A novel DNA-dependent protein kinase inhibitor, NU7026, potentiates the cytotoxicity of topoisomerase II poisons used in the treatment of leukemia

Elaine Willmore, Sarah de Caux, Nicola J. Sunter, Michael J. Tilby, Graham H. Jackson, Caroline A. Austin, and Barbara W. Durkacz

We report for the first time the use of a selective small-molecule inhibitor of DNA repair to potentiate topoisomerase II (topo II) poisons, identifying DNA-dependent protein kinase (DNA-PK) as a potential target for leukemia therapy. Topo II poisons form cleavable complexes that are processed to DNA double-strand breaks (DSBs). DNA-PK mediates nonhomologous end joining (NHEJ). Inhibition of this DSB repair pathway may sensitize cells to topo II poisons. We investigated the effects of a novel DNA-PK inhibitor, NU7026 (2-

(morpholin-4-yl)-benzo[h]chomen-4-one), on the response to topo II poisons using K562 leukemia cells. NU7026 (10 μ M) potentiated the growth inhibition of idarubicin, daunorubicin, doxorubicin, etoposide, amsacrine (mAMSA), and mitroxantrone with potentiation factors at 50% growth inhibition ranging from approximately 19 for mAMSA to approximately 2 for idarubicin (potentiation of etoposide was confirmed by clonogenic assay). In contrast, NU7026 did not potentiate camptothecin or cytosine arabinoside (araC). NU7026 did not affect the levels of etoposide-induced topo II α or β cleavable complexes. NU7026 alone had no effect on cell cycle distribution, but etoposide-induced accumulation in G2/M was increased by NU7026. A concentration-dependent increase in etoposide-induced DSB levels was increased by NU7026. The mechanism of NU7026 potentiation of topo II poisons involves inhibition of NHEJ and a G2/M checkpoint arrest. (Blood. 2004;103: 4659-4665)

© 2004 by The American Society of Hematology

Introduction

DNA topoisomerase II (topo II) is the target of some of the most important chemotherapeutic drugs used to treat leukemia. For example, induction therapies for adult acute myeloid leukemia (AML) include an anthracycline (eg, idarubicin or daunorubicin) in combination with cytosine arabinoside (araC), and consolidation protocols include other topo II poisons like mitoxantrone or etoposide.¹ Elevated expression of *P*-glycoprotein (*P-gp*), resulting in decreased cellular retention of topo II poisons, and decreased expression of topo II are important mechanisms of drug resistance in cell culture and in clinical specimens.²⁻⁵ Current therapies for AML are limited by associated toxicities (particularly neutropenia) and incomplete responses. Thus, acquisition of resistance to topo II poisons, incomplete responses, and dose-limiting toxicities are major clinical problems, and development of novel therapies remains an important goal.

Topo II is a nuclear enzyme that controls DNA topology. The enzymatic functions include relaxation of supercoiled DNA and decatenation of sister chromatids. There are 2 isoforms of topo II (α , 170 kDa; and β , 180 kDa) encoded by different genes. Although in vitro they perform the same catalytic function, evidence suggests that in vivo they possess distinct functions.⁶ During the normal catalytic cycle of topo II, the enzyme cleaves the phosphodiester backbone of DNA to generate a transient enzyme–bridged DNA double-strand break (DSB) through which another DNA duplex can pass, and then the DSB is religated. Topo II poisons trap the intermediate in this process, giving rise to a "cleavable complex".⁷

Cellular processing converts the protein-DNA cleavable complex into a DSB, which may be repaired by a DSB repair pathway.

Enzyme-mediated repair of DSBs is a major mechanism of resistance to both ionizing radiation (IR) and drugs that cause DSBs as intermediates in repair processes. Two complementary DSB repair pathways in eukaryotes are nonhomologous end joining (NHEJ) and homologous recombination repair (HRR).⁸ Important components in these repair pathways are the phosphatidylinositol 3-kinase (PI 3-K)–related protein kinase (PIKK) family of enzymes. These DNA damage–activated serine/threonine protein kinases include DNA-dependent protein kinase (DNA-PK), ataxia telangiectasia–mutated kinase (ATM), and ataxia telangiectasia and rad3–related kinase (ATR).⁹

The DNA-PK holoenzyme comprises a heterodimer of approximately 70- and 80-kDa polypeptides, known as Ku, that binds to DNA strand breaks, recruiting and activating the 470-kDa catalytic subunit, termed DNA-PKcs.¹⁰ Numerous studies have shown that cells lacking DNA-PK are hypersensitive to IR and cross-linking agents and defective in DSB repair.¹⁰⁻¹³ DNA-PK, together with the x-ray crosscomplementing (XRCC4)/DNA ligase IV complex and the recently identified cofactor Artemis, is specifically required for NHEJ,^{14,15} with about 80% of DNA DSBs repaired by this pathway.¹⁶

Studies using mutant cell lines and generic inhibitors of the PIKK family have implicated DNA-PK function in the survival of cells treated with topo II poisons. The xrs-1 cell line, which has a deletion mutation in the Ku80 gene rendering DNA-PK inactive, is

Supported by grant 0171 from the Leukaemia Research Fund,

United Kingdom (E.W.).

© 2004 by The American Society of Hematology

From the School of Cell and Molecular Biosciences, University of Newcastleupon-Tyne Medical School, Newcastle-upon-Tyne, United Kingdom; Northern Institute for Cancer Research, University of Newcastle-upon-Tyne Medical School, Newcastle-upon-Tyne, United Kingdom; and Department of Hematology, Royal Victoria Infirmary, Newcastle-upon-Tyne, United Kingdom.

Submitted July 29, 2003; accepted February 19, 2004. Prepublished online as *Blood* First Edition Paper, March 9, 2004; DOI 10.1182/blood-2003-07-2527.

Reprints: B. W. Durkacz, Northern Institute for Cancer Research, University of Newcastle-upon-Tyne Medical School, NE2 4HH, United Kingdom; e-mail: b.w.durkacz@ncl.ac.uk.

The publication costs of this article were defrayed in part by page charge payment. Therefore, and solely to indicate this fact, this article is hereby marked "advertisement" in accordance with 18 U.S.C. section 1734.

hypersensitive to a range of topo II poisons and partially defective in the repair of etoposide-induced DSBs.^{17,18} Etoposide resistance is restored by transfection of Ku80 cDNA into the mutant cell line.¹⁹ The first identified inhibitor of the PIKK enzyme family was the fungal metabolite wortmannin. Although primarily a PI 3-K inhibitor, it was also shown to potentiate IR-induced cytotoxicity and inhibit DSB repair at concentrations that inhibited cellular DNA-PK.²⁰⁻²² Similar results were also obtained with LY294002, another PIKK family inhibitor.²² Potentiation of etoposide cytotoxicity by wortmannin in a DNA-PK–proficient but not a DNA-PK– deficient cell line has also been demonstrated, indicating that this potentiation resulted specifically from DNA-PK inhibition by wortmannin.²³ These data support the idea that a specific DNA-PK inhibitor could prove a powerful tool for use in combination with topo II poisons in leukemia therapy.

NU7026 (2-(morpholin-4-yl)-benzo[h]chomen-4-one), a novel and specific inhibitor of DNA-PK, has been evaluated in this study. NU7026 is a competitive and highly selective inhibitor of DNA-PK, with a 60-fold greater potency against this enzyme than PI 3-K, and inactive against both ATM and ATR.²⁴ We have recently demonstrated that NU7026 is a potent radiosensitizer and inhibitor of DSB repair.²⁴ Here we describe the effects of NU7026 on the cytotoxic mechanisms of a range of topo II poisons in a human leukemia cell line. Cell growth and survival, cell cycle phase distribution, cleavable complex formation, and DSB repair were investigated.

Materials and methods

Cell culture

The K562 cell line was derived from a patient with chronic myelogenous leukemia (CML) in terminal blast phase and its metabolism is therefore similar to that of AML blasts. ML1 is an AML cell line. Both lines were maintained as a suspension culture in RPMI 1640 medium supplemented with 10% (vol/vol) fetal bovine serum, penicillin (50 U/mL), and streptomycin (50 µg/mL). Cells were grown at concentrations between 1×10^5 and 1×10^6 /mL and were free of *mycoplasma* contamination. The doubling time of the cells was approximately 24 hours (K562) and 30 hours (ML1). Cell culture reagents were obtained from Invitrogen (Paisley, United Kingdom).

Drugs

Idarubicin, daunorubicin, doxorubicin, mitoxantrone, and araC were made up as 1-mM stocks dissolved in water, stored in aliquots at -20° C, and diluted in water immediately prior to use. NU7026, mitoxantrone and amsacrine (mAMSA), and camptothecin were dissolved in anhydrous dimethyl sulfoxide (DMSO) as 1-mM stocks and stored at -20° C. Etoposide was dissolved in methanol as a 2-mM stock. Drugs were purchased from Sigma (Poole, United Kingdom). Idarubicin was kindly provided by Pharmacia-UpJohn (Cambridge, United Kingdom). NU7026 was synthesized by the Department of Chemistry, University of Newcastle-upon-Tyne, in collaboration with KuDOS Pharmaceuticals (Cambridge, United Kingdom).

XTT Growth inhibition assay

Exponentially growing cells were seeded (2×10^3 /well in 100 µL) into 96-well plates. Drug(s) (6 replicates) were added to plates 24 hours later. Following a 5-day incubation, growth inhibition was quantified using a sodium 3'-[1-(phenoaminocarbonyl)-3,4-tetrazolium] bis(4-methoxy-6-nitro)benzene sulfonic acid hydrate (XTT) cell proliferation kit (Roche, Lewes, United Kingdom), per manufacturer's instructions. Plates were read on a Bio-Rad 550 plate reader (Hercules, CA) at 450 nm, and results were expressed as a percentage of controls. NU7026 (10 µM) was found to have a small inhibitory effect on cell growth (~20% after 5 days) and

consequently the NU7026 alone samples were used as controls in calculations for drug combinations using a fixed concentration (10 μM) of inhibitor. Potentiation factors (PF_{50} values) at 50% growth inhibition (IC_{50}) were calculated from the ratios of the concentration of the drug alone that reduced growth by 50% (IC_{50} value) divided by the IC_{50} value obtained when the drug was used in combination with NU7026. Data were averaged from at least 3 independent experiments \pm SE.

Clonogenic assay

Exponentially growing cells were exposed to increasing concentrations of etoposide in the presence or absence of NU7026 (10 µM) for 16 hours. Alternatively, cells were exposed to increasing concentrations of NU7026 in the presence or absence of a fixed concentration (0.1 μ M) of etoposide. After exposure, drugs were removed by centrifugation and cell numbers were estimated by hemocytometer. A known number of cells were seeded in the absence of drug(s) into medium containing 0.125% (wt/vol) SeaKem agarose (Biowhittaker, Wokingham, United Kingdom) and 10% vol/vol fetal bovine serum in 35-mm culture dishes to allow colony formation. Under these conditions, K562 cells formed colonies with a plating efficiency of approximately 50% without the need for a feeder layer. Viable colonies were visualized by staining with MTT (3-[4,5-dimethylthiazol-2-yl]-2,5-diphenyltetrazolium bromide; 0.5 mg/mL in water). Colonies were counted using an Oxford Optronix colony counter (Oxford, United Kingdom). Survival was calculated as a percentage of control using the relative plating efficiency of control or NU7026-treated cells. Under the conditions used here, NU7026 had no effect on cell survival. PF50 values were calculated as for the XTT assays. Data were averaged from at least 3 independent experiments \pm SE.

TARDIS assay

The trapped in agarose DNA immunostaining (TARDIS) assay is used to quantitate cleavable complex formation and has been described in detail previously.²⁵ Briefly, cells were treated with drugs at the concentrations and for the times specified in the Figure legends and embedded in agarose on microscope slides. Proteins that were not covalently bound to the DNA were removed by placing the slides in 1M NaCl plus protease inhibitors followed by lysis buffer (1% sarkosyl; 80 mM phosphate, pH 6.8; 10 mM EDTA [ethylenediaminetetraacetic acid] plus protease inhibitors). Slides were stained with primary antibody (specific for either topo II α or β) and then a fluorescein isothiocyanate (FTTC)–conjugated secondary antibody before being counter-stained with Hoechst 33258. Images were captured using an epi-fluorescence microscope that separately visualizes blue (Hoescht-stained DNA) and green (FTTC-stained topo II) fluorescence. Imager 2 software (Astrocam, Cambridge, United Kingdom) was used to analyze and quantify levels of blue and green fluorescence. Data were averaged from at least 3 independent experiments ± SE.

Flow cytometric analysis

Cells were treated with drug(s) for 24 or 48 hours before permeabilization with 70% ethanol. Cells were treated with propidium iodide (400 μ g/mL) and RNAse (1 mg/mL) for 30 minutes at 37°C. Flow cytometry was performed on a Becton Dickinson FACScan (Heidelberg, Germany) equipped with an argon ion laser (excitation at 488 nm).

DNA strand break assay

DSB levels were assayed using the neutral filter elution technique.²⁶ The radiolabelling conditions used were exactly as previously described.²¹ Prelabelled cells were treated with etoposide (in the presence or absence of 10 μ M NU7026) for 1 hour before harvesting and preparing for neutral elution. Cells used as internal standards were exposed to IR (100 Gy) and placed immediately on ice. The internal standards were loaded onto the same filters as the experimental samples and eluted at pH 9.6. Data from elution profiles were summarized as relative elution (RE) values using the formula of Fornace and Little.²⁷ This calculates the amount of DNA retained on the filter when 50% of the internal standards have eluted. Data were averaged from 4 independent experiments ± SE.

Results

Growth inhibition

We have previously demonstrated that NU7026 (10 µM) completely inhibits purified human DNA-PK, is not cytotoxic per se in Chinese hamster ovary cells, and maximally potentiates IRinduced cytotoxicity.24 NU7026 was used at 10 µM in all experiments here, because, although it was slightly growth inhibitory (80% control growth), it maximally potentiated etoposide growth inhibition in K562 cells (results not shown) and was not cytotoxic in clonogenic survival assays (see "Cell survival"). Figure 1 shows the concentration-dependent effects of 4 topo II poisons, in the presence or absence of a fixed concentration (10 μ M) of NU7026, on cell growth. The IC_{50} and PF_{50} values for all the drugs evaluated are summarized in Table 1. NU7026 potentiated the growth inhibitory effects of idarubicin, daunorubicin, doxorubicin, etoposide, mAMSA, and mitoxantrone with PF₅₀ values ranging from approximately 19 for mAMSA to approximately 2 for idarubicin. In contrast, NU7026 did not potentiate the topo I poison, camptothecin, and, although NU7026 appeared to give some protection against the cytotoxicity of the nucleoside analog, araC $(PF_{50} = 0.45 \pm 0.29)$, this was not significant as assessed by the Student t test (P = .1532).

To test whether the effects of NU7026 were cell line specific, we used the AML cell line, ML1. Figure 2 shows that NU7026 also potentiated the growth inhibitory effect of etoposide in this leukemia cell line with a PF_{50} value of 10.53 ± 4.5 . Other data using clonogenic assays have demonstrated that NU7026 also potentiates etoposide- and doxorubicin-induced cytotoxicity in the human epithelial colorectal carcinoma cell lines, LoVo and SW620. Furthermore, NU7026 itself was shown to have little or no effect on the growth of the epithelial lines SW620 and HCT 116 (Yan Zhao, unpublished results, 2003).



Figure 1. Growth inhibitory effects of topo II poisons in the presence or absence of a fixed concentration of NU7026. K562 cells were treated with increasing concentrations of (A) etoposide, (B) idarubicin, (C) doxorubicin, or (D) mAMSA in the absence (III) or presence (\bigcirc) of 10 μ M NU7026. After 5 days of exposure, cells were stained with XTT and OD_{450nm} was quantified. Results are means of at least 4 experiments \pm SE.

Table 1. Comparison of IC_{50} and PF_{50} values derived from the growth inhibition curves

	Drug alone	Drug + NU7026		
Drug	IC ₅₀ , nM*	IC ₅₀ , nM*	PF50†	
mAMSA	32	1.6	19.2 ± 7.5	
Mitoxantrone	3	0.18	16.6 ± 6.6	
Etoposide	317	45	8.6 ± 1.0	
Doxorubicin	45	10	4.0 ± 0.38	
Daunorubicin	205	78	2.6 ± 0.11	
Idarubicin	9.5	5.4	1.7 ± 0.17	
Camptothecin	6.8	7.3	0.88 ± 0.17	
araC	33	66	0.45 ± 0.29	

*IC₅₀ values are the means of at least 3 independent experiments.

 \pm All mean PF₅₀ values (\pm SE) were calculated from the ratio of the individual IC₅₀ values (eg, the concentration of drug that caused 50% growth inhibition divided by the concentration of drug that caused 50% growth inhibition in the presence of NU7026).

Cell survival

The ability of NU7026 to potentiate cell killing by topo II poisons was evaluated by clonogenic survival assays. Figure 3A shows that treatment of K562 cells with etoposide resulted in a concentration-dependent decrease in survival, and this was potentiated by coincubation with NU7026 ($PF_{50} = 5.4 \pm 1.0$). It was observed that although 10 μ M NU7026 alone was slightly growth inhibitory in a 5-day exposure (see "Growth inhibition"), the 16-hour exposure used here prior to plating for colony formation was completely nontoxic.

In order to study the effects of NU7026 alone on cytotoxicity, K562 cells were exposed to increasing concentrations of NU7026 in the presence or absence of 0.1 μ M etoposide (a concentration that, used alone, reduced survival to ~ 65% of controls). Figure 3B demonstrates that NU7026 alone had no effect on cytotoxicity up to a concentration of 20 μ M. However, concentrations of 5 to 20 μ M all potentiated etoposide-induced cytotoxicity, reducing survival by, for example, 30% (5 μ M) and 70% (20 μ M).

Cleavable complex levels

An increase in topo II poison–induced cleavable complex formation by NU7026 is a possible mechanism that would enhance cytotoxicity. Therefore the effect of NU7026 on etoposide-induced cleavable complex levels was investigated using the TARDIS assay. Figure 4A shows the distinctive dose-dependent increase in green immunofluorescence in individual cells after etoposide treatment, corresponding to topo II α cleavable complex formation.²⁵ The scattergram in Figure 4B is a representative plot from a single experiment to demonstrate the distribution of immunofluorescence in individual cells after treatment with etoposide in the



Figure 2. Growth inhibitory effect of etoposide \pm NU7026 on ML1 cells. Cells were treated for 5 days before XTT staining. \blacksquare indicates etoposide alone; and \square , etoposide + 10 M NU7026. Results are means of 3 experiments \pm SE.



Figure 3. Clonogenic survival assay. (A) Cells were treated with increasing concentrations of etoposide in the presence or absence of NU7026 (10 μ M) for 16 hours. Indicates etoposide alone; and \Box , etoposide + NU7026. (B) cells were treated with increasing concentrations of NU7026 in the presence or absence of etoposide (0.1 μ M) for 16 hours. Indicates NU7026 alone; and \Box , NU7026 + etoposide. Cells were then plated in soft agar for colony formation in the absence of drugs. Results are the means of 4 experiments ± SE.

presence or absence of NU7026. Figure 4C (which shows the mean values derived from the scattergram data) demonstrates that coincubation with NU7026 had no significant effect on the levels of topo II α cleavable complexes. NU7026 also did not alter etoposide-induced topo II β cleavable complex levels (data not shown).

Cell cycle distribution

Flow cytometric analysis was used to assess the effects of the drugs on cell cycle phase. NU7026 by itself had no effect on cell cycle distribution. Following a 24-hour exposure (equivalent to one cell cycle) to 0.1 µM etoposide, cells accumulated in G2, and this effect was more pronounced after treatment with 0.4 µM etoposide (Figure 5). In the presence of NU7026, the etoposide-induced G2 blockade was enhanced. The percentage of cells in each stage of the cell cycle is summarized in Table 2. For example, at 24 hours the percentage of cells in G2 in NU7026-treated cells was 13% and this was increased to 51% in the presence of etoposide and to 62% when both drugs were present. The differences were even more marked at 48 hours, when cells attempted to progress into a second cell cycle. It was apparent that cells treated with 0.4 µM etoposide alone escaped the 24-hour G2 blockade and progressed into a second cell cycle where they arrested and accumulated in S phase (compare 11% in S phase at 24 h with 40% in S phase at 48 h; Table 2). In contrast, when etoposide and NU7026 were used in conjunction, the majority of cells remained arrested at the initial G2 checkpoint and were unable to progress into the second cell cycle (Figure 5; Table 2). It was also noted that by 48 hours a sub-G1 fraction of cells was accumulating, indicating that cells were undergoing apoptosis (Figure 5).



Figure 5. Flow cytometric analysis of cell cycle distribution. Cells were treated with 0.1 or 0.4 μ M etoposide in the presence or absence of NU7026 (10 μ M) for 24 or 48 hours before harvesting and preparing for flow cytometric analysis.

DSB assays

The effect of NU7026 on etoposide-induced DSB levels was assessed by the neutral elution technique. Cells were treated with etoposide for 1 hour in the presence or absence of NU7026. Figure 6A is a representative neutral elution profile showing a concentration-dependent increase in DSBs, represented by decreasing levels of [¹⁴C] thymidine (TdR) retained on the filter. In the presence of NU7026, levels of etoposideinduced DSB were higher. For example, (at 50% [³H] retention) 5 μ M etoposide resulted in an 82% retention of the [¹⁴C], whereas for etoposide + NU7026 this was reduced to 53%. These data were averaged and quantitated by conversion to RE values (Figure 6B). The same trend was observed over all the concentrations of etoposide tested (Figure 6B). For example, the RE value for 5 μ M etoposide was 0.056 \pm 0.013; in the presence of NU7026 this value was increased to 0.13 \pm 0.024, indicating that NU7026 inhibits the rejoining of etoposide-induced DSBs.

Discussion

Topo II poisons are a mainstay of leukemia therapy. Here we demonstrate for the first time that at noncytotoxic concentrations, the DNA-PK inhibitor NU7026 significantly potentiates the growth inhibitory effects of 6 topo II poisons. Furthermore, NU7026 potentiates the cytotoxicity of etoposide in a clonogenic survival assay. NU7026 does not affect the initial lesion of etoposide-induced topo II cleavable complex formation. NU7026 inhibits DSB repair and increases the etoposide-induced G2/M arrest. These data strongly implicate DNA-PK function in the cytotoxic



Figure 4. Tardis assay of cleavable complex levels. Cells were treated with etoposide in the presence or absence of 10 μ M NU7026 for 24 hours. Original magnification \times 10. (A) Immunofluorescent micrographs of control cells and cells treated with 1 and 10 μ M etoposide. The blue stain indicates DNA; green stain, topo II α cleavable complexes. (B) Scattergram. (C) Averaged data from scattergram (topo II α -specific antisera). \Box indicates etoposide alone; and \blacksquare , etoposide + NU7026. Results are means of 3 experiments \pm SE.

Table 2. Cell cycle distribution of cells treated with etoposide in the presence or absence of NU7026

	24 h			48 h	
Cell cycle phase	NU7026	Etoposide	Etoposide + NU7026	Etoposide	Etoposide + NU7026
G ₁ , %	71	34	16	32	14
S, %	12	11	10	40	37
G ₂ /M, %	13	51	62	16	44

Cells were treated with etoposide (0.4 $\mu M)$ in the presence or absence of NU7026 (10 $\mu M)$ for 24 or 48 hours, then harvested for flow cytometric analysis. Data are the means of duplicate samples from a representative experiment.

mechanism of topo II poisons downstream of cleavable complex formation. We believe that NU7026 potentiates topo II poisons by inhibiting the DNA-PK-mediated NHEJ of DSBs generated downstream from cleavable complexes. Failure to repair these breaks would lead to the arrest of cells at the DNA damage-induced G2 checkpoint mediated by ATM²⁸ as seen in Figure 5.

PF50 values for these 6 topo II poisons vary from 2 to 19 (Table 1). The wide variation in PF_{50} is intriguing since the main cytotoxic mechanism of all these agents is poisoning of topo II. There are a number of possibilities to explain this variability. We hypothesize that it is due to the differential targeting of these agents to topo $II\alpha$ and β . Human topoisomerase II exists in 2 isoforms (α and β), and the agents that give the highest PF_{50} directly correlate with those that target both topo II α and topo II β . The large PF₅₀ for mAMSA (19.2) and mitoxantrone (16.6) correlates with their known targeting of both isoforms of topo II as part of their cytotoxic mechanism.²⁹ Etoposide shows an intermediate PF_{50} (8.6) and this agent forms complexes with both isoforms in vivo²⁵ and both isoforms mediate cytotoxicity at high doses of etoposide.²⁹ The 3 topo II poisons that have the lowest PF50 are idarubicin, daunorubicin, and doxorubicin (1.7, 2.55, and 4, respectively) and these are all anthracyclines. We have previously demonstrated that these agents only form cleavable complexes with topo $II\alpha$ (and not with topo IIB) in K562 cells and that the sensitivity of mouse embryonic fibroblasts to the anthracyclines is not enhanced by topo IIB.³⁰ We propose that the dual targeting of both topo II α and β is the basis for the high levels of potentiation seen with mAMSA, mitoxantrone, and etoposide. However, we cannot exclude the possibility that other factors may play a role in the potentiation by NU7026. These could include the longevity of cleavable complexes³⁰ and



Figure 6. DNA DSB assay. (A) Representative neutral elution profile for K562 cells treated with 0 (**II**), 5 (**V**), 10 (**•**), or 20 (**•**) μ M etoposide in the absence (black symbols) or presence (white symbols) of 10 μ M NU7026 for 1 hour prior to neutral elution. (B) Mean relative elution values (calculated at 50% [³H] retention). indicates etoposide alone; and **II**, etoposide + NU7026. Results are means from 4 independent experiments \pm SD. *P* values were calculated using the 2-tailed unpaired Student *t* test in order to determine whether DSB levels were significantly different in the presence of NU7026 (*P* = .01, 5.0 μ M; *P* = .183, 10 μ M; *P* = .06, 20 μ M).



Model for NU7026 mechanism of action



Figure 7. Model for NU7026 mechanism of action. Schematic representing our model for the mechanism by which NU7026 potentiates topo II poison cytotoxicity. The topo II–DNA complex is an intermediate step in the catalytic cycle during which the enzyme passes one DNA duplex through another (via a covalent protein-bridged DNA break) and then religates the cleaved DNA. Treatment with detergent can disrupt or cleave these complexes, hence the term "cleavable complexes." However, treatment of cells with topo II poisons (eg, etoposide) results in stabilization of the normally transient cleavable complex. After stabilization by drugs, complexes may be processed into DNA DSB, stimulating activation of DNA repair enzymes. Recruitment of DNA-PK to DSB facilitates nonhomologous end joining and therefore repair of DNA. However, in the presence of NU7026, DNA-PK activity is inhibited, NHEJ is impeded, and thus persistence of DNA damage leads to cell death.

their location in the genome,³¹ both of which are likely to play a role in DNA damage responses. Therefore, for topo II poisons (anthracyclines in particular), it is not always possible to directly correlate levels of DNA strand breaks with cytotoxicity.^{32,33}

Our model (Figure 7) shows the proposed mechanism by which NU7026 potentiates topo II poison-induced cytotoxicity. Topo II cleavable complexes are transient protein-bridged DNA breaks that are normally religated after the enzyme has passed one DNA duplex through another. Upon drug treatment, these complexes are stabilized and can be processed into DNA DSBs. DNA DSBs can be repaired via recruitment of DNA-PK, and subsequent NHEJ results in cell survival. Alternatively, inhibition of DNA-PK activity by NU7026 results in decreased NHEJ and persisting DNA damage that leads to cell death.

Our data show that DNA-PK is involved in the repair of DSB following treatment of leukemia cells with topo II poisons. In vitro, DNA-PK is not activated by plasmid DNA bearing an etoposide-stabilized topo II–DNA cleavable complex.³⁴ However, removal of topo II (by proteinase K treatment) revealed a naked DNA break that does stimulate DNA-PK activity. These data are consistent with the requirement for cellular processing to remove topo II from the DNA break before DNA-PK activation can occur. Whether a novel tyrosyl-DNA phosphodiesterase (analogous to the one that acts on DNA topoisomerase I³⁵) or a known DNA repair enzyme is able to carry out this function remains to be elucidated.

Two common mechanisms leading to chemoresistance and therapy failure in relapsed leukemias are reduced topo II activity ("atypical drug resistance") and multidrug resistance (MDR), resulting in reduced intracellular levels of topoisomerase II poisons due to increased drug efflux.36 Increased expression of DNA-PKcs also contributes to chemoresistance and radioresistance by increasing the cellular capacity to carry out NHEJ. Increased DNA-PK activity has been widely demonstrated both in cell culture and in samples from patients with relapsed chronic lymphocytic leukemia (CLL) and correlates with the resistance of tumor cells to IR and bifunctional alkylating agents.³⁷⁻⁴⁰ An HL60 cell line selected for resistance to adriamycin was found to have a more than 15-fold increase in DNA-PKcs levels.⁴¹ DNA-PK activity and content correlated with sensitivity to doxorubicin and etoposide in clinical samples of cells from patients with CLL.42 Thus, increased DNA-PK activity can contribute to topo II poison-resistance mechanisms. Further research is required to confirm its significance in a clinical setting.

Leukemia-specific alterations in DNA-PK will be important determinants in designing therapeutic regimens for DNA-PK inhibitors. In leukemias where DNA-PK is overexpressed and tumors are radioresistant and chemoresistant, a selective DNA-PK inhibitor would resensitize these resistant cells. In 5 reports, antisense oligonucleotides, small interfering RNAs, and a C-terminal peptide that targets Ku80 and prevents DNA-PKcs binding to Ku have been used to deplete or inhibit DNA-PK function in human cell lines.⁴²⁻⁴⁷ Loss of DNA-PK activity correlated with radiosensitization, increased mutation, and inhibition of DNA damage repair.

This is the first report of a very selective small-molecule inhibitor of DNA repair in conjunction with topo II poisons. We believe that potentiation of the cytotoxicity of topo II poisons with DNA-PK inhibitors will be a powerful strategy for sensitizing cells to these agents. NU7026 is a specific DNA-PK inhibitor prototype, ideal for proof-of-principle experiments in cell line models, as reported in this study. The more recent identification of a DNA-PK inhibitor potent enough for pharmacologic evaluation and clinical use is the first step toward being able to test the potential of small-molecule inhibitors of DNA-PK in the clinic. The new compound is now the clinical candidate and is currently undergoing preclinical evaluation in human xenograft studies, cell line models, and leukemia cells derived from patients resistant to topo II poisons in order to test the hypothesis that inhibition of DNA-PK will prove a powerful strategy in overcoming therapeutic resistance.

References

- Lichtman MA, Liesveld JL. Acute myelogenous leukemia. In: Beutler E, Lichtman MA, Coller BS, Kipps TJ, Seligsohn U, eds. Williams Haematology. New York, McGraw-Hill; 2002:1047-1081.
- Nussler V, Pelka-Fleischer R, Zwierzina H, et al. *p*-Glycoprotein expression in patients with acute leukemia–clinical relevance. Leukemia. 1996;10: 23-31.
- Andoh T, Nishizawa M, Hida T, Ariyoshi Y, Takahashi T, Ueda R. Reduced expression of DNA topoisomerase II confers resistance to etoposide (VP-16) in small cell lung cancer cell lines established from a refractory tumor of a patient and by in vitro selection. Oncol Res. 1996;8:229-238.
- Fernandes DJ, Qiu J, Catapano CV. DNA topoisomerase II isozymes involved in anticancer drug action and resistance. Adv Enzyme Regul. 1995;35:265-281.
- Albertioni F, Gruber A, Arestrom I, Vitols S. Multidrug resistance gene (mdr1) RNA levels in relation to P-glycoprotein content of leukemic cells from patients with acute leukemia. Medical Oncol. 1995;12:79-86.
- Austin CA. Marsh KL. Eukaryotic DNA topoisomerase II beta. Bioessays. 1998;20:215-226.
- Kellner U, Sehested M, Jensen PB, Gieseler F, Rudolph P. Culprit and victim - DNA topoisomerase II. Lancet Oncol. 2002;3:235-243.
- Jackson SP. Sensing and repairing DNA double strand breaks. Carcinogenesis. 2002;23:687-696
- Durocher D, Jackson SP. DNA-PK, ATM and ATR as sensors of DNA damage: variations on a theme? Curr Opin Cell Biol. 2001;13:225-231.
- Smith GC, Jackson SP. The DNA-dependent protein kinase. Genes Dev. 1999;13:916-934.
- Caldecott J, Jeggo P. Cross sensitivity of gamma ray-sensitive hamster mutants to cross-linking agents. Mutat Res. 1991;255:111-121.
- Tanaka T, Yamagami T, Oka Y, Nomura T, Sugiyama H. The scid mutation in mice causes defects in the repair system for both double-strand DNA breaks and DNA cross-links. Mutat Res. 1993;288:277-280.
- Gu Y, Jin S, Gao Y, Weaver DT, Frederick W. Ku70-deficient embryonic stem cells have increased ionizing radiosensitivity, defective DNA end-binding activity, and inability to support V(D)J

recombination. Proc Natl Acad Sci U S A. 1997; 94:8076-8081.

- Critchlow SE, Jackson SP. DNA end joining: from yeast to man. Trends Biochem Sci. 1998;23:394-398.
- Pannicke MY, Schwarz K, Lieber MR. Hairpin opening and overhang processing by an Artemis/ DNA-dependent protein kinase complex in nonhomologous end joining and V(D)J recombination. Cell. 2002;108:781-794.
- DiBiase SJ, Zeng ZC, Chen R, Hyslop T, Curran WJ Jr, Iliakis G. DNA-dependent protein kinase stimulates an independently active, nonhomologous, end-joining apparatus. Cancer Res. 2000; 60:1245-1253.
- Jeggo PA, Caldecott K, Pidsley S, Banks GR. Sensitivity of Chinese hamster ovary mutants defective in DNA repair to topoisomerase II inhibitors. Cancer Res. 1989; 49:7057-7063.
- Caldecott K, Banks G, Jeggo P. DNA double strand break repair pathways and cellular tolerance to inhibitors of topoisomerase II. Cancer Res. 1990;50:5778-5783.
- He DM, Lee SE, Hendrickson EA. Restoration of X-ray and etoposide resistance, Ku-end binding activity and V(D)J recombination to the Chinese hamster sxi-3 mutant by a hamster Ku86 cDNA. Mutat Res. 1996;363:43-56.
- Izzard RA, Jackson SP, Smith GCM. Competitive and noncompetitive inhibition of the DNA-dependent protein kinase. Cancer Res. 1999;59:2581-2586.
- Boulton S, Kyle S, Yalçintepe L, Durkacz BW. Wortmannin is a potent inhibitor of DNA double strand break repair but not single strand break repair in Chinese hamster ovary cells. Carcinogenesis. 1996;17:2285-2290.
- Rosenzweig KE, Youmell MB, Palayoor ST, Price BR. Radiosensitization of human tumor cells by the phosphatidyl inositol 3-kinase inhibitors wortmannin and LY294002 correlates with inhibition of DNA-dependent protein kinase and prolonged G2-M delay. Clin Cancer Res. 1997;3:1149-1156.
- Boulton S, Kyle S, Durkacz BW. Mechanisms of enhancement of cytotoxicity in etoposide and ionising radiation treated cells by the protein kinase inhibitor wortmannin. Eur J Cancer. 2000;36:535-541.

- Downloaded from http://ashpublications.net/blood/article-pdf/103/12/4659/1698576/zh801204004659.pdf by guest on 02 June 202-
- Veuger SJ, Curtin, NJ, Richardson, CJ, Smith GC, Durkacz, BW. Radiosensitization and DNA repair inhibition by the combined use of novel inhibitors of DNA-dependent protein kinase and poly(ADP-ribose) polymerase-1. Cancer Res. 2003;63:6008-6015.
- Willmore E, Frank AJ, Padget K, Tilby MJ, Austin CA. Etoposide targets topoisomerase Ilalpha and Ilbeta in leukemic cells: isoform-specific cleavable complexes visualized and quantified in situ by a novel immunofluorescence technique. Mol Pharmacol. 1998;54:78-85.
- Bradley MO, Kohn KW. X-ray induced DNA double strand break production and repair in mammalian cells as measured by neutral filter elution. Nucleic Acids Res. 1979;7:793-804.
- Fornace AJ Jr, Little JB. DNA crosslinking induced by X-rays and chemical agents. Biochim Biophys Acta. 1977;477:343-355.
- Shiloh Y. ATM and ATR: networking cellular responses to DNA damage. Curr Opin Genet Dev. 2001;11:71-77.
- Errington F, Willmore E, Tilby MJ, et al. Murine transgenic cells lacking DNA topoisomerase IIbeta are resistant to acridines and mitoxantrone: analysis of cytotoxicity and cleavable complex formation. Mol Pharmacol. 1999;56:1309-1316.
- Willmore E, Errington, F, Tilby MJ, Austin CA. Formation and longevity of idarubicin-induced DNA topoisomerase II cleavable complexes in K562 human leukaemia cells. Biochem Pharmacol. 2002;63:1807-1815.
- Pommier Y, Leteutre F, Fesen MR, et al. Cellular determinants of sensitivity and resistance to DNA topoisomerase inhibitors. Cancer Invest. 1994; 12:530-542.
- Capranico G, Soranzo C, Zunino F. Single-strand DNA breaks induced by chromophore-modified anthracyclines in P388 leukemia cells. Cancer Res. 1986;46:5499-5503.
- Binaschi M, Capranico G, De Isabella P, et al. Comparison of DNA cleavage induced by etoposide and doxorubicin in two human small-cell lung cancer lines with different sensitivities to topoisomerase II inhibitors. Int J Cancer. 1990;45: 347-352.

- Martensson S, Nygren J, Osheroff N, Hammarsten O. Activation of the DNA-dependent protein kinase by drug-induced and radiation-induced DNA strand breaks. Radiat Res. 2003;160: 291-301.
- Yang SW, Burgin AB Jr, Huizenga BN, Robertson CA, Yao KC, Nash HA. A eukaryotic enzyme that can disjoin dead-end covalent complexes between DNA and type I topoisomerases. Proc Natl Acad Sci U S A. 1996;93:11534-11539.
- Larsen AK, Skladanowski A. Cellular resistance to topoisomerase-targeted drugs: from drug uptake to cell death. Biochim Biophys Acta. 1998; 1400:257-274.
- Xu W, Liu L, Smith GCM, Charles IG. Nitric oxide upregulates expression of DNA-PKcs to protect cells from DNA-damaging anti-tumour agents. Nat Cell Biol. 2000;2:329-345.
- Muller C, Chistodoulopoulos G, Salles B, Panasci L. DNA-dependent protein kinase activity correlates with clinical and in vitro sensitivity of chronic lym-

phocytic leukemia lymphocytes to nitrogen mustards. Blood. 1998;92:2213-2219.

- Sirzen F, Nilsson A, Zhivotovsky B, Lewensohn R. DNA-dependent protein kinase content and activity in lung carcinoma cell lines; correlation with intrinsic radiosensitivity. Eur J Cancer. 1999;35:111-116.
- Muller C, Calsou P, Salles B. The activity of the DNA-dependent protein kinase (DNA-PK) complex is determinant in the cellular response to nitrogen mustards. Biochimie. 2000;82:25-28.
- Tew KD, O'Brien M, Laing NM, Shen H. Coordinate changes in expression of protective genes in drug-resistant cells. Chem Biol Interact. 1998; 111-112:199-211.
- Eriksson A, Lewensoh R, Larsson R, Nilsson A. DNA-dependent protein kinase in leukaemia cells and correlation with drug sensitivity. Anticancer Res. 2002;22:1787-1793.
- 43. Sak A, Stuschke M, Wurm R, et al. Selective inactivation of DNA-dependent protein kinase with antisense oligodeoxynucleotides: consequences for the rejoining of radiation-induced DNA double-

strand breaks and radiosensitivity of human cancer cell lines. Cancer Res. 2002;62:6621-6624.

- 44. Peng Y, Zhang, Q, Nagasawa H, Okasayu R, Liber HL, Bedford JS. Silencing expression of the catalytic subunit of DNA-dependent protein kinase by small interfering RNA sensitizes human cells for radiationinduced chromosome damage, cell killing and mutation. Cancer Res. 2002;62:6400-6404.
- Kim C-H, Park S-J, Lee S-H. A targeted inhibition of DNA-dependent protein kinase sensitizes breast cancer cells following ionizing radiation. J Pharmacol Exp Ther. 2002;303:753-759.
- Omori S, Takiguchi Y, Suda A, et al. Suppression of a DNA double-strand break repair gene, *Ku70*, increases radio-and chemosensitivity in a human lung carcinoma cell line. DNA Repair. 2002;1: 299-310.
- Spencer CJ, Swartz MJ, Nelson WG, DeWeese TL. Enhanced radiation and chemotherapy-mediated cell killing of human cancer cells by small inhibitory RNA silencing of DNA repair factors. Cancer Res. 2003;63:1550-1554.