Brief report

High telomerase activity in granulocytes from clonal polycythemia vera and essential thrombocythemia

Anna Maria Ferraris, Rosa Mangerini, Natalija Pujic, Omar Racchi, Davide Rapezzi, Andrea Gallamini, Salvatore Casciaro, and Gian Franco Gaetani

Essential thrombocythemia (ET) and polycythemia vera (PV) are chronic myeloproliferative disorders that share the involvement of a multipotent progenitor cell and dominance of the transformed clone over normal hematopoiesis. On the other hand, the heterogeneity of these diseases with respect to clonal development from a common progenitor has been well established. To identify useful prognostic indicators, we analyzed telomerase activity (TA), a known marker of neoplastic proliferation, in granulocytes (PMNs) and mononuclear cells (MNCs) from 22 female patients with ET and PV. Clonality status was determined by investigation of X chromosome inactivation patterns (XCIPs). We found a statistically significant positive correlation between high TA and monoclonal pattern of XCIP. Therefore, our data suggest that the use of multiple tumor markers may contribute to a better understanding of the deregulated physiology of these disorders and provide useful prognostic factors. (Blood. 2005;105: 2138-2140)

© 2005 by The American Society of Hematology

Introduction

Telomeres are nonencoding regions of DNA capping the ends of chromosomes, essential for genetic stability and cell replication¹; telomerase is the ribonucleoprotein enzyme complex that maintains telomere integrity. Telomerase activity (TA) is generally undetectable in normal somatic cells, while it is expressed in approximately 85% of the common cancers.^{2,3} Hematopoietic stem cells express telomerase and contain long telomeres, which become shorter as cells differentiate and mature. While it is acknowledged that TA is elevated in the acute leukemias and aggressive lymphomas,⁴ exhaustive data on the level and prognostic significance of TA in other common hematopoietic diseases are still lacking.^{5,6}

Chronic myeloproliferative disorders (MPDs) are acquired diseases of the hematopoietic stem cell, characterized by clonal proliferation of one or more cell lineages. Several studies, however, have shown that a sizable proportion of patients with polycythemia vera (PV) and essential thrombocythemia (ET) have polyclonal hematopoiesis.⁷⁻⁹ The significance of this variable clonal expression and its relationships, if any, with other clinical and/or biologic characteristics remain elusive.

To investigate further the relevance and possible correlation(s) of these neoplastic markers, we decided to evaluate TA and the pattern of X chromosome inactivation (XCIP) as marker of clonality in a group of female patients with PV and ET.

Study design

Patients

Heparinized peripheral blood was obtained from 34 women with a diagnosis of PV or ET. Clinical findings of patients are summarized in Table

From the Ematologia Oncologica, Istituto Nazionale per la Ricerca sul Cancro; the Dipartimento di Medicina Interna, University of Genova; and the Struttura Complessa di Ematologia, Azienda Ospedaliera Santa Croce e Carle, Cuneo, Italy.

Supported by grants no. RBAU013W3J_010 from Fondo per gli Investimenti della Ricerca di Base (FIRB) and no. 3398 from Ministero della Salute.

Submitted July 2, 2004; accepted September 30, 2004. Prepublished online as *Blood* First Edition Paper, October 19, 2004; DOI 10.1182/blood-2004-06-2375.

1. Hematologic diagnosis was according to Polycythemia Vera Study Group (PVSG) criteria,¹⁰ and causes of secondary polycythemia (SP) or reactive thrombocytosis (RT) were carefully excluded. In no case was there a family history of MPD. All patients were in chronic phase without immature myeloid cells in the peripheral blood. Granulocytes (PMNs) and mono-nuclear cells (MNCs) were purified from peripheral blood according to standard methods.¹¹

All samples were obtained after informed consent following the guidelines of the Institutional Review Board of the University of Genova.

Estimation of telomerase activity

TA was determined in PMNs and MNCs with the TRAPeze telomerase detection kit (Intergen, Burlington, MA), based on the original method by Kim et al.² For each sample, the following controls were included: a heat-treated negative control, a positive control (HL-60 cell line),¹² and an internal standard, to monitor polymerase chain reaction (PCR) inhibition and for quantitative analysis of the reaction products. The amount of telomerase product for each reaction was expressed in total product generated units (TPGs). The assay has a linear range of 1 to 300 TPGs, equivalent to the TA from approximately 30 to 10 000 control cells. Thirty healthy age-matched females were included as controls.

XCIP PCR assay

To assess clonality, we took advantage of a highly polymorphic short tandem repeat (STR) within the X-linked human androgen receptor (HUMARA) gene. The HUMARA locus is very useful for X-chromosome inactivation studies because with a PCR-based method, approximately 85% of women are heterozygous, and results correlate well with those obtained by Southern blotting for phosphoglycerate kinase–hypoxanthine phosphoribosyltransferase (PGK-HPRT), and M27 β^{13} or transcriptional polymorphisms assay.^{7,14,15}

Reprints: Anna Maria Ferraris, Ematologia Oncologica Istituto Nazionale per la Ricerca sul Cancro, Largo Rosanna Benzi 10, 16132 Genova, Italy; e-mail: annamaria.ferraris@istge.it.

The publication costs of this article were defrayed in part by page charge payment. Therefore, and solely to indicate this fact, this article is hereby marked "advertisement" in accordance with 18 U.S.C. section 1734.

© 2005 by The American Society of Hematology

Table 1. Clinical and biologic features of informative patients

Patient no.	Diagnosis	Age at diagnosis (y)	Duration of disease (y)	Hgb g/L	Platelets 10³/μL	Splenomegaly	Therapy at time of analysis	Therapy prior to analysis	TA (PMN)	Clonality (PMN)
1	ET	95	1	131	736	N	HU	_	103	С
2	PV	60	9	127	200	N	HU	HU	1	Р
3	ET	68	14	126	794	N	—	HU	47.6	С
4	PV	69	1	158	749	Y	—	Ν	127.7	С
5	PV	72	3	193	311	N	HU	HU	89.6	С
6	ET	68	17	135	721	Y	HU	HU	212.5	С
7	PV	63	1	164	583	N	_	—	197	С
8	PV	78	3	184	603	Y	HU	P, HU	24.7	Р
9	ET	62	3	165	750	N	HU	HU	2.1	Р
10	ET	76	3	138	883	Y	HU	A, HU	123.7	C (LOH)
11	ET	61	9	158	769	N	HU	HU	19	Р
12	ET	76	9	168	900	N	HU	HU	105	С
13	ET	78	12	146	740	N	—	В	10.8	Р
14	PV	77	16	194	600	N	HU	P, HU	95	С
15	ET	79	6	143	795	N	_	В	70	С
16	PV	57	14	190	341	Y	P, HU	P, HU, A	83.7	С
17	ET	53	15	127	900	N	HU, A	HU	146.5	С
18	ET	63	14	135	480	N	HU	HU	10	Р
19	ET	88	15	125	800	Y	HU	HU	119	С
20	ET	62	1	128	950	N	HU	—	90	С
21	PV	77	0	180	280	Y	—	—	162	С
22	PV	28	1	176	821	Ν	—	_	33.3	Р

ET indicates essential thrombocythemia; PV, polycythemia vera; C, monoclonal; P, polyclonal; LOH, loss of heterozygosity; A, anagrelide; B, busulfan; HU, hydroxurea; P, phlebotomy; and TA, telomerase activity in TPG units.

indicates no therapy.

Analysis of X-chromosome inactivation was therefore performed according to Karasawa et al¹⁶ with some modifications. Primers sequences were as follows: F, 5'TCCAGAATCTGTTCAAGAGCG-TGC-3' and R, 5'GCTGTGAAGGTTGCTGTTCCTCAT-3'. The 5' end of the forward primer was labeled with 6-carboxyfluoroscein (FAM) fluorescent dye. The PCR products were separated by capillary electrophoresis on an ABI PRISM Genetic Analyzer (Applied Biosystems, Foster City, CA) and analyzed with a GeneScan software (Applied Biosystems).

Results and discussion

Thirty-four female patients with PV or ET were tested for heterozygosity at the HUMARA locus, and 27 (82%) were found informative. Of the 27, 5 were excluded from the study due to excess skewing, defined as greater than or equal to 90% of the cells with the same X-chromosome active.¹¹ Clonality status was therefore determined in PMNs and MNCs by XCIP analysis in 22 patients (Table 1). PMNs expressed a monoclonal XCIP in 9 (69%) of 13 subjects with ET and in 6 (67%) of 9 with PV. MNCs were always polyclonal. In patient 10 with ET (Table 1), the comparison of the undigested PMNs and MNCs after PCR amplification revealed loss of heterozygosity (LOH) at the HUMARA locus in PMNs. Although this finding prevented evaluation of XCIP, the presence of a LOH can be considered as evidence of a monoclonal process¹⁷ and will be counted as such.

TA was measured in all informative patients and then blindly compared with the results of XCIP analysis. Monoclonal PMNs displayed a TA ranging between 47.6 and 212.5 TPG units (mean, 118.2 \pm 45.5 SD), whereas polyclonal PMNs had a TA between 1.0 and 33.3 TPG units (mean, 14.4 \pm 11.9 SD) (Figure 1). The difference between TPG values in the 2 groups is highly significant (P < .001, unpaired Student *t* test). MNCs had a TA similar to that of polyclonal PMNs. TA of controls was 12.2 \pm 9.3 TPG units (range, 1.3-28.1). Our intention to analyze and determine the relations between 2 well-known markers of neoplastic proliferation has therefore been fulfilled by the finding of a direct correlation between TA and XCIP.

The stem cell origin and clonal development of MPDs are heterogeneous.^{7-9,18} One of the leading hypotheses is that the initial step of neoplastic proliferation can occur at different stages during the hemopoietic stem cell maturation process. On the other hand, controversies on the significance of clonality in MPDs are abundant in the literature.^{14,19,20} Several clinical and laboratory features have been evaluated in the literature, and a correlation between XCIP and vascular complications in ET was recently suggested,^{21,22} but most factors have shown no correlation with survival, thrombotic complications, or leukemic evolution.²²

The results of our study clearly indicate the existence of 2 distinct subgroups of chronic myeloproliferative diseases (CMPDs),



Figure 1. Distribution of PMN telomerase activity according to clonality expression.

one with monoclonal and the other with polyclonal granulopoiesis, each of them characterized by a specific level of TA, respectively, high and normal, but otherwise undistinguishable by conventional clinical parameters. Monoclonally restricted cells are thought to represent the real neoplastic tissue, but in CMPD they have never been found to express unambiguous morphologic or biomolecular markers.²³

Several questions remain unanswered, and the foremost concerns the relation, if any, between high or low TA and survival. The positive correlation of a high TA with monoclonal granulopoiesis may represent

References

- Blackburn EH. Structure and function of telomere. Nature. 1991;350:569-573.
- Kim NW, Piatyszek MA, Prowse KR, et al. Specific association of human telomerase activity with immortal cells and cancer. Science. 1994; 266:2011-2015.
- Broccoli D, Young JW, de Lange T. Telomerase activity in normal and malignant hematopoietic cells. Proc Natl Acad Sci U S A. 1995;92:9082-9086.
- Ohyashiki JH, Sashida G, Tauchi T, Ohyashiki K. Telomeres and telomerase in hematologic neoplasia. Oncogene. 2002;21:680-687.
- Engelhardt M, Mackenzie K, Drullinsky P, Silver RT, Moore MA. Telomerase activity and telomere length in acute and chronic leukemia, pre- and post-ex vivo culture. Cancer Res. 2000;60:610-617.
- Terasaki Y, Okumura H, Ohtake S, Nakao S. Accelerated telomere length shortening in granulocytes: a diagnostic marker for myeloproliferative diseases. Exp Hematol. 2002;30:1399-1404.
- El Kassar N, Hetet G, Li Y, Briere J, Grandchamp B. Clonal analysis of haemopoietic cells in essential thrombocythaemia. Br J Haematol. 1995;90: 131-137.
- Ferraris AM, Mangerini R, Racchi O, et al. Heterogeneity of clonal development in chronic myeloproliferative disorders. Am J Hematol. 1999;60: 158-160.
- Gilliland DG, Blanchard KL, Levy J, Perrin S, Bunn HF. Clonality in myeloproliferative disorders: analysis by means of the polymerase chain reaction. Proc Natl Acad Sci U S A. 1991;88: 6848-6852.

- Murphy S, Iland H, Rosenthal DS, Laszlo J. Essential thrombocythemia: an interim report from the Polycythemia Vera Study Group. Semin Hematol 1986;23:177-182.
- Racchi O, Mangerini R, Rapezzi D, Rolfo M, Gaetani GF, Ferraris AM. X chromosome inactivation patterns in normal females. Blood Cells Mol Dis. 1998;24:439-447.
- Ohyashiki JH, Ohyashiki K, Iwama H, Hayashi S, Toyama K, Shay JW. Clinical implications of telomerase activity levels in acute leukemia. Clin Cancer Res. 1997;3:619-625.
- Gale RE, Mein CA, Linch DC. Quantification of X-chromosome inactivation patterns in haematological samples using the DNA PCR-based HUMARA assay. Leukemia. 1996;10:362-367.
- Liu E, Jelinek J, Pastore YD, Guan Y, Prchal JF, Prchal JT. Discrimination of polycythemias and thrombocytoses by novel, simple, accurate clonality assays and comparison with PRV-1 expression and BFU-E response to erythropoietin. Blood. 2003;101:3294-3301.
- Kralovics R, Stockton DW, Prchal JT. Clonal hematopoiesis in familial polycythemia vera suggests the involvement of multiple mutational events in the early pathogenesis of the disease. Blood. 2003;102:3793-3796.
- Karasawa M, Tsukamoto N, Yamane A, et al. Analysis of the distribution of CAG repeats and X-chromosome inactivation status of HUMARA gene in healthy female subjects using improved fluorescence-based assay. Int J Hematol. 2001; 74:281-286.
- 17. D'Adda T, Candidus S, Denk H, Bordi C, Hofler H. Gastric neuroendocrine neoplasms: tumour

a powerful prognostic tool in PV and ET, but its significance will only be ascertained by a prospective study of clonality and TA on a larger cohort of patients over a long period of follow-up.

Although ET and PV are diseases with a relatively benign course, death due to progression to myelofibrosis and acute leukemia is not a rare event. Therefore, implications for future efforts to treat the subgroup of patients with monoclonal hematopoiesis and high TA with targeted chemotherapy and/or antitelomerase drugs²⁴ might not be excluded.

clonality and malignancy-associated large X-chromosomal deletions. J Pathol. 1999;189: 394-401.

- Klippel S, Strunck E, Temerinac S, et al. Quantification of PRV-1 mRNA distinguishes polycythemia vera from secondary erythrocytosis. Blood. 2003;102:3569-3574.
- Florensa L, Besses C, Zamora L, et al. Endogenous erythroid and megakaryocytic circulating progenitors, HUMARA clonality assay, and PRV-1 expression are useful tools for diagnosis of polycythemia vera and essential thrombocythemia. Blood. 2004;103:2427-2428.
- Streiff MB, Smith B, Spivak JL. The diagnosis and management of polycythemia vera in the era since the Polycythemia Vera Study Group: a survey of American Society of Hematology members' practice patterns. Blood. 2002;99:1144-1149.
- Shih LY, Lin TL, Lai CL, et al. Predictive values of X-chromosome inactivation patterns and clinicohematologic parameters for vascular complications in female patients with essential thrombocythemia. Blood. 2002;100:1596-1601.
- Fenaux P, Simon M, Caulier MT, Lai JL, Goudemand J, Bauters F. Clinical course of essential thrombocythemia in 147 cases. Cancer. 1990;66: 549-556.
- Harrison CN, Gale RE, Machin SJ, Linch DC. A large proportion of patients with a diagnosis of essential thrombocythemia do not have a clonal disorder and may be at lower risk of thrombotic complications. Blood. 1999;93:417-424.
- Shay JW, Wright WE. Telomerase: a target for cancer therapeutics. Cancer Cell. 2002;2:257-265.