

- gene are associated with lung cancer development. *Clin Cancer Res*. 2002;8:1115-1120.
21. Collins SJ, Ruscetti FW, Gallagher RE, Gallo RC. Normal functional characteristics of cultured human promyelocytic leukemia cells (HL-60) after induction of differentiation by dimethyl sulfoxide. *J Exp Med*. 1979;149:969-974.
 22. Lindmark A, Gullberg U, Olsson I. Processing and intracellular transport of cathepsin G and neutrophil elastase in the leukemic myeloid cell line U-937-modulation by brefeldin A, ammonium chloride, and monensin. *J Leukoc Biol*. 1994;55:50-70.
 23. Benson KF, Li F-Q, Person RE, et al. Mutations associated with neutropenia in dogs and humans disrupt intracellular transport of neutrophil elastase. *Nat Genet*. 2003;35:90-96.
 24. Borregaard N, Lollike K, Kjeldsen L, et al. Human neutrophil granules and secretory vesicles. *Eur J Haematol*. 1993;51:187-198.
 25. Gullberg U, Andersson E, Garwicz D, Lindmark A, Olsson I. Biosynthesis, processing and sorting of neutrophil proteins: insights into neutrophil granule development. *Eur J Haematol*. 1997;58:137-153.
 26. Yoshimura K, Crystal RG. Transcriptional and posttranscriptional modulation of human neutrophil elastase gene expression. *Blood*. 1992;79:2733-2740.
 27. Hunter MG, Druhan LJ, Massullo PR, Avalos BR. Proteolytic cleavage of granulocyte colony-stimulating factor and its receptor by neutrophil elastase induces growth inhibition and decreased cells surface expression of the granulocyte colony-stimulating factor receptor. *Am J Hematol*. 2003;74:149-155.
 28. El-Ouriaghli, Fujiwara H, Melenhorst JJ, Sconocchia G, Hensel N, Barrett AJ. Neutrophil elastase enzymatically antagonizes the in vitro action of G-CSF: implications for the regulation of granulopoiesis. *Blood*. 2003;101:1752-1758.
 29. Belaouaj A, McCarthy R, Baumabb M, et al. Mice lacking neutrophil elastase mutation derived from patients with severe congenital neutropenia have normal granulopoiesis. *Nat Med*. 1998;4:615-618.
 30. Grenda DS, Johnson SE, Mayer JR, et al. Mice expressing a neutrophil elastase mutation derived from patients with severe congenital neutropenia have normal granulopoiesis. *Blood*. 2002;100:3221-3228.
 31. Dillon SR, Mancini M, Rosen A, Schlissel MS. Annexin V binds to viable B cells and colocalizes with a marker of lipid rafts upon B cell receptor activation. *J Immunol*. 2000;164:1322-1332.
 32. Kuijpers TW, Maianski NA, Tool AT, et al. Neutrophils in Barth syndrome (BTSH) avidly bind annexin-V in the absence of apoptosis. *Blood*. 2004;103:3915-3923.
 33. Aprikyan AG, Liles WC, Person RE, Rodger E, Dale DC. Accelerated apoptosis of bone marrow progenitor cells in severe congenital neutropenia [abstract]. *Blood*. 1998;94:482.
 34. Carlsson G, Aprikyan AG, Tehranchi R, et al. Kostmann syndrome: severe congenital neutropenia associated with defective expression of Bcl-2, constitutive mitochondrial release of cytochrome c, and excessive apoptosis of myeloid progenitor cells. *Blood*. 2004;103:3355-3361.
 35. Aprikyan AG, Kutuyavin T, Stein S, et al. Cellular and molecular abnormalities in severe congenital neutropenia predisposing to leukemia. *Exp Hematol*. 2003;31:372-381.
 36. Mackey MC, Aprikyan AA, Dale DC. The rate of apoptosis in postmitotic neutrophil precursors of normal and neutropenic humans. *Cell Prolif*. 2003;31:372-381.
 37. Horwitz M, Benson KF, Duan Z, Li FQ, Person RE. Hereditary neutropenia: dogs explain human neutrophil elastase mutations. *Trends Mol Med*. 2004;10:163-170.
 38. Williams HR, Lin TY, Navia MA. Crystallization of human neutrophil elastase. *J Biol Chem*. 1987;262:17178-17181.
 39. Cai TQ, Wright SD. Human leukocyte elastase is an endogenous ligand for the integrin CR3 (CD11b/CD18, Mac-1, alpha M beta 2) and modulates polymorphonuclear leukocyte adhesion. *J Exp Med*. 1996;184:1213-1223.
 40. Allen DH, Tracy PB. Human coagulation factor V is activated to the functional cofactor by elastase and cathepsin G expressed at the monocyte surface. *J Biol Chem*. 1995;270:1408-1415.
 41. Kaup M, Dassler K, Reineke U, Weise C, Tauber R, Fuchs H. Processing of the human transferrin receptor at distinct positions within the stalk region by neutrophil elastase and cathepsin G. *Biol Chem*. 2002;383:1011-1020.
 42. Duan Z, Li F-Q, Wechsler J, et al. A novel notch protein, N2N, targeted by neutrophil elastase and implicated in hereditary neutropenia [abstract]. *Mol Cell Biol*. 2004;24:58.
 43. Pinkoski MJ, Waterhouse NJ, Heibein JA, et al. Granzyme B-mediated apoptosis proceeds predominantly through a Bcl-2-inhibitable mitochondrial pathway. *J Biol Chem*. 2001;276:12060-12067.
 44. Tehranchi R, Fadeel B, Forsblom A-M, et al. Granulocyte colony-stimulating factor inhibits spontaneous cytochrome c release and mitochondria-dependent apoptosis of myelodysplastic syndrome hematopoietic progenitors. *Blood*. 2003;101:1080-1086.
 45. Shelly LL, Fuchs C, Miele L. Notch-1 inhibits apoptosis in murine erythroleukemia cells and is necessary for differentiation induced by hybrid polar compounds. *J Cell Biochem*. 1999;73:164-178.

Erratum

In the article by Callén et al entitled “A common founder mutation in *FANCA* underlies the world’s highest prevalence of Fanconi anemia in Gypsy families from Spain,” which appeared in the March 1, 2005, issue of *Blood* (Volume 105:1946-1949), there is a mistake in the first sentence of the second paragraph of “Results and discussion.” The Afro-American FA patients of Ghana origin have a truncating mutation in exon 4 of *FANCC*, not in *FANCA*.

Therefore, the phenotypic effects of this mutation cannot be compared with the Spanish Gypsy truncating mutation in exon 4 of *FANCA*.