Brief report

Primary cutaneous follicle center lymphoma and primary cutaneous large B-cell lymphoma, leg type, are both targeted by aberrant somatic hypermutation but demonstrate differential expression of AID

Remco Dijkman, Cornelis P. Tensen, Maike Buettner, Gerald Niedobitek, Rein Willemze, and Maarten H. Vermeer

We assessed primary cutaneous large B-cell lymphoma, leg type (PCLBCL, leg type; n = 13), and primary cutaneous follicle center lymphoma (PCFCL; n = 19) for somatic hypermutation (SHM) of *BCL6*, and aberrant SHM of *MYC*, *RhoH/TTF*, and *PAX5*. We demonstrate SHM of *BCL6* in 8 PCLBCLs (62%), leg type, and 7 PCFCL patients (37%), and aberrant SHM in *PAX5*, *RhoH/TTF*, and/or *MYC* in 7 PCLBCLs (54%), leg type, and 10 PCFCL patients (53%). The majority of mutations consisted of single base-pair substitutions (n = 54) with rare deletions/insertions (n = 4), and displayed molecular features typical of the SHM process. Quantitative real-time PCR and immunohistochemical stainings for activation-induced cytidine deaminase, which is indispensable for SHM, demonstrated significantly higher expression in PCLBCL, leg type. Our results suggest that (aberrant) SHM may contribute to the pathogenesis of PCLBCL, leg type, and PCFCL and is not restricted to diffuse large B-cell lymphomas with an aggressive clinical behavior. (Blood. 2006;107:4926-4929)

© 2006 by The American Society of Hematology

Introduction

The recently published World Health Organization-European Organization for Research and Treatment of Cancer classification for primary cutaneous lymphomas distinguishes 2 main groups of primary cutaneous large B-cell lymphomas (PCLBCLs): primary cutaneous follicle center lymphomas (PCFCLs) with an indolent behavior (5-year survival > 95%) and PCLBCLs, leg type, with an intermediate prognosis (5-year survival = 50%).¹

Recent investigations identified (aberrant) somatic hypermutation (SHM) of proto-oncogenes as a novel mechanism of genetic lesion in diffuse large B-cell lymphomas (DLBCLs).² Normally, the SHM process introduces point mutations in the variable region of immunoglobulin genes (IgV) of germinal center (GC) B cells, allowing affinity maturation of the humoral immune response.³⁻⁵ Remarkably, the SHM process in normal GC B cells is not limited to the IgV sequences but can also target, albeit at a much lower rate, *BCL6*, *CD95*, and the B-cell receptor accessory proteins *CD79b* and *CD79a*.⁶⁻¹⁰

In more than half of the patients with DLBCL, AIDS-associated DLBCL, and Burkitt lymphomas, aberrant activity of the SHM process affects regulatory as well as coding sequences from multiple gene loci, including proto-oncogenes *PIM1*, *PAX5*, *RhoH/TTF*, and *MYC*.^{2,11} In contrast, mutations of *PIM1*, *PAX5*, *RhoH/TTF*, and *MYC* do not occur at a significant level in normal GC B cells.² Expression of AID (activation-induced cytidine deaminase) has been identified as an absolute requirement for SHM.^{12,13} Recent

From the Department of Dermatology, Leiden University Medical Center, Leiden, The Netherlands; and the Institute for Pathology, Friedrich-Alexander-University, Erlangen, Germany.

Submitted August 25, 2005; accepted February 11, 2006. Prepublished online as *Blood* First Edition Paper, February 28, 2006; DOI 10.1182/blood-2005-08-3443.

Supported by grants from The Netherlands Organisation for Health Research and Development (no. 907-00-066; M.H.V.).

R.D., C.P.T., R.W., and M.H.V. designed the study; R.D., C.P.T., and M.B. performed the research; R.D., C.P.T., G.N., M.B., R.W., and M.H.V. analyzed

studies demonstrated expression of AID mRNA and protein in Burkitt lymphomas and systemic DLBCL.¹⁴⁻¹⁹

These observations resulted in the concept that (aberrant) SHM plays a role in lymphomagenesis of aggressive B-cell lymphomas by mutating multiple proto-oncogenes.

We investigated the occurrence of (aberrant) SHM of *BCL6*, *PAX5*, *MYC*, and *RhoH/TTF* in PCFCL and PCLBCL, leg type. In addition, the expression of AID mRNA and protein was studied in both entities.

Study design

Patient material and RNA and DNA extraction

Frozen pretreatment skin biopsies of 19 patients with PCFCL and 17 patients with PCLBCL, leg type, with the histology of a DLBCL and containing 60% or more neoplastic cells were included in this study. Frozen lymph node samples from 2 patients with follicular hyperplasia were included as controls. RNA was extracted from $50 \times 20 \mu$ M sections using a RNeasy isolation kit (Qiagen, Hilden, Germany). DNA was extracted from $25 \times 20 \mu$ M sections using the Genome-tip 20/G isolation kit and Proteinase K solution (Qiagen). This study was performed in accordance with the Dutch code and Leiden University Medical Center guidelines on leftover material. Patients provided informed consent in accordance with the Declaration of Helsinki.

the data; R.D., C.P.T., R.W., and M.H.V. wrote the paper; and all authors checked the final version of the manuscript.

The online version of this article contains a data supplement.

Reprints: Maarten Vermeer, Department of Dermatology, B1-Q, Leiden University Medical Center, PO Box 9600, 2300 RC Leiden, The Netherlands; e-mail: m.h.vermeer@lumc.nl.

The publication costs of this article were defrayed in part by page charge payment. Therefore, and solely to indicate this fact, this article is hereby marked "advertisement" in accordance with 18 U.S.C. section 1734.

© 2006 by The American Society of Hematology

Check for updates

	Survival			Mutation, position*							
		Time, mo 28		BCL6		PAX5		RhoH/TTF		МҮС	
Case	Status		AID	No.	Mutations	No.	Mutations	No.	Mutations	No.	Mutations
1	D0		ND	1	g.751C>A	0	NA	0	NA	1	g.2480G>T
									g.744C>G, g.816G>A,		
2	A0	78	ND	1	g.842T>C	3	g.739C>T, g.846C>T, g.848C>T	3	g.901T>C	0	NA
3	A0	264	Occasional	0	NA	2	g.1082C>T, g.1187C>T		NA	0	NA
4	A+	86	Negative	0	NA	1	g.714_715insT	0	NA	0	NA
5	A0	35	ND	0	NA	0	NA	0	NA	0	NA
6	A0	30	ND	1	g.683-695del	1	g.839-855del	0	NA	1	g.2739G>T
									g.469A>G,		
7	A0	28	Negative	1	g.713G>A	0	NA	2	g.609G>A	1	g.2710C>T
8	A0	23	ND	0	NA	0	NA	0	NA	0	NA
9	A0	42	ND	1	g.798G>A	0	NA NA		NA	0	NA
							g.893C>T, g.1046C>T, g.1206G>A, g.1223G>A, g.1338C>G, g.1357C>G,				
10	A0	74	ND	2	g.766C>A, g.809G>A	7	g.1435G>C	0	NA	1	g.2541C>A
11	A0	63	Occasional	0	NA	0	NA	0	NA	0	NA
12	A0	57	Negative	0	NA	0	NA	0	NA	1	g.2466A>G
13	A0	62	ND	0	NA	0	NA	0	NA	0	NA
14	A0	69	ND	1	g.759G>A	0	NA	0	NA	1	g.2366C>T
15	D0	70	Negative	0	NA	0	NA	0	NA	0	NA
16	A0	52	ND	0	NA	0	NA	0	NA	0	NA
17	A0	40	Negative	0	NA	0	NA	0	NA	0	NA
18	D+	130	Negative	0	NA	0	NA	0	NA	0	NA
19	A0	113	ND	0	NA	0	NA	0	NA	1	g.2616C>G

Table 1. Overview of DCLO, FAXJ, RHOR/TTF, and MTC indiations in FCFCL
--

A0 indicates alive without disease; A+, alive with disease; D0, death unrelated to lymphoma; D+, death related to lymphoma; negative, no tumor cells stained; occasional, less than 10% of tumor cells stained; NA, not applicable; and ND, not determined.

*Sequences are aligned to the corresponding germline sequences. GenBank accession numbers are AY189709 (BCL6), AF386791 (PAX5), AF386789 (RhoH/TTF), and X00364 (MYC).

Detection of somatic mutations

PAX5, *RhoH/TTF*, and *MYC*.^{11,20} PCR products were purified and sequenced as described.²¹

Quantitative real-time PCR

In 13 cases of PCLBCL, leg type, and 19 cases of PCFCL, mutations in *BCL6*, *PAX5*, *RhoH/TTF*, and *MYC* were investigated. Mutation analysis was restricted to regions previously shown to contain more than 90% of mutations in systemic DLBCL.² Polymerase chain reaction (PCR) amplification of DNA was performed using previously described primer pairs covering a part of the promoter region, the first exon and intron of *BCL6*,

In 10 cases of PCLBCL, leg type, and 8 cases of PCFCL, quantitative real-time PCR for AID was performed. cDNA synthesis, real-time PCR, and data analysis were performed as described previously.²¹ For the

Table 2. Overview of DOLD, TAXS, THORY TT, and WTO mutations in TOLDOL, leg typ	Table 2. Overview of BCL6.	, PAX5, RhoH/TTF	and MYC mutations in	PCLBCL, leg typ	e
---	----------------------------	------------------	----------------------	-----------------	---

	Survival			Mutation, position*							
		Time, mo		BCL6		PAX5		RhoH/TTF		МҮС	
Case	Status		AID	No.	Mutations	No.	Mutations	No.	Mutations	No.	Mutations
20	A0	75	Double-positive	1	g.835T>C	2	g.883C>T, g.1070G>A	2	g.574T>G, g.589C>G	0	NA
21	D+	73	Double-positive	1	g.704G>C	1	g.1334C>T	1	g.912G>A	0	NA
22	D+	29	Double-positive	2	g.545G>T, g.661G>A	0	NA	0	NA	1	g.2405delC
23	A0	72	Negative	0	NA	0	NA	0	NA	0	NA
24	D+	20	Double-positive	1	g.799C>A	0	NA	0	NA	0	NA
25	D+	12	Occasional	2	g.552A>C, g.845C>T	0	NA	0	NA	0	NA
26	D0	10	Occasional	1	g.837T>G	0	NA	0	NA	1	g.4835G>C p.Ser146Leu
27	A0	31	Double-positive	0	NA	1	g.1347G>A	0	NA	0	NA
28	D+	26	ND	2	g.782C>G, g.794G>A	0	NA	2	g.368G>A, g.682C>T	0	NA
29	D+	3	ND	2	g.778C>T, g.779G>C	0	NA	0	NA	0	NA
30	D+	9	Negative	0	NA	0	NA	0	NA	0	NA
31	A0	76	Double-positive	0	NA	0	NA	0	NA	0	NA
32	D+	38	ND	0	NA	0	NA	1	g.306T>C	0	NA

A0 indicates alive without disease; A+, alive with disease; D0, death unrelated to lymphoma; D+, death related to lymphoma; negative, no tumor cells stained; occasional, less than 10% of tumor cells stained; positive, 10% to 50% of tumor cells stained; double-positive, more than 50% of tumor cells stained; and ND, not determined. *Sequences are aligned to the corresponding germline sequences. GenBank accession numbers are AY189709 (*BCL6*), AF386791 (*PAX5*), AF386789 (*RhoH/TTF*), and X00364 (*MYC*).



Figure 1. Expression of AID mRNA in PCFCL and PCLBCL, leg type. Expression of AID mRNA was measured in triplicate using quantitative real-time PCR and normalized to the expression of the housekeeping gene small nuclear ribonucleoprotein particle U1A (U1A). Error bars indicate standard deviations. Case numbers correspond to cases presented in Tables 1-2. dotted lines indicate the average AID mRNA expression within the group of PCFCL and PCLBCL, leg type; – indicates no tumor cells stained; +/- indicates less than 10% of tumor cells stained; LN indicates lymph nodes; IHC, immunohistochemical staining; and nd, not determined.

amplification of AID (*AID*, accession no. NM_020 661) and the housekeeping gene small nuclear ribonucleoprotein particle U1A (*U1A*, accession no. X06 347) the following primers were used: AID-F, 5'-AGA GGC GTG ACA GTG CTA CA-3'; AID-R, 5'-TGT AGC GGA GGA AGA GCA AT-3'; U1A-F, 5'-GCA GCT TAT GCC AGC ACA GAT-3'; and U1A-R, 5'-TTG GTG AGG AAC AAG ATG TGA TTC-3'.

Immunohistochemical stainings for AID

In 12 cases of PCLBCL, leg type, and 8 cases of PCFCL, immunohistochemical stainings for AID, using antibody Eh2-5G9, were performed as previously described.¹⁸

Results and discussion

This study demonstrates the presence of SHM in *BCL6* in 8 (62%) of 13 patients with PCLBCL, leg type, and 7 (37%) of 19 patients with PCFCL, and aberrant SHM in *PAX5*, *RhoH/TTF*, and/or *MYC* in 7 (54%) of 13 patients with PCLBCL, leg type, and 10 (53%) of 19 patients with PCFCL (Tables 1 and 2). The molecular profile of the detected mutations was similar in PCFCL and PCLBCL, leg



Figure 2. Expression of AID by tumor cells in PCLBCL, leg type. Neoplastic B cells of primary cutaneous large B-cell lymphoma, leg type, show positive staining for AID protein (original magnification \times 40). Images were captured with a Leica DMLB microscope (Leica, Bensheim, Germany), using a Leica PL FLUOTAR objective lens (40 \times /0.70) and a Leica DC200 camera. Images were directly imported into Powerpoint (Microsoft, Redmond, WA), using the Leica DC200 camera software (version 2.51).

type, and resembled the characteristics of mutations induced by the physiologic SHM process (Table S1, available on the *Blood* website; see the Supplemental Tables link at the top of the online article).³⁻⁵ These data provide additional evidence that PCLBCL, leg type, and PCFCL are derived from germinal center experienced B cells.

Similar to studies of systemic DLBCL, mutations were most frequently demonstrated in BCL6, and the mutation frequency for aberrant SHM in PAX5, MYC, and RhoH/TTF was generally lower than that observed for SHM of BCL6 sequences in the same cases (Tables S2-S3).^{2,11} A higher mutation frequency for BCL6 was observed in PCLBCL, leg type (P < .05, Student t test). Because one-third of patients with PCLBCL, leg type, harbor translocations affecting BCL6, whereas this chromosomal aberration is not observed in PCFCL, these data are in accordance with previous studies that suggested that SHM could play a role in the induction of chromosomal translocations.^{2,22} The functional consequences of the mutations detected in this study include potential deregulated gene transcription resulting from mutations in the 5' regulatory regions of BCL6, PAX5, RhoH/TTF, and MYC. In addition, in one patient with PCLBCL, leg type, a mutation in exon 2 of MYC, representing a mutational hotspot in translocated MYC alleles in Burkitt lymphoma and mouse plasmacytoma, was found.^{23,24}

Based on observations in systemic B-cell lymphomas, it was postulated that aberrant SHM is a selective feature of aggressive B-cell lymphomas.^{2,11} In our study, however, we demonstrate aberrant SHM in PCFCLs that are characterized by an indolent behavior and PCLBCL, leg type, with an intermediate prognosis. Also within the group of PCLBCL, leg type, the prevalence of aberrant SHM did not correlate with survival.

The mean expression level of AID mRNA, normalized to the housekeeping genes, was significantly higher in PCLBCL, leg type, than in PCFCL (P < .01, Student *t* test; Figure 1). Expression of AID protein was confined to the cytoplasm of neoplastic cells (Figure 2) and was demonstrated in 10 (83%) of 12 patients with PCLBCL, leg type (with more than 50% of tumor cells staining in 6 patients), whereas in PCFCL, AID protein was not detected in 6 of 8 patients and was expressed by less than 10% of tumor cells in 2 cases (Tables 1-2). A close correlation between AID transcripts and protein expression was found in the 10 patients in whom both quantitative real-time PCR and immunohistochemical stainings were performed (Figure 1). No correlation was found between expression of AID and occurrence of SHM, which is in line with previous observations.^{15,19}

Because PCLBCL, leg type, and PCFCL display an ABC-like and GCB-like gene expression profile, respectively,25 our observations are in line with studies in nodal DLBCL, which demonstrated that AID protein expression is tendentially higher in the ABC-like DLBCL.¹⁹

In conclusion, we demonstrate that in the majority of patients with PCFCL and PCLBCL, leg type, the proto-oncogenes BCL6, PAX5, RhoH/TTF, and, to a lesser extent, MYC are affected by (aberrant) SHM. In addition, we show that within the group of PCLBCLs, expression of AID is found in the vast majority of cases of PCLBCL, leg type, but is rarely detected in PCFCL. These results demonstrate that aberrant SHM is not restricted to B-cell non-Hodgkin lymphoma with an aggressive behavior, and suggest that persistent expression of AID may contribute to the lymphomagenesis of PCLBCL, leg type.

Acknowledgments

We thank Wim Zoutman and Jacqueline van Harmelen for their excellent technical assistance. We would like to thank Elisabeth Kremmer for providing the antibody EK2-5G9.

2005 205 541-547

651-659

1993:5:56-61.

18. Greiner A, Tobollik S, Buettner M, et al. Differen-

Cancer Res. 2004;64:5578-5586.

tions in vivo. Cell. 2004;118:431-438.

phoma and mouse plasmacytomas. Nat Genet.

ES, Raffeld M. Clustered mutations in the second

exon of the MYC gene in sporadic Burkitt's lym-

24. Yano T, Sander CA, Clark HM, Dolezal MV, Jaffe

25. Hoefnagel JJ, Dijkman R, Basso K, et al. Distinct

types of primary cutaneous large B-cell lym-

phoma identified by gene expression profiling.

phoma. Oncogene. 1993;8:2741-2748.

Blood. 2005;105:3671-3678.

tial expression of activation-induced cvtidine

deaminase (AID) in nodular lymphocyte-predomi-

nant and classical Hodgkin lymphoma. J Pathol.

References

- 1. Willemze R, Jaffe ES, Burg G, et al. WHO-EORTC classification for cutaneous lymphomas. Blood. 2005;105:3768-3785
- 2. Pasqualucci L, Neumeister P, Goossens T, et al. Hypermutation of multiple proto-oncogenes in B-cell diffuse large-cell lymphomas. Nature. 2001:412:341-346
- 3. Klein U, Goossens T, Fischer M, et al. Somatic hypermutation in normal and transformed human B cells. Immunol Rev. 1998;162:261-280.
- 4. Rajewsky K. Clonal selection and learning in the antibody system. Nature. 1996;381:751-758.
- 5. Jacobs H, Bross L. Towards an understanding of somatic hypermutation. Curr Opin Immunol. 2001;13:208-218.
- 6. Pasqualucci L, Migliazza A, Fracchiolla N, et al. BCL-6 mutations in normal germinal center B cells: evidence of somatic hypermutation acting outside Ig loci. Proc Natl Acad Sci U S A. 1998; 95:11816-11821.
- 7. Gordon MS, Kanegai CM, Doerr JR, Wall R. Somatic hypermutation of the B cell receptor genes B29 (Igbeta, CD79b) and mb1 (Igalpha, CD79a). Proc Natl Acad Sci U S A. 2003;100:4126-4131.
- 8. Shen HM, Peters A, Baron B, Zhu X, Storb U. Mutation of BCL-6 gene in normal B cells by the process of somatic hypermutation of Ig genes. Science. 1998;280:1750-1752.
- 9. Peng H-Z, Du M-Q, Koulis A, et al. Nonimmunoalobulin gene hypermutation in germinal center B cells. Blood. 1999;93:2167-2172.
- 10. Müschen M, Re D, Jungnickel B, Diehl V, Rajewsky K, Küppers R. Somatic mutation of the CD95 gene in human B cells as a side-effect of the ger-

minal center reaction. J Exp Med. 2000;18:1833-1840

- 11. Gaidano G, Pasqualucci L, Capello D, et al. Aberrant somatic hypermutation in multiple subtypes of AIDS-associated non-Hodgkin lymphoma. Blood. 2003;102:1833-1841.
- 12. Muramatsu M, Kinoshita K, Fagarasan S, Yamada S. Shinkai Y. Honio T. Class switch recombination and hypermutation require activation-induced cvtidine deaminase (AID), a potential RNA editing enzyme. Cell. 2000;102:553-563.
- 13. Revy P. Muto T. Levy Y. et al. Activation-induced cytidine deaminase (AID) deficiency causes the autosomal recessive form of the Hyper-IgM syndrome (HIGM2). Cell. 2000;102:565-575.
- 14. Smit LA, Bende RJ, Aten J, et al. Expression of activation-induced cytidine deaminase is confined to B-cell non-Hodgkin's lymphomas of germinalcenter phenotype. Cancer Res. 2003;63:3894-3898
- 15. Lossos IS, Levy R, Alizadeh AA. AID is expressed in germinal center B-cell-like and activated B-celllike diffuse large-cell lymphomas and is not correlated with intraclonal heterogeneity. Leukemia. 2004:18:1775-1779.
- 16. Hardianti MS, Tatsumi E, Syampurnawati M, et al. Expression of activation-induced cytidine deaminase (AID) in Burkitt lymphoma cells: rare AIDnegative cell lines with the unmutated rearranged VH gene. Leuk Lymphoma. 2004;45:155-160.
- 17. Greeve J, Philipsen A, Krause K, et al. Expression of activation-induced cytidine deaminase in human B-cell non-Hodgkin lymphomas. Blood. 2003;101:3574-3580.
- 19. Pasqualucci L, Guglielmino R, Houldsworth J, et al. Expression of the AID protein in normal and neoplastic B cells. Blood. 2004;104:3318-3325. 20. Capello D, Vitolo U, Pasqualucci L, et al. Distribution and pattern of BCL-6 mutations throughout the spectrum of B-cell neoplasia. Blood. 2000;95: 21. van Doorn R, Dijkman R, Vermeer MH, et al. Aberrant expression of the tyrosine kinase receptor EphA4 and the transcription factor twist in Sezary syndrome identified by gene expression analysis. 22. Ramiro AR, Jankovic M, Eisenreich T, et al, AID is required for c-myc/IgH chromosome transloca-23. Bhatia K, Huppi K, Spangler G, Siwarski D, Iyer R, Magrath I. Point mutations in the c-Myc transactivation domain are common in Burkitt's lym-