Expression of interleukin-3 receptor subunits on defined subpopulations of acute myeloid leukemia blasts predicts the cytotoxicity of diphtheria toxin interleukin-3 fusion protein against malignant progenitors that engraft in immunodeficient mice

Leman Yalcintepe, Arthur E. Frankel, and Donna E. Hogge

The interleukin-3 receptor (IL-3R) subunits are overexpressed on acute myeloid leukemia (AML) blasts compared with normal hematopoietic cells and are thus potential targets for novel therapeutic agents. Both fluorescence-activated cell sorter (FACS) analysis and quantitative real-time reverse transcriptionpolymerase chain reaction (QRT-PCR) were used to quantify expression of the IL-3R α and β_c subunits on AML cells. QRT-PCR for both subunits was most predictive of killing of AML colony-forming cells (AML-CFCs) by diphtheria toxinIL-3 fusion protein (DT₃₈₈IL3). Among 19 patient samples, the relative level of the IL-3R α was higher than the IL-3R β_c and highest in CD34⁺CD38⁻CD71⁻ cells, enriched for candidate leukemia stem cells, compared with cell fractions depleted of such progenitors. Overall, the amount of IL-3R β_c subunit did not vary among sorted subpopulations. However, expression of both subunits varied by more than 10-fold among different AML samples for all subpopulations studied. The level of IL-3R β_c expression versus glyceraldehyde-3-phosphate dehydrogenase (GAPDH) (set

at 1000) ranged from 0.14 to 13.56 in CD34⁺CD38⁻CD71⁻ cells from different samples; this value was correlated (r = .76, P = .05) with the ability of DT₃₈₈IL3 to kill AML progenitors that engraft in β_2 -microglobin–deficient nonobese diabetic/ severe combined immunodeficient (NOD/ SCID) mice (n = 7). Thus, quantification of IL-3R subunit expression on AML blasts predicts the effectiveness IL-3R-targeted therapy in killing primitive leukemic progenitors. (Blood. 2006;108:3530-3537)

© 2006 by The American Society of Hematology

Introduction

Acute myeloid leukemia (AML) is a clonal disorder of hematopoiesis characterized by the accumulation of large numbers of myeloid blast cells that fail to differentiate into functional blood cells.¹ In spite of these obvious abnormalities in cell maturation there are now considerable data demonstrating a hierarchy of malignant progenitor cells in AML similar to that found in normal hematopoiesis and the existence of a small subpopulation of leukemia stem cells (LSCs) that possesses extensive proliferative capacity and the potential for self-renewal.^{2,3} Leukemic cells that initiate long-term hematopoiesis in stromal coculture (AML LTC-ICs) and suspension cultures and engraft in nonobese diabetic/severe combined immunodeficient (NOD/SCID) mice (NOD/SCID leukemiainitiating cells [N/SL-ICs]) are typically present at a frequency of fewer than 1 in 10 000 malignant cells and express a cell-surface phenotype similar to that seen on normal multipotential long-term culture-initiating cells (LTC-ICs) and lymphomyeloid NOD/SCID mouse competitive repopulating units.⁴⁻⁷ Several recent studies have shown that N/SL-ICs can be phenotypically defined as CD34⁺, CD38⁻, human leukocyte antigen (HLA)-DR⁻, CD90⁻, CD71⁻, CD117⁻, and CD123⁺. This antigenic profile is intriguing in that it overlaps with normal stem cells (CD34⁺, CD38⁻, CD71⁻, HLA-DR⁻) but also has leukemia-specific features (CD90⁻, CD117⁻, and CD123⁺⁺).^{4,8-16} The frequent relapses observed in patients with AML who achieve an initial remission of their leukemia suggest that LSCs might be relatively resistant to conventional chemotherapy and also might be responsible for re-establishing the malignant clone in patients with leukemia. These cells are thus obvious targets for novel therapeutic approaches in AML.

The human interleukin-3 receptor (IL-3R) is a heterodimeric structure.¹⁷ The α subunit is essential for ligand binding and confers specificity on the receptor. The common β subunit (β_c), which is shared by the granulocyte-macrophage colony-stimulating factor (GM-CSF) and IL-5 receptors, is required for high-affinity ligand binding and signal transduction. The IL-3R is expressed on leukemic blasts from most patients with AML.^{14,18,19} The α subunit, in particular, is often expressed at much higher levels than are typically seen in normal hematopoietic cells and progenitors and is also detected on subpopulations of AML blasts enriched for N/SL-ICs.^{14,20} Thus, the IL-3R may be an appropriate target for cytotoxic drugs designed to selectively kill AML cells while sparing their normal hematopoietic-cell counterparts.

Fusion proteins in which the diphtheria toxin (DT) catalytic and translocation domains are genetically fused to a growth factor are a novel class of molecules which can selectively target and kill malignant cells expressing the relevant receptor.²¹⁻²⁶ After ligand binding and receptor-mediated endocytosis, the toxin translocates to the cytosol, where it ADP-ribosylates elongation factor 2,

From the Terry Fox Laboratory, British Columbia Cancer Agency, Vancouver, BC, Canada; and the Scott and White Memorial Hospital, Temple, TX.

The authors declare no competing financial interests.

© 2006 by The American Society of Hematology

Submitted April 3, 2006; accepted July 6, 2006. Prepublished online as *Blood* First Edition Paper, August 1, 2006; DOI 10.1182/blood-2006-04-013813.

Supported by the Canadian Institute for Health Research and National Cancer Institute of Canada (D.E.H.), National Institutes of Health (NIH) grants RO1CA76178and RO1CA90263, and Leukemia and Lymphoma Society grant 6006-05 (A.E.F.).

Reprints: Donna E. Hogge, Terry Fox Laboratory, BC Cancer Agency, 675 West 10th Ave, Vancouver, BC V5Z 1L3 Canada; e-mail: dhogge@ bccancer.bc.ca.

The publication costs of this article were defrayed in part by page charge payment. Therefore, and solely to indicate this fact, this article is hereby marked "advertisement" in accordance with 18 USC section 1734.

leading to inactivation of protein synthesis and cell death.²⁷⁻³⁰ $DT_{388}IL3$ consists of the truncated form of diphtheria toxin (DT_{388}) linked to human IL-3. This fusion protein has been shown to target and kill cells which express the IL-3R, including AML blasts and progenitors.³¹ Previous studies have shown that this molecule will kill AML colony forming cells (AML-CFCs), long-term cultureinitiating cells (AML LTC-ICs), and N/SL-ICs from many patient samples while showing little or no toxicity against analogous normal bone marrow progenitors.¹⁸ Although DT₃₈₈IL3 is effective against many AML samples, only partial eradication of malignant progenitors was seen with some leukemias in spite of detectable expression of high-affinity IL-3R on target blasts. More recently, we determined that the level of IL-3R subunit expression (particularly the β_c subunit) as detected by quantitative real-time reverse transcription-polymerase chain reaction (ORT-PCR) in AML blasts correlated with the effectiveness of DT388IL3 in killing AML-CFCs.32 However, in this study the relationship was not strong for a subset of patient samples and its usefulness in predicting the ability of DT₃₈₈IL3 to kill N/SL-ICs was not investigated thoroughly.

The current study examined the possibility that expression of the IL-3R subunits might be heterogeneous among AML cells and progenitors. We were particularly interested in the possibility that IL-3R expression on the most primitive AML cells (ie, the candidate LSCs that initiate long-term engraftment and proliferation in immunodeficient mice), might be different than the expression detected on the majority population of leukemic cells that have limited proliferative capacity. The level of IL-3R subunit expression was measured by QRT-PCR in subpopulations of leukemic cells enriched for AML progenitors with a multiparameter fluorescence-activated cell sorter (FACS) strategy. These levels were then compared with the ability of DT388IL3 to kill N/SL-ICs from many of the same AML samples. The results demonstrate that the level of expression of the IL-3R β_c (but not the IL-3R α) in unsorted blasts as well as subpopulations enriched for N/SL-ICs predicted the ability of DT₃₈₈IL3 to kill these progenitors from most patient samples. A phase 1/2 clinical trial testing DT₃₈₈IL3 in patients with relapsed or refractory AML is currently in progress. These data suggest that screening patient blast samples for expression of IL-3R subunits may allow prediction of therapeutic benefit with this agent.

Patients, materials, and methods

Blood and marrow samples

Peripheral-blood (PB) samples from 19 newly diagnosed patients with AML and bone marrow (BM) from 3 healthy allogeneic transplant donors were obtained with the approval of the Clinical Research Ethics Board of the University of British Columbia and after obtaining informed consent from patients with AML and from donors in accordance with the Declaration of Helsinki. Light-density mononuclear cells were isolated and cryopreserved as previously described.5 Diagnosis and classification of AML were based on the criteria of the French-American-British (FAB) group.33 Cytogenetic analysis was performed on diagnostic bone marrow samples of patients with AML. Thawed cells were washed twice in Iscove modified Dulbecco medium (IMDM) containing 10% fetal calf serum (FCS) before use in the experiments described below. Normal BM was enriched for CD34⁺ cells using an immunomagnetic separation technique (EasySep; StemCell Technologies, Vancouver, BC, Canada) prior to the FACS isolation of subpopulations described under "FACS analysis and subpopulation isolation of AML and normal cells."

DT₃₈₈IL-3 fusion protein

DT₃₈₈IL3 was constructed by joining the human IL-3 cDNA to a truncated diphtheria toxin (DT₃₈₈) sequence that lacks the native binding site via an intervening (G4S)₂ linker. Recombinant protein was prepared, purified, stored, and tested as described previously.³⁴ This material was found to kill IL-3R⁺ cell lines and AML-CFCs from primary AML samples expressing high-affinity IL-3Rs.^{19,31,34} The IL-3R binding affinity of this fusion toxin is similar to that of native human IL-3.²² In addition, the toxicity of DT₃₈₈IL3 against IL-3R–bearing TF1/Bcl2 cells was reduced more than 120-fold by coincubation with excess human IL-3.³¹

Cultures of AML cells

AML cells were incubated for 24 hours at 37°C at a concentration of 1×10^6 cells/mL with or without $DT_{388}IL3$. Equal fractions of the cells recovered from cultures with or without toxin were assayed in AML-CFC assay or in β_2 -microglobin–deficient NOD/SCID mice without regard to change in cell numbers.

AML-CFC assays were performed by plating cells at 0.2 to 1.0×10^5 cells/mL in methylcellulose-based medium (StemCell Technologies) as previously described. Cultures were scored after 14 days at 37°C in a humidified atmosphere with 5% CO₂ for the presence of clusters (4-20 cells) and colonies (more than 20 cells).⁵

FACS analysis and subpopulation isolation of AML and normal cells

Subpopulation analysis and isolation from AML PB and CD34⁺-enriched normal BM cells were performed with monoclonal antibodies directly coupled to the fluorochromes fluorescein isothiocyanate (FITC), phycoerythrin (PE), and cyanin5-succinimidylester (Cy5). Anti-CD34-Cy5 and anti-CD71-FITC were kind gifts from Dr Peter Lansdorp (Terry Fox Laboratory, Vancouver, BC, Canada); Anti-CD38-PE was obtained from Becton Dickinson (San Jose, CA). Biotinylated antibodies recognizing CD123 (IL-3R α) and CD131 (IL-3R β_c) were visualized by subsequent labeling with streptavidin-PE or -allophycocyanin (APC) (Becton Dickinson). Cells were analyzed on the basis of fluorescence intensity on a dual laser FACSaria instrument (Becton Dickinson). Gates were set to exclude nonviable propidium iodide+ (PI+) cells, and 100% of cells labeled with an irrelevant isotype control antibody. For analysis of IL-3R subunit expression the mean fluorescence intensity (MFI) and percentage of total AML blasts labeled with anti-CD123 and anti-CD131 were calculated using Cell Quest Software (Becton Dickinson).

Subpopulations of AML and normal BM cells labeled with anti-CD34, anti-CD38, and (for AML cells) anti-CD71 were sorted into IMDM (StemCell Technologies) with 20% FCS in microcentrifuge tubes at 4°C. Sort windows used to isolate the specific subfractions are shown in Figure 1. Sorted fractions were washed and used for RNA isolation and QRT-PCR analysis.

QRT-PCR analysis

Total RNA was extracted from sorted cells using the Absolutely RNA, Microprep, or Nanoprep kits (Stratagene, La Jolla, CA). The RT reaction



Figure 1. FACS strategy for isolation of AML cells based on expression of CD34, CD38, and CD71. (A) Cells labeled with anti-CD34–Cy5 (R2) were sorted on the basis of their expression of CD38-PE and CD71-FITC (B). Gate R1 cells were isolated as the CD34– population. Sort gates in panels A and B were set to exclude 100% of the cells labeled with an isotype-control antibody.

was performed in 20 µL with superscript II reverse transcriptase (Invitrogen, Burlington, ON, Canada) using random hexamer oligonucleotides (Amersham Pharmacia, Piscataway, NJ). Real-time PCR was performed using 12.5 μ L 2 × SYBR Green PCR Master Mix (Applied Biosystems, Foster City, CA), 1 µL of 20 pM-specific primers, 1 to 2 µL cDNA, and water to a final volume of 25 µL. Specific forward and reverse primers to produce approximately 100-bp amplicons for optimal amplification in real-time PCR of reverse-transcribed cDNA for human IL-3R α were 5'-GACCTGTACTTGAACGTTGCC and 5'-GAAACGACACCCGAT-ACGTGT, for human IL-3R β_c were 5'-GCAGCATGTCGGCCTTCACTA and 5'-GTCCCCGAATCCTACAGGGAA, and human glyceraldehyde-3phosphate dehydrogenase (GAPDH) were (5'-CCCATCACCATCTTCCAG-GAG and 5'-CTTCTCCATGGTGGTGAAGACG. Real-time PCR and data analysis were performed on an iCycler iQ system, using iCycler iQ Real-time Detection Software (Bio-Rad, Hercules, CA). The relative quantification data of human IL-3R α and IL-3R β_c compared with a reference gene (GAPDH) was generated on the basis of a mathematical model for relative quantification in real-time RT-PCR as described. 32,35,36

Transplantation and detection of AML cells in β_2 -microglobulin-deficient NOD/SCID mice

 $β_2$ -microglobulin–deficient NOD/SCID ($β_2m^{-/-}$ NOD/SCID) mice³⁷ were bred and maintained under sterile conditions in the British Columbia Cancer Research Center Joint Animal Facility according to protocols approved by the Animal Care Committee of the University of British Columbia. These animals were used based on our previous investigations demonstrating that they could allow more efficient engraftment of many AML patient samples than NOD/SCID animals.³⁸ Mice, 8 to 10 weeks of age, received 350 cGy from a ¹³⁷Cs source 16 to 24 hours before injection of AML cells. AML cells (1 to 3 × 10⁷) previously incubated for 24 hours with or without DT₃₈₈IL3 were injected via the tail vein. Femoral bone marrow aspirations were performed every 4 weeks after the injection of AML cells under isoflurane inhalation anesthesia.³⁹ At 16 weeks, mice were killed by CO₂ inhalation, and BM cells were obtained from the 4 long bones by flushing with α-MEM with 50% FCS.

Mouse BM cells were prepared for FACS analysis as previously described⁷ and analyzed for the expression of CD45, a human-specific panleukocyte marker, using an antibody prepared in our center from American Type Culture Collection (ATCC) (clone no. HB10508; Rockville, MD). FACS analysis was performed on a Becton Dickinson FACScan or FACSort flow cytometer. The percentage of CD45⁺ cells was determined after excluding 99.9% of cells labeled with the isotype control and

Age, y/sex

35/F

61/F

65/F

75/F

59/M

22/F

59/N

64/M

70/F

63/F

74/M

41/M

56/N

53/M

52/F

21/M

49/M

17/F

Table 1. Patient characteristics

Patient no.

1

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

nonviable cells. Nonspecific binding of CD45 on mouse BM cells is reliably less than 0.1% to 0.3%.⁷ Thus, a difference between the immunoglobulin G1–negative (IgG1⁻) control and the CD45 expression of the treated mice of more than 0.1% was regarded as evidence for engraftment of human cells. Values shown for engraftment of AML cells in mouse BM are the mean values (\pm SD) of CD45⁺ cells (%) obtained for all mice in the cohort that survived to the time of analysis.

Fluorescence in situ hybridization

Cells from mouse BM were cytocentrifuged onto glass microscope slides and then fixed in methanol–glacial acetic acid. The human-specific centromeric repeat probe used to detect the +8 abnormality (plasmid D8Z2; ATCC) and the yeast artificial chromosome clone containing 550 kb human DNA encompassing the breakpoint on 16p13 (CEPHy904E02 854; Max Planck Institute, Berlin, Germany) to detect the inv(16) were labeled by nick translation with digoxigenin (DIG; Boehringer Mannheim, Mannheim, Germany). Chromosomes 8 and 16 probes and slides were treated as previously described.⁵ For probe detection, slides were incubated with a sheep anti-DIG–FITC antibody (Boehringer Mannheim) and counterstained with PI as previously described.⁵ Slides were visualized on a Zeiss Axioplan (Oberkochen, Germany) fluorescence microscope using double or triple bandpass filters (Omega Optical, Battlebore, NC) to allow simultaneous visualization of FITC and PI signals.

Statistical analysis

Mean values of IL-3R subunit expression in subpopulations of AML and normal BM cells and correlation coefficients between variables were determined in Excel (Microsoft, Redmond, WA), and the significance of differences and correlations determined using a Student t test. A P value of less than .05 was considered significant.

Results

Table 1 shows the clinical characteristics of the 19 patients with AML patients whose cells were tested in this study. Their ages at diagnosis ranged from 17 to 75 years and the FAB subtype of these leukemias was M4 (n = 10), M1 (n = 5), or M5 (n = 3). Cytogenetic analysis of the diagnostic BM specimens revealed abnormalities associated with a good prognosis [inv(16)] in 3 patients and a

FAB	Marrow cytogenetics	WBC, ×10 ⁹ /L (% blasts in PB)
M5b	47, XY, +8	170 (71)
M1	46, XX	87 (94)
M4	44-45, XX, -3, -5, -7, -8 add(8), p23	24 (26)
M4	46,XX	205 (88)
M4	46,XX, del(11q)(q14;q25)	215 (90)
M1	46, XYinv(16)(p13;q22)	76 (90)
M4	46, XX,t(6;9)(p23;q34)	29 (38)
M4	46, XY	101 (90)
M1	49, XY, +10,+?13,+19	189 (75)
M4	46, XX	74 (16)
M1	Unsuccessful	295 (95)
M5	46, XY, t(8;16)(p11;p13)	85 (60)
M4eo	46, XY, inv(16)	238 (90)
ND	45, X, -Y	300 (95)
M4	47, XY, -4,del(5)(q12;q23)add(7)(p12),-12,-17	73 (46)
M4eo	46, XX	155 (45)
M1	46, XY, add(6)(p23)?+(6;11)(p27;q23)	320 (90)
M4eo	46, XY, inv(16)(p13;q22)idem,+15,+22	126 (60)
M5a	47, XX, +8,del(5)(q31;q?33)	205 (90)

WBC indicates white blood cell; M, male; F, female; and ND, not done.

poor prognosis in 6 of these 19 patients according to the Medical Research Council (MRC, United Kingdom) or Southwest Oncology Group (SWOG) criteria.^{40,41}

Comparison of QRT-PCR and FACS for detection of IL-3R subunit expression

In initial experiments the levels of expression of IL-3R subunits in AML blasts as detected by QRT-PCR was compared with detection by FACS analysis for 12 of the 19 patient samples shown on Table 1 using antibodies which detect IL-3R α (CD123) and IL-3R β_c (CD131) and primers and conditions for QRT-PCR described in "Patients, methods, and materials," and previously.32 As shown in Figure 2 there was a weak correlation between the level of IL-3R β_c expression detected by QRT-PCR and MFI for this molecule detected by FACS for CD131 (r = 0.40). No significant correlation was seen between IL-3Ra subunit expression detected by QRT-PCR and MFI detected for CD123. In addition, there was no significant correlation between the percentages of CD123⁺ or CD131⁺ cells quantified by FACS and the MFI or the value obtained by QRT-PCR for the same molecule. MFI observed for CD131 (but not CD123) correlated significantly (r = 0.74) with the percentage of AML-CFCs killed by DT₃₈₈IL3 while the percentages of CD131⁺ or CD123⁺ cells did not predict the AML-CFC killed. QRT-PCR levels for both the IL-3R α and β_c subunit expressions correlated significantly (r = 0.59 and 0.70, respectively) with AML-CFCs killed with DT₃₈₈IL3 (Table 2). Because QRT-PCR values for both the IL-3R α and β_c subunits from a given AML sample predicted DT₃₈₈IL3 cytotoxicity against AML-CFCs in both this and prior evaluations,32 while FACS analysis for only the IL-3R β_c subunit was predictive of the same endpoint, QRT-PCR was chosen for analysis of IL-3R expression in subsequent experiments on subpopulations of AML cells.

IL-3R subunit expression on subpopulations of AML and normal BM cells enriched for progenitors

Figure 1 shows a typical FACS profile of AML cells stained with antibodies against CD34, CD38, and CD71. Previous data had demonstrated that the CD34⁺CD38⁻ phenotype includes N/SL-ICs from most patient samples while addition of CD71 negativity to this phenotype increases the enrichment of sorted cells for these primitive progenitors.¹⁵ These studies have also demonstrated that AML-CFCs are largely CD34⁺ in most patient samples but are





IL-3 RECEPTORS ON AML STEM CELLS 3533

Table 2. Correlation between IL-3R subunit quantitation on AML blasts by FACS (MFI) and QRT-PCR and AML-CFC killing by $DT_{380}IL3$

	FACS-MFI		AML-CEC	
	CD123	CD131	kill	
QRT-PCR				
IL-3Rα	0.095	—	0.59†	
IL-3Rβ _c	_	0.40	0.70†	
AML-CFC* kill	0.23	0.74†	—	

Data expressed as correlation coefficients between the values indicated as determined for each of 12 of the 19 AML samples listed on Table 1 and according to "Patients, materials, and methods."

- indicates not done.

*Percentage of AML-CFCs killed after 24 hours of exposure to 50 ng/mL $\rm DT_{388}IL3.$

†P < .05, Student *t* test.

more heterogeneous in their expression of CD38^{8,15} (ie, most of these more mature progenitors express this antigen and thus are found in the CD34⁺CD38⁺ subpopulation, while N/SL-ICs are excluded). CD34⁻ cells from most AML samples have little progenitor activity. Thus, PB cells from the 19 patients with AML (Table 1) were sorted into 4 subpopulations defined as CD34⁺CD38⁻CD71⁻, CD34⁺CD38⁻, CD34⁺CD38⁺, and CD34⁻. As shown in Table 3, the large proportion of AML cells from 16 of these patients were CD34⁺. Overall, mean frequencies of the CD34⁺CD38⁻CD71⁻, CD34⁺CD38⁻, and CD34⁺CD38⁺ subpopulations were 3.2% (range, 0.05%-25.3%), 5.3% (range, 0.08%-38.4%), and 40.2% (range, 0.7%-90.9%), respectively, while 27% (range, 2.1%-80.9%) of cells were CD34⁻.

QRT-PCR was used to detect expression of the IL-3R α and β_c subunits in RNA isolated from the 4 sorted subpopulations of AML cells from these 19 patient samples. As shown in Figure 3, the median levels of expression of the α subunit were higher than that of the β_c subunit for each isolated subpopulation (range, 4- to 15-fold). The level of β_c expression detected among the subpopulations of cells enriched for primitive progenitors (CD34⁺CD38⁻

Table 3. Proportion of AML cells in FACS-isolated subpopulations

Patient no.	CD34+CD38-CD71-, %*	CD34+CD38-, %†	CD34+CD38+, %‡	CD34- %§						
1	1.5	1.9	44.5	12.3						
2	17.4	35.3	10.2	14.8						
3	0.9	2.4	56	11.8						
4	0.3	0.6	40	29.6						
5	2.9	3.8	64.9	8.6						
6	0.3	0.4	62.4	10.8						
7	0.4	0.7	43.1	26.3						
8	0.08	0.1	17.3	69.3						
9	2.6	4.1	71	5.8						
10	2.7	2.9	43.4	21.4						
11	0.05	0.08	0.7	67.1						
12	0.1	0.1	8.7	43						
13	1.6	1.8	62.6	6.1						
14	3.2	5.3	45.2	26.1						
15	25.3	38.4	24.3	19.5						
16	0.4	0.6	52.5	33.2						
17	0.6	0.6	90.9	2.1						
18	0.2	0.3	23.5	24.3						
19	0.7	0.8	2.1	80.9						

*Mean \pm SD = 3.2 \pm 6.6.

 \pm Mean \pm SD = 5.3 \pm 11.2.

‡Mean ± SD = 40.2 ± 25.3.

 $\Delta = 27 \pm 22.9.$





Figure 3. IL-3R subunit expression on AML cells as detected by QRT-PCR. RNA was extracted from AML cells sorted into the subpopulations indicated as described in "Patients, materials, and methods." Values shown are the levels of expression of β_c (A) and α (B) subunits of the IL-3R relative to the expression of GAPDH arbitrarily set at 1000 for each of 19 patient samples. Each data point represents the value for an individual patient sample. The horizontal bars and adjacent numbers indicate the mean level of expression for each of the α and β_c subunits in the different subpopulations.

cells or such cells lacking CD71) was not significantly different from that detected in the subpopulations depleted of such cells (CD34⁺CD38⁺ or CD34⁻ cells). However, expression of the IL-3R α subunit was significantly higher among CD34⁺CD38⁻CD71⁻ and CD34⁺CD38⁻ cells than among CD34⁺CD38⁺ or CD34⁻ cells (P < .002, paired *t* test).

RNA also was isolated from CD34⁺CD38⁻ and CD34⁺CD38⁺ cells from 3 normal BM samples. QRT-PCR demonstrated that the mean (\pm SD) level of expression of the IL-3R α and β_c subunits (relative to GAPDH set at 1000) was 21.6 (\pm 8.6) and 9.0 (\pm 5.1), respectively, for CD34⁺CD38⁻ cells, and 8.2 (\pm 1.8) and 26.2 (\pm 3.8), respectively, for CD34⁺CD38⁺ cells. These values for the α subunit were significantly ($P \leq .002$) lower than those obtained for the same subpopulations of AML cells. In contrast, the β_c subunit was expressed at similar (in CD34⁺CD38⁻ cells) or higher (P = .003 in CD34⁺CD38⁺ cells) levels among normal than among malignant cells.

In our previous studies we had defined an expression of the IL-3R β_c subunit relative to GAPDH of 3 or less as being associated with poor killing of AML-CFCs.³² To assess the value of this measurement in predicting cytotoxicity of DT₃₈₈IL3 against N/SL-ICs from different patient samples, AML blasts from 7 samples, which had levels of IL-3R β_c expression above or below that level in the sorted subpopulations were selected. Three of these AML samples had low levels of IL-3R β_c expression levels for samples 1, 16, and 13 were 1.8, 0.14, and 0.3, respectively), while the remaining 4 patient samples had higher levels of expression

(expression for samples 8, 17, 10, and 11 was 8, 5, 11.8, and 13.5, respectively). Comparison of this level of expression with that seen in the other analyzed subpopulations and unsorted AML blasts from the same sample showed no consistent differences (Figure 4). Among these 7 samples expression of the IL-3R α subunit was higher among CD34⁺CD38⁻ cells compared with cells expressing CD38 or lacking CD34 as described for the entire patient group (Figure 3). However, there were no significant differences in the expression of IL-3R α between the 3 samples with low expression and the 4 samples with high expression of IL-3R β_c in any subpopulation studied.

AML blasts from these same samples were incubated for 24 hours with or without DT₃₈₈IL3 at 50 or 250 ng/mL and then injected into cohorts of sublethally irradiated $\beta_2 m^{-/-}$ NOD/SCID mice. For the 3 samples with low levels of IL-3R β_c expression, exposure to either DT₃₈₈IL3 concentration did not change the level of engraftment detected in mouse bone marrow 12 weeks later indicating that N/SL-ICs from these samples had survived this treatment (Figure 5A). Fluorescence in situ hybridization (FISH) analysis of BM cells harvested from mice injected with $DT_{388}IL3$ -treated or -untreated AML cells from patients 1 and 13 demonstrated the presence of the +8 and inv(16) chromosome abnormalities, respectively, in a similar proportion of cells to that found in the patients' diagnostic BM sample (data not shown). In contrast, with 3 samples that expressed high levels of IL-3R β_c complete eradication of N/SL-ICs from samples 8 and 17 and near-complete eradication from sample 10 was achieved with 250 ng/mL of DT₃₈₈IL3. Although sample 11 expressed the IL-3R β_c subunit at levels similar to these 3 samples, only a modest (but statistically significant [P < .05]) reduction in engraftment of AML cells was observed (Figure 5B). The percentage of reduction in AML cell engraftment achieved with prior DT₃₈₈IL3 treatment of these 7 patient samples correlated most strongly with the level of IL-3R β_c expression in CD34⁺CD38⁻CD71⁻ cells (r = 0.76) compared with the other FACS-sorted subpopulations, but a correlation was also seen with the expression of the same molecule on unsorted blast cells (r = 0.56).

Discussion

DT₃₈₈IL3 is a novel fusion protein which is currently undergoing clinical evaluation as a possible new therapeutic agent for the treatment of AML. Although data from tissue culture and mouse



Figure 4. Comparison of the relative level of expression of the IL-3R β_c subunit between different subpopulations of AML cells from 7 patient samples tested for DT₃₈₈IL3 sensitivity in β_2 -microglobin–deficient NOD/SCID mice. QRT-PCR was performed on FACS-sorted cells of the indicated cell-surface phenotype as described for Figure 3.



Figure 5. IL-3R β_c subunit expression detected by QRT-PCR predicts cytotoxicity of DT₃₈₈IL3 against AML progenitors which engraft in β_2 -microglobin-deficient NOD-SCID mice. (A) AML cells from 3 patients (nos. 1, 16, and 13 from Table 1) expressing low levels of IL-3R β_c and (B) 4 patients (nos. 8, 17, 10, and 11 from Table 1) who had high levels of IL-3R β_c expression were incubated for 24 hours with or without DT₃₈₈IL-3 at 50 and 250 ng/mL and then injected intravenously into sublethally irradiated mice as described in "Patients, materials, and methods." Data are shown as the mean level of engraftment of human AML cells in cohorts of mice receiving treated or untreated cells 8 to 12 weeks previously as detected by FACS analysis of mouse BM cells for CD45⁺ human cells. Indicates untreated cells; S, cells treated with 50 ng/mL DT₃₈₈IL-3. and \Box , cells treated with 250 ng/mL DT₃₈₈IL-3. Error bars indicate SEM level engraftment for all mice in the cohort. The level of engraftment was significantly (P < .05; Student *t* test) less than that obtained for the untreated cells for the 4 patient samples treated with both doses of DT₃₃₈IL3 shown in panel B.

models had suggested that this drug would be active against many human AMLs, it also suggested that some leukemias would be much more sensitive than others.^{18,19,31} This was particularly true when DT₃₈₈IL3 was tested against AML progenitors that engraft in NOD/SCID mice (N/SL-ICs).18 Thus, for some samples DT₃₈₈IL3 was highly cytotoxic when tested against AML-CFCs but much less so against N/SL-ICs, while for other samples the drug had little effect against any class of AML progenitors. Evaluation of the expression of high- and low-affinity IL-3-binding sites on AML blasts demonstrated that the presence of IL-3Rs was necessary but not always sufficient to ensure leukemic progenitor kill by the fusion protein.^{19,34} Previous studies had also demonstrated that AML blast populations highly enriched for or depleted of progenitor activity could be obtained by isolating cells that varied in their expression of CD34, CD38, and CD71.8,15 N/SL-ICs are largely restricted to the CD34⁺CD38⁻ subpopulation of AML blasts from most patient samples and can be further enriched by excluding CD71⁺ cells, while CD34⁻ cells are typically depleted of all progenitor activity.8,15,16

Using FACS analysis other investigators have demonstrated that the IL-3R α subunit is typically highly expressed on AML blasts, including the CD34⁺CD38⁻ subpopulation.^{14,20} More recently this finding has been confirmed and extended by using MFI to quantify expression of both the IL-3R subunits on AML blasts labeled with anti-CD123 and anti-CD131.⁴² This measurement was also shown to correlate with the percentage of AML cells undergoing apoptosis when exposed to DT₃₈₈IL3. As an alternative to flow cytometry we have used QRT-PCR to show that the level of IL-3R subunit expression in AML blasts predicted the killing of AML-CFCs by DT₃₈₈IL3 and its variants.³² However, with both techniques some patient samples showed relative resistance to DT₃₈₈IL3 cytotoxicity in spite of high-level IL-3R expression.

In initial experiments described here, flow cytometry was compared with QRT-PCR for its ability to detect and quantify the IL-3R subunits. Both techniques provided quantitative and reproducible data. However, there was no correlation between the FACS data (either MFI or percentage of positive cells) and QRT-PCR for the IL-3R α subunit. While a correlation was seen between MFI detected by FACS and ORT-PCR for the IL-3RB_c subunit, it was relatively weak and did not reach statistical significance (Table 2). These data indicate that RNA and protein levels for these molecules do not necessarily correspond in AML blasts. Our data differ from the recent report from Testa et al, who found that MFI detected by anti-CD123 antibodies predicted AML blast apoptosis induced by DT₃₈₈IL3.⁴² This difference might be explained by the different endpoint in our studies (AML-CFC killing), the use of different antibodies, and the larger number of samples tested in the earlier investigation. Nevertheless, in the current study we were able to demonstrate a correlation between expression of both IL-3R subunits as detected by QRT-PCR and AML-CFC killing with DT₃₈₈IL3. These data confirm those in our previous report and are also consistent with our previous conclusion that the relationship is strongest for the IL-3Rβ_c subunit.³²

Since LSCs are the most relevant target population for novel antileukemic therapy such as DT₃₈₈IL3 we used QRT-PCR to study the expression of IL-3R subunits on subpoulations of AML cells known to be enriched for or depleted of leukemic progenitors that engraft in immunodeficient ($\beta_2 m^{-/-}$ NOD/SCID) mice. Among 19 AML patient samples no consistent difference in expression of the IL-3R β_c subunit was detected between the most primitive CD34⁺CD38⁻CD71⁻ cell fraction and any other subpopulation (Figure 3A). On the other hand, the IL-3Ra subunit was most highly expressed among cells enriched for candidate LSCs which engraft in mice (Figure 3B). This and the generally much higher level of expression of the IL-3R α subunit compared with the β_c subunit across all cell populations studied suggest that expression of the latter molecule is more likely than the former to be a limiting factor in the formation of high-affinity IL-3R-binding sites on N/SL-ICs.

Although N/SL-ICs are a small proportion of the total cells present in even the most enriched population studied here,¹⁵ and the number of AML samples studied in $\beta_2 m^{-/-}$ NOD/SCID mice was limited to 7, the correlation between IL-3R β_c expression in CD34⁺CD38⁻CD71⁻ cells and DT₃₈₈IL3 cytotoxicity against cells engrafting in mice was strong. In fact, a correlation between expression of this molecule in unsorted blast cells and reduced engraftment of DT₃₈₈IL3-treated AML cells was also seen. This and the overall expression data shown on Figure 3 suggest that there is little difference between the expression of the IL-3R β_c among AML blasts and progenitors. On the other hand, while expression of the IL-3R α subunit may vary considerably among different AML-cell fractions, it does not appear to limit the cytotoxicity of DT₃₈₈IL3 against candidate LSCs.

In initial studies investigating the relative cytotoxicity of $DT_{388}IL3$ against normal and leukemic progenitor cells, a striking lack of toxicity was seen against primitive normal multipotential progenitors, including those with lymphomyeloid potential, that are detected in NOD/SCID mice.¹⁸ This was somewhat surprising given the fact that expression of both IL-3R subunits has been demonstrated on CD34⁺CD38⁻ cells from normal BM as well as cord blood and fetal liver using RT-PCR.⁴³ We have confirmed the findings of this earlier study. However, using the QRT-PCR technique described in the current report the relative level of expression of the IL-3R α subunit among both CD34⁺CD38⁻ and CD34⁺CD38⁺ normal BM cells was found to be generally lower than that detected in the same population of AML cells. This

difference may partly explain the low toxicity observed with $DT_{388}IL3$ against primitive normal hematopoietic cells. Other investigators have also shown that expression of the IL-3R α subunit is generally higher on primitive AML cells than on comparable normal cell populations.^{14,20} Interestingly, the expression of the β_c subunit was similar among normal and malignant cells. Thus, it remains possible that the clinical experience with $DT_{388}IL3$ will uncover more myelosuppressive activity for this fusion protein than has so far been suggested by the preclinical data. Alternatively, events that take place after $DT_{388}IL3$ binding to the receptor may differ between normal and malignant cells. It seems feasible that enhanced association of $DT_{388}IL3$ with the IL-3R α subunits on AML cells (which must also express sufficient β_c subunits to form the high-affinity binding complex) might

accelerate the formation and internalization of the α/β_c receptor complex,⁴⁴ resulting in more efficient delivery of DT molecules to the cytoplasm of AML progenitors.

A phase 1/2 clinical trial of DT₃₈₈IL3 is currently under way in patients with relapsed or refractory AML.⁴⁵ Data presented here suggest that it may be possible to use quantification of target molecules on AML blasts to predict which patients are likely to benefit from therapy with this fusion protein. The use of QRT-PCR for the IL-3R subunits is currently being tested in this context. Furthermore, these results suggest that detailed evaluation of subpopulations of malignant cells with different functional properties may reveal similarities and differences in gene expression that are important for the rational development of targeted antileukemic therapy.

References

- Fialkow PJ, Singer JW, Raskind WH, et al. Clonal development, stem-cell differentiation, and clinical remissions in acute nonlymphocytic leukemia. N Engl J Med. 1987;317:468-473.
- 2. McCulloch EA. Stem cells in normal and leukemic hemopoiesis. Blood. 1983;62:1-13.
- Griffin JD, Lowenberg B. Clonogenic cells in acute myeloblastic leukemia. Blood. 1986;68: 1185-1195.
- Sutherland HJ, Blair A, Zapf RW. Characterization of a hierarchy in human acute myeloid leukemia progenitor cells. Blood. 1996;87:4754-4761.
- Ailles LE, Gerhard B, Hogge DE. Detection and characterization of primitive malignant and normal progenitors in patients with acute myelogenous leukemia using long-term coculture with supportive feeder layers and cytokines. Blood. 1997;90:2555-2564.
- Hogge DE, Lansdorp PM, Reid D, Gerhard B, Eaves CJ. Enhanced detection, maintenance and differentiation of primitive human hematopoietic cells in cultures containing murine fibroblasts engineered to produce human Steel factor, interleukin-3 and granulocyte colony-stimulating factor. Blood. 1996;88:3765-3773.
- Ailles LE, Gerhard B, Kawagoe H, Hogge DE. Growth characteristics of acute myelogenous leukemia progenitors that initiate malignant hematopoiesis in nonobese diabetic/severe combined immunodeficient mice. Blood. 1999;34:1761-1772.
- Kawagoe H, Humphries RK, Blair A, Sutherland HJ, Hogge DE. Expression of Hox genes, HOX cofactors, and MLL in phenotypically and functionally defined subpopulations of leukemic and normal human hematopoietic cells. Leukemia. 1999;13:687-698.
- Sauvageau G, Lansdorp PM, Eaves CJ, et al. Differential expression of homeobox genes in functionally distinct CD34⁺ subpopulations of human bone marrow cells. Proc Natl Acad Sci U S A. 1994;91:12223-12227.
- Bhatia M, Wang JCY, Kapp U, Bonnet D, Dick JE. Purification of primitive human hematopoietic cells capable of repopulating immune-deficient mice. Proc Natl Acad Sci U S A. 1997;94:5320-5325.
- Conneally E, Cashman J, Petzer A, Eaves C. Expansion in vitro of transplantable human cord blood stem cells demonstrated using a quantitative assay of their lympho-myeloid repopulating activity in nonobese diabetic-*scid/scid* mice. Proc Natl Acad Sci U S A. 1997;94:9836-9841.
- Blair A, Hogge DE, Ailles LE, Lansdorp PM, Sutherland HJ. Lack of expression of Thy-1 (CD90) on acute myeloid leukaemia cells with long-term proliferative ability in vitro and in vivo. Blood. 1997; 89:3104-3112.
- 13. Blair A, Sutherland HJ. Primitive acute myeloid

leukemia cells with long-term proliferative ability in vitro and in vivo lack surface expression of ckit. Exp Hematol. 2000;28:660-671.

- Jordan CT, Upchurch D, Szilvassy SJ, et al. The interleukin-3 receptor alpha chain is a unique marker for human acute myelogenous leukemia stem cells. Leukemia. 2000;14:1777-1784.
- Blair A, Hogge DE, Sutherland HJ. Most acute myeloid leukaemia progenitor cells with long-term proliferative ability in vitro and in vivo have the phenotype CD34⁺/CD71⁻/HLA-DR⁻. Blood. 1998;92:4325-4335.
- Bonnet D, Dick JE. Human acute myeloid leukemia is organized as a hierarchy that originates from a primitive hematopoietic cell. Nat Med. 1997;3:730-736.
- Chang F, Steelman LS, Lee JT, et al. Signal transduction mediated by the Ras/Raf/MEK/ERK pathway from cytokine receptors to transcription factors: potential targeting for therapeutic intervention. Leukemia. 2003;17:1263-1293.
- Feuring-Buske M, Frankel AE, Alexander RL, Gerhard B, Hogge DE. A diphtheria toxin-interleukin 3 fusion protein is cytotoxic to primitive acute myeloid leukemia progenitors but spares normal progenitors. Cancer Res. 2002;62:1730-1736.
- Alexander RL, Ramage J, Kucera GL, Caligiuri MA, Frankel AE. High affinity interleukin-3 receptor expression on blasts from patients with acute myelogenous leukemia correlates with cytotoxicity of a diphtheria toxin/IL-3 fusion protein. Leuk Res. 2001;25:875-881.
- Testa U, Riccioni R, Militi S, et al. Elevated expression of IL-3Ra in acute myelogenous leukemia is associated with enhanced blast proliferation, increased cellularity, and poor prognosis. Blood. 2002;100:2980-2988.
- Fitzgerald D, Pastan I. Targeted toxin therapy for the treatment of cancer. J Natl Cancer Inst. 1989; 81:1455-1463.
- Frankel AE, Ramage J, Kiser M, Alexander R, Kucera G, Miller MS. Characterization of diphtheria fusion proteins targeted to the human interleukin-3 receptor. Protein Eng. 2000;13:575-581.
- Murphy JR, Bishai W, Borowski M, Miyanohara A, Boyd J, Nagle S. Genetic construction, expression, and melanoma-selective cytotoxicity of a diphtheria-toxin-related alpha-melanocyte-stimulating hormone fusion protein. Proc Natl Acad Sci U S A. 1986;83:8258-8262.
- Williams DP, Parker K, Bacha P, et al. Diphtheria toxin receptor binding domain substitution with interleukin-2: genetic construction and properties of a diphtheria toxin-related interleukin-2 fusion protein. Protein Eng. 1991;1:493-498.
- Chadwick DE, Williams DP, Niho Y, Murphy JR, Minden MD. Cytotoxicity of a recombinant diphtheria toxin-granulocyte colony-stimulating factor fusion protein on human leukemic blast cells. Leuk Lymphoma. 1993;11:249-262.

- Feuring-Buske M, Frankel A, Gerhard B, Hogge D. Variable cytotoxicity of diphtheria toxin 388granulocyte-macrophage colony-stimulating factor fusion protein for acute myelogenous leukemia stem cells. Exp Hematol. 2000;28:1390-1400.
- Sandvig K, Olsnes S. Rapid entry of nicked diphtheria toxin into cell at low pH: characterization of the entry process and effects of low pH on the toxin molecule. J Biol Chem. 1981;256:9068-9076.
- Draper RK, Simon MI. The entry of diphtheria toxin into the mammalian cell cytoplasm: evidence for lysosomal involvement. J Biol Chem. 1980;87:849-854.
- Honjo T, Nishizuka Y, Kato I, Hayaishi O. Adenosine diphosphate ribosylation of aminoacyl transferase II and inhibition of protein synthesis by diphtheria toxin. J Biol Chem. 1971;246:4251-4260.
- Yamaizumi M, Mekada E, Uchida T, Okada Y. One molecule of diphtheria toxin fragment A introduced into a cell can kill the cell. Cell. 1978;15: 245-250.
- Frankel AE, McCubrey JA, Miller MS, et al. Diphtheria toxin fused to human interleukin-3 is toxic to blasts from patients with myeloid leukemias. Leukemia. 2000;14:576-585.
- Hogge DE, Yalcintepe L, Wong S-H, Gerhard B, Frankel AE. Variant diphtheria toxin-interleukin-3 fusion proteins with increased receptor affinity have enhanced cytotoxicity against acute myeloid leukemia progenitors. Clin Cancer Res. 2006;12: 1284-1291.
- Bennett JM, Catovsky D, Daniel MT, et al. Proposed revised criteria for the classification of acute myeloid leukemia: a report of the French-American-British Cooperative Group. Ann Intern Med. 1985;103:620-625.
- Alexander RL, Kucera GL, Klein B, Frankel AE. In vitro interleukin-3 binding to leukemia cells predicts cytotoxicity of a diphtheria toxin/IL-3 fusion protein. Bioconjug Chem. 2000;11:564-568.
- Pfaffl MW. A new mathematical model for relative quantification in real-time RT-PCR. Nucleic Acids Res. 2001;29:e45.
- Bustin SA. Absolute quantification of mRNA using real-time reverse transcription polymerase chain reaction assays. J Mol Endocrinol. 2000;25:169-193.
- 37. Christianson SW, Greiner DL, Hesselton R, et al. Enhanced human CD4⁺ T cell engraftment in β_2 microglobulin-deficient NOD-*scid* mice. J Immunol. 1997;158:3578-3586.
- Feuring-Buske M, Gerhard B, Cashman J, Humphries RK, Eaves CJ, Hogge DE. Improved engraftment of human acute myeloid leukemia progenitor cells in beta 2-microglobulin-deficient

NOD/SCID mice and in NOD/SCID mice transgenic for human growth factors. Leukemia. 2003; 17:760-763.

- Verlinden SF, van Es HH, van Bekkum DW. Serial bone marrow sampling for long-term follow up of human hematopoiesis in NOD/SCID mice. Exp Hematol. 1998;26:627-630.
- Grimwade D, Walker H, Oliver F, et al. The importance of diagnostic cytogenetics on outcome in AML: analysis of 1,612 patients entered into the MRC AML 10 trial. Blood. 1998;92:2322-2333.
- 41. Slovak ML, Kopecky KJ, Cassileth PA, et al.

Karyotypic analysis predicts outcome of pre- and post-remission therapy in adult acute myeloid leukemia (AML): A SWOG/ECOG intergroup study [abstract]. Blood. 1998;92:678a.

- Testa U, Riccioni R, Biffoni M, et al. Diphtheria toxin fused to variant human interleukin-3 induces cytotoxicity of blasts from patients with acute myeloid leukemia according to the level of interleukin-3 receptor expression. Blood. 2005;106:2527-2529.
- Oh I-H, Lau A, Eaves CJ. During ontogeny primitive (CD34⁺CD38⁻) hematopoietic cells show altered expression of a subset of genes associ-

ated with early cytokine and differentiation responses of their adult counterparts. Blood. 2000; 96:4160-4168.

- Sato N, Caux C, Kitamura T, et al. Expression and factor-dependent modulation of the interleukin-3 receptor subunits on human hematopoietic cells. Blood. 1993;82:752-761.
- Frankel AE, Weir MA, Hall PD, Hogge DE, Rizzieri DA. Diphtheria toxin-interleukin 3 fusion protein therapy of patients with elderly or relapsed/ refractory acute myeloid leukemia (AML) [abstract]. J Clin Oncol [suppl Jun 20]. 2006;24:6569.