

Brief report

Frequent antibody production against RAR α in both APL mice and patients

Marie Robin, Juliette Andreu-Gallien, Marie-Helene Schlageter, Djaouida Bengoufa, Isabelle Guillemot, Katerina Pokorna, Carine Robert, Jerome Larghero, Philippe Rousselot, Emmanuel Raffoux, Herve Dombret, Pierre Fenaux, Marika Pla, Dominique Charron, Rose-Ann Padua, and Christine Chomienne

In an acute promyelocytic leukemia (APL)–transplantable mouse model, we previously reported the presence of antibodies recognizing PML-RAR α and RAR α in the sera of ATRA-treated mice. To evaluate this immune response, we determined the prevalence of anti-RAR α antibodies in a cohort of 48 APL mice, treated by ATRA (n = 24) or by placebo pellets (n = 24), and in a preliminary subset of 9 patients with APL using a specific enzyme-linked

immunosorbent assay (ELISA). In APL mice, significantly higher antibody levels were observed at the latest time points (day 48 to 58 levels superior to day 15 to 18 or day 28 to 38 levels). Antibody levels were higher in ATRA-treated mice than in placebo-treated mice and were also predictive of better survival. In the patients with APL, anti-RAR α antibodies were detected at diagnosis and after maintenance therapy, reminiscent of the ATRA-

treated APL mice. Antinuclear or antineutrophil cytoplasmic autoantibodies were also detected. These data reveal for the first time that in patients with APL an immune response may be detected at diagnosis and enhanced after maintenance therapy. (Blood. 2006;108:1972-1974)

© 2006 by The American Society of Hematology

Introduction

Acute promyelocytic leukemia (APL) is characterized by a differentiation block at the promyelocytic stage and by a reciprocal chromosomal translocation, t(15;17)(q24;q21), fusing the promyelocytic leukemia gene (*PML*) with the retinoic acid receptor alpha (*RARA*) gene.¹ Treatment with all-trans retinoic acid (ATRA) and chemotherapy induces complete remissions in 90% of PML-RAR α -positive patients with APL through ATRA-induced differentiation of the leukemic cells and their subsequent elimination by apoptosis.² Apart from their known effects on cell differentiation, retinoids also play an important role in infection and immune functions.³ In hypovitaminosis A animal models and in vitamin A-deficient children, alterations in antiviral, antibacterial, or antiparasite responses have been shown.³⁻⁵ In an APL-transplantable mouse model that shares the promyelocytic features and response to ATRA of human APL,⁶ we have previously reported the presence of antibodies in ATRA-treated mouse sera.⁷ By Western blot analysis, we showed that these antibodies recognized mouse and human RAR α from spleen and HL-60 cells, respectively, as well as PML-RAR α from mouse spleen and NB4 leukemic cells. We hypothesized that such adjuvant effects of ATRA might also participate in the efficacy of ATRA in patients with APL, especially during maintenance therapy when the tumor burden is low and immunocompetency restored.²

The aim of our study was therefore to further characterize anti-RAR α antibody production in APL mice as well as in patients with APL. In patients with APL, antibodies to RAR α and autoanti-

bodies like antinuclear autoantibodies (ANAs) or antineutrophil cytoplasmic autoantibodies (ANCA) were analyzed.

Study design

The transplantable APL mouse model of mice undergoing APL transplantation and ATRA therapy was set up as already described.^{6,7} Data from 3 protocols that did not differ in their response to ATRA were pooled. Placebo (n = 24) and ATRA-treated APL (n = 24) mice died from day 18 to 29 (median, 26 days) and from day 28 to 95 (median, 55 days), respectively, as already reported.^{6,7} Antibodies against RAR α were detected by an enzyme-linked immunosorbent assay (ELISA) at different time intervals as previously described.⁷ For each mouse serum, specific absorbance (SA) was calculated as the difference between duplicates of mean absorbance with and without GST-RAR α . To normalize results between experiments, arbitrary units (AUs) were obtained by dividing the SA obtained in serum dilution by the SA obtained with 1:200 000 dilution of positive control monoclonal anti-RAR α antibody (9 α F).

An ELISA was performed in sera from 9 patients with APL to detect the presence of anti-RAR α antibodies. ANAs and ANCA were measured by indirect immunofluorescence. We analyzed sera from 9 patients for whom stored sera were available at diagnosis and after 2-year maintenance therapy.² Patients were enrolled in APL trials: patients 3, 5, and 8 from APL93 and patients 1, 2, 4, 6, 7, and 9 from the APL2000 trial. In these trials informed consent was obtained to use stored samples. For all patients sera were stored in similar conditions at -30°C and thawed once. Eighteen healthy subjects served as controls.

From the Institut National de la Santé et de la Recherche Médicale (INSERM), Unité mixte de recherche (UMR) S 718, Paris; Université Paris 7, Institut Universitaire d'Hématologie (IUH), UMR S 718, Paris; Laboratoire d'Immunologie et d'Histocompatibilité, Assistance Publique-Hopitaux de Paris (AP-HP), Hôpital Saint-Louis, Paris; Service d'Hématologie, AP-HP, Hôpital Saint-Louis, Paris; Service d'Hématologie, AP-HP, Hôpital Avicenne, Bobigny; INSERM, UMR S 662, Paris; Université Paris 7, IUH, UMR S 662, Paris, France.

Submitted March 28, 2006; accepted May 2, 2006. Prepublished online as *Blood* First Edition Paper, May 25, 2006; DOI 10.1182/blood-2006-03-013177.

Supported by the French Fondation pour la Recherche Médicale, the Fondation de France, the French Association de Recherche contre le Cancer,

the French Ligue Nationale contre le Cancer, the Fondation Saint-Louis, Eli Lilly, and INSERM.

M.R. and J.A.-G. contributed equally to this work.

Reprints: Christine Chomienne, INSERM UMR S 718, Institut Universitaire d'Hématologie (IUH), Hôpital Saint-Louis, 1, Avenue Claude Vellefaux, Paris Cedex 10, F-75475 France; e-mail: lbch.secretariat@sls.aphp.fr.

The publication costs of this article were defrayed in part by page charge payment. Therefore, and solely to indicate this fact, this article is hereby marked "advertisement" in accordance with 18 U.S.C. section 1734.

© 2006 by The American Society of Hematology




Figure 1. Anti-RAR α antibody production in APL mice and patients. (A) Comparative anti-RAR α ELISA dose-response curves with a positive mouse serum and an anti-RAR α monoclonal antibody control. SA measured by ELISA of a serum from an ATRA-treated mouse serially diluted from 1:50 to 1:10 000 (Δ) and from the anti-RAR α monoclonal antibody 9 α F diluted from 1:10 000 to 1:200 000 (reference curve, \square). (B) Time-dependent presence of anti-RAR α antibodies in mice with APL. Anti-RAR α antibodies were measured by ELISA in 48 mice with APL. On days 15 to 18, all mice were alive, and sera were obtained from only 46 mice, including 23 mice treated by ATRA and 23 treated by placebo. On days 28 to 38, 30 mice were alive, including 24 mice treated by ATRA and 6 treated by placebo. On days 48 to 58, only 13 mice, treated by ATRA, were alive. The box plot in combination with dot plot displays a statistical summary of the data (quartiles and median). Data are expressed as arbitrary units. (C) Time course of the IgG and IgM SA in the anti-RAR α ELISA of 1 ATRA-treated APL mouse. For IgG and IgM detection, secondary antibodies specific for mouse IgG and IgM were used in ELISA tests. In 1 ATRA-treated mouse the anti-RAR α ELISA was performed in serum collected on days 18, 28, and 38 using specific secondary antibodies for mouse IgG (\bullet) and mouse IgM (\blacksquare). Results are expressed as SA. (D) Time-dependent presence of anti-RAR α antibodies in APL patient sera. Anti-RAR α antibodies (expressed as SA) were measured by ELISA in the serum of 9 patients with APL at diagnosis and after maintenance therapy. ELISA tests were performed in 96-well plates (Nunc; Merck Eurolab, VWR, Fontenay-Sous-Bois, France) coated with either GST or GST-RAR α (obtained by fusion of a GST tag to the full-length RAR α). After overnight saturation with PBS1X-BSA5%, sera (diluted 1:200 in PBS1X-BSA5%) were incubated 1 hour at room temperature. Peroxidase-conjugated goat anti-human IgG antibody (Sigma-Aldrich, Lyon, France) was added and incubated 1 hour at room temperature, followed by TMB substrate revelation (BD Pharmingen, San Diego, CA). Absorbance was measured at 450 nm (reference filter at 630 nm) using an optical densitometer (Dynatech MR5000; Labsystems, Cergy, France). For each serum, SA was calculated as the difference between duplicates of mean absorbance between GST-RAR α and GST. Anti-RAR α antibodies were also measured in sera from 18 healthy persons (controls). The box plot in combination with dot plot displays a statistical summary of the data (quartiles and median).

Results and discussion

Anti-RAR α antibody production in APL mice

As we have previously reported, anti-RAR α antibodies were detected in ATRA-treated APL mice.⁷ Specificity of the ELISA signal defined previously on immunoblotting experiments with recombinant GST-RAR α ⁷ was further assessed in this study by ELISA competition experiments. Preincubation of an anti-RAR α -positive serum overnight at 4°C with GST-RAR α in a 20-fold excess resulted in an 87% binding inhibition (SA of 0.158 compared with 1.23 with preincubation with buffer alone) (data not shown). The dilution of a positive anti-RAR α mouse serum shows a dose-dependent SA similar to that obtained with a known anti-RAR α monoclonal antibody, 9 α F⁸ (Figure 1A). In this large cohort of 48 APL mice, it is clear that anti-RAR α antibody levels progressively increase with time, from a median AU level of 0.26 on days 15 to 18 (D“18”) to median AU levels of 0.88 on days 28 to 38 (D“38”) and 6.33 on days 48 to 58 (D“58”) (*t* test, *P* < .001 between D“18” and D“58”). On D“38,” among 6 surviving placebo-treated mice, only 1 mouse had antibody levels reaching a threshold of 4-fold that of the D“18” median, whereas in ATRA-treated mice 9 of 24 mice reached this level on D“38” and 12 of 13 on D“58,” when all placebo-treated mice were dead (Figure 1B). AUs reached a maximal value of 16.8 on D“58” in ATRA-treated mice. This antibody appearance followed the classic scheme of antibody production, with a time-decreased production of IgM and a time-increased production of IgG from day 18 to day 58 (Figure 1C). To investigate the prognostic value of these antibodies independently of time, we asked whether antibody levels obtained at D“18,” when all mice were alive, were predictive of survival. At this time point, a higher level of antibody was associated with increased survival: Median survival was 28 days in mice with a level lower than the median D“18” AUs (0.26), versus 49 days for mice with a level higher than the median D“18” AUs (log-rank test, *P* < .001).

Anti-RAR α antibody production in patients with APL

To provide relevance to the preclinical data obtained in APL mice, anti-RAR α antibodies were retrospectively analyzed by ELISA in 9 patients with APL. All patients were in complete remission at the time of postmaintenance treatment study. For the first time, anti-RAR α antibodies were detected both at diagnosis and after maintenance therapy in patients with APL, and the antibody production followed a time-dependent pattern similar to the kinetics of anti-RAR α antibodies

Table 1. Anti-RAR α antibodies, ANAs, and ANCAs in patients with APL

Patient no.	Months after treatment†	Anti-RAR α antibody		ANA		ANCA	
		Diagnosis*	After treatment†	Diagnosis*	After treatment†	Diagnosis*	After treatment†
1	6	+	+	-	++	-	-
2	6	+++	+++	++	++	-	-
3	36	++	++++	-	+	-	-
4	4.7	++++	+++	++	+	-	-
5	67	++++	++++	++	++	-	-
6	6	++	++++	-	-	+	++
7	4	+	+++	+	++	-	-
8	31	+	+++	-	+	++	+
9	10	+	+++	++	++	+	+

Results of anti-RAR α antibodies are expressed as semiquantitative values according to specific absorbance (SA) value: SA less than 0.2 (+), SA 0.21 to 0.4 (++), SA 0.41 to 0.6 (+++), SA more than 0.61 (++++). Results of ANAs and ANCAs are expressed as negative (-) if no fluorescence was observed and as weakly positive (+) or strongly positive (++) according to the intensity of the signal. Both ANA and ANCA tests were performed according to standard routine procedures of the Laboratoire d’Immunologie by 2 independent investigators. Briefly, ANAs were detected in the same sera (diluted 1:80) by indirect immunofluorescence with commercial Hep2000 cells (ImmunoConcepts; BMD, Marne la Vallée, France).⁹ ANCAs were tested in sera (diluted 1:20) by indirect immunofluorescence with commercial human neutrophils (Inova Diagnostics, San Diego, CA).¹⁰

*Day 0 of therapy for all patients.

†After maintenance therapy (delay in months is given for each patient).

observed in APL mice (Figure 1D and Table 1). In 5 of 9 patients anti-RAR α antibody levels increased with time. Of note, among the 4 patients in whom no increase in anti-RAR α antibody levels was seen, 3 already had high levels of antibodies at diagnosis. Interestingly, other autoimmunity manifestations were observed in these patients. At diagnosis, 7 (78%) were either positive for ANAs or ANCAs, and all patients were either ANA or ANCA positive after maintenance therapy (Table 1). This proportion was higher than that expected in the general population (13.3% and 0% to 1.8% of healthy persons may present ANAs⁹ or ANCAs,¹⁰ respectively). Thus, patients with APL present frequent autoantibodies, especially after maintenance therapy.

Antibodies against multiple tumor-associated antigens like onco-gene proteins (myc,¹¹ ras¹²), gene suppressor proteins (p53¹³), tumor antigens (MUC1,¹⁴ erbB2,¹⁵ WT1¹⁶), or normal proteins (SPAN-Xb¹⁷) have been described in patients with solid tumors or hematologic malignancies.¹⁸⁻²⁰ More recently, autoantibodies have been shown to be markers of better prognosis in melanoma patients treated by interferon- α .²¹ Our results show that in patients with APL, antibodies to RAR α and autoantibodies like ANCA, and especially ANA, are detected. An increase in anti-RAR α antibodies and ANAs was seen after maintenance therapy that includes ATRA and chemotherapy. In mice,²² as in humans,¹⁴ immunotherapy approaches have induced specific antibody production, and in an APL patient vaccinated with an HLA-A2–restricted peptide PR1 derived from proteinase 3, PR1 tetramer–sorted cytotoxic T lymphocytes (CTLs) were obtained after vaccination and showed lysis of the patient's bone marrow cells,²³ suggesting that an antitumor response may be elicited in APL. The data of our study support the notion that an immune response elicited by APL cells is enhanced by ATRA therapy and may well be implicated in its efficacy, because immunocompromised APL mice treated by ATRA have re-

duced survival.²⁴ The immunogenic role played by ATRA in APL is not known and may be related to at least 2 mechanisms. One relates to the now recognized role of vitamin A and its derivatives in the immune response, well seen in vitamin A deficiency³ as in studies on the T helper-2 (Th2) response²⁵ and dendritic cell activation triggered by retinoids.²⁶ The other may relate to the direct effect of ATRA on the leukemic cell itself, where apoptosis and degradation of the oncoprotein PML-RAR could participate in cross-presentation. Ongoing prospective immunomonitoring of patients with APL at diagnosis and throughout treatment will allow us to establish whether the achievement of an efficient immune response is of prognostic value in APL and which criteria may be determinant. If the latter study demonstrates that antibody response correlates with disease-free survival, it will provide, in the setting of minimal residual disease, the basis for vaccine-boosting strategies²⁵⁻²⁷ that enhance a humoral and/or a cellular immune response.

Acknowledgments

We thank Cécile Rochette-Egly, UMR-S 596, Strasbourg, France, for providing monoclonal anti-RAR α antibody; Dr Felix Agbalika and Dr Catherine Scieux, Laboratoire de Virologie, Hôpital Saint-Louis, Paris, for providing some of the stored sera of patients with APL; Scott Kogan and J. Michael Bishop, University of California, San Francisco, for the APL mouse cells; and Bernard Boursin, Service d'Infographie, IUH, Hôpital Saint-Louis, Paris, for excellent graphic work.

Katerina Pokorna was on leave from the Institute of Molecular Genetics, Academy of Sciences of the Czech Republic, Prague, at the time of this article's writing.

References

- de The H, Chomienne C, Lanotte M, Degos L, Dejean A. The t(15;17) translocation of acute promyelocytic leukaemia fuses the retinoic acid receptor alpha gene to a novel transcribed locus. *Nature*. 1990;347:558-561.
- Fenaux P, Chastang C, Chevret S, et al. A randomized comparison of all transretinoic acid (ATRA) followed by chemotherapy and ATRA plus chemotherapy and the role of maintenance therapy in newly diagnosed acute promyelocytic leukemia. The European APL Group. *Blood*. 1999;94:1192-1200.
- Stephensen CB. Vitamin A, infection, and immune function. *Annu Rev Nutr*. 2001;21:167-192.
- Cantorna MT, Nashold FE, Hayes CE. In vitamin A deficiency multiple mechanisms establish a regulatory T helper cell imbalance with excess Th1 and insufficient Th2 function. *J Immunol*. 1994;152:1515-1522.
- Semba RD, Muhilal, Scott AL, et al. Depressed immune response to tetanus in children with vitamin A deficiency. *J Nutr*. 1992;122:101-107.
- Brown D, Kogan S, Lagasse E, et al. A PML-RAR α transgene initiates murine acute promyelocytic leukemia. *Proc Natl Acad Sci U S A*. 1997;94:2551-2556.
- Padua RA, Larghero J, Robin M, et al. PML-RARA-targeted DNA vaccine induces protective immunity in a mouse model of leukemia. *Nat Med*. 2003;9:1413-1417.
- Gaub MP, Rochette-Egly C, Lutz Y, et al. Immunodetection of multiple species of retinoic acid receptor alpha: evidence for phosphorylation. *Exp Cell Res*. 1992;201:335-346.
- Tan EM, Feltkamp TE, Smolen JS, et al. Range of antinuclear antibodies in "healthy" individuals. *Arthritis Rheum*. 1997;40:1601-1511.
- Maillefert JF, Pfitzenmeyer P, Thenet M, et al. Prevalence of ANCA in a hospitalized elderly French population. *Clin Exp Rheumatol*. 1997;15:603-607.
- Zhang JY, Casiano CA, Peng XX, Koziol JA, Chan EKL, Tan EM. Enhancement of antibody detection in cancer using panel of recombinant tumor-associated antigens. *Cancer Epidemiol Biomarkers Prev*. 2003;12:136-143.
- Takahashi M, Chen W, Byrd DR, et al. Antibody to ras proteins in patients with colon cancer. *Clin Cancer Res*. 1995;1:1071-1077.
- Soussi T. p53 Antibodies in the sera of patients with various types of cancer: a review. *Cancer Res*. 2000;60:1777-1788.
- Adluri S, Gilewski T, Zhang S, Ramnath V, Ragupathi G, Livingston P. Specificity analysis of sera from breast cancer patients vaccinated with MUC1-KLH plus QS-21. *Br J Cancer*. 1999;79:1806-1812.
- Visco V, Bei R, Moriconi E, Gianni W, Kraus MH, Muraro R. ErbB2 immune response in breast cancer patients with soluble receptor ectodomain. *Am J Pathol*. 2000;156:1417-1424.
- Gaiger A, Reese V, Disis ML, Cheever MA. Immunity to WT1 in the animal model and in patients with acute myeloid leukemia. *Blood*. 2000;96:1480-1489.
- Wang Z, Zhang Y, Liu H, Salati E, Chiriva-Internati M, Lim SH. Gene expression and immunologic consequence of SPAN-Xb in myeloma and other hematologic malignancies. *Blood*. 2003;101:955-960.
- Boon T, van der Bruggen P. Human tumor antigens recognized by T lymphocytes. *J Exp Med*. 1996;183:725-729.
- Moldrem JJ, Lee PP, Wang C, Champlin RE, Davis MM. A PR1-human leukocyte antigen-A2 tetramer can be used to isolate low-frequency cytotoxic T lymphocytes from healthy donors that selectively lyse chronic myelogenous leukemia. *Cancer Res*. 1999;59:2675-2681.
- Robin M, Schlageter MH, Chomienne C, Padua RA. Targeted immunotherapy in acute myeloblastic leukemia: from animals to humans. *Cancer Immunol Immunother*. 2005;54:933-943.
- Gogas H, Ioannovich J, Dafni U, et al. Prognostic significance of autoimmunity during treatment of melanoma with interferon. *N Engl J Med*. 2006;354:709-718.
- Rinaldi M, Ria F, Parrella P, et al. Antibodies elicited by naked DNA vaccination against the complementary-determining region 3 hypervariable region of immunoglobulin heavy chain idiotype determinants of B-lymphoproliferative disorders specifically react with patients' tumor cells. *Cancer Res*. 2001;61:1555-1562.
- Moldrem JJ, Kant S, Lu S, et al. Peptide vaccination with PR1 elicits active T cell immunity that induces cytogenetic remission in acute myelogenous leukemia. Paper presented at the Annual Meeting of the American Society of Hematology. December 6-10, 2002. Philadelphia, PA.
- Westervelt P, Pollock JL, Oldfather KM, et al. Adaptive immunity cooperates with liposomal all-trans-retinoic acid (ATRA) to facilitate long-term molecular remissions in mice with acute promyelocytic leukemia. *Proc Natl Acad Sci U S A*. 2002;99:9468-9473.
- Yu S, Xia M, Xu W, Chu Y, Wang Y, Xiong S. All-trans retinoic acid biases immune response induced by DNA vaccine in a Th2 direction. *Vaccine*. 2005;23:5160-5167.
- Geissmann F, Revy P, Brousse N, Lepelletier Y, et al. Retinoids regulate survival and antigen presentation by immature dendritic cells. *J Exp Med*. 2003;198:623-634.
- Lollini PL, Cavallo F, Nanni P, Forni G. Vaccines for tumour prevention. *Nat Rev Cancer*. 2006;6:204-216.