

Role of CD28 in fatal autoimmune disorder in *scurfy* mice

Nagendra Singh,¹ Phillip R. Chandler,¹ Yoichi Seki,¹ Babak Baban,¹ Mayuko Takezaki,¹ David J. Kahler,¹ David H. Munn,¹ Christian P. Larsen,² Andrew L. Mellor,¹ and Makio Iwashima¹

¹Immunotherapy Center, Medical College of Georgia, Augusta; and ²Emory Transplant Center, Department of Surgery, and Yerkes Regional Primate Research Center, Emory University School of Medicine, Atlanta, Georgia

***Scurfy* mice develop CD4 T-cell-mediated lymphoproliferative disease leading to death within 4 weeks of age. The *scurfy* mutation causes loss of function of the *foxp3* gene (*foxp3^{sh}*), which is essential for development and maintenance of naturally occurring regulatory CD4 T cells (nTregs). In humans, mutations of the *foxp3* gene cause immune dysregulation, polyendocrinopathy, enteropathy, and X-linked syndrome (IPEX). In most patients with IPEX and also in *scurfy* mice, T cells**

show hyperreactivity and levels of Th1- and Th2-associated cytokines are substantially elevated. We report that removal of CD28 expression rescued *scurfy* mice from early death. Longer-term surviving CD28-deficient *scurfy* mice still had lymphoproliferative disorder, but their CD4 T cells showed decreased interferon- γ and no sign of interleukin-4 or interleukin-10 hyperproduction. Furthermore, injection of CTLA4-Ig to block CD28-B7 interactions substantially improved the survival of

***scurfy* mice by blocking effector T-cell differentiation. These data support the hypothesis that CD28-B7 interactions play a critical role in the etiology of lethal autoimmune disease in *scurfy* mice by stimulating the differentiation of antigen-activated naive T cells into effector T cells. (Blood. 2007;110:1199-1206)**

© 2007 by The American Society of Hematology

Introduction

The *scurfy* mutation causes loss of function of the *foxp3* gene, which is essential for development and maintenance of naturally occurring regulatory CD4⁺CD25⁺ T cells (nTregs).¹ Disease in *scurfy* mice is characterized by lymphocytic infiltration of lymph nodes, spleen, liver, and skin.² CD4⁺ T cells from these mice were hyperresponsive to T-cell receptor (TCR) stimulation and produced highly elevated levels of cytokines, including interleukin (IL)-2, -4, -5, -10, -13, interferon (IFN)- γ , granulocyte-macrophage colony-stimulating factor, and tumor necrosis factor- α .^{3,4} These hyperreactive T cells were relatively resistant to immunosuppressive drugs.^{4,5} When adoptively transferred into nu/nu recipients, CD4 T cells from *scurfy* mice induced a rapid wasting disease indicating that these cells play a central role in *scurfy* disease development.⁶

CD28 plays a pivotal role in antigen-peptide-induced proliferation of naive T cells,⁷ whereas CD28 costimulation increases the level of IL-2 mRNA by transcriptional and posttranscriptional mechanisms,⁸⁻¹⁰ up-regulates expression of the antiapoptotic protein Bcl-XL that promotes cell survival,¹¹⁻¹³ and elevates antigen receptor proximal events by strengthening immunologic synapse.^{14,15} CD28 dependency decreases under conditions of high antigenic load or in the presence of high avidity peptide agonists.¹⁶ In contrast to naive T cells, proliferation and cytokine production by recently activated T cells and memory T cells are less dependent on CD28 costimulation.¹⁷ CD28/B7 costimulation plays a critical role also in the control of Th1/Th2 balance.⁷ Differentiation of naive CD4⁺T cells toward Th2, but not Th1 phenotype, is dependent on CD28/B7 costimulation both in vitro and in vivo.^{18,19}

CTLA-4 is an inhibitory cell surface receptor of the CD28 family and down-regulates T cell responses. Although CD28 is

constitutively expressed on resting and activated T cells, CTLA-4 is not expressed by resting T cells but is induced on activation.²⁰ CD28 and CTLA-4 compete for binding to CD80 (B7.1) and CD86 (B7.2). Studies suggest that CD86 preferentially interacts with CD28, in contrast to CD80, which binds to CTLA-4 more strongly than to CD28.²¹ Thus, a fusion protein consisting of the extracellular domain of CTLA-4 and the Fc part of IgG, CTLA4Ig, through its binding to CD80 and CD86 inhibits CD28-mediated positive signals and blocks T-cell responses to prevent autoimmune diseases and transplant rejections (reviewed in ²²). Disruption of the CTLA-4 gene leads to uncontrolled T cell lymphoproliferation in mice leading to death by 3 to 4 weeks after birth. Lymphoproliferation in CTLA-4-deficient mice is mediated by CD28 signaling, because mice lacking all 3 molecules, CTLA-4, B7.1, and B7.2 (*ctla-4^{-/-}*, *b7.1^{-/-}*, *7.2^{-/-}*), or both CTLA-4 and CD28 (*ctla-4^{-/-}cd28^{-/-}*) showed no hyperproliferation.^{23,24} It should be noted that nTregs constitutively express higher amounts of CTLA-4 than do naive conventional T cells, and previous studies have implicated a role for CTLA-4 in the suppressive mechanism used by nTregs.^{25,26}

CD28 also plays a critical role in autoimmunity. Studies using antigen-induced experimental autoimmune encephalomyelitis, a model of human multiple sclerosis, showed that disruption of CD28/B7 interactions at the time of immunization was associated with a reduction in the severity of the autoimmune disease.²⁷⁻²⁹ Blockade of CD28/B7 interactions also reduced disease severity in mouse models of myocarditis,³⁰ arthritis,³¹ thyroiditis,³² and myasthenia gravis.³³ In the New Zealand black/New Zealand white mouse strain, which develops an antibody-dependent lupus-like autoimmune syndrome, treatment with CTLA4Ig or a combination

Submitted October 27, 2006; accepted April 23, 2007. Prepublished online as *Blood* First Edition paper, April 26, 2007; DOI 10.1182/blood-2006-10-054585.

The online version of this article contains a data supplement.

The publication costs of this article were defrayed in part by page charge payment. Therefore, and solely to indicate this fact, this article is hereby marked "advertisement" in accordance with 18 USC section 1734.

© 2007 by The American Society of Hematology

of anti-B7-1 and anti-B7-2 antibodies prevented the development and progression of the disease.^{34,35}

We examined the role of CD28 in the pathogenesis of autoimmune disease development in *scurfy* mice. The data demonstrate that CD28 plays a pivotal role in the early death of mice with the *scurfy* mutation. However, lymphoproliferative disorders of CD28-deficient *scurfy* mice remained as severe as those of CD28-sufficient *scurfy* mice, showing that aberrant T-cell proliferation caused by loss of nTregs is not sufficient to cause the early death of *scurfy* mice.

Materials and methods

Mice

Female *scurfy* mice, CD28 knockout mice (backcrossed to C57BL/6 for either 8 or 12 generations, respectively), and C57BL/6 mice were purchased from Jackson Laboratory (Bar Harbor, ME). Female *foxp3^{sf/+}* mice were crossed with *cd28^{-/-}* males to obtain *foxp3^{sf/y}cd28^{+/-}* and *foxp3^{sf/+}cd28^{+/-}* progeny. Female *foxp3^{sf/+}cd28^{+/-}* mice were then backcrossed with *cd28^{-/-}* males to obtain *foxp3^{sf/y}cd28^{-/-}* males. From this mating, we screened 4 litters and obtained 3 *cd28^{-/-}scurfy* males, which were mated with *cd28^{-/-}foxp3^{sf/+}* females to generate *cd28^{-/-}scurfy* males and females. Mice carrying the *scurfy* mutation or CD28 gene disruption were identified by polymerase chain reaction using primers and conditions as suggested by Jackson laboratory. All procedures have been reviewed and approved by the Medical College of Georgia Institutional Animal Care and Use Committee (IACUC).

Antibodies and flow cytometry

Fluorochrome- and biotin-coupled monoclonal antibodies directed against CD4 (RMA 4-5), CD8 (53-6.7), CD62L (MEL-14), CD25 (7D4), CD3 (17A2), and DX5 were purchased from BD Biosciences (San Jose, CA). Anti-CD3 (2c11) and CD28 (37-51) were from eBiosciences (San Diego, CA). Stained cells were analyzed on a FACS Calibur (BD Biosciences).

T-cell proliferation and cytokine assay

Defined T-cell populations were sorted by Mo-Flo (Dako Colorado, Fort Collins, CO) or FACSaria (BD Biosciences) cell sorter. Lymph node T cells were purified by panning or a Mo-Flo cell sorter and were cultured with plate-bound anti-CD3 and soluble anti-CD28 antibody (0.5 mg/mL) in RPMI 1640 supplemented with 5% fetal calf serum (FCS), penicillin/streptomycin, 2 mM L-glutamine, and 50 μ M 2-mercaptoethanol in 96-well plates. After 48 hours culture, supernatant (50 μ L) was collected for cytokine analysis and plates were then pulsed with 0.5 μ Ci/well of [³H]-thymidine to measure proliferation. Cytokine production was analyzed by enzyme-linked immunosorbent assay as described.³⁶

Histology

Tissues were fixed in 10% buffered formalin and embedded in paraffin. Hematoxylin and eosin-stained sections (5 μ m) were mounted with Cytoseal60 (Richard Allan Scientific, Kalamazoo, MI) and analyzed by light microscopy. Images were acquired at room temperature using an Olympus Provis (Model AX70TRF; Tokyo, Japan) microscope equipped with Spot Camera (model 1.3.0; Diagnostic Instruments, Sterling Heights, MI) and Spot software.

CTLA4-IgG2a treatment

One hundred micrograms CTLA4-IgG2a in 50 μ L phosphate-buffered saline (PBS) was injected intraperitoneally on days 7, 11, 15, and 19 postpartum.

Results

Loss of CD28 expression rescues *scurfy* mice from early death

To examine the role of CD28 in the pathogenesis of *scurfy* mice, we bred female (*foxp3^{sf/+}*) mice with *cd28* gene knockout (*cd28^{-/-}*) mice. Male F1 mice with a *cd28^{+/-}foxp3^{sf}* genotype (n = 9) survived longer than parental *scurfy* mice (n = 9) (mean survival time 63 days and 24 days, respectively). From F1 female *cd28^{+/-}foxp3^{sf/+}* mice, we derived male *cd28^{-/-}foxp3^{sf}* mice (n = 10) 50% of which survived longer than 200 days and were fertile (Figure 1A). These data demonstrate that CD28 is a component of the lethal autoimmune disease caused by the *scurfy* mutation.

Despite the increased survival, 3-week-old *cd28^{-/-}* male *scurfy* mice exhibited splenomegaly and lymphadenopathy similar to those of age-matched *cd28^{+/+}* male *scurfy* mice (Figure 1B; Figure S1, available on the *Blood* website; see the Supplemental Materials link at the top of the online article). Substantially increased cellularities were observed in the lymph nodes of *cd28^{-/-}scurfy* mice relative to *cd28^{+/+}scurfy* mice, mainly as a result of an increase in the numbers of T and B cells. Although lower percentages of CD4⁺ cells were present in *cd28^{-/-}scurfy* mice, cell numbers of CD4⁺ cells were equivalent between *cd28^{+/+}* and *cd28^{-/-}scurfy* mice (Figure 1B). Similar patterns of elevation in cell numbers also occurred in the spleens (data not shown). These data show that prolonged survival did not correlate with decreased T-cell expansion in *cd28^{-/-}scurfy* mice.

In contrast, the number of thymocytes in 3-week-old *cd28^{-/-}scurfy* mice showed a profound difference compared with *cd28^{+/+}scurfy* mice (Figure 1C). *cd28^{+/+}scurfy* mice thymi were involuted (Figure S1) and had 40-fold lower numbers of cells than in wild-type and *cd28^{-/-}* mice (Figure 1D). Few CD4⁺CD8⁺ cells were present in *cd28^{+/+}scurfy* mouse thymi (Figure 1C), whereas thymi from *cd28^{-/-}scurfy* mice displayed the same levels of surface antigen expression as thymi from wild-type mice (Figure 1D). A recent report showed a potential defect in thymic stroma cells of *scurfy* mice that caused impairment of CD4⁺CD8⁻ cell thymopoiesis.³⁷ We found similar percentages and numbers of CD4⁺CD8⁺ cells in 5-day-old *cd28^{+/+}scurfy* mice as well as in *cd28^{+/+}* mice (Figure 2). Thus, the data presented here suggest that FoxP3 does not itself play a role in thymopoiesis and the absence of CD4⁺CD8⁺ cells in 3-week-old *scurfy* mice may be the result of an effect of the thymic and/or peripheral environment of *scurfy* mice with advanced autoimmune disease.

Impaired effector cytokine production by *-/-*CD4 T cells from *cd28^{-/-}scurfy* mice

In the periphery of *cd28^{+/+}scurfy* mice, approximately 50% of CD4 T cells showed downmodulation of CD3, whereas in *cd28^{-/-}scurfy* mice, this was only 10% of CD4 T cells (Figure 3A). Moreover, more than 50% of *cd28^{+/+}scurfy* mice CD4 T cells were CD44^{high}CD62L⁻, consistent with an effector/memory phenotype. In contrast, the CD4 T-cell compartment in *cd28^{-/-}scurfy* mice contained only 16% of CD44^{high}CD62L⁻ cells, whereas the majority (54%) were found to be CD44^{low}CD62L⁻. More than 50% of CD4 T cells from wild-type mice were CD44^{low}CD62L⁺ (Figure 3B). Thus, a majority of the population of CD4⁺ T cells in *cd28^{-/-}scurfy* mice T cells displayed a phenotype intermediate between activated T cells in *cd28^{+/+}scurfy* mice and resting naive T cells in wild-type mice.




Figure 1. State of T cells from CD28 deficient *scurfy* mice. (A) Lifespan of *scurfy* (*sf*), CD28 heterozygous *scurfy* [*sf/CD28(+/-)*], and CD28-deficient *scurfy* [*sf/CD28(-/-)*] mice ($P < .001$). (B) Total number of live cells present in inguinal lymph nodes (left panel) and subpopulations of lymph node cells (right panel). CD4⁺ cells: □, CD8⁺ cells: ■, B220⁺ cells: ■ (mean ± SD, n = 3). (C) CD4/CD8 phenotypes of lymph node cells (top row) and thymocytes (bottom row). Cells from 25-day-old C57BL/6 (WT), CD28-deficient *scurfy* [CD28(-/-) *scurfy*], and *cd28^{+/+} scurfy* (*scurfy*) mice were stained for CD4 and CD8 and analyzed by flow cytometry. Numbers in each quadrant indicate the percentage of cells in each region against the total. (D) Total number of live cells present in each thymic lobe from C57BL/6 (WT), CD28-deficient (CD28-), CD28-deficient *scurfy* [CD28- *scurfy*], and CD28-sufficient *scurfy* (*scurfy*) mice (mean ± SD, n = 3).

Despite the surface antigen phenotypes of mild activation, T cells from *cd28^{-/-} scurfy* mice responded to anti-CD3 stimulation *in vitro* as vigorously as those from age-matched *cd28^{+/+} scurfy* mice (Figure 3C). Addition of anti-CD28 antibody did not enhance the proliferation of T cells from either *cd28^{-/-}* or *cd28^{+/+} scurfy* mice but did increase the proliferation of T cells from wild-type mice. Moreover, when stimulated with anti-CD3, CD4 T cells from *cd28^{-/-} scurfy* mice produced much higher levels of IL-2 compared with CD4 T cells from *cd28^{+/+} scurfy* mice (Figure 3D). Elevation of IL-2 production was also observed in CD8 and natural killer T (NKT) cells from *cd28^{-/-} scurfy* mice (Figure S2).

Most notably, anti-CD3 Ab-stimulated CD4 T cells from *cd28^{-/-} scurfy* mice produced levels of IL-4 and IL-10 comparable to those from wild-type mice but exhibited a more than 500-fold reduction compared with CD4 T cells from *cd28^{+/+} scurfy* mice. Engagement of CD28 on *cd28^{+/+} scurfy* mice T cells did not change the level of IL-4 production and only slightly enhanced IL-10 production, indicating that these cells are already CD28-independent in terms of IL-4 and IL-10 production. CD4 T cells from *cd28^{-/-} scurfy* mice also produced much less IFN- γ compared with CD4 T cells from *cd28^{+/+} scurfy* mice.

To determine whether subpopulations of CD4 T cells have differences in their ability to produce effector cytokines, we compared TCR-induced responses of CD62L⁺CD44⁻ (naive phenotype) and CD62L⁻CD44⁺ (effector/memory phenotype) CD4⁺CD25⁻ T cells from wild-type, *scurfy*, and *cd28^{-/-} scurfy* mice (Figure 4). We used groups of *cd28^{+/+}*, *cd28^{+/-} scurfy*, and *cd28^{-/-} scurfy* mice because they can be obtained as littermates. Anti-CD3 stimulation induced substantial proliferation of naive (CD62L⁺CD44⁻) CD4 T cells from all genotypes and we observed no hyperreactivity in cells from *cd28^{-/-} scurfy* mice (Figure 4A). CD28 costimulation augmented the anti-CD3 induced proliferation of naive CD4 T cell from both wild-type and *scurfy* mice by 6- to 10-fold, whereas it had no effect on the proliferation of *cd28^{-/-} scurfy* naive T cells (Figure 4A). In contrast, effector/memory type CD4⁺ T cells (CD62L⁻CD44⁺) from *cd28^{+/-}* and *cd28^{-/-} scurfy* mice were hyperresponsive to anti-CD3 stimulation compared with their wild-type counterparts. Addition of anti-CD28 induced a 2.2-fold increase in proliferation of CD4 T cells from *scurfy* mice in the presence of 0.1 μ g/mL anti-CD3 and a 3.1-fold increase was observed in CD4 T cells from wild-type mice activated with 1 μ g/mL anti-CD3 antibody (Figure 4A).




Figure 2. Thymocytes from 5- and 27-day-old *scurfy* and wild-type mice were analyzed for CD4 and CD8 expression. Numbers in each quadrant indicate the percentage of cells in each region against the total.




Figure 3. Reactivity of T cells from CD28-deficient *scurfy* mice. (A,B) In vivo activation of *cd28*^{-/-} *scurfy* mouse T cells. Lymph node cells from mice indicated above each panel were stained for CD4, CD8, CD3, CD44, and CD62L. Shown are (A) CD3 and (B) CD44 and CD62L staining on CD4⁺CD8⁻-gated cells. In panel A, numbers in each subpanel indicate the percentage of cells in each region against the total. In panel B, numbers in or above each quadrant indicate the percentage of cells in each region against the total. (C) Proliferative response of T cells from *cd28*^{-/-} *scurfy* mouse. Purified T cells from *cd28*^{-/-} and *cd28*^{+/+} mice were stimulated with anti-CD3 (left panel) or anti-CD3+anti-CD28 (right panel) antibodies. Proliferation of T cells was measured in response to titrated doses of anti-CD3, respectively, by ³H-thymidine incorporation (mean ± SD, n = 3). (D) Cytokine production by T cells from *cd28*^{-/-} *scurfy* mice. Mo-Flo sorted TCR⁺CD4⁺DX5⁻ cells (2 × 10⁴ cells/well in flat-bottomed 96-well plate) were activated with 1 μg/mL anti-CD3-coated plates in the absence □, or presence ■, of anti-CD28 antibody (0.5 μg/mL). Forty-eight hours later, a portion of culture supernatant was harvested for analysis of cytokines as indicated on the top of the panels and culture supernatant from stimulated T cells were analyzed for cytokines denoted above each panel by enzyme-linked immunosorbent assay. CD4 T cells from wild-type (WT), *cd28*^{-/-} [28(-)sf] and *cd28*^{+/+} *scurfy* (sf) mice were analyzed (mean ± SD, n = 2).

Next we analyzed cytokine production by naive and effector/memory CD4 T cells. When stimulated in vitro with anti-CD3, naive CD4 T cells from *cd28*^{-/-} *scurfy* mice produced less IL-2 than those from *cd28*^{+/+} wild-type or *cd28*^{+/+} *scurfy* mice (Figure

4B). CD28 costimulation clearly augmented IL-2 production by *cd28*^{+/+} *scurfy* mouse T cells indicating that CD28 plays a critical role in IL-2 production even with the *scurfy* background. In contrast, effector/memory CD4 T cells from *cd28*^{-/-} *scurfy* mice made significantly higher levels of IL-2 compared with CD4 T cells from *cd28*^{+/+} *scurfy* and *cd28*^{+/+} wild-type mice. Although the levels of IL-2 production by CD4 T cells from *cd28*^{+/+} *scurfy* and wild-type mice were augmented by anti-CD28 stimulation, they did not reach the levels of IL-2 production by CD4 T cells from *cd28*^{-/-} *scurfy* mice.

As expected, naive CD4 T cells from all genotypes failed to produce IL-4, and effector/memory CD4 T cells from *cd28*^{+/+} *scurfy* mice produced significantly higher amounts of IL-4 (Figure 4B) and IL-10 (Figure S2) than CD4 T cells from *cd28*^{+/+} or *cd28*^{-/-} *scurfy* mice. A small elevation was observed in the level of IFN-γ production by naive and effector/memory T cells from *cd28*^{+/+} *scurfy* mice compared with that by T cells from *cd28*^{+/+} wild-type and *cd28*^{-/-} *scurfy* mice (Figure S2). Taken together, effector/memory (CD62L⁻CD44⁺) CD4 T cells from *cd28*^{-/-} *scurfy* mice showed patterns of cytokine production clearly distinct from those of T cell from *cd28*^{+/+} *scurfy* mice. These data indicate that T cells from *cd28*^{-/-} mice are defective in the differentiation process from the IL-2-producing stage into the Th2 (and Th1 to a lesser degree) type cytokine-producing cells.

NK T cells from *cd28*^{-/-} *scurfy* mice also produced less than 10% IL-4 and IL-10 compared with NK T cells from *cd28*^{+/+} *scurfy* mice (Figure S3). On the other hand, IFN-γ production by CD8 T cells and NK T cells was similar between *cd28*^{-/-} and *cd28*^{+/+} *scurfy* mice compared with wild-type mice.

Differences between CD28-deficient and -sufficient *scurfy* mice in vivo

To test whether differences in cytokine production reflected the state of T cells in vivo, we examined sera from *cd28*^{-/-} and *cd28*^{+/+} *scurfy* mice. Whereas sera from *cd28*^{+/+} *scurfy* mice showed extremely high levels of IL-4 and IFN-γ, these cytokines were undetectable in sera from both wild-type and *cd28*^{-/-} *scurfy* mice (Figure 5A). Moreover, as observed in patients with immune




Figure 4. Responses of naive and activated CD4 T cells from CD28^{-/-} *scurfy* mice. CD4⁺CD25⁻ NK1.1⁻ populations from mice as indicated were sorted into CD62L⁺CD44⁻ and CD62L⁻CD44⁺ populations. Sorted cells (1.5 × 10⁴ cells/well in flat-bottomed 96-well plate) were activated in triplicate wells coated with 0.1 or 1 μg/mL of anti-CD3 in the presence or absence of anti-CD28 antibody (0.5 μg/mL). (A) Proliferation (mean ± SD, n = 3) and (B) cytokine production (mean ± SD, n = 2) was determined as in Figure 3.




Figure 5. Analysis of in vivo lymphocyte functions of *cd28*^{-/-} scurfy mice. Serum from *cd28*^{-/-} and *cd28*^{+/+} scurfy mice were analyzed for (A) interleukin-4 and interferon-γ, and (B) immunoglobulin isotypes. WT: C57BL/6, 28(-): *cd28*^{-/-} mice with C57BL/6 background, 28(-) sf: *cd28*^{-/-} scurfy mice, sf: *cd28*^{+/+} scurfy mice (mean ± SD, n = 3).

dysregulation, polyendocrinopathy, enteropathy, X-linked syndrome (IPEX),⁴ IgE and IgG₁ levels were highly elevated in the sera of *cd28*^{+/+} scurfy mice compared with *cd28*^{-/-} scurfy and wild-type mice (Figure 5B). Although IgM levels were significantly higher in *cd28*^{-/-} scurfy mice than in *cd28*^{+/+} scurfy mice, levels of total Ig were comparable.

In addition to elevated cytokine production, the disease of scurfy mice is characterized by multiorgan tissue destruction by

infiltrating lymphocytes.¹ Mononuclear infiltrates were present in the lungs (Figure S4A) and livers (Figure S4B) of 3-week-old *cd28*^{+/+} scurfy mice, whereas no major infiltrates were detected in either of these organs from *cd28*^{-/-} scurfy mice.

CTLA4-Ig injection rescues scurfy mice from early death

The data presented here demonstrated that CD28 plays a pivotal role in disease development in scurfy mice. This predicts that blockade of CD28 signaling in scurfy mice will protect them from lethal disease. To test this prediction, we examined whether injection of soluble CD152 (CTLA4)-Ig into scurfy mice blocks progression of disease. CTLA4-Ig is a fusion protein that has been used successfully to block CD28 function by inhibiting interactions between B7 molecules and CD28 both in vitro and in vivo.³⁸ Litters of scurfy mice of unknown phenotype were injected with CTLA4-Ig (100 μg) or PBS every 4 days from day 7 until day 19. After treatment, litters of mice were either killed at day 22 for analysis or monitored further without any additional injections.

Scurfy mice (n = 5) injected with CTLA4-Ig showed a markedly increased lifespan (Figure 6A). No visible symptoms were observed at day 22 and the first scurfy mouse to become moribund was killed at day 42. Two of the remaining mice became moribund at day 48 and 2 others remained healthy past day 70. Because CTLA4-Ig injection was stopped at day 19, the data suggest that continuous costimulatory blockade is important to prevent lethality of scurfy disease.

CTLA4-Ig-injected scurfy mice killed on day 22 showed phenotypes in their T-cell compartment comparable with those of *cd28*^{-/-} scurfy mice. Lymphoid organs of CTLA4-Ig-injected scurfy mice (day 22) were enlarged and the cell numbers were highly elevated compared with wild-type (Figure 6B). Analysis of the surface antigens showed that CTLA4-Ig injection did not alter the CD4/CD8 ratio but reduced the numbers of CD44^{high}CD62L⁻ T cells in scurfy mice compared with PBS-injected scurfy mice (Figures 6C,D). At day 48, however, T cells from CTLA4-Ig-injected scurfy mice were predominantly CD44^{high} CD62L⁻ as was also observed in PBS-injected scurfy mice, indicating that self-reactive T cells are fully activated by this time.




Figure 6. Effect of CTLA4-Ig injection in scurfy mice. (A) Lifespan of CTLA4-Ig-injected (dotted line, n = 5) and phosphate-buffered saline-injected (solid line, n = 3) scurfy mice. Intraperitoneal CTLA4-Ig injection took place every 4 days from day 7 until day 19 after birth (shown above the panel) (P < .001). (B) Total cell number of inguinal lymph nodes from phosphate-buffered saline- and CTLA4-Ig-injected scurfy (sf) and wild-type (wt) mice. (C,D) Surface antigen analysis of T cells from CTLA4-Ig-injected scurfy mice. Numbers in each quadrant indicate the percentage of cells in each region against the total. (C) CD4 and CD8 and (D) CD44 and CD62L (CD4 gated) surface phenotypes of inguinal lymph node cells from wild-type and scurfy mice. Cells were isolated from CTLA4-Ig-injected scurfy mice at day 22 (d22) and day 48 (d48) after birth. As a control, data from phosphate-buffered saline-injected scurfy mice at day 22 are shown.




Figure 7. Functional analysis of T cells from CTLA4-Ig-injected mice. CD4 T cells were isolated from CTLA4-Ig-injected *scurfy* mice at day 22 (d22) and day 48 (d48) after birth. As a control, data from a phosphate-buffered saline-injected *scurfy* mouse at day 22 are shown. T cells were stimulated with anti-CD3 (□) or anti-CD3 and CD28 (■). (A) ³H-thymidine uptake of T cells after 72 hours of stimulation (proliferation) (mean ± SD, n = 3). (B) Interleukin-2, interleukin-4, interleukin-10, and interferon-γ production after 48 hours are shown. Results are representative of 3 CTLA4-Ig-injected mice (mean ± SD, n = 2).

T cells from CTLA4-Ig-injected and PBS-injected *scurfy* mice were equally reactive to anti-CD3 stimulation and proliferated vigorously without CD28 stimulation (Figure 7A). Yet cytokine production by T cells from day 22 CTLA4-injected *scurfy* mice showed clear differences from those of PBS-injected *scurfy* mouse T cells (Figure 7B). IL-2 production was elevated significantly compared with PBS-injected *scurfy* mice. Conversely, IL-4, IL-10, and IFN-γ production was clearly reduced in CTLA4-Ig-injected *scurfy* mice compared with PBS-injected mice. In addition, histology showed a clear absence of infiltration by mononuclear cells in the lungs of CTLA4-Ig-injected *scurfy* mice at day 22 (Figure S5).

At day 48, CTLA4-Ig-injected mice showed overt symptoms of weight loss, skin lesions, and lack of mobility. CD4 T cells from 48-day-old CTLA4-Ig-injected *scurfy* mice produced elevated levels of IL-10 compared with their 22-day-old counterparts, whereas the levels of IL-4 and IFN-γ were comparable between day 22 and 48 CTLA4-Ig-injected *scurfy* mice (Figure 7B). CD4 T cells from CTLA4-Ig-treated *scurfy* mice, regardless of their age, produced less IL-4 and IFN-γ than those from PBS-treated *scurfy* mice (Figure 7B). However, the amounts of IL-4 and IL-10 produced by CD4 T cells from CTLA4-Ig-treated *scurfy* mice were still several-fold higher than those produced by CD4 T cells from CTLA4-Ig-injected littermate wild-type mice (data not shown). In contrast, IFN-γ production by CD4 T cells from CTLA4-Ig-injected *scurfy* mice was mildly increased compared with CTLA4-Ig-treated wild-type mice (data not shown). Both the PBS-injected *scurfy* mice (day 22) and CTLA4-Ig-injected *scurfy* mice (day 48) showed significant mononuclear cell infiltration in their lungs (Figure S5). These data suggest that production of IL-4, IL-10, and/or infiltration by mononuclear cells may contribute to the progression of *scurfy* disease.

Discussion

Our data show that CD28 plays a pivotal role in the progression of lethal autoimmune disease in *scurfy* mice and that survival of

cd28^{-/-} *scurfy* mice is unrelated to aberrant T-cell proliferation. Indeed, CD62L⁻CD44⁺ CD4 T cells from *cd28*^{-/-} *scurfy* mice produced more IL-2 than *cd28*^{+/-} *scurfy* mouse CD4 T cells and lack of CD28 did not impair their ability to proliferate after CD3 stimulation. The data demonstrated that in *scurfy* mice, CD4 T cells expand without committing to become effectors when CD28 is absent. Moreover, CD28 is also required for the process leading to tissue infiltration by lymphocytes.

Importantly, naive CD4 T cells (CD62L⁺CD44⁻) from *scurfy* and wild-type mice responded in a similar manner to anti-CD3 and anti-CD28 stimulation, whereas naive CD4 T cells from *cd28*^{-/-} *scurfy* mice showed reduced ability to produce IL-2. In contrast, effector/memory CD4 T cells (CD62L⁻CD44⁺) from *cd28*^{+/-} and *cd28*^{-/-} *scurfy* mice were hyperproliferative in response to anti-CD3 stimulation. Collectively, these data suggest that hypersensitivity of *scurfy* CD4 T cells is not a cell intrinsic property of the *scurfy* background, but is most likely the result of the presence of large number of preactivated T cells caused by lack of nTregs.

In *scurfy* mice, activated CD4 T cells differentiate into effectors and produce IL-4, IL-10, and IFN-γ. Surprisingly, activated CD4 T cells from *cd28*^{-/-} *scurfy* mice produced increased amounts of IL-2, decreased IFN-γ, and virtually no IL-4 and IL-10. The data demonstrate that even in the absence of nTregs, CD28 signaling plays a critical role in commitment of activated CD4 T cells into Th2 or, to a lesser extent, Th1 effectors. Failure of CD4 T cells from *cd28*^{-/-} *scurfy* mice to produce effector cytokines is consistent with published reports (reviewed in Salomon and Bluestone⁷).

In contrast to IL-4 and IL-10, IL-2 production was much up-regulated in *cd28*^{-/-} *scurfy* mice. Increased amount of IL-2 production by CD4 T cells from *cd28*^{-/-} *scurfy* mice is not caused by a cell intrinsic abnormality of naive T cells in *scurfy* mice because they show no sign of enhanced IL-2 production. The reason why *cd28*^{-/-} effector T cells make enhanced levels of IL-2 is currently unclear. Although beyond the scope of this article, one question remains to be answered: how naive CD4 T cells become activated in *scurfy* mice to produce IL-2 in the absence of CD28.

Previous reports showed that CD28 costimulation is not always required for T-cell proliferation and IL-2 production, especially when strong and/or sustained antigen receptor signal is provided. Thus, one potential explanation for CD28-independent IL-2 production is that naive T cells receive prolonged antigenic stimulation in the absence of nTregs. Another mutually nonexclusive possibility is the contribution from homeostatic expansion. During the neonatal period, T cells undergo homeostatic proliferation in response to lymphopenia.³⁹ Homeostatic expansion of T cells is less dependent on CD28 than antigen-induced proliferation of T cells and can promote polyclonal proliferation (reviewed in Khoruts and Fraser⁴⁰). Thus, the lymphopenic environment of the neonate may initiate CD28-independent activation of self-reactive CD4 T cells in *scurfy* mice.

The data presented here exhibit interesting similarities and differences between *scurfy* and *ctla4*^{-/-} mice.^{23,24,41,42} Like *scurfy* mice, *ctla4*^{-/-} mice develop a lymphoproliferative disorder by 3 weeks of age and die around the same time.⁴³ *cd28*^{-/-}*ctla4*^{-/-} mice, *b7-1*^{-/-}*b7-2*^{-/-}*ctla4*^{-/-} mice, and *ctla4*^{-/-} mice injected with CTLA4-Ig all show dramatically extended lifespans and reduced levels of cytokine production by T cells.^{23,24,41,42} However, CD28-deficient and CTLA4-Ig-treated *ctla4*^{-/-} mice show striking differences from CD28-deficient and CTLA4-Ig-treated *scurfy* mice. For example, when CD28-B7 interactions are blocked, T cells from *ctla4*^{-/-} mice do not show any lymphoproliferative disorders and exhibited decreased T-cell responsiveness in vivo and

in vitro. *cd28^{-/-}ctla4^{-/-}* T cells are hypoproliferative at levels comparable with those of *cd28^{-/-}* mouse T cells.²⁴ Thus, hyperproliferation of T cells caused by loss of CTLA4 requires CD28, but hyperproliferation of T cells from *scurfy* mice is independent of CD28. It was also shown that nTregs from CTLA4-deficient mice have uncompromised suppressive activity.⁴⁴ Thus, in the absence of CD28, these nTregs may dominate the reactivity of nonregulatory T cells, whereas lack of Foxp3 may cause complete loss of suppressive function.

Our data show a potential connection between effector T-cell response and lethality in *scurfy* disease. CD4 T cells from CD28-deficient *scurfy* mice showed no sign of IL-4, IL-10, and IFN- γ hyperproduction. Similarly, a potential connection between Th2 response and lethality of *scurfy* mice was previously reported with *cd4^{-/-} scurfy* mice.⁶ Moreover, levels of IL-4, IL-10, and IFN- γ are highly elevated in patients with IPEX.^{3,4} Thus, uncontrolled cytokine production by *scurfy* T cells may be responsible for lethal disease development in *scurfy* mice.

An intriguing observation was made in the CTLA4-Ig-injected *scurfy* mice when killed on day 48. These mice showed clear symptoms of *scurfy* disease. Because CTLA4-Ig injection was discontinued at day 19 after birth, the data suggest that continuous blockade of B7 molecules may be required to attenuate progression of *scurfy* disease. Alternatively, it may reflect the difference of the mode by which CTLA4-Ig works from deficiency of CD28 expression. Several reports of B7-CTLA-4 pathways of immunoregulation have been published and not all depend on B7-CD28-mediated costimulation (reviewed in Alegre and Fallarino⁴⁵). We are currently in the process of distinguishing among the potential pathways that may lead to the observed phenotype.

Except for IL-10 levels in 48-day-old mice (which were elevated), the amounts of IL-4, IL-10, and IFN- γ produced by CD4 T cells from CTLA4-Ig-treated *scurfy* mice were reduced compared with CD4 T cells from PBS-injected *scurfy* mice. However, cytokine levels were still much higher than those produced by CD4 T cells from CTLA4-Ig-injected wild-type mice. This may be enough over time to induce disease in CTLA4-Ig-injected *scurfy* mice at a later time point.

References

- Ochs HD, Ziegler SF, Torgerson TR. FOXP3 acts as a rheostat of the immune response. *Immunol Rev.* 2005;203:156-164.
- Godfrey VL, Wilkinson JE, Russell LB. X-linked lymphoreticular disease in the *scurfy* (*sf*) mutant mouse. *Am J Pathol.* 1991;138:1379-1387.
- Kanangat S, Blair P, Reddy R, et al. Disease in the *scurfy* (*sf*) mouse is associated with overexpression of cytokine genes. *Eur J Immunol.* 1996;26:161-165.
- Wildin RS, Smyk-Pearson S, Filipovich AH. Clinical and molecular features of the immunodysregulation, polyendocrinopathy, enteropathy, X linked (IPEX) syndrome. *J Med Genet.* 2002;39:537-545.
- Baud O, Goulet O, Canioni D, et al. Treatment of the immune dysregulation, polyendocrinopathy, enteropathy, X-linked syndrome (IPEX) by allogeneic bone marrow transplantation. *N Engl J Med.* 2001;344:1758-1762.
- Blair PJ, Bultman SJ, Haas JC, Rouse BT, Wilkinson JE, Godfrey VL. CD4+CD8- T cells are the effector cells in disease pathogenesis in the *scurfy* (*sf*) mouse. *J Immunol.* 1994;153:3764-3774.
- Salomon B, Bluestone JA. Complexities of CD28/B7: CTLA-4 costimulatory pathways in autoimmunity and transplantation. *Annu Rev Immunol.* 2001;19:225-252.
- Jenkins MK, Taylor PS, Norton SD, Urdahl KB. CD28 delivers a costimulatory signal involved in antigen-specific IL-2 production by human T cells. *J Immunol.* 1991;147:2461-2466.
- Lindstein T, June CH, Ledbetter JA, Stella G, Thompson CB. Regulation of lymphokine messenger RNA stability by a surface-mediated T cell activation pathway. *Science.* 1989;244:339-343.
- Fraser JD, Irving BA, Crabtree GR, Weiss A. Regulation of interleukin-2 gene enhancer activity by the T cell accessory molecule CD28. *Science.* 1991;251:313-316.
- Noel PJ, Boise LH, Green JM, Thompson CB. CD28 costimulation prevents cell death during primary T cell activation. *J Immunol.* 1996;157:636-642.
- Radvanyi LG, Shi Y, Vaziri H, et al. CD28 costimulation inhibits TCR-induced apoptosis during a primary T cell response. *J Immunol.* 1996;156:1788-1798.
- Boise LH, Minn AJ, Noel PJ, et al. CD28 costimulation can promote T cell survival by enhancing the expression of Bcl-XL. *Immunity.* 1995;3:87-98.
- Bromley SK, Iaboni A, Davis SJ, et al. The immunological synapse and CD28-CD80 interactions. *Nat Immunol.* 2001;2:1159-1166.
- Wetzel SA, McKeithan TW, Parker DC. Live-cell dynamics and the role of costimulation in immunological synapse formation. *J Immunol.* 2002;169:6092-6101.
- Kundig TM, Shahinian A, Kawai K, et al. Duration of TCR stimulation determines costimulatory requirement of T cells. *Immunity.* 1996;5:41-52.
- London CA, Lodge MP, Abbas AK. Functional responses and costimulator dependence of memory CD4+ T cells. *J Immunol.* 2000;164:265-272.
- McAdam AJ, Schweitzer AN, Sharpe AH. The role of B7 co-stimulation in activation and differentiation of CD4+ and CD8+ T cells. *Immunol Rev.* 1998;165:231-247.
- Van Gool SW, Vandenberghe P, de Boer M, Ceuppens JL. CD80, CD86 and CD40 provide accessory signals in a multiple-step T-cell activation model. *Immunol Rev.* 1996;153:47-83.
- Alegre ML, Noel PJ, Eisfelder BJ, et al. Regulation of surface and intracellular expression of CTLA4 on mouse T cells. *J Immunol.* 1996;157:4762-4770.
- Linsley PS, Brady W, Urnes M, Grosmaire LS, Damsle NK, Ledbetter JA. CTLA-4 is a second receptor for the B cell activation antigen B7. *J Exp Med.* 1991;174:561-569.

In addition, tissue infiltration by lymphocytes may play a critical role in the disease development. Unlike IL-4 and IFN- γ production, *scurfy* mice injected with CTLA4-Ig showed significant mononuclear cell infiltration in the lung and liver at day 48. The data indicate that CD28 is required for T cells to initiate tissue invasion. This mononuclear cell infiltration may be the direct effect of a CD28 signal to activated T cells or an indirect effect through up-regulation of cytokine/chemokine signaling and/or induction of other coactivators such as ICOS.

Together, the data demonstrate the significance of CD28 for critical development of autoimmune disorder in *scurfy* mice. Although the mechanisms underlying the autoimmune pathology caused by mutations of human and mouse Foxp3 may be different, intervention of CD28-B7 interaction may prove beneficial in treating patients with IPEX. Such possibilities need to be explored in future analysis of the human systems.

Acknowledgments

This study was supported in part by grants from NIH (AI055022 and AI049398) and JDRF (5-2006-388) (M.I.) and from the Arthritis Foundation (Y.S.).

We thank Dr Jeff Lee for advice, MCG Flow Cytometry Core and Histology Core for support, and Joyce Wilson and Doris McCool for assistance.

Authorship

Contribution: N.S., A.M., and M.I. designed the research; N.S., P.C., Y.S., B.B., M.T., and D.K. performed the experiments; N.S. and M.I. analyzed the data; D.H.M. and C.P.L. contributed vital reagents; and N.S. and M. I. wrote the paper.

Conflict-of-interest disclosure: The authors declare no competing financial interests.

Correspondence: Makio Iwashima, CN4155, Immunotherapy Center, MCG, 1120 15th St, Augusta, GA 30912; e-mail: miwashima@mcg.edu.

22. Bluestone JA, St Clair EW, Turka LA. CTLA4Ig: bridging the basic immunology with clinical application. *Immunity*. 2006;24:233-238.
23. Mandelbrot DA, McAdam AJ, Sharpe AH. B7-1 or B7-2 is required to produce the lymphoproliferative phenotype in mice lacking cytotoxic T lymphocyte-associated antigen 4 (CTLA-4). *J Exp Med*. 1999;189:435-440.
24. Mandelbrot DA, Oosterwegel MA, Shimizu K, et al. B7-dependent T-cell costimulation in mice lacking CD28 and CTLA4. *J Clin Invest*. 2001;107:881-887.
25. Read S, Greenwald R, Izcue A, et al. Blockade of CTLA-4 on CD4+CD25+ regulatory T cells abrogates their function in vivo. *J Immunol*. 2006;177:4376-4383.
26. Takahashi T, Tagami T, Yamazaki S, et al. Immunologic self-tolerance maintained by CD25(+)CD4(+) regulatory T cells constitutively expressing cytotoxic T lymphocyte-associated antigen 4. *J Exp Med*. 2000;192:303-310.
27. Karandikar NJ, Vanderlugt CL, Bluestone JA, Miller SD. Targeting the B7/CD28:CTLA-4 costimulatory system in CNS autoimmune disease. *J Neuroimmunol*. 1998;89:10-18.
28. Racke MK, Ratts RB, Arredondo L, Perrin PJ, Lovett-Racke A. The role of costimulation in autoimmune demyelination. *J Neuroimmunol*. 2000;107:205-215.
29. Anderson DE, Sharpe AH, Hafler DA. The B7-CD28/CTLA-4 costimulatory pathways in autoimmune disease of the central nervous system. *Curr Opin Immunol*. 1999;11:677-683.
30. Matsui Y, Inobe M, Okamoto H, et al. Blockade of T cell costimulatory signals using adenovirus vectors prevents both the induction and the progression of experimental autoimmune myocarditis. *J Mol Cell Cardiol*. 2002;34:279-295.
31. Tada Y, Nagasawa K, Ho A, et al. CD28-deficient mice are highly resistant to collagen-induced arthritis. *J Immunol*. 1999;162:203-208.
32. Peterson KE, Sharp GC, Tang H, Braley-Mullen H. B7.2 has opposing roles during the activation versus effector stages of experimental autoimmune thyroiditis. *J Immunol*. 1999;162:1859-1867.
33. Shi FD, He B, Li H, Matusevicius D, Link H, Ljunggren HG. Differential requirements for CD28 and CD40 ligand in the induction of experimental autoimmune myasthenia gravis. *Eur J Immunol*. 1998;28:3587-3593.
34. Finck BK, Linsley PS, Wofsy D. Treatment of murine lupus with CTLA4Ig. *Science*. 1994;265:1225-1227.
35. Nakajima A, Azuma M, Kodera S, et al. Preferential dependence of autoantibody production in murine lupus on CD86 costimulatory molecule. *Eur J Immunol*. 1995;25:3060-3069.
36. Koike T, Yamagishi H, Hatanaka Y, et al. A novel ERK-dependent signaling process that regulates interleukin-2 expression in a late phase of T cell activation. *J Biol Chem*. 2003;278:15685-15692.
37. Chang X, Gao JX, Jiang Q, et al. The *scurfy* mutation of FoxP3 in the thymus stroma leads to defective thymopoiesis. *J Exp Med*. 2005;202:1141-1151.
38. Dall'Era M, Davis J. CTLA4Ig: a novel inhibitor of costimulation. *Lupus*. 2004;13:372-376.
39. Min B, McHugh R, Sempowski GD, Mackall C, Foucras G, Paul WE. Neonates support lymphopenia-induced proliferation. *Immunity*. 2003;18:131-140.
40. Khoruts A, Fraser JM. A causal link between lymphopenia and autoimmunity. *Immunol Lett*. 2005;98:23-31.
41. Tivol EA, Boyd SD, McKeon S, et al. CTLA4Ig prevents lymphoproliferation and fatal multiorgan tissue destruction in CTLA-4-deficient mice. *J Immunol*. 1997;158:5091-5094.
42. Chambers CA, Sullivan TJ, Allison JP. Lymphoproliferation in CTLA-4-deficient mice is mediated by costimulation-dependent activation of CD4+ T cells. *Immunity*. 1997;7:885-895.
43. Waterhouse P, Penninger JM, Timms E, et al. Lymphoproliferative disorders with early lethality in mice deficient in Ctl4-4. *Science*. 1995;270:985-988.
44. Tang Q, Boden EK, Henriksen KJ, Bour-Jordan H, Bi M, Bluestone JA. Distinct roles of CTLA-4 and TGF-beta in CD4+CD25+ regulatory T cell function. *Eur J Immunol*. 2004;34:2996-3005.
45. Alegre ML, Fallarino F. Mechanisms of CTLA-4-Ig in tolerance induction. *Curr Pharm Des*. 2006;12:149-160.