

## B-lineage commitment prior to surface expression of B220 and CD19 on hematopoietic progenitor cells

Robert Mansson,<sup>1-3</sup> Sasan Zandi,<sup>1</sup> Kristina Anderson,<sup>2,3</sup> Inga-Lill Martensson,<sup>4</sup> Sten Eirik W. Jacobsen,<sup>2,3</sup> David Bryder,<sup>5</sup> and Mikael Sigvardsson<sup>1-3</sup>

<sup>1</sup>Department for Biomedicine and Surgery, Linköping University, Linköping, Sweden; <sup>2</sup>Hematopoietic Stem Cell Laboratory and <sup>3</sup>Lund Strategic Research Center for Stem Cell Biology and Cell Therapy, Lund University, Lund, Sweden; <sup>4</sup>Laboratory of Lymphocyte Signaling and Development, The Babraham Institute, Cambridge, United Kingdom; and <sup>5</sup>Department of Immunology, Lund University, Lund, Sweden

**Commitment of hematopoietic progenitor cells to B-lymphoid cell fate has been suggested to coincide with the development of PAX5-expressing B220<sup>+</sup>CD19<sup>+</sup> pro-B cells. We have used a transgenic reporter mouse, expressing human CD25 under the control of the B-lineage-restricted *Igll1* ( $\lambda$ 5) promoter to investi-**

**gate the lineage potential of early progenitor cells in the bone marrow. This strategy allowed us to identify a reporter-expressing LIN<sup>-</sup>B220<sup>-</sup>CD19<sup>-</sup>CD127<sup>+</sup>FLT3<sup>+</sup>SCA1<sup>low</sup>KIT<sup>low</sup> population that displays a lack of myeloid and a 90% reduction in in vitro T-cell potential compared with its reporter-negative counterpart. Gene ex-**

**pression analysis demonstrated that these lineage-restricted cells express B-lineage-associated genes to levels comparable with that observed in pro-B cells. These data suggest that B-lineage commitment can occur before the expression of B220 and CD19. (Blood. 2008;112:1048-1055)**

### Introduction

The development of mature blood cells can be viewed as a process where multipotent progenitors gradually lose lineage potential as they progress through maturation to become functional hematopoietic cells. One of the first steps in the progression of lymphoid differentiation involves the development of lineage marker (LIN)<sup>-</sup>negative SCA1<sup>+</sup>KIT<sup>+</sup>CD34<sup>+</sup>FLT3<sup>+</sup> lymphoid primed multipotent progenitor (LMPP) cells that display a reduced megakaryocyte and erythroid lineage potential but a preserved ability to differentiate into myeloid and lymphoid cells.<sup>1,2</sup> A subfraction of the LMPPs express a number of lymphoid-associated genes, including *Rag1* and *Rag2* as well as *Dntt* (*TdT*) and *CD127* (*IL7 $\alpha$* ),<sup>2</sup> and although the expression of a *Rag1* or *Rag2* reporter gene in early progenitors does not mark complete loss of myeloid lineage potential,<sup>3,4</sup> such cells are primed to lymphoid development and further differentiation into common lymphoid progenitors (CLPs).<sup>4</sup> The surface phenotype of the CLP largely resembles that of the LMPP,<sup>1</sup> however, with lower expression of SCA1 and KIT and robust expression of CD127.<sup>5</sup> CLPs have been shown to support B-, T-, and NK-cell development, although their myeloid potential is largely reduced compared with more immature cells.<sup>5-7</sup> The regulation of B-lymphoid fate has been suggested to depend on an interplay between transcription factors including PU.1 (*Sfpi1*), early B-cell factor 1 (EBF1), and E2A,<sup>6,8,9</sup> with extracellular signals such as those mediated by FLT3-ligand (FL), interleukin 7 (IL7),<sup>10</sup> and the ligands of the NOTCH signaling pathway.<sup>11,12</sup> However, even if these factors are able to prime cells for or against development along the B-lymphoid pathway, the definitive lock of lineage potential has been suggested to depend on the expression of the transcription factor PAX5. Loss-of-function models have shown that in the absence of PAX5, development along the B-lymphoid pathway is initiated, but although the cells express a

large number of B-lineage genes, such cells retain the ability to develop into alternative lymphoid or myeloid pathways.<sup>13-15</sup> Hence B-cell commitment may occur at the B220<sup>+</sup>CD19<sup>+</sup> stage, where robust expression of *Pax5* can be detected.<sup>3,16</sup> However, the finding that *Pax5* is a direct target for EBF1,<sup>17</sup> and vice versa,<sup>18</sup> complicates the understanding of commitment in normal B-cell development. In addition, the fact that *Ebf1* is expressed already at the CLP stage<sup>19,20</sup> suggests that the issue of B-lineage commitment needs further investigations.

To investigate the temporal regulation of B-cell development and the link between functional activation of early B-lineage genes and lineage commitment, we have used a transgenic mouse where the expression of a human CD25 (hCD25) reporter gene is controlled by the EBF1-regulated *Igll1* ( $\lambda$ 5) promoter.<sup>21,22</sup> Expression of this reporter could be detected on a fraction of the LIN<sup>-</sup>B220<sup>-</sup>CD19<sup>-</sup>CD127<sup>+</sup>FLT3<sup>+</sup>SCA1<sup>low</sup>KIT<sup>low</sup> cells, commonly referred to as CLPs, with a highly reduced ability to develop into T cells and an inability to develop into myeloid cells. This was accompanied by increased expression of B-lymphoid-associated genes including *Pax5*, leading us to suggest that B-lineage commitment can occur before surface expression of B220 and CD19.

### Methods

#### Quantitative RT-PCR

Quantitative reverse transcriptase-polymerase chain reaction (Q-RT-PCR) analysis of sorted cells was performed as previously described.<sup>1,2</sup> Assays-on-Demand probes (Applied Biosystems, Foster City, CA) used were as follows: *Cd19*; Mm00515420\_m1, *CD79a* (*Mb1*); Mm00432423\_m1,

Submitted November 20, 2007; accepted April 6, 2008. Prepublished online as *Blood* First Edition paper, May 21, 2008; DOI 10.1182/blood-2007-11-125385.

The online version of this article contains a data supplement.

The publication costs of this article were defrayed in part by page charge payment. Therefore, and solely to indicate this fact, this article is hereby marked "advertisement" in accordance with 18 USC section 1734.

© 2008 by The American Society of Hematology

*Pax5*; Mm00435501\_m1, *Hprt*; Mm00446968\_m1, *Ebf1*; and Mm00432948\_m1. Sequences for the Assay-by-Design probe used for  $\lambda 5$  (*Igll1*) detection were as follows: fw, 5'-GGAACAACAGGCCTAGC-TATGG; rev, 5'-CTCCCCGTGGGATGATCTG; and probe, 5'-CCG-GCAGCTCCTGTTC. All experiments were performed in triplicates and differences in cDNA input were compensated by normalizing against *Hprt* expression levels.

### Fluorescence-activated cell sorting of CLPs and B-lineage cells

Bone marrow (BM) and spleen cells were harvested from 10- to 15-week-old C57Bl/6 wild-type or heterozygous hCD25 ( $\lambda 5$ ) transgenic mice<sup>21</sup> (on C57Bl/6 background) and single-cell suspensions were prepared.

For CLP isolation, BM cells were subjected to magnetic-activated cell sorting (MACS) column (Miltenyi Biotec, Bergisch Gladbach, Germany) enrichment of KIT<sup>+</sup> cells using anti-CD117 immunomagnetic beads (Miltenyi Biotec). KIT<sup>+</sup> cells were subsequently stained with Fc-block (CD16/CD32, 2.4G2) followed by CD19 (6D5) fluorescein isothiocyanate (FITC), FLT3 (A2F10) phycoerythrin (PE), CD11b (M1/70) PE-cyanin-5 (PE-Cy5), GR1 (RB6-8C5) PE-Cy5, TER119 (Ter119) PE-Cy5, CD3 (145-2c11) PE-Cy5, B220 (RA3-6B2) PE-Cy7, hCD25 (BC96) allophycocyanin (APC), KIT (2B8) APC-Alexa750, SCA1 (D7) PacificBlue, IL7r (A7R34) biotin (visualized with streptavidin QDot605 or streptavidin QDot655), and propidium iodide (PI).

For isolation of B-lineage cells, BM/spleen cells were stained with purified TER119 (Ter119), GR1 (RB6-8C5), CD11c (M1/70) CD4 (GK1.5), and CD8a (53-6.7) (visualized with goat antirat QDot605). Cells were subsequently stained with Fc-block (CD16/CD32, 2.4G2) followed by IgM (R6-60.2) FITC, CD43 (S7) PE, B220 (RA3-6B2) APC, CD19 (1D3) PE-Cy7, AA4.1 (AA4.1) biotin (visualized with streptavidin QDot655), and PI.

Analysis and cell sorting was done on a BD FACSAria (BD Biosciences, San Jose, CA).

### Affymetrix gene expression and data analysis

RNA was extracted from purified adult BM subsets as described for Q-RT-PCR. RNA was labeled and amplified according to the Affymetrix GeneChip Expression Analysis Technical Manual.<sup>23</sup> Chips were scanned using a GeneChip Scanner 3000 (Affymetrix, Santa Clara, CA) and scaled to a median intensity of 100. RNA from cells sorted on separate occasions was separately hybridized for each investigated population. Probe level expression values were calculated using robust multichip average (RMA) and further analysis was done using dChip (Dana-Farber Cancer Institute and Harvard School of Public Health, <http://www.dchip.org>). Array data used are accessible through the Gene Expression Omnibus (GEO; GSE7302 and GSE11110).<sup>24</sup>

### In vitro evaluation of B- and T-lymphoid potential by OP9/OP9DL1 coculture

For evaluating B-cell and T-cell potential, cells were clone sorted (using a FACSAria) directly into 96-well plates containing preplated (2000 cells/well) OP9 (supplemented with 10 ng/mL FLT3L and IL7) and OP9DL1 (supplemented with 10 ng/mL FLT3L and IL7 or only 10 ng/mL FLT3L) stroma layers, respectively. Cultures were substituted with new cytokines every 7 days. OptiMEM supplemented with 10% fetal calf serum, 50  $\mu$ g/mL gentamicin, and 50  $\mu$ M  $\beta$ -mercaptoethanol was used for maintaining the OP9/OP9DL1 stroma cell lines as well as for cocultures.

Cocultures were evaluated by flow cytometry staining with CD19 (1D3) PE, B220 (RA3-6B2) APC, and PI for OP9 cocultures and with CD25 (7D4) FITC, CD19 (1D3) PE, CD90.2/Thy1.2 (53-2.1) APC, and PI for OP9DL1 cocultures.

OP9 and OP9DL1 cocultures of CLPs were in initial experiments evaluated at day 14 and day 21, respectively, and in later experiments at days 7 to 8 and days 14 to 15, respectively, with indistinguishable results. Samples were analyzed on a BD FACSCalibur (BD Biosciences).

### In vitro evaluation of myeloid potential

Myeloid potential was evaluated as previously described<sup>1,2</sup> but with a modified cytokine combination. In brief, 150 cells were sorted in 3 mL medium (OptiMEM supplemented with 10% fetal calf serum, 50  $\mu$ g/mL gentamicin, 50  $\mu$ M  $\beta$ -mercaptoethanol, 25 ng/mL KIT ligand [KL], 25 ng/mL fms-like tyrosine kinase 3 ligand [FLT3L], 5 ng/mL interleukin 3 [IL-3], 5 ng/mL colony-stimulating factor 1 [CSF1], 5 ng/mL colony stimulating factor 2 [CSF2], and 10 ng/mL colony-stimulating factor 3 [CSF3]) and 20  $\mu$ L was plated into each well of 2 60-well plates (Nunc Minitrays; Rochester, NY). Wells were scored after 6 days, with an inverted light microscope, for clonal growth and size of the clones. At least 600 cells per population were plated in 2 independent experiments and at least 16 clones picked randomly from each population for evaluation by RT-PCR. Nested RT-PCRs were essentially done as described for single-cell RT-PCRs (for primers used, see "Gene expression analysis of single cells by multiplex RT-PCR" and Mansson et al<sup>2</sup>). For morphologic evaluation, May-Grünwald Giemsa-stained cytospin preparations were made from 3 to 4 pooled clones to make 16 slides per population from 2 independent experiments.

### Gene expression analysis of single cells by multiplex RT-PCR


Multiplex single-cell RT-PCR analysis was performed as previously described.<sup>1,2,25</sup> Primers used were as follows: *Hprt*: (1) 5'-GGGGCTATA-AGTTCTTTGC; (2) 5'-GTTCTTTGCTGACCTGCTGG; (3) 5'-TGGGGCTGACTGCTTAACC; (4) 5'-TCCAACTTCGAGAGGTCC. *Rag1*: (1) 5'-CCAAGCTGCAGACATTCTAGCACTC; (2) 5'-CAGACAT-TCTAGCACTCTGG; (3) 5'-GCTTGACTTCCCATCAGCATGGA; (4) 5'-CAACATCTGCCTTCACGTCGATCC. *Il7r*: (1) 5'-CTCTCTCAGAAT-GATGGC; (2) 5'-TGACTCAGACATCAACACAGC; (3) 5'-AAGATA-CATCGTCCAGTTGC; (4) 5'-GTCGTAGTTTTCTCTGTGG. *Pax5*: (1) 5'-CTACAGGCTCCGTGACGCAG; (2) 5'-ATGGCCACTCACTTC-CGGGC; (3) 5'-GTCATCCAGGCCTCCAGCCA; (4) 5'-TCTCGGCCT-GTGACAATAGG. *Igll1* ( $\lambda 5$ ): (1) 5'-AGTTCTCTCCTGCTGCTGC; (2) 5'-GGGTCTAGTGGATGGTGTCC; (3) 5'-CAAACTGGGGCTTA-GATGG; (4) 5'-CCCACCACAAAGACATACC. *Ebf1*: (1) 5'-CCCTCT-TATCTGGAACATGC; (2) 5'-CTACTCCCTGTATCAAAGCC; (3) 5'-TGTACGACAGTGTGACTTCC; (4) 5'-TAAGGATCACTTCCTTTGGC. *Mb-1*: (1) 5'-CCTCCTTCTTGTGCATACG; (2) 5'-AAACAATGGCAG-GAACCC; (3) 5'-TGATGATGCGGTTCTTGG; (4) 5'-GAACAGTCAT-CAAGGTTCCAGG. Primers 1 plus 4 and 2 plus 3 are outer and inner primer pairs, respectively.

### Institutional review board approval

Animal procedures were performed with consent from the local ethics committee at Lund University (Lund, Sweden) and Linköping University (Linköping, Sweden).

## Results

Although CLPs have been reported to express significant amounts of B-lineage-associated transcripts,<sup>3,26</sup> it has not been reported that this is associated with B-cell commitment. Rather, such transcripts have been assigned to lineage priming, associated with the activation of chromatin, making progenitor cells permissive for development along the B-lymphoid pathway. However, single-cell PCR has suggested that approximately one-third of CLPs coexpress the B-lineage genes *Igll1* and *Pax5* without the expression of the T-lineage genes *Pta* or *CD3e*,<sup>26</sup> a finding that might suggest that the CD19<sup>+</sup>B220<sup>+</sup> CLP compartment contains a B cell-committed subpopulation. To investigate if the early lineage marker-negative compartment contains committed progenitors, we adopted an approach using an *Igll1* ( $\lambda 5$ ) promoter-controlled human CD25 reporter gene (hCD25)<sup>21</sup> to trace cells with an active B-lineage-



**Figure 1. The B220<sup>-</sup>CD19<sup>-</sup> compartment contains lineage-restricted B-cell progenitors marked by transgenic expression of hCD25.** (A) Representative FACS plots and purification scheme used to identify hCD25<sup>+</sup> and hCD25<sup>-</sup> LIN<sup>-</sup>B220<sup>-</sup>CD19<sup>-</sup>CD127<sup>+</sup>FLT3<sup>+</sup>SCA1<sup>low</sup>KIT<sup>low</sup> fractions. Numbers indicate percentage of gated cells. (B) The frequency (%) of cells composing colonies generated by coculture of single cells on OP9/OP9DL1 cells in the presence of indicated cytokines. The data are collected from 2 or more independent experiments. (C) FACS plots of the output from representative clones generated on OP9 (top) and OP9DL1 (bottom).

associated transcription program. We speculated that the use of this reporter mouse should allow us to identify cells that have initiated the expression of B-lineage-restricted genes, possibly reflecting lineage commitment. The fact that the *Igll1* promoter is a direct target for EBF1<sup>22</sup> and that functional expression of this transcription factor is crucial for the expression of the *Igll1*<sup>27</sup> gene make it useful as a reporter for functional EBF1 activity. As expected, the reporter gene was active in a majority of the B220<sup>+</sup>CD19<sup>+</sup> bone marrow cells (data not shown and Hu et al<sup>25</sup>), but we could also detect a minor fraction (~5%) of the LIN<sup>-</sup>B220<sup>-</sup>CD19<sup>-</sup>CD127<sup>+</sup>FLT3<sup>+</sup>SCA1<sup>low</sup>KIT<sup>low</sup> cells (often referred to as CLPs) that expressed hCD25 on the surface (Figure 1A). This would correspond to a population size comparable with that of the CD34<sup>-</sup>FLT3<sup>-</sup>LSK compartment<sup>1</sup> (in the range of 1/20 000 bone marrow cells). To verify that the reporter gene expression overlapped with that of the endogenous *Igll1* gene, we performed Q-PCR analysis of sorted CD25<sup>+</sup> and CD25<sup>-</sup> cells, revealing that all the detectable *Igll1* expression was found in the CD25<sup>+</sup> progenitor fraction, supporting the notion that the reporter gene provides correct information about expression of the endogenous gene (Figure 3C).

Knowing that the LIN<sup>-</sup>B220<sup>-</sup>CD19<sup>-</sup>CD127<sup>+</sup>FLT3<sup>+</sup>SCA1<sup>low</sup>KIT<sup>low</sup> fraction contains a subpopulation of cells with a functional transcription factor network that allows for the activation of the *Igll1* gene, we wanted to investigate whether the hCD25<sup>+</sup> progenitors were functionally different from the hCD25<sup>-</sup>LIN<sup>-</sup>B220<sup>-</sup>CD19<sup>-</sup>CD127<sup>+</sup>FLT3<sup>+</sup>SCA1<sup>low</sup>KIT<sup>low</sup> cells. To gain the highest possible resolution and study lineage potential at a single-cell level, we sorted hCD25<sup>+</sup> or hCD25<sup>-</sup> LIN<sup>-</sup>B220<sup>-</sup>CD19<sup>-</sup>CD127<sup>+</sup>FLT3<sup>+</sup>SCA1<sup>low</sup>KIT<sup>low</sup> single cells and cultured

them on OP9 stroma cells (Figure 1B). Of 294 hCD25<sup>+</sup> cells plated, 181 (62%) developed into CD19<sup>+</sup>B220<sup>+</sup> B-lineage progenitors (B cells, Figure 1B), whereas 3% of the wells contained a portion of cells with only B220 and no CD19 expression, indicating an incomplete differentiation process. Plating of hCD25<sup>-</sup> cells revealed that 18% of the 180 analyzed wells contained only B220<sup>+</sup>CD19<sup>+</sup> cells, whereas 20% contained a portion of cells lacking the expression of CD19, indicating that these cells develop with a slower kinetics than the hCD25<sup>+</sup> cells. To investigate this further, we seeded hCD25<sup>-</sup> or hCD25<sup>+</sup> LIN<sup>-</sup>B220<sup>-</sup>CD19<sup>-</sup>CD127<sup>+</sup>FLT3<sup>+</sup>SCA1<sup>low</sup>KIT<sup>low</sup> cells on OP9 feeders and investigated the cellular output after 3, 4, 5, 6, and 10 days after initiation of the cultures (Figure S1). At day 3, the cultures initiated with hCD25<sup>+</sup> cells contained more than 95% CD19<sup>+</sup> cells, whereas the cultures from the reporter-negative cells contained mainly CD19<sup>-</sup> cells at this time after seeding. The overall cell numbers were modestly different, but because the fraction of CD19<sup>+</sup> cells was approximately 5 times higher in cells from the hCD25<sup>+</sup> cells, these data support the idea that hCD25<sup>+</sup> cells are more potent as short term B-cell progenitors. This difference was somewhat reduced at later time points after seeding, and after 10 days of differentiation the contents of the cultures were comparable although the number of cells generated from hCD25<sup>-</sup> cells were higher (Figure S1). We also investigated the presence of hCD25<sup>-</sup> cells in these cultures, revealing that a low number of B220<sup>-</sup>hCD25<sup>-</sup> cells were generated from the hCD25<sup>-</sup> cells and the LMPPs (Figure S2). This rapid kinetics makes in vivo analysis complicated due to the short time given for expansion and the short time between radiation and analysis. However, to ensure that the

hCD25<sup>+</sup> progenitors were able to give rise to CD19<sup>+</sup> cells *in vivo*, we transplanted 1000 hCD25<sup>+</sup> or hCD25<sup>-</sup> progenitors into irradiated mice (800 rad). Analysis of spleen and bone marrow suggested that both populations gave rise to a comparable amount of CD19<sup>+</sup> cells 7 days after transplantation (Figure S3). The apparent lack of functional differences *in vivo* could reflect the *in vitro* finding that the difference in B-cell output is less apparent already after 6 days. We were also able to detect a low amount of lineage-negative cells selectively in the spleens of mice that received a transplant of hCD25<sup>-</sup> CLPs, indicating that residual progenitors were present at the time of analysis. However, we have been unable to investigate *in vivo* reconstitution at earlier time points, and we are therefore limited to the conclusion that the reporter-positive cells have *in vivo* B-cell potential. This suggests that the LIN<sup>-</sup>B220<sup>-</sup>CD19<sup>-</sup>CD127<sup>+</sup>FLT3<sup>+</sup>SCA1<sup>low</sup>KIT<sup>low</sup> (CLP) progenitor population contains a subpopulation of cells prone to development into B-lineage cells *in vitro*.

To investigate if this increased B-cell potential was associated with any changes in T-cell potentials, we sorted single cells on OP9DL1 cells known to be highly efficient to stimulate development of T-lineage cells (Figure 1B). Thirty-two percent of the hCD25<sup>+</sup> cells generated clones under these conditions, but these were composed mainly of B220<sup>+</sup>CD19<sup>+</sup> cells (26%), and only 4% of the 192 plated cells generated T-cell progenitors (CD25<sup>+</sup>CD90.2<sup>+</sup>; Figure 1B). In contrast 51% of the 192 seeded hCD25<sup>-</sup> cells gave rise to colonies composed of CD25<sup>+</sup>CD90.2<sup>+</sup> cells, and only 1% of the cells gave rise to B220<sup>+</sup>CD19<sup>+</sup> cells. The protocol used for OP9DL1 cocultures includes the addition of IL7, known to be a highly potent stimulator of B-lineage development. Thus, to investigate the role of IL7 in this apparent loss of T-lineage potential in hCD25<sup>+</sup> LIN<sup>-</sup>B220<sup>-</sup>CD19<sup>-</sup>CD127<sup>+</sup>FLT3<sup>+</sup>SCA1<sup>low</sup>KIT<sup>low</sup> cells, we incubated the OP9DL1 cocultures with FL alone (Figure 1B). Under these conditions, the cloning frequency dropped so that only 11% of the 213 plated hCD25<sup>+</sup> cells developed into detectable clones. Among these, 7% were composed of B-lineage and 3.5% of T-lineage cells. Of the 288 plated hCD25<sup>-</sup> cells, 31% developed into T-lineage cells, whereas only one clone contained B-lineage cells. These data suggest that expression of the *Igll1* promoter-controlled reporter is linked to a reduced ability of CD19<sup>-</sup>B220<sup>-</sup> cells to develop into T-lymphoid cells, supporting that B- versus T-lineage restriction may occur already at this lineage marker-negative stage.

#### Expression of hCD25 on LIN<sup>-</sup>B220<sup>-</sup>CD19<sup>-</sup>CD127<sup>+</sup>FLT3<sup>+</sup>SCA1<sup>low</sup>KIT<sup>low</sup> cells marks a loss of residual myeloid potential in early progenitors


Although the previously defined CLPs, which functionally have a pronounced preference to develop into lymphoid cells,<sup>5,7</sup> evidence suggesting that these cells maintain a myeloid potential<sup>3,28</sup> (which may be lost as late as after the B- versus T-lineage choice<sup>2</sup>) has been presented. Therefore, to investigate whether we could reveal a myeloid potential in LIN<sup>-</sup>B220<sup>-</sup>CD19<sup>-</sup>CD127<sup>+</sup>FLT3<sup>+</sup>SCA1<sup>low</sup>KIT<sup>low</sup> compartment of cells, we plated hCD25<sup>+</sup> and hCD25<sup>-</sup> LIN<sup>-</sup>B220<sup>-</sup>CD19<sup>-</sup>CD127<sup>+</sup>FLT3<sup>+</sup>SCA1<sup>low</sup>KIT<sup>low</sup> cells into 60-well plates and incubated them for 6 days in a cytokine cocktail (KL, FL, IL3, CSF1, CSF2, and CSF3) to promote the development of myeloid cells. As a control population, we sorted LMPPs and cultured them under the same conditions for 6 days. Analysis of the colonies generated suggested that although the general cloning frequency was comparable among the hCD25<sup>-</sup> and hCD25<sup>+</sup> populations, the hCD25<sup>-</sup> cells generated a few large

clones of cells resembling those generated by the LMPPs (Figure 2A). An even more striking difference was seen in the distribution of the cells in the wells and in the morphology of the cells. The clones generated from the hCD25<sup>+</sup> cells were small, focused at a small area of the well, and the cells were rounded and nonadherent (Figure 2B). In contrast, a majority of the hCD25<sup>-</sup> cells gave rise to colonies composed of a mix of cells that were spread over the well, attached to the plastic, and also in some cases appeared to have pseudopodias (Figure 2B). To investigate this further, we randomly pooled the content of 3 to 4 wells to perform cytopins and May-Grünwald Giemsa stainings. Investigating the morphology of the cells revealed that all 16 slides obtained from 2 independent experiments using hCD25<sup>+</sup> progenitors contained only rounded cells with a small cytoplasm, indicative that these cells are lymphoid related (Figure 2B). hCD25<sup>-</sup> cells did, however, generate a mix of cells on all of the 16 slides investigated. Among these were granular cells with large cytoplasm, a morphology clearly indicative of myeloid cells (Figure 2B). To obtain molecular support for the morphologic analysis, we picked cells from at least 16 randomly chosen clones and performed nested RT-PCR in search for myeloid and lymphoid transcripts in the generated cells (Figure 2C). Using cells generated from hCD25<sup>+</sup> LIN<sup>-</sup>B220<sup>-</sup>CD19<sup>-</sup>CD127<sup>+</sup>FLT3<sup>+</sup>SCA1<sup>low</sup>KIT<sup>low</sup> progenitors, 15 of 16 samples contained detectable amounts of both *Pax5* and *Igll1* transcripts, whereas none of the samples contained either *Mpo* or *Csf3r* message normally found in myeloid cells. This was in contrast to the cells obtained from the LIN<sup>-</sup>B220<sup>-</sup>CD19<sup>-</sup>CD127<sup>+</sup>FLT3<sup>+</sup>SCA1<sup>low</sup>KIT<sup>low</sup> hCD25<sup>-</sup> progenitors, because 12 of the 16 samples analyzed contained transcripts encoding either *Mpo*, *Csf3r*, or both. *Igll1* expression could be detected in 2 of the samples and *Pax5* in 1 sample. We were also able to detect *CD3e* expression, indicative of T-lineage cells, in 5 of these samples. As expected LMPPs gave rise mainly to myeloid cells and all 16 samples contained *Mpo* and *Csf3r* message, although lymphoid transcripts could be detected in some samples. These data indicate that although the hCD25<sup>-</sup> LIN<sup>-</sup>B220<sup>-</sup>CD19<sup>-</sup>CD127<sup>+</sup>FLT3<sup>+</sup>SCA1<sup>low</sup>KIT<sup>low</sup> cells had lost most of their potential to generate large number of myeloid cells, the conditions we provided allowed them to reveal a residual myeloid potential that was lost in the hCD25<sup>+</sup> cells. Thus, hCD25<sup>+</sup> LIN<sup>-</sup>B220<sup>-</sup>CD19<sup>-</sup>CD127<sup>+</sup>FLT3<sup>+</sup>SCA1<sup>low</sup>KIT<sup>low</sup> cells display a dramatically reduced ability to adopt other cell fates than B-lineage even under highly permissive conditions, supporting the idea that B-lineage commitment may occur before the expression of B220 and CD19.

#### Increased level of *Ebf1* expression is associated with the activation of a B-lymphoid regulatory program in CD19<sup>-</sup>B220<sup>-</sup> progenitor cells

Although our data strongly support the idea that the expression of the *Igll1*-controlled hCD25 reporter in early progenitors marks B-lineage commitment, we wanted to investigate molecular changes associated with this event. To this end, we performed microarray experiments to analyze gene expression patterns in hCD25<sup>+</sup> and hCD25<sup>-</sup> LIN<sup>-</sup>B220<sup>-</sup>CD19<sup>-</sup>CD127<sup>+</sup>FLT3<sup>+</sup>SCA1<sup>low</sup>KIT<sup>low</sup> cells. To put the data into context, expression patterns were compared with those in LMPPs, CD19<sup>+</sup>AA4.1<sup>+</sup>CD43<sup>low</sup> pro-B cells,<sup>29</sup> CD19<sup>+</sup>B220<sup>+</sup>CD43<sup>-</sup>IgM<sup>-</sup> pre-B cells, and IgM<sup>+</sup>CD19<sup>+</sup> mature spleen B cells. This revealed a significant difference in overall expression patterns: the hCD25<sup>+</sup> cells clustered with the committed B-lineage cells, whereas the hCD25<sup>-</sup> cells rather resembled the more immature LMPP cells (Figure 3A). This was a result mainly






**Figure 2. Activation of the *Igll1* promoter is associated with a reduced myeloid potential in progenitor cells.** (A) The cloning frequency and the size of the colonies obtained after culture of single cells in a mix of myeloid cytokines. Error bars represent SEM. (B) Colonies detected (50 $\times$  magnification) and the morphology of the cells generated during in vitro incubation with cytokines as visualized by May-Grünwald Giemsa staining (MGG; 500 $\times$  magnification). Cells displayed as one picture are collected from the same slide, but the images of individual cells have been moved to fit the picture format. Images of tissue culture plate wells were taken with an Olympus IX70 microscope (with a LCPlanF1 20 $\times$ /0.40 Ph1 objective; Olympus Optical, Tokyo, Japan) using a Nikon E4500 camera (Nikon Nordic AB, Solna, Sweden). Images of MGG-stained slides were acquired with an Olympus BX51 microscope (with a UPlanF1 100 $\times$ /0.30 oil objective; Olympus Optical) using an Olympus DP70 camera and acquisition software (DP Controller 1.1.1.65; Olympus, Tokyo, Japan). (C) The number of PCR-positive samples generated after analysis of 16 randomly chosen clones by multiplex RT-PCR. Each horizontal line of dots indicates the gene expression pattern observed in a single investigated clone. Data presented are collected from 2 independent experiments.

of a reduction in the expression of a large set of genes, including *Notch1*, comparably low in the hCD25<sup>+</sup> cells (Figure S4). hCD25<sup>+</sup> cells also expressed higher levels of *Igll1*, *Vpreb1*, *Vpreb3*, *Cd19*, *Cd79a*, *Cd79b*, and *Ebfl* (Figure 3B), all indicative of B-lineage cells. We could also see an increased expression of several B-lineage-associated transcription factors such as *Ocab*, *Lef1*, and interestingly *Foxo1*, possibly linking *Ebfl* directly into the network of Fox proteins that play an important role in early B-cell development.<sup>30</sup> To verify some of these data, we analyzed the expression of a set of these genes by Q-RT-PCR (Figure 3C) and compared the relative expression levels with those observed in IgM<sup>-</sup>B220<sup>+</sup>CD43<sup>low</sup>CD19<sup>+</sup> pro-B cells,<sup>29</sup> IgM<sup>-</sup>B220<sup>+</sup>CD43<sup>-</sup>CD19<sup>+</sup> pre-B cells, and IgM<sup>+</sup>CD19<sup>+</sup> splenic B cells. The hCD25<sup>+</sup> cells expressed 4 times as much *Ebfl* message as the hCD25<sup>-</sup> cells. This level of *Ebfl* expression was comparable with that observed in the pro-B cells and only 2-fold lower than in the pre-B cells. *Igll1* transcripts were not detectable in the hCD25<sup>-</sup> cells, whereas levels comparable with those in the pro-B cell could be found in the hCD25<sup>+</sup> cells, providing evidence that the reporter gene expression is perfectly correlated to the expression of the endogenous *Igll1* gene. In addition, both *Pax5* and *Mbl* message levels were increased to levels comparable with that in CD19<sup>+</sup>B220<sup>+</sup> pro-B cells in the hCD25<sup>+</sup> cells. These cells also expressed significant amounts of *Cd19* mRNA, and although this level was low compared to what was observed in the pro-B cells, it was higher than in the

hCD25<sup>-</sup> cells. LMPPs expressed low or undetectable levels of all the analyzed transcripts (Figure 3C). These data support the idea that the expression of the reporter gene allows us to identify two molecular and functionally distinct LIN<sup>-</sup>B220<sup>-</sup>CD19<sup>-</sup>CD127<sup>+</sup>FLT3<sup>+</sup>SCA1<sup>low</sup>KIT<sup>low</sup> cell populations in the mouse bone marrow.

To investigate the modulation in gene expression patterns at the single-cell level, we sorted hCD25<sup>-</sup> and hCD25<sup>+</sup> LIN<sup>-</sup>B220<sup>-</sup>CD19<sup>-</sup>CD127<sup>+</sup>FLT3<sup>+</sup>SCA1<sup>low</sup>KIT<sup>low</sup> cells and analyzed their gene expression patterns by multiplex RT-PCR analysis (Figure 3D). To put the data into context, we also sorted LIN<sup>-</sup>KIT<sup>+</sup>SCA1<sup>+</sup>FLT3<sup>+</sup> LMPPs, LIN<sup>-</sup>KIT<sup>Low</sup>SCA1<sup>low</sup>CD127<sup>+</sup>FLT3<sup>+</sup> CLPs, as well as CD19<sup>+</sup>AA4.1<sup>+</sup>CD43<sup>low</sup> pro-B cells. Among the LMPP cells for which we obtained HPRT signal, 4 (6%) contained detectable levels of *Ebfl* transcripts, whereas 3 (4%) expressed *Il7r $\alpha$*  (CD127) transcripts. Seven (10%) expressed *Rag1* transcripts, whereas none of the cells contained detectable levels of *Pax5*, *Igll1*, or *Mbl* (CD79 $\alpha$ ) mRNA. In the CLP compartment, we were able to detect *Rag1* expression in 48 (55%) of the 87 analyzed cells and *Ebfl* expression in more than 51% of the analyzed cells, whereas *Pax5*, *Igll1*, and *Mbl* transcripts were found in 14%, 9%, and 17% of the cells, respectively. Notably, the majority of cells expressed *Pax5*, *Igll1*, and *Mbl* in a coordinated manner, supporting the idea that the lineage program is initiated already at the level of the currently defined CLP. A similar gene expression pattern was



**Figure 3. Expression of the *Igll1* promoter-controlled hCD25-transgenic marker in LIN<sup>-</sup> progenitors is associated with induced expression of genes linked to B-cell development.** (A) A diagram of the relative relationship between the hCD25<sup>+</sup> and hCD25<sup>-</sup> LIN-B220<sup>-</sup>CD19<sup>-</sup>CD127<sup>+</sup>FLT3<sup>+</sup>SCA1<sup>low</sup>KIT<sup>low</sup> progenitors to LMPs and B-lineage cells. The diagram is based on genes differentially expressed between the hCD25<sup>+</sup> and hCD25<sup>-</sup> cells. The complete analysis is presented in Figure S4 (available on the *Blood* website; see the Supplemental Materials link at the top of the online article). (B) A dChip analysis of hCD25<sup>+</sup> and hCD25<sup>-</sup> LIN-B220<sup>-</sup>CD19<sup>-</sup>CD127<sup>+</sup>FLT3<sup>+</sup>SCA1<sup>low</sup>KIT<sup>low</sup> cells in the context of LMPs and other stages of B-cell development. Clustering shows genes being expressed in either pro-B cells or LMPs (100 + expression units in either) and being up-regulated 1.5-fold from hCD25<sup>-</sup> to hCD25<sup>+</sup>. Red represents high; white, intermediate; and blue, low expression of the gene indicated to the far right. Superimposed values show RMA-modeled array expression values. (C) Q-RT-PCR data from CD25<sup>+</sup> and CD25<sup>-</sup> cells as well as control cell populations as indicated (data from one representative experiment). The error bars indicate standard deviation. (D) The collected result of multiplex single-cell PCR on LMPs, CLPs, pro-B cells, and hCD25<sup>+/+</sup> LIN-B220<sup>-</sup>CD19<sup>-</sup>CD127<sup>+</sup>FLT3<sup>+</sup>SCA1<sup>low</sup>KIT<sup>low</sup> cells as indicated. Each horizontal line of dots indicates the gene expression pattern observed in a single investigated cell. The values on top of the data panel indicate the number of positive cells and the values below the panels indicate the total number of cells analyzed. Error bars indicate standard deviation.

seen upon analysis of hCD25<sup>-</sup> LIN-B220<sup>-</sup>CD19<sup>-</sup>CD127<sup>+</sup>FLT3<sup>+</sup>SCA1<sup>low</sup>KIT<sup>low</sup> cells. Of 140 analyzed cells, 85 (61%) expressed *Ebf1* message, whereas in 99 cells (71%) we were able to detect *Il7ra*, and in 60 (43%) we detected *Rag1* transcripts. Six cells (4%) expressed *Igll1*; 5 (4%), *Pax5*; and 19 (14%), *Mb1*. In the hCD25<sup>+</sup> LIN-B220<sup>-</sup>CD19<sup>-</sup>CD127<sup>+</sup>FLT3<sup>+</sup>SCA1<sup>low</sup>KIT<sup>low</sup> cells, a similar pattern was seen when analyzing the expression of *Il7ra* (60%) and *Rag1* (61%) in the 145 investigated cells. A minor difference could be observed in the percentage of *Ebf1*-positive (90%) and *Mb1*-positive (26%) cells, whereas the number of *Igll1* and *Pax5*-positive cells increased to 57% and 52%, respectively. This was more comparable with the data obtained using the pro-B cells: 88% expressed detectable levels of *Ebf1* message, whereas 55 (66%)

of 93 contained *Pax5* mRNA. *Igll1* was detected in 92% of the cells; *Mb1*, in 68%; and *Rag1*, in 34% of the pro-B cells. These data support the idea that B-cell commitment occurs already in the phenotypically defined CLP compartment and indicate that the expression of *Ebf1* is a precommitment event, not directly linked to the expression of target genes or lineage restriction. Furthermore, because the expression level is increased 4-fold in the hCD25<sup>+</sup> progenitors compared with the hCD25<sup>-</sup> fraction, although the number of cells with detectable expression is increased only from 68% to 92%, this taken together suggests an increase of *Ebf1* mRNA at the single-cell level. Thus, investigations of transcriptional programs at a single-cell level support the idea that B-lineage fate can be determined before the expression of B220 and KIT19.

## Discussion

We here report data suggesting that B-lineage commitment of hematopoietic progenitors can occur at an early progenitor stage before the expression of B220 and CD19 in LIN<sup>-</sup>CD127<sup>+</sup>FLT3<sup>+</sup>SCA1<sup>low</sup>KIT<sup>low</sup>hCD25<sup>+</sup> B-lymphoid committed progenitor (BLCP). These finding does not support the currently dominating view that commitment occurs at a later stage of development with surface expression of both CD19 and B220. The idea that CD19 expression would coincide with commitment is to a substantial degree based on data from *Pax5*-deficient mice or cell lines where robust expression of *Pax5* has been shown to result in expression of *Pax5* target genes such as *CD19* and repression of alternative cell fates.<sup>13-16</sup> This view is also supported by studies in wild-type mice where CD19<sup>-</sup>B220<sup>-</sup> CLPs have been reported to give rise to CD11<sup>+</sup> myeloid cells as well as lymphoid cells, whereas B220<sup>+</sup> cells maintain their T-cell potential and only B220<sup>+</sup>CD19<sup>+</sup> cells appear to have undergone lineage commitment.<sup>3</sup> However, our experiments using a transgenic reporter provide a possibility to investigate these phenomena with a higher resolution. Our data do not contradict the idea that PAX5 is involved in lineage commitment because *Pax5* expression is increased in the BLCP. However, they add new information concerning the temporal regulation of events suggested from these previous observations. Several transgenic mice models have been used to identify lineage-committed or -restricted progenitors in the early compartments, and although the use of *Rag1* reporter mice<sup>4</sup> or staining for TDT<sup>31</sup> has allowed for a dramatic enrichment of cells primed for lymphoid development, the cells committed to B-lymphoid fate have been identified mainly in the lineage marker-positive fractions in the bone marrow.<sup>3</sup> Experiments using a hCD25 reporter under the regulation of the T-lineage-restricted *pTα* promoter have allowed for the identification of lineage-restricted T-cell progenitors in the blood,<sup>32</sup> whereas reporter-positive cells in the bone marrow appeared to retain a substantial B-lineage potential.<sup>33</sup> To our knowledge, there are no previous reports of a committed B220<sup>-</sup>CD19<sup>-</sup> cell, however, our data are supported by observations in other transgenic models. First, *Rag* expression and DJ recombination of the immunoglobulin heavy chain locus are dramatically impaired in CLPs from mice lacking the B-lineage-restricted E-Rag enhancer.<sup>34,35</sup> This suggests that B-lineage-restricted control elements may be active already in the CLP. Second, mice carrying a GFP reporter gene under the control of the *Pax5* locus present a population of GFP<sup>+</sup> cells within the LIN<sup>-</sup>CD127<sup>+</sup>SCA1<sup>low</sup>KIT<sup>low</sup> compartment.<sup>16</sup> The lineage potential of these cells was not investigated in the published report, but the size of this population is comparable with that of the hCD25<sup>+</sup> progenitor population and may well represent the same cells. Furthermore, multiplex single-cell PCR analysis of sorted CLPs suggested that approximately one third of the cells contained only B-lineage-associated transcripts.<sup>26</sup> Thus, we believe that our data find support in the existing literature.

In addition to an increased insight into the temporal regulation of B-lineage commitment, our results allow for alternative models for the molecular regulation of lineage choices in the early progenitors. Single-cell PCR analysis suggests that, as predicted from transgenic models, the transcription of lymphoid-associated genes such as *CD127* or *Rag1* is not linked to B- versus T-lineage decision event.<sup>4</sup> This also appears to be true for the expression of *Ebfl* because the number of progenitors that express EBF1 in hCD25<sup>+</sup> and hCD25<sup>-</sup> cells are comparable. However, in hCD25<sup>+</sup>

cells, the level of *Ebfl* message is increased 4-fold and the expression of EBF1 target genes is induced in a coordinated manner. The extremely low transcript levels (Figure 3C) and the low number of cells with detectable levels of transcripts from lineage-restricted genes in the noncommitted progenitors (Figure 3D) argue against a lineage priming model. Rather, our data suggest that the transcripts found in the CLP compartment are generated from B-lineage-restricted cells. This is also supported by the finding that the levels of *Ebfl*, *Pax5*, and *Mb1* mRNA found in the BLCPs are fully comparable with those found in IgM<sup>-</sup>B220<sup>+</sup>CD43<sup>low</sup>CD19<sup>+</sup> pro-B cells. The finding that a reporter gene controlled by the *Igll1* promoter—a direct target gene for EBF1—allows for tracing of B-lineage commitment points to a direct role of functional EBF1 activity in lineage restriction events, as previously suggested from overexpression studies.<sup>36</sup> The finding that *Ebfl* message can be detected in the majority (61%) of the CD127<sup>+</sup>FLT3<sup>+</sup>SCA1<sup>low</sup>KIT<sup>low</sup> hCD25<sup>-</sup> cells able to efficiently generate T-cell clones, and before the detectable expression of either *Pax5* or *Igll1*, suggests that the activation of the *Ebfl* gene is a precommitment event but that further increased levels of *Ebfl* have a striking effect on target gene activation and lineage commitment. This is also in line with the finding that *Pax5*-deficient pro-B cells could be made B-lineage restricted by ectopic expression of EBF, suggesting that the key role of PAX5 in lineage commitment is to enhance EBF expression.<sup>37</sup> Although EBF1 has been proposed to act in a dose-dependent manner<sup>17</sup> and the effect observed may be a direct response to increased level of EBF1, other posttranscriptional regulatory mechanisms including reduction of EBF1 activity by active NOTCH signaling<sup>38</sup> might be involved.

Although our data do not directly contradict the idea that PAX5 is the crucial factor in B-lineage commitment, the low level of CD19 transcripts in the BLCPs argues against the idea that these cells have established a fully developed PAX5 regulatory network. We rather believe that our findings fit a model where lineage restriction initially is mediated by EBF1, whereas the role of PAX5 is to ensure the establishment of a regulatory circuit involving the control of EBF1 expression levels in the committed B-lineage cells.<sup>37</sup> Although we currently are unable to say how large a part of the B cells that develop through a BLCP pathway, as opposed to those that undergo commitment at a CD19<sup>+</sup>B220<sup>+</sup> stage,<sup>3</sup> our findings suggest that this process can occur in lineage-negative cells—presenting new possibilities to study early events in B-cell development.

## Acknowledgments

The authors thank A. Cumano and J. C. Zuniga-Pflucker for providing the OP9 and OP9DL1 stromal cell lines; Liselotte Lenner and Gerd Sten for expert advice and technical assistance; and the SweGene Affymetrix facility for its assistance with the microarrays.

These studies were generously supported by grants from the Swedish Cancer Society, the Swedish Research Council, Tobias Stiftelsen, the Swedish Childhood Cancer Foundation, and Swedish Foundation for Strategic Research (all Stockholm, Sweden). The Lund Stem Cell Center is supported by a Center of Excellence grant from the Swedish Foundation for Strategic Research.



## Authorship

Contribution: R.M. and S.Z. performed the majority of the *in vitro* differentiation experiments and single-cell PCR; D.B. performed cell sorting experiments; K.A. performed and analyzed Giemsa stainings; M.S. performed Q-PCR experiments; R.M., S.Z., D.B., S.E.W.J., and M.S. were involved in the design

of the experiments; I.-L.M. provided crucial reagents; all authors have read and given comments to the paper written by R.M. and M.S.

Conflict-of-interest disclosure: The authors declare no competing financial interests.

Correspondence: Mikael Sigvardsson, Faculty of Health Sciences, Linköping University Lab1, Level 13, Linköping, Sweden; e-mail: mikael.sigvardsson@ibk.liu.se.

## References

- Adolfsson J, Mansson R, Buza-Vidas N, et al. Identification of Flt3+ lympho-myeloid stem cells lacking erythro-megakaryocytic potential a revised road map for adult blood lineage commitment. *Cell*. 2005;121:295-306.
- Mansson R, Hultquist A, Luc S, et al. Molecular evidence for hierarchical transcriptional lineage priming in fetal and adult stem cells and multipotent progenitors. *Immunity*. 2007;26:407-419.
- Rumfelt LL, Zhou Y, Rowley BM, Shinton SA, Hardy RR. Lineage specification and plasticity in CD19- early B cell precursors. *J Exp Med*. 2006;203:675-687.
- Igarashi H, Gregory SC, Yokota T, Sakaguchi N, Kincade PW. Transcription from the RAG1 locus marks the earliest lymphocyte progenitors in bone marrow. *Immunity*. 2002;17:117-130.
- Kondo M, Weissman IL, Akashi K. Identification of clonogenic common lymphoid progenitors in mouse bone marrow. *Cell*. 1997;91:661-672.
- DeKoter RP, Singh H. Regulation of B lymphocyte and macrophage development by graded expression of PU.1. *Science*. 2000;288:1439-1441.
- Kondo M, Scherer DC, Miyamoto T, et al. Cell-fate conversion of lymphoid-committed progenitors by instructive actions of cytokines. *Nature*. 2000;407:383-386.
- Medina KL, Pongubala JM, Reddy KL, et al. Assembling a gene regulatory network for specification of the B cell fate. *Dev Cell*. 2004;7:607-617.
- Seet CS, Brumbaugh RL, Kee BL. Early B cell factor promotes B lymphopoiesis with reduced interleukin 7 responsiveness in the absence of E2A. *J Exp Med*. 2004;199:1689-1700.
- Sitnicka E, Brakebusch C, Martensson IL, et al. Complementary signaling through flt3 and interleukin-7 receptor alpha is indispensable for fetal and adult B cell genesis. *J Exp Med*. 2003;198:1495-1506.
- Pui JC, Allman D, Xu L, et al. Notch1 expression in early lymphopoiesis influences B versus T lineage determination. *Immunity*. 1999;11:299-308.
- Radtke F, Wilson A, Stark G, et al. Deficient T cell fate specification in mice with an induced inactivation of Notch1. *Immunity*. 1999;10:547-558.
- Nutt SL, Heavey B, Rolink AG, Busslinger M. Commitment to the B-lymphoid lineage depends on the transcription factor Pax5 [see comments]. *Nature*. 1999;401:556-562.
- Rolink AG, Schaniel C, Bruno L, Melchers F. *In vitro* and *in vivo* plasticity of Pax5-deficient pre-B1 cells. *Immunol Lett*. 2002;82:35-40.
- Rolink AG, Nutt SL, Melchers F, Busslinger M. Long-term *in vivo* reconstitution of T-cell development by Pax5-deficient B-cell progenitors [see comments]. *Nature*. 1999;401:603-606.
- Fuxa M, Busslinger M. Reporter gene insertions reveal a strictly B lymphoid-specific expression pattern of Pax-5 in support of its B cell identity function. *J Immunol*. 2007;178:3031-3037.
- O'Riordan M, Grosschedl R. Coordinate regulation of B cell differentiation by the transcription factors EBF and E2A [in process citation]. *Immunity*. 1999;11:21-31.
- Roessler S, Gyory I, Imhof S, et al. Distinct promoters mediate the regulation of Ebf1 gene expression by interleukin-7 and Pax5. *Mol Cell Biol*. 2007;27:579-594.
- Kikuchi K, Lai AY, Hsu CL, Kondo M. IL-7 receptor signaling is necessary for stage transition in adult B cell development through up-regulation of EBF. *J Exp Med*. 2005;201:1197-1203.
- Dias S, Silva H Jr, Cumano A, Vieira P. Interleukin-7 is necessary to maintain the B cell potential in common lymphoid progenitors. *J Exp Med*. 2005;201:971-979.
- Martensson IL, Melchers F, Winkler TH. A transgenic marker for mouse B lymphoid precursors. *J Exp Med*. 1997;185:653-661.
- Sigvardsson M, O'Riordan M, Grosschedl R. EBF and E47 collaborate to induce expression of the endogenous immunoglobulin surrogate light chain genes. *Immunity*. 1997;7:25-36.
- Affymetrix. GeneChip Expression Analysis Technical Manual (version 701021 rev. 5). [http://www.affymetrix.com/support/downloads/manuals/expression\\_print\\_manual.zip](http://www.affymetrix.com/support/downloads/manuals/expression_print_manual.zip). Accessed March 1, 2007.
- National Center for Biotechnology Information. Gene Expression Omnibus. <http://www.ncbi.nlm.nih.gov/geo/>. Accessed March 9, 2008.
- Hu M, Krause D, Greaves M, et al. Multilineage gene expression precedes commitment in the hemopoietic system. *Genes Dev*. 1997;11:774-785.
- Miyamoto T, Iwasaki H, Reizis B, et al. Myeloid or lymphoid promiscuity as a critical step in hematopoietic lineage commitment. *Dev Cell*. 2002;3:137-147.
- Lin H, Grosschedl R. Failure of B-cell differentiation in mice lacking the transcription factor EBF. *Nature*. 1995;376:263-267.
- Balciunaite G, Ceredig R, Massa S, Rolink AG. A B220+ CD117+ CD19- hematopoietic progenitor with potent lymphoid and myeloid developmental potential. *Eur J Immunol*. 2005;35:2019-2030.
- Li YS, Wasserman R, Hayakawa K, Hardy RR. Identification of the earliest B lineage stage in mouse bone marrow. *Immunity*. 1996;5:527-535.
- Hu H, Wang B, Borde M, et al. Foxp1 is an essential transcriptional regulator of B cell development. *Nat Immunol*. 2006;7:819-826.
- Tudor KS, Payne KJ, Yamashita Y, Kincade PW. Functional assessment of precursors from murine bone marrow suggests a sequence of early B lineage differentiation events. *Immunity*. 2000;12:335-345.
- Krueger A, von Boehmer H. Identification of a T lineage-committed progenitor in adult blood. *Immunity*. 2007;26:105-116.
- Martin CH, Aifantis I, Scimone ML, et al. Efficient thymic immigration of B220+ lymphoid-restricted bone marrow cells with T precursor potential. *Nat Immunol*. 2003;4:866-873.
- Borghesi L, Hsu LY, Miller JP, et al. B lineage-specific regulation of V(D)J recombinase activity is established in common lymphoid progenitors. *J Exp Med*. 2004;199:491-502.
- Hsu LY, Lauring J, Liang HE, et al. A conserved transcriptional enhancer regulates RAG gene expression in developing B cells. *Immunity*. 2003;19:105-117.
- Zhang Z, Cotta CV, Stephan RP, deGuzman CG, Klug CA. Enforced expression of EBF in hematopoietic stem cells restricts lymphopoiesis to the B cell lineage. *EMBO J*. 2003;22:4759-4769.
- Pongubala JM, Northrup DL, Lancki DW, et al. Transcription factor EBF restricts alternative lineage options and promotes B cell fate commitment independently of Pax5. *Nat Immunol*. 2008;9:203-215.
- Smith EM, Akerblad P, Kadesch T, Axelson H, Sigvardsson M. Inhibition of EBF function by active Notch signaling reveals a novel regulatory pathway in early B-cell development. *Blood*. 2005;106:1995-2001.