

# Iron regulates phosphorylation of Smad1/5/8 and gene expression of *Bmp6*, *Smad7*, *Id1*, and *Atoh8* in the mouse liver

\*Léon Kautz,<sup>1,2</sup> \*Delphine Meynard,<sup>1,2</sup> Annabelle Monnier,<sup>3,4</sup> Valérie Darnaud,<sup>1,2</sup> Régis Bouvet,<sup>5</sup> Rui-Hong Wang,<sup>6</sup> Chiuxia Deng,<sup>6</sup> Sophie Vaulont,<sup>7,8</sup> Jean Mosser,<sup>3,5,9</sup> †Hélène Coppin,<sup>1,2</sup> and †Marie-Paule Roth<sup>1,2</sup>

<sup>1</sup>Institut National de la Santé et de la Recherche Médicale (Inserm), U563, Centre de Physiopathologie de Toulouse Purpan, Toulouse, France; <sup>2</sup>Université Toulouse III Paul-Sabatier, Institut Fédératif de Recherche 30 (IFR 30), Toulouse, France; <sup>3</sup>Centre National de la Recherche Scientifique (CNRS), Unité Mixte de Recherche 6061 (UMR6061), Institut de Génétique et Développement, and <sup>4</sup>CNRS (UMR6553), Ecobio, Université de Rennes 1, Rennes, France; <sup>5</sup>Service de Génétique Médicale, Centre Hospitalier Universitaire (CHU) de Rennes, Rennes, France; <sup>6</sup>Genetics of Development and Disease Branch, National Institute of Diabetes and Digestive and Kidney Diseases, National Institutes of Health, Bethesda, MD; <sup>7</sup>Institut Cochin, Université Paris Descartes, CNRS (UMR 8104), Paris, France; <sup>8</sup>Inserm, U567, Paris, France; and <sup>9</sup>Plate-forme transcriptome OUEST-genopole, Rennes, France

Although hepcidin expression was shown to be induced by the BMP/Smad signaling pathway, it is not yet known how iron regulates this pathway and what its exact molecular targets are. We therefore assessed genome-wide liver transcription profiles of mice of 2 genetic backgrounds fed iron-deficient, -balanced, or -enriched diets. Among 1419 transcripts significantly modulated by the dietary iron content, 4 were regulated similarly to the hepcidin genes *Hamp1* and *Hamp2*. They

are coding for *Bmp6*, *Smad7*, *Id1*, and *Atoh8* all related to the Bmp/Smad pathway. As shown by Western blot analysis, variations in *Bmp6* expression induced by the diet iron content have for functional consequence similar changes in Smad1/5/8 phosphorylation that leads to formation of heteromeric complexes with Smad4 and their translocation to the nucleus. Gene expression variations induced by secondary iron deficiency or iron overload were compared with those

consecutive to *Smad4* and *Hamp1* deficiency. Iron overload developed by *Smad4*- and *Hamp1*-deficient mice also increased *Bmp6* transcription. However, as shown by analysis of mice with liver-specific disruption of *Smad4*, activation of *Smad7*, *Id1*, and *Atoh8* transcription by iron requires Smad4. This study points out molecules that appear to play a critical role in the control of systemic iron balance. (Blood. 2008;112:1503-1509)

## Introduction

Iron, by virtue of its ability to accept or donate electrons, is essential for many of the biologic reactions carried out by living systems. This same characteristic, however, allows free iron in solution to form highly reactive oxygen species that can oxidize lipids, DNA, and proteins, and lead to cell damage. As there is no physiological means for excreting iron in mammals, systemic iron homeostasis must be maintained by tight regulation of intestinal iron absorption, as well as macrophage and hepatocyte iron release.

Hepcidin has a key role in coordinating the use and storage of iron with iron acquisition.<sup>1</sup> It acts by binding to ferroportin, an iron exporter present on the surface of enterocytes, macrophages, and hepatocytes, and induces its internalization and lysosomal degradation.<sup>2</sup> The loss of ferroportin from the cell surface prevents iron efflux from intestinal enterocytes, recycling of iron from senescent erythrocytes by macrophages, and iron mobilization from hepatic stores. Hepcidin production is inhibited by iron deficiency anemia and tissue hypoxia,<sup>3</sup> which increases iron availability for use in erythropoiesis. By contrast, hepcidin expression is enhanced by dietary or parenteral iron loading,<sup>4</sup> thus providing a feedback mechanism to limit intestinal iron absorption. Individuals with mutations in any of the hepcidin (*HAMP*),<sup>5</sup> hemojuvelin (*HJV*),<sup>6</sup> transferrin receptor 2 (*TFR2*),<sup>7</sup> or *HFE*<sup>8</sup> genes have low hepcidin

levels despite excessive iron stores. As a consequence, they are unable to effectively repress iron absorption.

Recently, Babitt et al<sup>9,10</sup> demonstrated that hepcidin expression was induced by the bone morphogenetic protein (BMP) signaling pathway. This pathway involves members of the transforming growth factor beta (TGF- $\beta$ ) superfamily of ligands that act by binding 2 type I and 2 type II BMP receptors (BMPR-I and BMPR-II). This induces the phosphorylation of BMPR-I by BMPR-II and the activated complex, in turn, phosphorylates a subset of Smad proteins (Smad1, Smad5, and Smad8). The receptor-activated Smads then form heteromeric complexes with the common mediator Smad4, and these translocate to the nucleus where they regulate the transcription of specific targets.<sup>11</sup> Hemojuvelin (HJV) was shown to act as a BMP coreceptor in vitro and to facilitate the activation of the BMPR-I/II complex.<sup>9</sup> Interestingly, stimulation of BMP signaling by HJV is paralleled by an increase in *HAMP* expression. Support of the involvement of the BMP signaling pathway in the regulation of hepcidin also comes from the observation that liver-specific disruption of the *Smad4* gene in mice leads to reduced hepcidin expression and systemic iron loading.<sup>12</sup> These mice are unable to synthesize hepcidin in response to inflammatory stimuli or to iron load. The BMP/Smad4 pathway therefore is critical to hepcidin expression.

Submitted March 6, 2008; accepted May 19, 2008. Prepublished online as Blood First Edition paper, June 6, 2008; DOI 10.1182/blood-2008-03-143354.

\*L.K. and D.M. contributed equally to this work

†H.C. and M.-P.R. contributed equally to this work

The online version of this article contains a data supplement.

The publication costs of this article were defrayed in part by page charge payment. Therefore, and solely to indicate this fact, this article is hereby marked "advertisement" in accordance with 18 USC section 1734.

There are, however, many open questions. First, there are several BMP molecules that can bind HJV in vitro and regulate hepcidin expression,<sup>10,13</sup> but further work is needed to determine which of these BMP molecules is the endogenous regulator in vivo. Second, it is not yet known how iron regulates hepcidin. In particular, whether iron is sensed by molecules forming the hepcidin-regulating BMP/HJV/BMPR complex remains to be elucidated. Third, it is still not known whether hepcidin is a direct or indirect target for BMP and the activated Smad/Smad4 complex. To decipher these important issues, we assessed the liver transcription profiles of 2 strains of wild-type mice fed an iron-deficient or an iron-enriched diet and identified 4 genes that were regulated similarly to hepcidin in that context. Analysis of their expression in *Smad4*- or *Hamp1*-deficient mice showed that they were all participating in the Bmp/Smad signaling pathway.

## Methods

### Wild-type mice fed diets with different iron contents

Male mice of the C57BL/6 (B6) and DBA/2 (D2) backgrounds were purchased from the Center d'Élevage Robert Janvier (Le Genest St-Isle, France) and maintained at the IFR 30 animal facility. Dietary iron deficiency was induced by feeding 5 4-week-old-mice of each strain a diet with virtually no iron content (< 3 mg iron/kg; SAFE, Augy, France) and demineralized water for a period of 3 weeks. Five control animals of each strain received the iron-balanced diet of the same composition (200 mg iron/kg). Dietary iron loading was obtained by feeding five 4-week-old animals of each strain the iron-balanced diet supplemented with carbonyl iron (8.3 g/kg) for 3 weeks. All mice were analyzed at 7 weeks and fasted for 14 hours before they were killed. Experimental protocols were approved by the Midi-Pyrénées Animal Ethics Committee. A total of 30 livers were dissected for RNA isolation, rapidly frozen, stored in liquid nitrogen, and used for genome-wide expression profiling. Serum iron parameters and nonheme iron were quantified as previously described.<sup>14</sup>

### *Smad4*- and *Hamp1*-deficient mice

Liver samples from six 4-month-old males with liver-specific disruption of *Smad4* and 5 10-week-old *Hamp1*-deficient males that were subjected to detailed analysis in previous publications<sup>12,15</sup> were used here to compare gene expression variations consecutive to lack of functional *Smad4* or *Hamp1* with those induced by secondary iron deficiency or iron overload. Liver samples from males of the same genetic backgrounds and the same ages as *Smad4*- and *Hamp1*-deficient mice, respectively, were used as controls.

### RNA isolation, preparation of labeled cRNA, microarray hybridization, and analysis of expression data

Total RNA was extracted and purified using the RNeasy Lipid Tissue kit (Qiagen, Courtaboeuf, France). RNA quality was checked on RNA 6000 Nano chips using a Bioanalyzer 2100 (Agilent Technologies, Palo Alto, CA). RNA samples used for chip experiments all had RNA integrity numbers (RINs)<sup>16</sup> greater than 9. Agilent's Low RNA Input Linear Amplification Kit PLUS (One-color) was used to generate fluorescent cRNA. The amplified cyanine 3-labeled cRNA samples were then purified using Qiagen's RNeasy mini spin columns and hybridized to Agilent Whole Mouse Genome Microarrays, 4 × 44 K. Microarray slides were washed and scanned with an Agilent scanner, according to the standard protocol of the manufacturer. Information from probe features was extracted from microarray scan images using the Agilent Feature Extraction software v.9.5.1. Expression data were submitted to the National Center for Biotechnology Information's (NCBI's) Gene Expression Omnibus (GEO) repository and are available under the accession number (GenBank: GSE10421).<sup>17</sup> All the analyses were

performed using Bioconductor, an open source software for the analysis of genomic data rooted in the statistical computing environment R (Fred Hutchinson Cancer Research Center, Seattle, WA).<sup>18</sup> Normalization between arrays was performed using the quantile method. Genes for which the background-corrected signal intensities were not greater than 2.6 standard deviations above the average background in at least 3 B6 and 3 D2 mice were assumed not to be expressed in the liver and were excluded from further analysis. The R/MAANOVA package implemented in Bioconductor<sup>19</sup> was used to perform a 2-way analysis of variance in which the log<sub>2</sub>-transformed expression level was considered to be a function of diet, strain, and the effects of the interaction between these 2 factors. F tests, based on the James-Stein shrinkage estimates of the error variance, were computed on a gene-by-gene basis and *P* values obtained by permutation analysis. The proportion of false positives among all the genes initially identified as being differentially expressed (FDR) was assessed using the procedure described by Storey et al.<sup>20</sup> When influence of diet on expression levels was significant, *t* tests were performed to investigate specific effects of iron deficiency and iron enrichment.

### Real-time quantitative PCR

All primers (Table S1, available on the *Blood* website; see the Supplemental Materials link at the top of the online article) were designed using the Primer Express 2.0 software (Applied Biosystems, Foster City). Real-time quantitative polymerase chain reaction (Q-PCR) reactions were prepared with Moloney murine leukemia virus (M-MLV) reverse transcriptase (Promega, Charbonnières-les-Bains, France) and LightCycler 480 DNA SYBR Green I Master reaction mix (Roche Diagnostics, Mannheim, Germany) as previously described<sup>14</sup> and run in duplicate on a LightCycler 480 Instrument (Roche Diagnostics). For each mouse and transcript, an expression measure was calculated as  $2^{Ct_{ToI} - Ct_{HPRT}}$ , where ToI is the transcript of interest; HPRT, a transcript with stable level between strains, diets, and genotypes, quantified to control for variation in cDNA amounts; and Ct, the cycle number where fluorescence reaches a given threshold. Expression values were log<sub>2</sub> transformed, and the contribution of diet (iron deficient, standard, or iron enriched) or genotype (wild type, knockout) in each strain was assessed by analysis of variance followed by Scheffe posthoc tests or by Student *t* tests, respectively, using SAS version 9.1.3 (SAS Institute, Cary, NC). The transcript fold changes following the different dietary conditions or *Smad4/Hamp1* disruption were calculated on values obtained by raising 2 to the power *n*, where *n* is the mean of the log<sub>2</sub>-transformed expression measures within each group. Correlations between liver iron and expression values were assessed by nonparametric Spearman rank-order correlation tests.

### Western blot analysis

Livers were homogenized in a FastPrep-24 Instrument (MP Biomedicals Europe, Illkirch, France) for 40 seconds at 4 m/s. The lysis buffer (50 mM Tris-HCl, pH 8, 150 mM NaCl, 5 mM EDTA, pH 8, 1% NP-40) included inhibitors of proteases (1 mM PMSF, 10 μg/mL leupeptin, 10 mg/mL pepstatin A, and 1 mg/mL antipain) and of phosphatases (10 μL/mL phosphatase inhibitor cocktail 2; Sigma-Aldrich, Saint-Quentin Fallavier, France). Proteins were quantified using the Bio-Rad Protein Assay kit (Bio-Rad Laboratories, Hercules, CA) based on the method of Bradford. Protein extracts (20 μg for Smad7, 30 μg for phosphorylated Smad1/5/8 and Smad1, and 40 μg for Id1) were diluted in Laemmli buffer (Sigma-Aldrich), incubated for 5 minutes at 95°C, and subjected to SDS-polyacrylamide gel electrophoresis (PAGE). Proteins were then transferred to Western Clear Signal PVDF membranes (Schleicher and Schuell, Maidstone, United Kingdom). Membranes were blocked with either 5% of dry milk in TBS-T buffer (10 mM Tris-HCl, pH 7.5, 150 mM NaCl, 0.15% Tween 20) for phosphorylated Smad1/5/8 and Id1, or a more stringent buffer (10% dry milk, 1% BSA, 1% NP40, 0.5% Tween 20 in PBS) for Smad1 and Smad7. They were incubated with rabbit polyclonal antibodies to phosphorylated Smad1/5/8 (1/1000; Cell Signaling Technology, Danvers, MA), Id1 (1/200; Santa Cruz Biotechnology, Santa Cruz, CA) and Smad7 (1/500; Santa

**Table 1. Iron parameters in B6 and D2 mice fed diets with different iron contents**

Diet	Serum iron, $\mu\text{M}$	TIBC, $\mu\text{M}$	Transferrin saturation	Liver iron concentration, $\mu\text{g iron/g dry weight}$
<b>B6 strain</b>				
Iron deficient	7.12 $\pm$ 1.90	61.12 $\pm$ 4.63	11 $\pm$ 3%	211 $\pm$ 24
Iron adequate	11.28 $\pm$ 1.63	39.36 $\pm$ 1.98	29 $\pm$ 4%	441 $\pm$ 85
Iron supplemented	30.46 $\pm$ 1.69	30.74 $\pm$ 0.94	99 $\pm$ 7%	3569 $\pm$ 502
<b>D2 strain</b>				
Iron deficient	7.32 $\pm$ 1.31	66.78 $\pm$ 8.35	11 $\pm$ 3%	224 $\pm$ 44
Iron adequate	22.18 $\pm$ 3.37	52.20 $\pm$ 3.66	42 $\pm$ 4%	502 $\pm$ 69
Iron supplemented	31.38 $\pm$ 2.80	43.62 $\pm$ 3.97	72 $\pm$ 3%	2353 $\pm$ 163

Results are expressed as mean  $\pm$  SD (n=5 mice per group). Analyses of variance show that, within each strain, the dietary iron content significantly affects serum iron, total iron binding capacity (TIBC), transferrin saturation, as well as hepatic iron concentration (all *P* values are  $<$  .001).

Cruz Biotechnology), or with a mouse polyclonal antibody to Smad1 (1/250; Santa Cruz Biotechnology) at 4°C overnight, and washed with TBS-T buffer. Following incubation with a goat anti-rabbit or anti-mouse IgG antibody (1/5000; Cell Signaling Technology) conjugated to horseradish peroxidase, enzyme activity was visualized by an enhanced chemiluminescence (ECL)-based detection system (Amersham Pharmacia Biotech, Orsay, France). Blots were then stripped and reprobed with mouse monoclonal antibody to  $\beta$ -actin (1/20 000; Sigma-Aldrich). Quantification of proteins was calculated by normalizing the specific probe band to  $\beta$ -actin using the Quantity One software (Bio-Rad Laboratories).

## Results


### Genome-wide mRNA expression profiling identified only 4 genes regulated similarly to hepcidin by the dietary iron content

As shown in Table 1, serum iron parameters and liver iron concentration were all significantly affected by the dietary iron content. Whereas, as previously observed, D2 mice fed an iron-balanced diet have slightly higher serum iron parameters than have B6 mice,<sup>21</sup> the latter accumulated significantly more iron when fed the iron-enriched diet than did D2 mice. This is particularly noteworthy in the liver (Table 1) and seems to be characteristic of the strain.<sup>22</sup> *Hamp1* and *Hamp2* gene expression was assessed by real-time quantitative PCR. As previously reported, B6 mice fed an iron-balanced diet have lower *Hamp2* gene expression than D2 mice fed the same diet.<sup>23</sup> As expected, *Hamp1* gene expression was virtually abolished by iron deficiency, but was increased approximately 7-fold by iron overload in both B6 and D2 mice. Similarly, *Hamp2* gene expression was abolished by iron deficiency and increased by iron overload approximately 5-fold in D2 mice and 13-fold in B6 mice (Figure 1). A total of 30 Agilent microarray hybridizations were performed using mRNA extracted from the livers of 15 B6 and 15 D2 mice (5 mice per diet/strain combination). Two-factor (diet and strain) ANOVA identified 1419 transcripts significantly modulated by the diet iron content (FDR 1%/<sub>oo</sub>). These transcripts were classified according to their variations similar or opposite to hepcidin mRNAs in iron deficiency and/or iron overload (Table S2). The DAVID annotation tool<sup>24,25</sup> was used to search for overrepresentation of functional categories within the different gene groups (Tables S2,S3). Interestingly, there was a significant excess of genes involved in the TGF- $\beta$ /BMP signaling pathway among those down-regulated by iron deficiency and up-regulated by iron overload as were *Hamp1* and *Hamp2*. Three genes in this functional category had fold changes induced by both diets of 1.5 or more. Two are coding


for the bone morphogenetic protein Bmp6 and for an inhibitory Smad (Smad7). Of note, Smad7 is important for the tight regulation of Bmp/Smad signal transduction.<sup>26</sup> The third gene, *Id1*, whose promoter is strongly activated by Bmp in a Smad-dependent manner,<sup>27</sup> encodes a negative inhibitor of basic helix-loop-helix (bHLH) proteins. Besides these genes involved in the TGF- $\beta$ /BMP signaling pathway, the gene encoding Atoh8, a bHLH protein known to promote neuronal versus glial fate determination in the nervous system,<sup>28</sup> was also down-regulated by iron deficiency and up-regulated by iron overload with a fold-change of 1.5 or more.

### Real-time quantitative PCR confirms microarray data and shows that *Bmp6*, *Smad7*, *Id1*, and *Atoh8* mRNA levels are highly correlated with the liver iron burden

As shown in Figure 2, real-time quantitative PCR confirms that expression of *Bmp6*, *Smad7*, *Id1*, and *Atoh8* is up-regulated in mice fed an iron-enriched diet. This up-regulation is significantly larger in B6 mice that have accumulated more iron in their liver than D2 mice. Conversely, the expression of these genes is down-regulated in response to iron deficiency. Pearson correlation coefficients of *Bmp6*, *Smad7*, *Id1*, and *Atoh8* expression with liver iron content are highly significant (0.90, 0.86, 0.91, and 0.86, respectively; all *P*  $<$  .001). Recently, Lin et al showed that iron transferrin regulates hepcidin synthesis in primary hepatocyte culture through hemojuvelin and BMP2/4.<sup>29</sup> Although expression of *Bmp2* was found modulated by iron overload but not by iron deficiency (Table S3), we did not observe any in vivo modulation of the *Bmp4* mRNA by



**Figure 1. Expression of hepcidin1 (*Hamp1*) and hepcidin2 (*Hamp2*) in the liver of mice fed diets with different iron contents.** The expression of hepcidin1 and hepcidin2 transcripts was assessed in mice of 2 strains (B6 and D2) fed an iron-deficient (ID), iron-balanced (IB), or iron-enriched (IE) diet. Expression levels were normalized to HPRT and log<sub>2</sub> transformed, and are expressed as means of 5 samples plus or minus standard deviation. The contribution of diet was assessed by ANOVA, and statistical significant differences relative to the iron-balanced diet were determined by Scheffe posthoc tests. \**P*  $<$  .05; \*\*\**P*  $<$  .001.




**Figure 2. Expression of *Bmp6*, *Smad7*, *Id1*, and *Atoh8* in the liver of mice fed diets with different iron contents.** The expression of the 4 transcripts was assessed in mice of 2 strains (B6 and D2) fed an iron-deficient (ID), iron-balanced (IB), or iron-enriched (IE) diet. Expression levels were normalized to HPRT and log<sub>2</sub> transformed, and are expressed as means of 5 samples plus or minus standard deviation. The contribution of diet was assessed by ANOVA, and statistical significant differences relative to the iron-balanced diet were determined by Scheffe posthoc tests. \* $P < .05$ ; \*\* $P < .01$ ; \*\*\* $P < .001$ .

the dietary iron content in our microarray data. Nevertheless, we assessed their expression by real-time quantitative PCR. Interestingly, although expression of *Bmp4* was not significantly affected when mice of either the B6 or the D2 genetic background were made iron overloaded or iron deficient ( $P = .33$  and  $P = .38$ , respectively), a small but significant increase in the expression of *Bmp2* was observed in mice of the B6 strain fed the iron-enriched diet ( $P = .001$ ). No such increase was observed in the D2 mice that have accumulated less iron in their liver than B6 mice (Figure S1). In addition, no decrease in the expression of *Bmp2* was observed in mice of either strain fed the iron-deficient diet. We quantified the Id1 and Smad7 proteins to which antibodies were available in the liver of the 2 mouse strains and observed that Id1 (Figure S2) and Smad7 were modulated by iron not only at the transcript level but also at the protein level.

#### Smad1/5/8 phosphorylation is increased by iron overload and decreased by iron deficiency


Since *Bmp6* belongs to the TGF- $\beta$  superfamily and transmits signal through phosphorylation of Smads,<sup>30</sup> we tested whether phosphorylation of Smad1/5/8 was significantly modulated in liver extracts of mice fed an iron-deficient or an iron-enriched diet. Total protein lysates from 18 mice were prepared (3 mice per strain/diet combination) and the amount of the phosphorylated forms of Smad1/5/8 in each condition was determined by Western blot analysis. As shown in Figure 3 and Figure S3, the iron-enriched diet induced Smad1/5/8 phosphorylation. This induction was stronger in B6 mice than in D2 mice, which correlates with the higher liver iron burden and *Bmp6* expression observed in these mice. Of note, a significant decrease of Smad1/5/8 phosphorylation was also observed in mice fed an iron-deficient diet. The amount of Smad1/5/8 phosphorylation in the liver of these mice is therefore correlated with body iron stores and the quantity of *Bmp6* transcripts.



**Figure 3. Smad1/5/8 phosphorylation is increased by iron overload and decreased by iron deficiency.** (A) Liver lysates from mice fed an iron-deficient (ID), iron-balanced (IB), or iron-enriched (IE) diet ( $n = 3$  in each group) were analyzed by Western blot with an antibody to phosphorylated Smad1/5/8. Blots were stripped and reprobed with an antibody to  $\beta$ -actin as loading control. A representative experiment is shown. (B) Chemiluminescence was quantified using Quantity One software to calculate the ratio of phosphorylated Smad1/5/8 (p-Smad) relative to  $\beta$ -actin. Mean ratios of 3 samples ( $\pm$  SD) are represented on this figure, relative to the mean ratio of the B6 group of mice fed the iron-balanced diet. Statistical significant differences relative to the iron-balanced diet were determined by Student  $t$  tests. \*\*\* $P < .001$ .

#### Smad4 knockout mice have increased *Bmp6* but decreased *Smad7*, *Id1*, and *Atoh8* gene expression

The receptor-activated Smad1, Smad5, and Smad8 form heteromeric complexes with Smad4 and these translocate to the nucleus where they regulate the transcription of specific targets. To investigate whether the variations in the mRNA expression of some of these genes could be induced by the observed variations in Smad1/5/8 phosphorylation, we assessed *Bmp6*, *Smad7*, *Id1*,



**Figure 4. Expression of *Bmp6*, *Smad7*, *Id1*, *Atoh8*, and *Hamp1* in the livers of mice with genetic iron overload.** Mice with liver-specific knockout of *Smad4* (A) or *Hamp1* disruption (B) were compared with their respective controls. Expression levels were normalized to HPRT and log<sub>2</sub> transformed for the statistical analyses. Statistically significant differences relative to the controls were determined by Student  $t$  tests. \* $P < .05$ ; \*\* $P < .01$ ; \*\*\* $P < .001$ . Geometric means ( $2^n$ , where  $n$  is the mean of the log<sub>2</sub>-transformed expression measures within each group) ( $\pm$  SD) are represented on this figure, relative to the geometric mean of the appropriate control group.

*Atoh8*, and *Hamp1* gene expression in the liver of 6 mutant mice with Cre-loxP-mediated liver-specific disruption of *Smad4* and controls of the same genetic background at 4 months of age. Liver-specific disruption of *Smad4* was shown to result in accumulation of iron in many organs, in particular the liver.<sup>12</sup> As shown in Figure 4A, although liver-specific knockout of *Smad4* resulted in markedly decreased expression of *Hamp1*, it led to increased *Bmp6* expression as did the iron-enriched diet. Interestingly, despite the accumulation of iron observed in these mice, *Smad4* deficiency resulted in decreased *Smad7*, *Id1*, and *Atoh8* mRNA expression, as did the iron-deficient diet. These results show that the binding partner for the receptor-activated Smad1, Smad5, and Smad8 is necessary for activation of *Smad7*, *Id1*, and *Atoh8* transcription by iron in response to Bmp signaling.

#### **Hamp1 knockout mice have increased Bmp6, Smad7, Id1, and Atoh8 gene expression**

Targeted disruption of the hepcidin1 gene also results in early and severe multivisceral iron overload.<sup>15</sup> To check whether the Bmp signaling pathway is still functional in these mice, we assessed the expression of the same genes in the liver of 5 *Hamp1* knockout mice and controls of comparable genetic background at 10 weeks of age. As shown in Figure 4B, the *Hamp1*<sup>-/-</sup> mice behave like mice fed an iron-enriched diet. Iron overload induced by *Hamp1* deficiency is indeed able to increase the expression not only of *Bmp6* but also of *Smad7*, *Id1*, and *Atoh8*.

## **Discussion**

Analysis of liver transcription profiles in C57BL/6 and DBA/2 mice fed an iron-deficient or an iron-enriched diet allowed us to identify 4 genes that were regulated by iron similarly to hepcidin in both strains. These results were confirmed by quantitative real-time PCR and point out important molecules participating in the BMP/Smad signaling pathway that regulates hepcidin expression in vivo.

We therefore tested whether the dietary iron content effectively activated/repressed the Smad signaling pathway in these mice through phosphorylation of the receptor-regulated Smads. Activation of Smad1, Smad5, and Smad8 indeed is considered to be the major signaling pathway for BMPs.<sup>31</sup> Interestingly, we observed that Smad1/5/8 phosphorylation was induced by iron overload, and that this induction was proportional to the liver iron burden of the mice. Conversely, it was reduced by iron deficiency. This is the first demonstration that dietary iron regulates hepcidin synthesis in vivo through Smad1/5/8 phosphorylation. Of note, injection of iron dextran was also shown to lead to an increase in Smad1/5/8 phosphorylation in liver extracts from both zebrafish and mouse.<sup>32</sup>


Which of the BMP ligands is (are) responsible for the observed activation of the Smad pathway by iron in vivo is still unknown. Recent studies have shown that many ligands of the TGF- $\beta$ /BMP superfamily could positively regulate hepcidin expression in vitro<sup>9,12,13</sup> and in vivo.<sup>10</sup> However, in these studies, hepatic cells or mice were exposed to high levels of exogenous ligands and the observed signals may not reflect the physiological response to iron exposure. Our data suggest that *Bmp6*, which is highly expressed in the liver,<sup>33</sup> has a preponderant role in the activation of the Smad signaling pathway in vivo. We indeed showed in the present study that liver expression of *Bmp6* was transcriptionally regulated by iron (ie, induced by iron overload and repressed by iron deficiency) and that its induction paralleled the liver iron burden of the mice.

Of note, expression of *Bmp2* was also slightly increased in mice with the most severe iron burden (ie, the B6 mice fed the iron-enriched diet) which indicates that, in addition to *Bmp6*, *Bmp2* can also participate in the activation of the Smad signaling pathway. No other ligand of the TGF- $\beta$ /BMP superfamily, including *Bmp4*, had its gene expression significantly modulated by iron in vivo. What exactly triggers *Bmp6* and *Bmp2* modulation by the dietary iron content remains to be determined, but may well be serum transferrin saturation.<sup>34,35</sup>

The composition of BMP ligands after exogenous challenges has been observed to be modulated both at the mRNA and at the protein levels.<sup>36</sup> It is therefore likely that *Bmp6* and *Bmp2* protein levels in iron-overloaded and in iron-deficient mice reflect *Bmp6* and *Bmp2* transcript levels. Whereas hemojuvelin (HJV) acts as a BMP coreceptor to enhance cellular responses to BMP ligands and increase hepcidin expression,<sup>9</sup> soluble HJV seems to bind and sequester BMP ligands and thus to inhibit BMP signaling and hepcidin expression both in vitro and in vivo.<sup>10</sup> Interestingly, the ability of soluble HJV to inhibit BMP induction of hepcidin promoter activation in Hep3B cells is selective and much more effective for BMP6 than for other ligands.<sup>10</sup> siRNA inhibition of endogenous BMP6 in HepG2 cells significantly reduces basal hepcidin expression,<sup>10</sup> confirming that regulation of hepcidin by BMP6 is bidirectional, that is, not only induced when BMP6 is added to cells, but also repressed when endogenous BMP6 expression is decreased. Altogether, these results converge to suggest that BMP6 is the ligand of the TGF- $\beta$ /BMP superfamily that has a key role in the maintenance of systemic iron homeostasis, although *BMP2* can also function as an endogenous regulator of hepcidin expression in vivo, especially when body iron stores are abundant and require strong activation of the Smad signaling pathway.

Our data also show that *Id1*, *Smad7*, and *Atoh8* gene expression in the liver of mice fed an iron-enriched or an iron-deficient diet follows *Bmp6* gene expression and the level of Smad1/5/8 phosphorylation. *Id1* is known to be a direct target gene for BMPs that strongly activate its promoter in a Smad-dependent manner. Indeed, Smad-binding elements (SBEs) and a GC-rich region, which bind both Smad1 and Smad5, confer BMP responsiveness to the *Id1* promoter.<sup>27</sup> Receptor-activated Smads have also been shown to regulate *Smad7* transcription through direct binding to SBEs on its promoter.<sup>37</sup> However, no information on the regulation of *Atoh8* transcription is yet available. To check that the regulation of these genes by body iron stores relies on the formation of heteromeric complexes between phosphorylated Smad1/5/8 and Smad4 and the translocation of these complexes into the nucleus, we measured their expression in mice with liver-specific knockout of *Smad4*. These mice have compromised Bmp signaling and develop severe iron overload.<sup>12</sup> Interestingly, *Bmp6* gene expression is increased in these mice, which shows that its up-regulation by iron is not restricted to dietary iron overload. However, the expression of *Id1*, *Smad7*, and *Atoh8* is decreased, suggesting that *Atoh8*, like *Id1* and *Smad7*, is a direct target gene for BMP ligands. Further analyses are conducted to check whether receptor-activated Smads also bind to the *Atoh8* promoter. Of note, in mice with genetic iron overload due to targeted disruption of *Hamp1*,<sup>15</sup> Bmp signaling is functional, and the expression of *Bmp6*, *Smad7*, *Id1*, and *Atoh8* is increased as it is in mice with dietary iron overload.

It is not clear yet how *Id1*, *Smad7*, and *Atoh8*, whose expression is regulated by body iron providing that *Smad4* is



**Figure 5. A working model for hepcidin regulation by iron in the mouse.** (A) Bmp6, whose expression is induced by iron, binds to type I and II receptors and to the coreceptor, hemojuvelin (Hjv). The constitutively active kinase domains of type II receptors phosphorylate type I receptors, and this in turn activates the Smad signaling pathway through phosphorylation of receptor Smads (Smad1, Smad5, and Smad8). These associate with co-Smads (Smad4) to form a heteromeric complex that translocates to the nucleus and stimulates the expression of a wide range of target genes, including the genes coding for Id1, Smad7, and possibly also Atoh8. Hypothetical roles for Id1, Smad7, and Atoh8 are presented on panels B through E. (B) When dietary iron availability is low, hepcidin transcription is repressed by specific basic-loop-helix (bHLH) repressors. (C) When the diet iron content increases, inhibitor of differentiation proteins Id1 are synthesized. These are known to associate with bHLH proteins and prevent them from binding DNA. By sequestering the bHLH repressors, Id1 proteins may stop them from achieving their inhibiting functions on the hepcidin promoter. (D) Iron overload also increases Smad7, which could enhance Atoh8 activity as it does for another bHLH transcription activator, MyoD. (E) The hepcidin promoter contains several E-boxes known to interact with bHLH transcription factors and further investigation will determine whether Atoh8 binds to and activates the hepcidin promoter.

functional, can act on hepcidin regulation. Inhibitor of differentiation proteins Id1 associate with tissue-specific basic helix-loop-helix (bHLH) transcription factors that can either activate<sup>38</sup> or repress transcription.<sup>39</sup> In this study, we observed that activation of the BMP/Smad signaling pathway by iron leads to synthesis of Id1 that seems proportional to the needs for hepcidin. It should therefore be explored whether Id1 could overcome the inhibitory effect of a transcription factor and allow hepcidin transcription to proceed. The expression of *Smad7* mRNA is known to be quickly induced by members of the TGF- $\beta$  family.<sup>40</sup> Smad7 interferes with binding to type I receptors and thus activation of receptor-regulated Smads or recruits the E3 ubiquitin ligase Smurf to receptors and thus targets them for degradation.<sup>26,41</sup> Although Smad7 could participate in a negative feedback loop to control the intensity and duration of responses to ligands such as Bmp6,<sup>26,41</sup> it could also enhance transcriptional activity of Atoh8, a bHLH protein<sup>28</sup> whose role in the maintenance of iron balance was unsuspected so far, as was shown recently for another bHLH protein, MyoD.<sup>42</sup> The hepcidin promoter contains several E-boxes<sup>43</sup> known to interact with bHLH transcription factors and it would be interesting to test whether Atoh8 could activate the hepcidin promoter. This would establish a link among iron, the different proteins identified in this study, and hepcidin, as depicted in our working model (Figure 5).

In summary, our data underscore a critical role of Bmp6 in the maintenance of systemic iron homeostasis in 2 different strains of mice and suggest that Smad7, Id1, and Atoh8 may participate in this maintenance. They also underline that Smad1/5/8 phosphorylation in the liver is controlled by body iron. It is now important to

understand how exactly these molecules are involved in the complex process of regulation of hepcidin synthesis by iron.

## Acknowledgments

The authors thank the "Service de Zootechnie" (Institut Fédératif de Recherche 30) and the platform "Génomique" (Génopole Toulouse) for assistance and skilled advice.

This work was supported in part by grants from the Agence Nationale pour la Recherche, Paris, France (ANR, program IRONGENES), the European Commission, Brussels, Belgium (LSHM-CT-2006-037296: EUROIRON1), and the Association pour la Recherche sur le Cancer (ARC), Villejuif, France.

## Authorship

Contribution: L.K. and D.M. performed research, analyzed data, and reviewed the paper; A.M. and J.M. performed research and reviewed the paper; V.D. and R.B. performed research; R.-H.W. and C.D. contributed *Smad*-deficient mice and reviewed the paper; S.V. contributed *Hamp1*-deficient mice, discussed the experiments, and reviewed the paper; H.C. and M.-P.R. designed research, analyzed and interpreted data, performed statistical analysis, and wrote the paper.

Conflict-of-interest disclosure: The authors declare no competing financial interests.

Correspondence: Marie-Paule Roth, Inserm U563, CHU Purpan, BP 3028, F-31024 Toulouse Cedex 3, France; e-mail: marie-paule.roth@inserm.fr.

## References

- Ganz T, Nemeth E. Regulation of iron acquisition and iron distribution in mammals. *Biochim Biophys Acta*. 2006;1763:690-699.
- Nemeth E, Tuttle MS, Powelson J, et al. Hepcidin regulates cellular iron efflux by binding to ferroportin and inducing its internalization. *Science*. 2004;306:2090-2093.
- Nicolas G, Chauvet C, Viatte L, et al. The gene encoding the iron regulatory peptide hepcidin is regulated by anemia, hypoxia, and inflammation. *J Clin Invest*. 2002;110:1037-1044.
- Pigeon C, Ilyin G, Courselaud B, et al. A new mouse liver-specific gene, encoding a protein homologous to human antimicrobial peptide hepcidin, is overexpressed during iron overload. *J Biol Chem*. 2001;276:7811-7819.
- Roetto A, Papanikolaou G, Politou M, et al. Mutant antimicrobial peptide hepcidin is associated with severe juvenile hemochromatosis. *Nat Genet*. 2003;33:21-22.
- Papanikolaou G, Samuels ME, Ludwig EH, et al. Mutations in HFE2 cause iron overload in chromosome 1q-linked juvenile hemochromatosis. *Nat Genet*. 2004;36:77-82.

7. Nemeth E, Roetto A, Garozzo G, Ganz T, Camaschella C. Heparin is decreased in TFR2 hemochromatosis. *Blood*. 2005;105:1803-1806.
8. Piperno A, Girelli D, Nemeth E, et al. Blunted hepcidin response to oral iron challenge in HFE-related hemochromatosis. *Blood*. 2007;110:4096-4100.
9. Babitt JL, Huang FW, Wrighting DM, et al. Bone morphogenetic protein signaling by hemojuvelin regulates hepcidin expression. *Nat Genet*. 2006;38:531-539.
10. Babitt JL, Huang FW, Xia Y, Sidis Y, Andrews NC, Lin HY. Modulation of bone morphogenetic protein signaling in vivo regulates systemic iron balance. *J Clin Invest*. 2007;117:1933-1939.
11. Anderson GJ, Frazer DM. Iron metabolism meets signal transduction. *Nat Genet*. 2006;38:503-504.
12. Wang RH, Li C, Xu X, et al. A role of SMAD4 in iron metabolism through the positive regulation of hepcidin expression. *Cell Metab*. 2005;2:399-409.
13. Truksa J, Peng H, Lee P, Beutler E. Bone morphogenetic proteins 2, 4, and 9 stimulate murine hepcidin 1 expression independently of Hfe, transferrin receptor 2 (Tfr2), and IL-6. *Proc Natl Acad Sci U S A*. 2006;103:10289-10293.
14. Dupic F, Fruchon S, Bensaid M, et al. Inactivation of the hemochromatosis gene differentially regulates duodenal expression of iron-related mRNAs between mouse strains. *Gastroenterology*. 2002;122:745-751.
15. Lesbordes-Brion JC, Viatte L, Bennoun M, et al. Targeted disruption of the hepcidin 1 gene results in severe hemochromatosis. *Blood*. 2006;108:1402-1405.
16. Schroeder A, Mueller O, Stocker S, et al. The RIN: an RNA integrity number for assigning integrity values to RNA measurements. *BMC Mol Biol* (<http://www.biomedcentral.com/bmcmolbiol/>). 2006;7:3.
17. National Center for Biotechnology Information. Gene Expression Omnibus. <http://www.ncbi.nlm.nih.gov/geo/>. Accessed June 30, 2008.
18. Gentleman RC, Carey VJ, Bates DM, et al. Bioconductor: open software development for computational biology and bioinformatics. *Genome Biol*. (<http://genomebiology.com>). 2004;5:R80.
19. Cui X, Churchill GA. Statistical tests for differential expression in cDNA microarray experiments. *Genome Biol*. (<http://genomebiology.com>). 2003;4:210.
20. Storey JD, Xiao W, Leek JT, Tompkins RG, Davis RW. Significance analysis of time course microarray experiments. *Proc Natl Acad Sci U S A*. 2005;102:12837-12842.
21. Dupic F, Fruchon S, Bensaid M, et al. Duodenal mRNA expression of iron related genes in response to iron loading and iron deficiency in four strains of mice. *Gut*. 2002;51:648-653.
22. Rodriguez A, Hilvo M, Kytomaki L, et al. Effects of iron loading on muscle: genome-wide mRNA expression profiling in the mouse. *BMC Genomics* (<http://www.biomedcentral.com/bmcgenomics>). 2007;8:379.
23. Courselaud B, Troade MB, Fruchon S, et al. Strain and gender modulate hepatic hepcidin 1 and 2 mRNA expression in mice. *Blood Cells Mol Dis*. 2004;32:283-289.
24. National Institute of Allergy and Infectious Diseases. Database for Annotation, Visualization and Integrated Discovery (DAVID). <http://david.abcc.ncifcrf.gov/>. Accessed June 30, 2008.
25. Dennis G Jr, Sherman BT, Hosack DA, et al. DAVID: Database for Annotation, Visualization, and Integrated Discovery. *Genome Biol*. (<http://genomebiology.com>). 2003;4:P3.
26. Itoh S, ten Dijke P. Negative regulation of TGF-beta receptor/Smad signal transduction. *Curr Opin Cell Biol*. 2007;19:176-184.
27. Korchynski O, ten Dijke P. Identification and functional characterization of distinct critically important bone morphogenetic protein-specific response elements in the Id1 promoter. *J Biol Chem*. 2002;277:4883-4891.
28. Inoue C, Bae SK, Takatsuka K, Inoue T, Bessho Y, Kageyama R. Math6, a bHLH gene expressed in the developing nervous system, regulates neuronal versus glial differentiation. *Genes Cells*. 2001;6:977-986.
29. Lin L, Valore EV, Nemeth E, Goodnough JB, Gabayan V, Ganz T. Iron transferrin regulates hepcidin synthesis in primary hepatocyte culture through hemojuvelin and BMP2/4. *Blood*. 2007;110:2182-2189.
30. Kersten C, Sivertsen EA, Hystad ME, Forfang L, Smeland EB, Myklebust JH. BMP-6 inhibits growth of mature human B cells: induction of Smad phosphorylation and upregulation of Id1. *BMC Immunol* (<http://www.biomedcentral.com/bmcmimmunol/>). 2005;6:9.
31. Miyazawa K, Shinozaki M, Hara T, Furuya T, Miyazono K. Two major Smad pathways in TGF-beta superfamily signalling. *Genes Cells*. 2002;7:1191-1204.
32. Yu PB, Hong CC, Sachidanandan C, et al. Dorsomorphin inhibits BMP signals required for embryogenesis and iron metabolism. *Nat Chem Biol*. 2008;4:33-41.
33. Xia Y, Babitt JL, Sidis Y, Chung RT, Lin HY. Hemojuvelin regulates hepcidin expression via a selective subset of BMP ligands and receptors independently of neogenin. *Blood*. 2008;111:5195-5204.
34. Nemeth E, Rivera S, Gabayan V, et al. IL-6 mediates hypoferrremia of inflammation by inducing the synthesis of the iron regulatory hormone hepcidin. *J Clin Invest*. 2004;113:1271-1276.
35. Schmidt PJ, Toran PT, Giannetti AM, Bjorkman PJ, Andrews NC. The transferrin receptor modulates Hfe-dependent regulation of hepcidin expression. *Cell Metab*. 2008;7:205-214.
36. Rosendahl A, Pardali E, Speletas M, Ten Dijke P, Heldin CH, Sideras P. Activation of bone morphogenetic protein/Smad signaling in bronchial epithelial cells during airway inflammation. *Am J Respir Cell Mol Biol*. 2002;27:160-169.
37. Nagarajan RP, Zhang J, Li W, Chen Y. Regulation of Smad7 promoter by direct association with Smad3 and Smad4. *J Biol Chem*. 1999;274:33412-33418.
38. Ruzinova MB, Benezra R. Id proteins in development, cell cycle and cancer. *Trends Cell Biol*. 2003;13:410-418.
39. Jogi A, Persson P, Grynfeld A, Pahlman S, Axelson H. Modulation of basic helix-loop-helix transcription complex formation by Id proteins during neuronal differentiation. *J Biol Chem*. 2002;277:9118-9126.
40. Nakao A, Afrakhte M, Moren A, et al. Identification of Smad7, a TGFbeta-inducible antagonist of TGF-beta signalling. *Nature*. 1997;389:631-635.
41. Valdimarsdottir G, Goumans MJ, Itoh F, Itoh S, Heldin CH, ten Dijke P. Smad7 and protein phosphatase 1alpha are critical determinants in the duration of TGF-beta/ALK1 signaling in endothelial cells. *BMC Cell Biol* (<http://www.biomedcentral.com/bmccellbiol/>). 2006;7:16.
42. Kollias HD, Perry RL, Miyake T, Aziz A, McDermott JC. Smad7 promotes and enhances skeletal muscle differentiation. *Mol Cell Biol*. 2006;26:6248-6260.
43. Bayele HK, McArdle H, Srai SK. Cis and trans regulation of hepcidin expression by upstream stimulatory factor. *Blood*. 2006;108:4237-4245.