

Gene expression profile of the third pharyngeal pouch reveals role of mesenchymal MafB in embryonic thymus development

*Dil Afroz Sultana,¹ *Shuhei Tomita,¹ Michito Hamada,² Yasuyuki Iwanaga,^{1,3} Yuki Kitahama,¹ Nguyen Van Khang,¹ Shuichi Hirai,^{1,3} Izumi Ohigashi,¹ Sachiko Nitta,¹ Takashi Amagai,⁴ Satoru Takahashi,² and Yousuke Takahama¹

¹Division of Experimental Immunology, Institute for Genome Research, University of Tokushima, Tokushima; ²Department of Anatomy and Embryology, Graduate School of Comprehensive Human Sciences, University of Tsukuba, Ibaraki; ³Department of Anesthesiology and Emergency Medicine, Faculty of Medicine, Kagawa University, Kagawa; and ⁴Department of Immunology and Microbiology, Meiji University of Oriental Medicine, Kyoto, Japan

The thymus provides a microenvironment that induces the differentiation of T-progenitor cells into functional T cells and that establishes a diverse yet selftolerant T-cell repertoire. However, the mechanisms that lead to the development of the thymus are incompletely understood. We report herein the results of screening for genes that are expressed in the third pharyngeal pouch, which contains thymic primordium. Polymerase chain reaction (PCR)-based cDNA subtraction screening for genes expressed in microdissected tissues of the third pharyngeal pouch rather than the second pharyngeal arch yielded one transcription factor, MafB, which was predominantly expressed in CD45⁻IA⁻PDGFR α^+ mesenchymal cells and was detectable even in the third pharyngeal pouch of FoxN1-deficient *nude* mice. Interestingly, the number of CD45⁺ cells that initially accumulated in the embryonic thymus was significantly decreased in MafB- deficient mice. Alterations of gene expression in the embryonic thymi of MafBdeficient mice included the reduced expression of Wnt3 and BMP4 in mesenchymal cells and of CCL21 and CCL25 in epithelial cells. These results suggest that MafB expressed in third pharyngeal pouch mesenchymal cells critically regulates lymphocyte accumulation in the embryonic thymus. (Blood. 2009;113:2976-2987)

Introduction

A functionally competent T-cell pool with a diverse repertoire of T-cell antigen receptors (TCRs) is essential in mounting immune responses to invading pathogens.^{1,2} Most peripheral T cells bearing $\alpha\beta$ TCRs are generated in the thymus.^{2,3} The migration of hematopoietic stem cell–derived T-lymphoid progenitor cells into the thymus and the subsequent interactions with thymic stromal microenvironments are essential for T-cell development in the thymus.^{4,5} The entry of hematopoietic cells into thymus primordium is initiated during embryogenesis as early as embryonic day 11.5 (E11.5) in mice.⁶

The initial formation of thymic primordium occurs before the migration of hematopoietic cells and involves interactions between epithelial cells of the third pharyngeal pouch (3PP) endoderm and mesenchymal cells derived from neural crest at earlier stages (E9.5 to E11.5) of embryogenesis.^{7,8} This initial thymus development is governed by several transcription factors, including Tbx1, Hoxa3, Pax1, Pax9, Eya1, and Six1.9-18 Tbx1 is expressed in pharyngeal endoderm^{9,10} and is required for pharyngeal segmentation; its deficiency causes various pharyngeal defects, including impaired generation of the thymus and the parathyroid glands.^{11,12} Hoxa3 is expressed in the 3PP endoderm and neural crest mesenchyme;13,14 the lack of Hoxa3 reduces the expression of Pax1 and Pax9, which in turn causes defective formation of the thymus and the parathyroid glands.¹⁵⁻¹⁷ Eya1 and Six1 are expressed in pouch endoderm and neural crest mesenchyme of the third pharyngeal clefts, and are required for the development of the thymus and the parathyroid glands.¹⁸ Thus, transcriptional regulations of epithelialmesenchymal interactions during this prelymphocyte stage of thymus development are mostly shared between epithelial and mesenchymal cells and between thymus and parathyroid development.

Subsequently, at E11.5, FoxN1 is detectable in the ventral aspect of the 3PP.19 FoxN1 is specifically expressed in epithelial cells (but not in mesenchymal cells) of the thymus but not the parathyroid glands.¹⁹ The formation of thymic primordium before the entry of T-lymphoid progenitor cells does not seem to require FoxN1, but the subsequent differentiation of thymic primordium into functional thymus is dependent on FoxN1.20 Thus, FoxN1-dependent development of thymic epithelial cells is implicated in the interactions between thymic epithelial cells and developing lymphoid cells.²⁰⁻²² Indeed, FoxN1 is required for thymic epithelial cells to optimally produce CCL25, a chemokine involved in attracting T-lymphoid progenitor cells to the thymus, 23,24 and to produce DLL1 and DLL4, the Notch ligands involved in supporting T-cell development in the thymus.²⁵ However, FoxN1-dependent epithelial-lymphocyte interactions are not sufficient for thymus development. Rather, the contribution of mesenchymal cells appears crucial for the optimal generation of the thymus even at this stage. Several investigators have reported that neural crest-derived mesenchymal cells critically regulate the growth and development of thymic epithelial cells²⁶ by secreting molecules, including Wnts, bone morphogenetic proteins (BMPs), and fibroblast growth factors (FGFs).²⁷⁻³⁰ However, the transcriptional mechanisms that govern these epithelial-mesenchymal interactions at this lymphocyte accumulation stage have remained elusive.

The online version of this article contains a data supplement.

© 2009 by The American Society of Hematology

Submitted June 24, 2008; accepted November 22, 2008. Prepublished online as *Blood* First Edition paper, January 22, 2009; DOI 10.1182/blood-2008-06-164921.

^{*}D.A.S. and S. Tomita contributed equally to this work.

The publication costs of this article were defrayed in part by page charge payment. Therefore, and solely to indicate this fact, this article is hereby marked "advertisement" in accordance with 18 USC section 1734.

Table 1. PCR primers

Gene	Primer	Primer	PCR product size, bp
GAPDH	5'-CCGGTGCTGAGTATGTCGTG-3'	5'-CAGTCTTCTGGGTGGCAGTG-3'	349
CCL21	5'-GGGGAACCTCTAAGTCTGGAA-3'	5'-CTCATTCCCTGGGAGACACT-3'	341
PTH	5'-GCTGGCAGTCTGTCTTCTTACCC-3'	5'-TGTCAGTGCCCTGCACTGTC-3'	343
Casr	5'-GGTATACAGCGCACACGTGA-3'	5'-CCTTTGCCTCTCCTGCACTT-3'	282
Spink8	5'-AGTTGCAGTCCTTGTTCTGG-3'	5'-CTGTCTCCCAAACTGGGTATC-3'	331
Pla2g7	5'-CAGACTCCAAAGGACATCGCAA-3'	5'-GTCAAAGGGTGACCCAGGAA-3'	303
Elovl2	5'-CTACTACGGCCTGTCTGTG-3'	5'-GGGCTTTGGGGAAACCATTC-3'	281
MafB	5'-GTTATAGGGGAGGTCTAGGTGT-3'	5'-AAGCTCGTTTCCGATGCAG-3'	292
lrs4	5'-AGACAGGGTACAGAAGGTAGAG-3'	5'-GCAACTCAAAGCAGGTCTGTGAAG-3'	407
IL7	5'-CTGGTGAACTGCACAAGTAAGG-3'	5'-AGCTGCATTTCTGTGTCCATC-3'	287
Nedd4	5'-TTGAGGAACAGCCTACACTTCC-3'	5'-TCCAGTGATTGCCACGTTCT-3'	329
Pax1	5'-TGGAGCTAGCACAGCTGGGTATC-3'	5'-GCCCCTGTTTGCTCCATAAACGTC-3'	440
Gcm2	5'-GCCAGCCATCTGTGACAAGG-3'	5'-GCCCTGGAATAGGGAAGCTG-3'	383
Angpt2	5'-AATGTTCCGTGGGAGTTCAG-3'	5'-GTACAGTCTCCGCATTCACC-3'	388
Eeflg	5'-AGCCCCAGTTCAGGGCTATC-3'	5'-AGCGGTACTCGGCATACCAC-3'	394
2610528KiiRik	5'-GAGCACCAACAGGCTACGAC-3'	5'-ATGTGCCTGGGAAGGTTTGG-3'	500
Gpsn2	5'-CCCATACCCCACCAAGAACC-3'	5'-GGTAGTCGCGGAACTCCTTC-3'	206
Hnrpf	5'-GGGCTGCCCTACAAAGCAAC-3'	5'-CCCCGTAACAGCCACTCACT-3'	325
PD-L1	5'-GAGAGCCTCGCGTCCAAAG-3'	5'-GTGGTTTTGCCCTGGCTGTGATCT-3'	409
FGF7	5'-CTCTACAGGTCATGCTTCCACC-3'	5'-ACAGAACAGTCTTCTCACCCT-3'	174
FGF10	5'-CAGCGGGACCAAGAATGAAG-3'	5'-TGACGGCAACAACTCCGATTT-3'	77
IGF1	5'-CAGGCTATGGCTCCAGCATTCGG-3'	5'-CAGATCACAGCTCCGGAAGC-3'	77
IGF2	5'-GAGCTTGTTGACACGCTTCAGTTTGTC-3'	5'-GTTTGGCCTCTCTGAACTCTTTGAG-3'	357
IGF1R	5'-GACATCCGCAACGACTATCAG-3'	5'-GTAGTTATTGGACACCGCATC-3'	393
Wnt3	5'-ACCTGGAGAAGGCTGGAAGT-3'	5'-CTTGTCCTTGAGGAAGTCGC-3'	280
Wnt4	5'-CTCAAAGGCCTGATCCAGAG-3'	5'-TCACAGCCACACTTCTCCAG-3'	293
Wnt7a	5'-GGCTTCGCCAAGGTCTTCG-3'	5'-CATGAGGTCACAGCCACTGG-3'	440
Wnt7b	5'-TGCCCGTGAGATCAAAAAG-3'	5'-CTGCGTTGTACTTCTCCTTG-3'	200
Wnt10a	5'-CCTGGAGACTCGGAACAAAG-3'	5'-AACCGCAAGCCTTCAGTTTA-3'	157
Wnt10b	5'-GATACCCACAACCGCAACTC-3'	5'-GGCTCACCTTCATTTACACACA-3'	323
Wnt11	5'-GCTCCATCCGCACCTGTT-3'	5'-CGCTCCACCACTCTGTCC-3'	331
FGFR2IIIb	5'-AACGGTCACCACACCGGC-3'	5'-AGGCAGACTGGTTGGCCTG-3'	311
BMP2	5'-TGTGACCAGACTATTGGACACC-3'	5'-AGTTCAGGTGGTCAGCAAGG-3'	460
BMP4	5'-CGAGGCGACACTTCTACAG-3'	5'-TGGGGGCTTCATAACCT-3'	394
FoxN1	5'-CTCGTCGTTTGTGCCTGAC-3'	5'-TGCCTCTTGTAGGGGTGGAAA-3'	243
DLL1	5'-GTTAGCATCATTGGGGCTACCCAG-3'	5'-CGCCTCTGCTAACTCTGAGAGAACC-3'	519
DLL4	5'-CTGTCCTTATGGCTTTGTGG-3'	5'-GCTCCTTCTTCTGGTTTGTG-3'	260
CCL25	5'-GTTTTGTTGGGGCCTGGATG-3'	5'-ACTCCTCACGCTTGTACTGTTG-3'	345
PDGFRa	5'-TCGTCCTTTTCTCTGAGATGCGTT-3'	5'-CCCACGCTGAAGGTTCCGTT-3'	428
RARb	5'-ACCGGCATACTGCTCAATCC-3'	5'-ACTGGCATCGGTTCCTAGTG-3'	264
LCK	5'-ACGGTGGCTTCTACATCTCC-3	5'-GGCCGTGTACTCATTGTCCT-3	626

By screening for genes that are expressed in E11.5 3PPs, we have identified that the transcription factor MafB is strongly expressed in thymic mesenchymal cells rather than thymic epithelial cells or thymocytes. We show that MafB-deficient mice generate thymic primordium containing FoxN1-expressing epithelial cells by E11.5. However, subsequent generation of the thymus with developing thymocytes is significantly impaired in MafB-deficient mice. The E11.5 3PP of MafB-deficient mice exhibits reduced expression of several mesenchymal molecules, including Wnt3 and BMP4, and several epithelial molecules, including CCL21 and CCL25, suggesting that MafB expressed in thymic mesenchymal cells critically regulates embryonic thymus development at the lymphocyte accumulation stage.

Methods

Mice

All mice were maintained under specific pathogen-free conditions in accordance with intramural guidelines. C57BL/6 (B6), BALB/c-*nu/nu*, and BALB/c-*nu/+* mice were obtained from SLC (Shizuoka, Japan). MafB-

deficient mice³¹ were described previously. The day when a vaginal plug was first observed was designated as gestation day 0.5 (E0.5). All experiments were carried out under the approval of the Institutional Animal Care and Use Committee of the University of Tokushima.

Laser-capture microdissection

E11.5 embryos were embedded in OCT compound (Sakura Finetek, Tokyo, Japan), sliced into 5- μ m-thick sections, and attached to glass slides for staining with Histogene reagent (Arcturus Engineering, Mountain View, CA). Sections were placed on a thin-laser pressure-catapulting membrane (Arcturus). Epithelial linings of 3PPs, second pharyngeal arches (2PAs), and several other tissues were dissected with a PixCell II laser capture microdissection system (Arcturus).

cDNA amplification, subtraction, and cDNA library construction

3PP and 2PA tissues were microdissected from 34 sections of E11.5 embryos. RNA was extracted from the microdissected tissues using CapSure HS LCM Caps and PicoPure RNA Isolation Kit (Arcturus). DNaseI-treated RNA was polymerase chain reaction (PCR)–amplified using the Super SMART PCR cDNA Synthesis Kit (Clontech, Mountain View, CA). Amplified cDNA was digested with *Rsa*I and then subjected to

the subtraction using the PCR-Select cDNA Subtraction Kit (Clontech) for genes expressed in 3PPs rather than 2PAs. Subtracted cDNA was cloned in TA cloning vector. A total of 1000 randomly picked clones were sequenced.

Isolation of thymic stromal cells

Thymic stromal cells were prepared from E14.5 embryonic thymi and 6-week-old adult thymi by digestion with collagenase, dispase, and DNaseI, as described.³² For isolation of fetal thymic stromal cells, cells were stained with allophycocyanin-conjugated anti-CD45 antibody, FITC-conjugated anti–I-A antibody, and biotinylated anti-PDGFR α antibody (eBioscience, San Diego, CA) followed by phycoerythrin-conjugated streptavidin. Cells were isolated with FACSVantage cell sorter (BD Biosciences, San Jose, CA) as described.³³ For isolation of adult thymic stromal cells, CD45⁻ cells enriched with a magnetic cell sorter (Miltenyi Biotec, Auburn, CA) were stained with allophycocyanin-conjugated anti-CD45 antibody, FITC-conjugated anti–I-A antibody, and fibroblast-specific MTS15 antibody,³⁴ followed by biotinylated goat anti–rat antibody and phycoerythrin-conjugated streptavidin. Propidium iodide was used to exclude dead cells. Sorted cells with at least 80% purity were used for further analysis.

RNA amplification for quantitative **RT-PCR** analysis

RNA was extracted from microdissected E11.5 embryonic tissues using the PicoPure RNA Isolation Kit (Arcturus). DNaseI-treated RNA was amplified by the Message Amp II aRNA Kit (Ambion, Austin, TX), which consisted of reverse transcription (RT) with an oligo-dT primer containing the T7 promoter sequence and in vitro transcription of resulting cDNA with T7 RNA polymerase to amplify RNA copies. RNA purity and quantification was determined with a Nanodrop spectrophotometer (Nanodrop Technologies).

Quantitative RT-PCR analysis

Amplified RNA or total cellular RNA extracted using Isogen (Wako Pure Chemical Industries, Osaka, Japan) was reverse-transcribed with oligo-dT primer and Superscript III reverse transcriptase (Invitrogen, Carlsbad, CA). Real-time PCR was performed with SYBER Premix Ex Taq (Takara, Otsu, Japan) and ABI Prism 7900HT Sequence Detection System with SDS 2.0 software (Applied Biosystems, Foster City, CA). Amplified signals were confirmed to be single bands over gel electrophoresis. Gene expression was normalized to GAPDH mRNA using the Δ cycle threshold (Ct) method.³⁵ Primers for PCR are listed in Table 1.

Multicolor confocal microscopy analysis

Fresh tissues were embedded in OCT compound (Sakura Finetek). Frozen sections measuring 5-µm thick were fixed with either acetone or paraformaldehyde and stained with the following antibodies: anti-mouse FoxN1 antibody,24 ER-TR5 monoclonal antibody specific for medullary epithelial cells,³⁶ biotinylated monoclonal antibodies specific for mouse CCL21 and CCL25 (R&D Systems, Minneapolis, MN), rabbit anti-mouse AIRE antibody (M-300; Santa Cruz Biotechnology, Santa Cruz, CA), rabbit anti-pancytokeratin polyclonal antibody (Dako Cytomation, Carpinteria, CA), Alexa 488-conjugated anti-rabbit IgG antibody, and Alexa 633-conjugated anti-mouse IgG antibody. To analyze green fluorescent protein (GFP) expression derived from the MafB- allele, in which the coding sequence of MafB was replaced with GFP,31 of MafB+/heterozygous mice, E15.5 thymus was fixed in 4% paraformaldehyde, sectioned, and stained with anti-pancytokeratin antibody. Stained sections were mounted with a fluorescence mounting medium (Dako). Images were acquired with a TCS SP2 laser scanning microscope (Leica, Mannheim, Germany) equipped with argon and helium-neon lasers (excitation at 488, 546, and 633 nm), $20 \times$ 1.25-0.75 NA and $40 \times 1.25-0.75$ CS oil objectives, and Leica confocal software version 2.0.

Bioinformatics

Sequences were analyzed using the public databases of the National Center for Biotechnology Information (NCBI),³⁷ JAX–Mouse Genome Informatics (MGI),³⁸ and European Molecular Biology Laboratory (EMBL)–



Figure 1. Subtraction screening for genes expressed in the 3PP. (A) Sagittal sections of E11.5 C57BL/6 mouse embryos were 2-color stained with anti-CD45 (green) and anti-keratin (red) antibodies. Regions of the 3PP and the 2PA are marked with blue lines. (B) Regions of 3PP (top panels, marked with dashed lines) and 2PA (bottom panels, marked with dashed lines) in histogene-stained sagittal sections of E11.5 C57BL/6 mouse embryos were microdissected for RNA preparation. Left and right panels show the sections before and after microdissection, respectively. (C) Schematic diagrams of the screening strategy for genes expressed in E11.5 3PPs.

InterPro.³⁹ Domains and motifs were identified using the NCBI–Conserved Domain Database (CDD)⁴⁰ and the EMBL-SMART tool.⁴¹

Statistics

Statistical comparison was performed by the Student t test using Excel software (Microsoft, Redmond, WA).

Results

Subtraction screening for genes expressed in the 3PP

To identify genes that are selectively expressed in the 3PP that contains thymic primordium, tissues of the 3PP and the 2PA were isolated from E11.5 mouse embryos by laser capture microdissection (Figure 1A,B). Total RNA isolated from these tissues was reverse-transcribed and amplified. The 3PP-derived cDNA was hybridized with an excess amount of 2PA-derived cDNA, and subtracted cDNA was used for the preparation of a plasmid library (Figure 1C). The representative size of library inserts was 0.5 to 3 kb. Analysis of 1000 randomly picked clones yielded sequences of 678 inserts. A total of 217 genes were found in these inserts: 17 genes appeared no less than 3 times (Table 2; group 1); 50 genes appeared twice (Table 2; group 2); and 150 genes appeared only once (Table 2; group 3). Evaluation of gene expression in the 3PP and the 2PA by quantitative RT-PCR confirmed that 12 of the 17 group 1 genes were expressed at higher levels in the 3PP than in the 2PA, whereas none of the 8 genes randomly chosen from groups 2 and 3 showed any apparent differences in expression levels between the 2 tissues (Table 2). Table 3 lists the 12 genes that were expressed at higher levels in the 3PP than in the 2PA. Among these genes, 6 encoding Pax1, Gcm2, calcium-sensing receptor

Table 2. Screening for genes expressed in E11.5 3PPs

	Gene from subt	raction screening		Verification of expression specificity	
Group/No.	Name	Accession no.	Frequency	3PP/2PA (SE)	
1					
1	CCL21	NM_011124	254	407.7 (86)	*
2	PTH	NM_020623	104	2.1 (1.1)	*
3	Casr	NM_013803	10	19 088.5 (1963)	*
4	Spink8	NM_183136	8	53.9 (19)	*
5	Pla2g7	NM_013737	7	5.3 (3.8)	*
6	Elovl2	NM_019423	6	0.5 (0.3)	
7	MafB	NM_010658	6	1 842.7 (542)	*
8	Irs4	NM_010572	5	4 166.0 (958)	*
9	IL7	NM_008371	4	658.7 (32)	*
10	Nedd4	NM_010890	3	210.3 (94)	*
11	Pax1	NM_008780	3	2.5 (2.1)	*
12	Gcm2	NM_008104	3	8.2 (4.6)	*
13	Angpt2	NM_007426	3	3.2 (0.7)	*
14	Eeflg	NM_026007	3	0.1 (0.1)	
15	2610528KiiRik	NM_175184	3	0.8 (0.1)	
16	Gpsn2	NM_134118	3	0.3 (0.1)	
17	Hnrnpf	NM_133834	3	0.5 (0.1)	
	17 genes				12 genes
2					
1	Dtprp	NM_010088	2	0.1 (0.0)	
2	Pbx1	NM_008783	2	1.6 (0.7)	
3	Atp5a1	NM_007505	2	0.4 (0.1)	
4	Zfp9	NM_011763	2	1.1 (0.5)	
	50 genes				
3					
1	Ndufv1	NM_133666	1	0.1 (0.0)	
2	Neurod	NM_010894	1	0.3 (0.1)	
3	Cks2	NM_025415	1	0.3 (0.1)	
4	Ptpn18	NM_011206	1	0.1 (0.0)	
	150 genes				
Total	217 genes				12 genes

Subtraction screening was carried out as described in "Methods." Of the 217 genes identified in the screening, 17 that appeared more than twice are categorized in group 1. Groups 2 and 3 consist of genes that appeared twice (50 genes) and once (150 genes), respectively. Ratios of expression levels in the 3PP to expression levels in the 2PA were verified by quantitative RT-PCR, and the values indicate averages and standard errors (SEs) of 3 independent measurements.

*Twelve of the 17 group 1 genes were confirmed to be expressed at higher levels in 3PP than in 2PA. None of 4 randomly selected genes from group 2 and 4 randomly selected genes from group 3 showed significantly higher expression in 3PP than in 2PA.

(Casr), parathyroid hormone (PTH), CCL21, and IL7 are known for their roles in the development and function of the thymus and the parathyroid glands,^{15,16,19,23,42-46} suggesting that this library screening effectively profiled genes that are involved in the development and function of the 3PP and its descendant organs. The rest of the genes were *MafB*, *Irs4*, *Pla2g7*, *Nedd4*, *Agpt2*, and *Spink8*, which have not been previously identified for their roles in the development or function of the thymus or the parathyroid glands (Table 3). Therefore, we further analyzed their expression and function.

Expression profiles of 3PP genes

We next examined the expression of these functionally unclear 3PP genes (*MafB*, *Irs4*, *Pla2g7*, *Nedd4*, *Agpt2*, and *Spink8*) in different

Table 3. Summary of gen	es expressed	in E11.	5 3PPs
-------------------------	--------------	---------	--------

	, , ,			
Gene	Synonym	Category	Pharyngeal functions	References
Pax1	Paired box 1	Transcription factor	Pharyngeal development	15,16
Gcm2	Glial cells missing homolog 2	Transcription factor	Parathyroid development	19,42
Casr	Calcium-sensing receptor	Receptor	Parathyroid function	43
PTH	Parathyroid hormone	Hormone	Parathyroid function	44
CCL21	CC chemokine ligand 21	Chemokine	Parathyroid/thymus function	23,45
IL7	Interleukin 7	Cytokine	Thymus function	46
MafB		Transcription factor	Unknown	
lrs4	Insulin receptor substrate 4	Receptor signaling	Unknown	
Pla2g7	PAF acetylhydrolase	Phospholipase	Unknown	
Nedd4		Ubiquitin ligase	Unknown	
Agpt2	Angiopoietin 2	Angiogenesis	Unknown	
Spink8	Tof2	Peptidase inhibitor	Unknown	

The 12 genes selectively expressed in E11.5 3PPs are divided into two groups. The 6 genes in the top group are known for their roles in the development or function of parathyroid glands or thymus. The involvement in pharyngeal region of the 6 genes in the bottom group has not been identified.



Figure 2. Expression profiles of 3PP genes. Quantitative RT-PCR analysis of total cellular RNA from indicated tissues of E11.5, E14.5, and 8-week-old (8W) C57BL/6 mice. mRNA levels of indicated genes were normalized to GAPDH mRNA levels, and are indicated as the ratio to the amount expressed in E11.5 2PAs (■). Shown are geometric means (bars) and standard errors (lines) of 3 independent measurements.

tissues isolated from different ontogenies of mice. Total RNA isolated from indicated tissues of E11.5, E14.5, and 8-week-old mice was reverse-transcribed (Figure 2). Quantitative RT-PCR confirmed that all of these 6 genes were expressed at higher levels in the 3PP than in the 2PA at E11.5 (Figure 2), in good agreement with the results of initial screening (Table 2). However, none of them was expressed strictly in the 3PP or specifically in the thymus; rather, all 6 genes were expressed in various organs throughout the ontogeny (Figure 2). It may be interesting to note that *Irs4* and *Pla2g7* were undetectable in several organs of E11.5 mice, and *Irs4* was undetectable in several organs of 8-week-old mice (Figure 2), suggesting that *Irs4* and *Pla2g7* show higher tissue expression specificity than the other 4 genes.

We also examined the expression of these 6 genes in various cell subsets in embryonic thymus. Freshly isolated E14.5 fetal thymus lobes were digested with collagenase, dispase, and DNase, and single-cell suspensions were multicolor-stained for flow cytometry sorting of CD45⁺ leukocytes and CD45⁻ stromal cells. CD45⁻ stromal cells were additionally fractionated into IA⁺ thymic epithelial cells and IA⁻ nonthymic epithelial cells. CD45⁻IA⁻ nonthymic epithelial cells were further fractionated into PDGFRa⁺ mesenchymal cells and PDGFRa⁻ nonmesenchymal cells⁴⁷ (Figure 3). Quantitative RT-PCR analysis of the thymic epithelial cell–specific gene *FoxN1*, T-lymphoid cell–specific gene *Lck*, and mesenchymal cell–specific gene *PDGFRa* verified the successful fractionation of sorted cell populations (Figure S1, available on the Blood website; see the Supplemental Materials link at the top of the online article). In this analysis of sorted cell populations, all 6 genes were more strongly expressed in CD45⁻ stromal cells than CD45⁺ leukocytes (Figure 3). Of the 6 genes, 5 (MafB, Pla2g7, Nedd4, Agpt2, and Spink8) were more strongly expressed in CD45⁻IA⁻ nonthymic epithelial cells than CD45⁻IA⁺ thymic epithelial cells, whereas Irs4 was more strongly expressed in thymic epithelial cells than nonepithelial thymic stromal cells. MafB, Irs4, Pla2g7, Nedd4, and Agpt2 were more strongly expressed in CD45⁻IA⁻PDGFRa⁺ thymic mesenchymal cells than PDGFR α^{-} nonmesenchymal thymic stromal cells, whereas Spink8 was more strongly expressed in nonmesenchymal, nonepithelial thymic stromal cells than thymic mesenchymal cells (Figure 3). These results indicate that among the 6 genes that are expressed in the 3PP but whose functions are unclear, MafB, Pla2g7, Nedd4, and Agpt2 are predominantly expressed in mesenchymal cells in the embryonic thymus, whereas Irs4 and Spink8 are predominantly expressed in thymic epithelial cells and nonepithelial, nonmesenchymal stromal cells, respectively, in the embryonic thymus.

MafB is expressed in thymic mesenchymal cells

Because MafB is the only transcription factor in the 6 3PP genes whose roles are unclear in the development and function of **Figure 3. Expression of 3PP genes in fetal thymus cells.** Quantitative PCR analysis of total cellular RNA from indicated cell fractions purified from E14.5 C57BL/6 mice. mRNA levels of indicated genes were normalized to GAPDH mRNA levels and are indicated as the ratio to the amount expressed in CD45⁺ leukocytes (left panels), CD45⁻IA⁺ epithelial cells (middle panels), or CD45⁻IA⁻PDGFRα⁻ nonepithelial, nonmesenchymal stromal cells (right panels). Means and standard errors of the results obtained from 3 independent measurements are shown.



3PP-derived organs, we further analyzed the expression and function of MafB in thymus development. To do so, we initially examined MafB expression in cell subpopulations isolated from embryonic and adult thymi. As described above, MafB in E14.5 embryonic thymus was predominantly expressed in mesenchymal cells (Figure 3). To examine the expression of MafB in postnatal thymus, adult thymus lobes were digested with collagenase, dispase, and DNase, and the single-cell suspension was fractionated into CD45⁺ leukocytes and CD45⁻ stromal cells. CD45⁻ stromal cells were further fractionated into CD45⁻IA⁺ thymic epithelial cells, CD45⁻IA⁺MTS15⁺ thymic mesenchymal cells, and CD45⁻IA⁺MTS15⁻ nonepithelial, nonmesenchymal thymic stromal cells. Quantitative RT-PCR analysis of MafB mRNA levels showed that MafB was most strongly expressed in CD45⁻IA⁺MTS15⁺ thymic mesenchymal cells in the adult thymus (Figure 4A). These results indicate that MafB in the thymus is most strongly expressed in mesenchymal cells in either the embryonic or postnatal period.

MafB expression in microdissected tissues from E11.5 3PPs of FoxN1-deficient *nude* mice was comparable with that of control heterozygotes, whereas the expression of PD-L1, which was previously shown to be dependent on thymic epithelial cell–specific transcription factor FoxN1,⁴⁸ was undetectable in E11.5 3PPs of *nude* mice (Figure 4B). These results indicate that MafB expression in E11.5 3PPs is independent of FoxN1.

In situ expression analysis of MafB in the developing thymus by monitoring GFP expression derived from the MafB⁻ allele, in which the coding sequence of MafB was replaced with GFP,³¹ of MafB^{+/-} heterozygous mice showed that MafB expression in embryonic thymus was most prominently detected at the capsules and also detectable within the thymus (Figure 4C). This distribution of MafB-expressing cells agrees with the reported distribution of neural crest–derived mesenchymal cells.⁴⁹⁻⁵¹

Thymus development in MafB-deficient mice

MafB is essential for central respiratory control, possibly involving the specification of rhythmogenic neurons. MafB-deficient mice die from central apnea at birth and are defective for respiratory rhythmogenesis.31,52 MafB-deficient newborn mice also display renal dysgenesis.³¹ We thus examined thymus development in MafB-deficient mice during embryogenesis before death at birth. We found that thymus containing keratin-expressing thymic epithelial cells was formed in MafB-deficient mice at E12.5 (Figure 5A). However, the number of CD45⁺ lymphoid cells that accumulated within E12.5 thymus of MafB-deficient mice was significantly reduced and approximately half of that of wild-type mice (Figure 5A, D). The number of CD45⁺ cells in the E12.5 thymus of MafB^{+/-} heterozygous mice was less severely reduced and approximately 75% of that of wild-type mice (Figure 5A,D), suggesting that haploid dosage of the MafB gene affects the number of E12.5 thymocytes. Similarly, the numbers of CD45⁺ cells in E13.5 thymus, which were 4-fold larger than those in E12.5 thymus, were reduced in MafB^{-/-} mice (approximately 50% of wild-type mice) and MafB^{+/-} mice (approximately 75% of wild-type mice; Figure 5B,E). On the other hand, the numbers of thymic epithelial cells identified by the coexpression of FoxN1 and keratin were comparable and not significantly different among MafB^{-/-}, MafB^{+/-}, and wild-type mice (Figure 5C,F). These results indicate that the number of thymocytes that initially accumulate in the fetal thymus



Figure 4. Expression of MafB in adult thymus cells and *nude* 3PPs. (A) Quantitative PCR analysis of MafB transcript levels in indicated cell fractions isolated from adult C57BL/6 mice. MafB mRNA levels were normalized to GAPDH mRNA levels and are indicated as the ratio to the amount expressed in CD45⁺ leukocytes. Means and standard errors of the results obtained from 4 independent measurements are shown. (B) Quantitative PCR analysis of MafB and PD-L1 transcript levels in microdissected 3PPs from E11.5 *nul*+ mice or E11.5 *nulnu* mice. mRNA levels were normalized to GAPDH mRNA levels and are indicated as the ratio to the amount expressed in *nul*+ 3PPs. Means and standard errors of the results obtained from 3 independent measurements are shown. (C) In situ expression analysis of MafB in developing thymus by monitoring GFP expression (green) derived from the MafB⁻ allele, in which the coding sequence of MafB was replaced with GFP, of MafB^{+/-} heterozygous mice.³¹ Paraformaldehyde-fixed E15.5 thymus sections were stained with anti-pancytokeratin antibody (red). Scale bars indicate 100 µm.

is significantly reduced in MafB-deficient mice, and that MafB affects the optimal cellularity of thymocytes but not thymic epithelial cells.

Later in development at E18.5, the thymi of MafB-deficient mice contained ER-TR5–expressing medullary epithelial cells that included AIRE⁺ cells (Figure 6A) and generated unimpaired CD4/CD8 profiles of thymocytes that included CD4⁺CD8⁺ thymocytes (Figure 6B,C). The number of thymocytes was not significantly impaired in E18.5 MafB-deficient mice (Figure 6D). These results indicate that unlike initial thymus development at E12.5 and E13.5, thymus development at E18.5 is restored to normalcy in MafB-deficient mice, and therefore suggest that MafB is involved in embryonic thymus development only during initial ontogeny.

Reduced chemokine expression in embryonic thymus of MafB-deficient mice

Finally, we examined the expression of genes that were previously shown to participate in thymus development^{6,8} (Figure 7) to understand how MafB regulates initial thymus development. Using quantitative RT-PCR, the genes were first examined for the ratios of



Figure 5. Early development of embryonic thymus in MafB-deficient mice. (A-C) Sagittal sections of indicated mice at E12.5 (A) or E13.5 (B) were 2-color–stained for CD45 (green) and keratin (red). Sagittal E13.5 sections of frozen embryos from indicated mice were also 2-color–stained for FoxN1 (red) and keratin (blue; C). Scale bars indicate 50 μ m in panel A and 150 μ m in panels B and C. (D-F) Means and standard errors (n = 10 for panel D, n = 4 for panel E, and n = 2 for panel F) of the numbers of CD45⁺ leukocytes (D,E) or FoxN1⁺ cells (F) in keratin⁺ thymic areas per section of E12.5 (D) and E13.5 (E,F) embryos are shown. ****P* < .001; ***P* < .01; N.S., not significant.

expression between CD45⁻IA⁻PDGFR α^+ thymic mesenchymal cells and CD45-IA+ thymic epithelial cells isolated from the embryonic thymus (Figure 7A). Wnt11, FGF7, and PDGFRa genes were more than 10 times more strongly detectable in mesenchymal cells than in epithelial cells (Figure 7A mesenchymal genes). In contrast, the expression levels of DLL4, Wnt7a, and BMP2 genes showed a less than 10-fold difference between mesenchymal and epithelial cells (Figure 7A mesenchymal and epithelial genes). Wnt4, FGFR2IIIb, Pax1, and FoxN1 genes were more than 10 times more strongly detectable in epithelial cells than in mesenchymal cells (Figure 7A epithelial genes). We then isolated cellular RNA from microdissected tissues of E11.5 3PPs of MafB-deficient mice as well as wild-type mice (Figure 7B). Quantitative RT-PCR analysis of mesenchymal and/or epithelial genes in the 3PPs of MafB-deficient mice showed significantly reduced expression of mesenchymal genes, such as Wnt11, PDGFR α , BMP4, RAR β , and Wnt3; mesenchymal and epithelial genes, such as DLL4, Wnt7a, BMP2, Wnt7b, Wnt10a, IGF1R, and Wnt10b; and epithelial genes, such as CCL21 and CCL25 (Figure 7B). Other genes in the 3PPs of MafB-deficient mice, including mesenchymal FGF7 and FGF10 and epithelial IL7, Pax1, Gcm2, and FoxN1, showed no expression reduction (Figure 7B). These results indicate that MafB, which in the thymus is most strongly expressed in mesenchymal cells, regulates the expression of several epithelial genes, including CCL21 and CCL25, in addition to



Figure 6. Late development of embryonic thymus in MafB-deficient mice. (A,B) Frozen sections of embryonic thymus isolated from indicated mice at E18.5 were multicolor-stained either for AIRE (green) and ER-TR5 (a marker for medullary thymic epithelial cells; red) or for CD4 (green), CD8 (blue), and ER-TR5 (red). Representative results of 4 independent experiments are shown. Scale bars indicate 50 μ m in panel A and 100 μ m in panel B. (C) Two-color flow cytometry analysis of CD4 and CD8 in E18.5 thymocytes from indicated mice. Numbers indicate percentage of cells in quadrant. Representative results of 4 independent experiments are shown. (D) Means and standard errors (n = 3-4) of the numbers of thymocytes from indicated mice at E18.5 are shown. N.S. indicates not significant.

several mesenchymal genes, including *Wnt3* and *BMP4*, in E11.5 3PPs.

It was previously shown that FoxN1 essentially promotes the expression of CCL25, DLL1, and DLL4,¹⁹⁻²⁵ whereas Gcm2 is essential for the expression of CCL21, PTH, and Casr.^{19,23,42-44} Our results demonstrate that the expression of some FoxN1-dependent genes, including CCL25 and DLL4, and some Gcm2-dependent genes, including CCL21, PTH, and Casr, was reduced in the 3PPs of MafB-deficient mice, whereas both FoxN1 and Gcm2 were normally detectable in the absence of MafB (Figure 7B). These results suggest that FoxN1 and Gcm2 expressed in the 3PP are not sufficient for the expression of CCL21 and CCL25 as well as several other epithelial genes, and that mesenchymal MafB, in addition to epithelial FoxN1 and Gcm2, critically regulates the development of the thymus and the parathyroid glands.

We further found that the expression of some parathyroid epithelial genes, such as *PTH* and *Casr*, was significantly reduced in IA⁺ thymic epithelial cells of MafB-deficient mice even at E15.5 (Figure 7C), suggesting that MafB affects the maturation of the epithelial compartment in the common primordia for the thymus and the parathyroid glands. Interestingly, the numbers of PDGFRa⁺ mesenchymal cells, IA⁺ epithelial cells, and PDGFRa⁻IA⁻ stromal cells were not significantly reduced in the thymus of E15.5 MafB-deficient embryos (Figure S2), whereas the expression of some mesenchymal genes, such as *Wnt3* and *BMP4*, was significantly reduced in PDGFR α^+ thymic mesenchymal cells of E15.5 MafB-deficient embryos (Figure 7D), suggesting that the reduced expression of such mesenchymal genes as *Wnt3* and *BMP4* reflects reduced gene expression per mesenchymal cell rather than fewer mesenchymal cells in the embryonic thymus.

We previously identified that the coordination between CCL21 and CCL25 is essential for chemotactic guidance of T-lymphoid progenitor cells to fetal thymic primordium before, but not after, vascularization of fetal thymus.^{23,45,53} It was therefore interesting to know whether or not CCL21 and CCL25 protein expression might be also decreased in the E11.5 3PPs of MafB-deficient mice. As shown in Figure 8, both CCL21 and CCL25 proteins in E11.5 3PPs were less detectable in MafB-deficient mice than in wild-type mice. These results indicate that the 3PP of MafB-deficient mice is defective in the expression of CCL21 and CCL25. The reduction of these 2 chemokines can explain why thymocyte cellularity is reduced only during initial development, rather than late development, of the embryonic thymus, because these 2 chemokines are required for lymphoid progenitor colonization before, but not after, vascularization of fetal thymus.

Discussion

Mesenchymal cells essentially contribute to thymus development before and after the entry of hematopoietic cells. In particular, their interactions with epithelial cells pivotally regulate the maturation of thymic epithelial cells via such mediators as Wnts, BMPs, and FGFs.²⁶⁻³⁰ However, the transcriptional mechanisms underlying the development of thymic mesenchymal cells were unclear. Our results show that the transcriptional factor MafB in the thymus is predominantly expressed in mesenchymal cells and is essential for optimum accumulation of thymocytes during embryonic thymus development. The expression of Wnt family genes, including Wnt3 and Wnt11, as well as BMP family genes, including BMP4, which were predominantly produced by thymic mesenchymal cells, was decreased in embryonic thymus of MafB-deficient mice. Coincident with the decrease in expression of these mesenchymal genes, the expression of several genes, including CCL21 and CCL25, which were predominantly produced by thymic epithelial cells, was reduced in the embryonic thymi of MafB-deficient mice. These results suggest that MafB expressed in mesenchymal cells of the embryonic thymus governs the production of several molecules, such as Wnt3, Wnt11, and BMP4, which in turn act on thymic epithelial cells to promote the production of CCL21 and CCL25, which attract lymphoid progenitor cells into the thymus. Thus, this study proposes a novel molecular pathway that mediates mesenchymal contribution to thymus development.

MafB is a basic leucine zipper transcription factor belonging to the large Maf family. It forms dimers with various basic leucine zipper proteins, thereby regulating a wide variety of genes.⁵⁴ MafB is expressed in various tissues, including kidney, pancreas, lens, and hindbrain, as well as in hematopoietic cells.^{31,49,54,55} A deficiency in MafB results in death at birth from the defect of respiratory neurons in the hindbrain and subsequent central breathing failure.⁵² MafB-deficient mice also display renal dysgenesis.³¹ A missense mutation in MafB, called *kreisler*, causes segmentation abnormalities in the caudal hindbrain and defective inner ear development.⁵⁵ The expression and function of MafB in mesenchymal cells of embryonic thymus revealed in the present study agree



Figure 7. Gene expression profiles of thymic stromal cells in MafB-deficient mice. (A) Quantitative RT-PCR analysis of indicated genes in CD45⁻IA⁻ PDGFRa⁺ thymic mesenchymal cells (Ms) and CD45⁻IA⁺ thymic epithelial cells (Ep) isolated from E14.5 C57BL/6 embryonic thymus. mRNA levels were normalized to GAPDH mRNA levels and are indicated as the ratios of the levels in Ms cells to the levels in Ep cells. Shown are geometric means (bars) and standard errors (lines) of 3 independent measurements. The genes are aligned according to the values of the ratios. In this list, genes that are more than 10 times more strongly detectable in Ms cells than in Ep cells are categorized as Ms and Ep genes. Genes that are not more than 10 times different in Ms cells and Ep cells are categorized as Ms and Ep genes. Genes that are more than 10 times more strongly detectable in Ep cells than in Ms cells are categorized as Ep genes. (B) Quantitative RT-PCR analysis of indicated genes in microdissected 3PP tissues from either E11.5 MafB-deficient (MafB-KO) or wild-type mice (WT). mRNA levels were normalized to GAPDH mRNA levels and are indicated genes in CD45⁻IA⁺ thymic epithelial cells (C) and CD45⁻PDGFRa⁺ thymic mesenchymal cells (D) isolated from E15.5 MafB-deficient (MafB-KO) or wild-type mice (WT). mRNA levels in MafB-KO samples to those in WT samples. Data are means and standard errors of 4 separate measurements. ****P* < .001; ***P* < .01; ***P* < .05.

with the previous detection of MafB in rhombomere 6 of the hindbrain,⁵⁵ which gives rise to neural crest cells that migrate to the third pharyngeal region.^{49,56} We also showed that MafB in the thymus is barely detectable in other cell types, including thymic epithelial cells, and that MafB deficiency does not result in a decrease in the number of FoxN1-expressing thymic epithelial cells. These results suggest that MafB regulates thymus development by specifically controlling mesenchymal functions that map downstream of and/or in parallel with FoxN1 expression in thymic epithelial cells.

Interestingly, a MafB ortholog in *Drosophila melanogaster* encoded by *traffic jam* (*tj*) is expressed in somatic gonadal cells that are in direct contact with germline cells, and in *tj* mutant gonads, somatic cells fail to intermingle and properly envelop germline cells, causing early blockade of germ cell differentiation.⁵⁷ Thus, the possible role of MafB in mesenchymal cells and in cell-to-cell interactions for thymus development may be analogous to the role of *tj* in gonad morphogenesis in the fruit fly.

Regarding the mechanisms underlying thymus development regulation by MafB in mesenchymal cells, our results showed that the 3PPs of MafB-deficient mice express significantly reduced levels of *Wnt3*, *Wnt11*, and *BMP4* genes among the genes that are expressed in thymic mesenchymal cells rather than thymic epithelial cells. Wnt7a, Wnt7b, Wnt10a, Wnt10b, and BMP2, which are expressed by both thymic mesenchymal cells and thymic epithelial cells, are also decreased in the 3PPs of MafB-deficient mice. Previous reports described that Wnts regulate thymocyte cellularity^{58,59} and can act on thymic epithelial cells²⁷ and thymocytes.⁶⁰ It was also reported that BMPs are expressed in thymic mesenchymal cells and act to regulate thymus development by controlling the epithelial-mesenchymal interaction that shapes thymic stroma.^{28,61-63} BMPs in the thymus may also directly act on immature thymocytes.^{64,65} On the other hand, FGF7 and FGF10, which are also expressed in thymic mesenchymal cells rather than thymic epithelial cells and regulate the development of thymic epithelial cells, 26,29,66,67 are normally detectable in the 3PPs of MafB-deficient mice. Thus, we think it is possible that MafB in thymic mesenchymal cells regulates the expression of these Wnts and BMPs rather than FGFs, thereby regulating thymus development by acting on thymic epithelial cells and thymocytes. It is interesting to note that MafB does not affect FGF7 and FGF10 gene expression, in agreement with the finding that the number of FoxN1-expressing thymic epithelial cells is not reduced in MafB-deficient mice.

Genes that are severely affected in the 3PPs of MafB-deficient mice include *CCL21* and *CCL25*. The expression of CCL21 and CCL25 proteins in thymic primordium is indeed reduced in MafB-deficient

Figure 8. Expression of CCL21 and CCL25 proteins in E11.5 3PPs of MafB-deficient mice. Sagittal sections of frozen embryos from indicated mice were 2-color–stained either for CCL21 (pink) and keratin (blue) or for CCL25 (pink) and keratin (blue). Anterior-posterior (A-P) and dorsal-ventral (D-V) orientations of the sections are indicated. Scale bars indicate 100 μ m. Two representative datasets (nos. 1 and 2) of 10 to 14 different sections are shown.



mice. CCL21 and CCL25 are expressed in Gcm2-dependent parathyroid epithelial cells and FoxN1-dependent thymic epithelial cells, respectively, rather than thymic mesenchymal cells and are involved in attracting T-lymphoid progenitor cells to the fetal thymus before vascularization.^{23,45,53} Thus, we think it is possible that MafB in 3PP mesenchymal cells regulates the expression of Wnts and BMPs, and these molecules, in turn, act on 3PP epithelial cells to produce CCL21 and CCL25. Later in embryonic development, thymocyte cellularity recovers to normalcy in MafB-deficient mice, in agreement with the possibility that the reduced thymocyte cellularity in early embryogenesis is due to impaired chemokine production by fetal thymus without MafB. These results indicate that unlike initial thymus development at E12.5 and E13.5, thymus development at E18.5 is restored to normalcy in MafB-deficient mice, therefore suggesting that MafB is involved in embryonic thymus development only during initial ontogeny. Indeed, the numbers of V γ 3-expressing TCR- $\gamma\delta^+$ T cells, which represent the first wave of T cells in the ontogeny,^{23,68} in the thymus and the skin at E18.5 were not reduced in the embryos of MafB-deficient mice (Figure S3).

Itoi et al reported that in PDGFR α -deficient *patch* mutant mice, whose mesenchymal cells are primarily affected, epithelial cells in thymic primordium show a reduction in number and are defective in the expression of SCF, DLL4, and class II MHC molecules.⁶⁹ Interestingly, thymic mesenchymal cells in *patch* mutant embryos are defective in producing FGF7 and FGF10, whereas the genes that exhibit reduced expression in thymic epithelial cells of *patch* mutant mice do not include *CCL21* or *CCL25*.⁶⁹ These results indicate that spectra of molecules affected in *patch* mutant mice and MafB-deficient mice are unequal either in thymic mesenchymal cells or in thymic epithelial cells. Thus, it is likely that epithelial-mesenchymal interactions during thymus development are mediated by multiple molecular pathways in parallel rather than a single molecular stimulus.

In this study, we used PCR-based cDNA subtraction screening for the genes expressed in microdissected tissues of the 3PP rather than the 2PA. Among the 12 genes extracted, 6 (*Pax1*, *Gcm2*, *Casr*, *PTH*, *CCL21*, and *IL7*) are known for their roles in the development and function of the thymus and the parathyroid glands,^{15,16,19,23,42-46} suggesting that our gene expression profiling has a good fidelity in extracting genes that are involved in the development and function of the 3PP and its derivative organs. However, it should be noted that our screening did not extract such genes as *Tbx1*, *Hoxa3*, and *FoxN1*, which are also specifically expressed in the 3PP and are also crucial for its development.^{8-15,19-21} These genes may be expressed in the 2PA and thus subtracted during the procedure (eg, *Tbx1*) or expressed in E11.5 3PPs at very low levels and thus undetectable in the present screening (eg, *FoxN1*).

In conclusion, the present study examining the gene expression profile of the mouse 3PP has identified the role of MafB in thymus development. The results have revealed a novel transcriptional regulation of thymus development by mesenchymal cells and offer a novel molecular pathway that mediates mesenchymal contribution to thymus development via Wnts, BMPs, and chemokines. Because it is shown that transcription factors belonging to the large Maf family, including MafB, MafA, and c-Maf, can exert partly overlapping roles in organ development and cellular functions,^{70,71} it is possible that MafA and c-Maf may also be involved in controlling thymus development. An improved understanding of the molecular mechanisms of thymus development should aid in the formulation of strategies to enhance thymus regeneration in the elderly or those following chemotherapy and/or radiotherapy.

Acknowledgments

We thank Drs N. Iwanami, T. Nitta, and T. Ueno for critically reading the manuscript.

This study was supported by a Ministry of Education, Culture, Sports, Science, and Technology (MEXT; Tokyo, Japan) Grant-in-Aid for Scientific Research.

Authorship

Contribution: D.A.S., S. Tomita, and Y.T. designed the research and wrote the paper; D.A.S., S. Tomita, M.H., Y.I., Y.K., N.V.K., S.H., I.O., and S.N. performed research; D.A.S. and S. Tomita analyzed data; and T.A. and S. Takahashi contributed vital materials.

Conflict-of-interest disclosure: The authors declare no competing financial interests.

The current address for Dr Tomita is Department of Pharmacol-

Immunology, Institute for Genome Research, University of To-

kushima, 3-18-15 Kuramoto, Tokushima, Japan; e-mail: takahama@

ogy, Institute of Health Biosciences, University of Tokushima, Tokushima 770-8503, Japan.

Correspondence: Yousuke Takahama, Division of Experimental

References

- Miller JFAP. The thymus and the development of immunologic responsiveness. Science. 1964; 144:1544-1551.
- Kisielow P, von Boehmer H. Development and selection of T cells: facts and puzzles. Adv Immunol. 1995;58:87-209.
- Scollay R, Wilson A, D'Amico A, et al. Developmental status and reconstitution potential of subpopulations of murine thymocytes. Immunol Rev. 1988;104:81-120.
- Petrie HT. Cell migration and the control of postnatal T-cell lymphopoiesis in the thymus. Nat Rev Immunol. 2003;3:859-866.
- Takahama Y. Journey through the thymus: stromal guides for T-cell development and selection. Nat Rev Immunol. 2006;6:127-135.
- Hollander G, Gill J, Zuklys S, Iwanami N, Liu C, Takahama Y. Cellular and molecular events during early thymus development. Immunol Rev. 2006;209:28-46.
- Cordier A, Haumont S. Development of thymus, parathyroids and ultimo-branchial bodies in NMR1 and nude mice. Am J Anat. 1980;157:227-263.
- 8. Manley NR. Thymus organogenesis and molecular mechanisms of thymic epithelial cell differentiation. Sem Immunol. 2000;12:421-428.
- Yamagishi H, Maeda J, Hu T, et al. Tbx1 is regulated by tissue-specific forkhead proteins through a common Sonic hedgehog-responsive enhancer. Genes Dev. 2003;17:269-281.
- Xu H, Cerrato F, Baldini A. Timed mutation and cell-fate mapping reveal reiterated roles of Tbx1 during embryogenesis, and a crucial function during segmentation of the pharyngeal system via regulation of endoderm expansion. Development. 2005;132:4387-4395.
- Lindsay EA, Vitelli F, Su H, et al. Tbx1 haploinsufficiency in the DiGeorge syndrome region causes aortic arch defects in mice. Nature. 2001;410:97-101.
- Jerome LA, Papaioannou VE. DiGeorge syndrome phenotype in mice mutant for the T-box gene, Tbx1. Nat Genet. 2001;27:286-291.
- Manley NR, Capecchi MR. The role of Hoxa-3 in mouse thymus and thyroid development. Development. 1995;121:1989-2003.
- Manley NR, Capecchi MR. Hox group 3 paralogs regulate the development and migration of the thymus, thyroid and parathyroid glands. Dev Biol. 1998;195:1-15.
- Su D, Ellis S, Napier A, Lee K, Manley NR. Hoxa3 and pax1 regulate epithelial cell death and proliferation during thymus and parathyroid organogenesis. Dev Biol. 2001;236:316-329.
- Wallin J, Eibel H, Neubuser A, Wilting J, Koseki H, Balling R. Pax1 is expressed during development of the thymus epithelium and is required for normal T cell maturation. Development. 1996; 122:23-30.
- Peters H, Neubuser A, Kratochwil K, Balling R. Pax-9-deficient mice lack pharyngeal pouch derivatives and teeth and exhibit craniofacial and limb abnormalities. Genes Dev. 1998;12:2735-2747.
- Zou D, Silvius D, Davenport J, Grifone R, Maire P, Xu PX. Patterning of the third pharyngeal pouch into thymus/parathyroid by Six and Eya1. Dev Biol. 2006;293:499-512.
- Gordon J, Bennett AR, Blackburn CC, Manley NR. Gcm2 and Foxn1 mark early parathyroidand thymus-specific domains in the developing

third pharyngeal pouch. Mech Dev. 2001;103: 141-143.

genome.tokushima-u.ac.jp.

- Nehls M, Kyewski B, Messerle M, et al. Two genetically separable steps in the differentiation of thymic epithelium. Science. 1996;272:886-889.
- Su DM, Navarre S, Oh WJ, Condie BG, Manley NR. A domain of Foxn1 required for crosstalkdependent thymic epithelial cell differentiation. Nat Immunol. 2003;4:1128-1135.
- Itoi M, Kawamoto H, Katsura Y, Amagai T. Two distinct steps of immigration of hematopoietic progenitors into the early thymus anlage. Int Immunol. 2001;13:1203-1211.
- Liu C, Saito F, Liu Z, et al. Coordination between CCR7- and CCR9-mediated chemokine signals in prevascular fetal thymus colonization. Blood. 2006;108:2531-2539.
- Itoi M, Tsukamoto N, Amagai T. Expression of DII4 and CCL25 in Foxn1-negative epithelial cells in the post-natal thymus. Int Immunol. 2007;19:127-132.
- Tsukamoto N, Itoi M, Nishikawa M, Amagai T. Lack of Delta like 1 and 4 expressions in nude thymus anlages. Cell Immunol. 2005;234:77-80.
- Jenkinson WE, Jenkinson EJ, Anderson G. Differential requirement for mesenchyme in the proliferation and maturation of thymic epithelial progenitors. J Exp Med. 2003;198:325-332.
- Balciunaite G, Keller MP, Balciunaite E, et al. Wnt glycoproteins regulate the expression of FoxN1, the gene defective in nude mice. Nat Immunol. 2002;3:1102-1108.
- Bleul CC, Boehm T. BMP signaling is required for normal thymus development. J Immunol. 2005; 175:5213-5221.
- Erickson M, Morkowski S, Lehar S, et al. Regulation of thymic epithelium by keratinocyte growth factor. Blood. 2002;100:3269-3278.
- Rossi SW, Jeker LT, Ueno T, et al. Keratinocyte growth factor (KGF) enhances postnatal T-cell development via enhancements in proliferation and function of thymic epithelial cells. Blood. 2007;109:3803-3811.
- Moriguchi T, Hamada M, Morito N, et al. MafB is essential for renal development and F4/80 expression in macrophages. Mol Cell Biol. 2006;26: 5715-5727.
- Gray DH, Chidgey AP, Boyd RL. Analysis of thymic stromal cell populations using flow cytometry. J Immunol Methods. 2002;260:15-28.
- Ueno T, Liu C, Nitta T, Takahama Y. Development of T-lymphocytes in mouse fetal thymus organ culture. Methods Mol Biol. 2005;290:117-133.
- Gray DH, Tull D, Ueno T, et al. A unique thymic fibroblast population revealed by the monoclonal antibody MTS-15. J Immunol. 2007;178:4956-4965.
- Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative PCR and the 2(-ΔΔC_T) method. Methods. 2001; 25:402-408.
- Vilet V, Melis EM, van Ewijk W. Monoclonal antibodies to stromal cell types of the mouse thymus. Eur J Immunol. 1984;14:524-529.
- National Center for Biotechnology Information. BLAST database. Available at: http://www.ncbi. nih.gov. Accessed January 2008.
- JAX Laboratories. Mouse Genome Informatics (MGI) database. Available at: http://www.informatics.jax.org. Accessed September 2008.
- 39. European Molecular Biology Laboratory. InterPro

Database. Available at: http://www.ebi.ac.uk/interpro. Accessed November 2007.

- National Center for Biotechnology Information. Conserved Domain Database (CDD). Available at: http://www.ncbi.nlm.nih.gov/Structure/cdd/ wrpsb.cgi. Accessed January 2008.
- European Molecular Biology Laboratory. SMART tool. Available at: http://smart.embl-heidelberg.de. Accessed December 2006.
- Günther T, Chen ZF, Kim J, et al. Genetic ablation of parathyroid glands reveals another source of parathyroid hormone. Nature. 2000;406:199-203.
- Ho C, Conner DA, Pollak MR, et al. A mouse model of human familial hypocalciuric hypercalcemia and neonatal severe hyperparathyroidism. Nat Genet. 1995;11:389-394.
- Miao D, He B, Karaplis AC, Goltzman D. Parathyroid hormone is essential for normal fetal bone formation. J Clin Invest. 2002;109:1173-1182.
- Liu C, Ueno T, Kuse S, et al. The role of CCL21 in recruitment of T-precursor cells to fetal thymi. Blood. 2005;105:31-39.
- Von FJU, Vieira P, Lucian LA, McNeil T, Burdach SE, Murray R. Lymphopenia in interleukin (IL)-7 gene-deleted mice identifies IL-7 as a nonredundant cytokine. J Exp Med. 1995;181:1519-26.
- Jenkinson WE, Simona WR, Sonia MP, Jenkinson EJ, Anderson G. PDGFRa-expressing mesenchyme regulates thymus growth and the availability of intrathymic niches. Blood. 2007; 109:954-960.
- Bleul CC, Boehm T. Laser capture microdissection-based expression profiling identifies PD1ligand as a target of the nude locus gene product. Eur J Immunol. 2001;31:2497-2503.
- Le Douarin NM, Jotereau FV. Tracing of cells of the avian thymus through embryonic life in interspecific chimeras. J. Exp. Med. 1975;142:17-40.
- Yamazaki H, Sakata E, Yamane T, et al. Presence and distribution of neural crest-derived cells in the murine developing thymus and their potential for differentiation. Int Immunol. 2005;17:549-558.
- Foster K, Julie Sheridan J, Veiga-Fernandes H, et al. Contribution of neural crest-derived cells in the embryonic and adult thymus. J Immunol. 2008; 180:3183-3189.
- Blanchi B, Kelly LM, Viemari JC, et al. MafB deficiency causes defective respiratory rhythmogenesis and fatal central apnea at birth. Nat Neurosci. 2003;6:1091-1100.
- Jenkinson WE, Simona WR, Parnell SM, et al. Chemokine receptor expression defines heterogeneity in the earliest thymic migrants. Eur J Immunol. 2007;37:2090-2096.
- Kataoka K, Fujiwara KT, Noda M, Nishizawa M. MafB, a new Maf family transcription activator that can associate with Maf and Fos but not with Jun. Mol Cell Biol. 1994;14:7581-7591.
- Cordes SP, Barsh GS. The mouse segmentation gene kr encodes a novel basic domain-leucine zipper transcription factor. Cell. 1994;79:1025-1034.
- Lumsden A, Sprawson N, Graham A. Segmental origin and migration of neural crest cells in the hindbrain region of the chick embryo. Development. 1991;113:1281-1291.
- Li MA, Alls JD, Avancini RM, Koo K, Godt D. The large Maf factor Traffic Jam controls gonad morphogenesis in Drosophila. Nat Cell Biol. 2003;5: 994-1000.
- 58. Mulroy T, McMahon JA, Burakoff SJ, McMahon

AP, Sen J. Wnt-1 and Wnt-4 regulate thymic cellularity. Eur J Immunol. 2002;32:967-971.

- Liang H, Coles AH, Zhu Z, et al. Noncanonical Wnt signaling promotes apoptosis in thymocyte development. J Exp Med. 2007;204:3077-3084.
- Pongracz J, Hare K, Harman B, Anderson G, Jenkinson EJ. Thymic epithelial cells provide Wnt signals to developing thymocytes. Eur J Immunol. 2003;33:1949-1956.
- Tsai PT, Lee RA, Wu H. BMP4 acts upstream of FGF in modulating thymic stroma and regulating thymopoiesis. Blood. 2003;102:3947-3953.
- Patel SR, Gordon J, Mahbub F, Blackburn CC, Manley NR. Bmp4 and Noggin expression during early thymus and parathyroid organogenesis. Gene Expr Patterns. 2006;6:794-799.
- 63. Anderson G, Jenkinson WE, Jones T, et al. Es-

tablishment and functioning of intrathymic microenvironments. Immunol Rev. 2006;209:10-27.

- Hager-Theodorides AL, Outram SV, Shah DK, et al. Bone morphogenetic protein 2/4 signaling regulates early thymocyte differentiation. J Immunol. 2002;169:5496-5504.
- Graf D, Nethisinghe S, Palmer DB, Fisher AG, Merkenschlager M. The developmentally regulated expression of Twisted gastrulation reveals a role for bone morphogenetic proteins in the control of T cell development. J Exp Med. 2002;196: 163-171.
- Revest JM, Suniara RK, Kerr K, Owen JJ, Dickson C. Development of the thymus requires signaling through the fibroblast growth factor receptor R2-IIIb. J Immunol. 2001;167:1954-1961.
- 67. Rossi S, Blazar BR, Farrell CL, et al. Keratinocyte growth factor preserves normal thymopoiesis and

thymic microenvironment during experimental graft-versus-host disease. Blood. 2002;100:682-691.

- Havran WL, Allison JP. Developmentally ordered appearance of thymocytes expressing different T-cell antigen receptors. Nature. 1988;335:443-445.
- Itoi M, Tsukamoto N, Yoshida H, Amagai T. Mesenchymal cells are required for functional development of thymic epithelial cells. Int Immunol. 2007;19:953-964.
- Matsuoka TA, Zhao L, Artner I, et al. Members of the large Maf transcription family regulate insulin gene transcription in islet beta cells. Mol Cell Biol. 2003;23:6049-6062.
- Tsuchiya M, Taniguchi S, Yasuda K, et al. Potential roles of large mafs in cell lineages and developing pancreas. Pancreas. 2006;32: 408-416.