

Vascular complications after splenectomy for hematologic disorders

Shelley E. Crary¹ and George R. Buchanan¹

¹Department of Pediatrics, University of Texas Southwestern Medical Center at Dallas

The most widely recognized long-term risk of splenectomy is overwhelming bacterial infection. More recently, thrombosis has become appreciated as another potential complication of the procedure. Because of these long-term risks, the indications for and timing of splenectomy are debated in the medical community. Ac-

cordingly, the adverse effects and benefits of splenectomy for hematologic disorders and other conditions demand further study. This comprehensive review summarizes the existing literature pertaining to vascular complications after splenectomy for hematologic conditions and attempts to define the potential patho-

physiologic mechanisms involved. This complex topic encompasses diverse underlying conditions for which splenectomy is performed, diverse thrombotic complications, and multiple pathophysiologic mechanisms. (*Blood*. 2009;114:2861-2868)

Introduction

The spleen was once considered unnecessary for life; however, it clearly serves extremely important hematologic and immunologic functions. The spleen is separated into 2 major functional compartments: the white pulp and the red pulp. The white pulp contains a large mass of lymphoid tissue and serves a vital role in the recognition of antigens and production of antibodies. The red pulp of the spleen consists of a tight meshwork of sinusoids, the cords of Billroth, which primarily serve hematologic functions, especially filtration of the blood. The milieu of the red pulp is relatively acidic and hypoglycemic. Aged or damaged red cells not able to tolerate this harsh environment are ultimately removed by splenic macrophages. Antibody-coated cells and bacteria are also recognized and ingested by these phagocytic cells lining the sinusoids. Therefore, persons without a functioning spleen have a severe impairment in their ability to clear encapsulated organisms from the bloodstream. Particulate matter is also removed from red cells as they pass through the splenic sinusoids, and “polished” or “conditioned” red cells, free of surface imperfections, are returned to the bloodstream. The red pulp also acts as a reservoir for approximately one-third of the total platelet mass and a smaller proportion of granulocytes.

Asplenia and hyposplenism

Congenital asplenia can occur in isolation or may be associated with certain forms of congenital heart defects or heterotaxy syndromes.¹ Children with sickle cell disease have acquired hyposplenism that begins at several months of age and progresses to splenic infarction. Young patients with sickle cell disease may also develop recurrent and even life-threatening splenic sequestration requiring surgical splenectomy. Moreover, many immunologic and rheumatic disorders are associated with impairment of the spleen’s phagocytic and immunologic functions. Transient functional hyposplenism may also occur during therapy with cortico-

steroids and after pharmaceutical reticuloendothelial blockade with intravenous IgG or anti-D immunoglobulin.

Surgical splenectomy

According to the National Hospital Discharge Survey, approximately 22 000 total splenectomies were performed for all causes in the United States during 2005.² In most institutions, trauma and incidental splenectomy are the primary indications,³ although splenectomy for trauma is becoming less common than in years past, resulting from more conservative nonoperative management of splenic injury.⁴ The most frequent medical indication for splenectomy is a hematologic disorder (Table 1). Splenectomy is performed in patients having hemolytic anemia (eg, hereditary spherocytosis [HS] and autoimmune hemolytic anemia) because the intrinsically abnormal or antibody-coated red blood cells are prematurely destroyed by splenic macrophages. Because splenectomy can ameliorate the underlying anemia, it is often considered the treatment of choice for such conditions. Sick cell disease may be complicated by splenic sequestration requiring surgical splenectomy, and patients with β -thalassemia may undergo splenectomy to relieve splenomegaly resulting in increased destruction of red blood cells. Splenectomy is also performed in patients with immune thrombocytopenic purpura (ITP), especially when chronic or severe.

Septic risk of asplenia and hyposplenism

For decades, it has been known that in persons with asplenia the major long-term complication is overwhelming bacterial sepsis.^{5,6} These infections occur in persons after surgical splenectomy as well as in conditions predisposing to hyposplenism or asplenia. This complication is less frequent than in years past as a result of pneumococcal vaccination, prophylactic penicillin, and prompt administration of parenteral antibiotics when fever occurs.

Table 1. Current indications for splenectomy in hematologic disorders

Hereditary spherocytosis
Other hereditary hemolytic anemias (eg, stomatocytosis, pyropoikilocytosis, pyruvate kinase deficiency)
Sickle cell disease
Thalassemia major or intermedia
Immune thrombocytopenia
Myeloproliferative disorders
Autoimmune hemolytic anemia
Hypersplenism
Lymphoproliferative disorders
Thrombotic thrombocytopenic purpura

Potential vascular complications after splenectomy

An increased risk of vascular complications involving both the venous and the arterial sides of the circulation may result from splenectomy. A vascular complication is defined here as any condition that causes narrowing or occlusion of a blood vessel. This can be a consequence of in situ thrombosis, thromboembolism, vascular smooth muscle remodeling, vasospasm, or atherosclerosis. The risk of thromboembolic events and pulmonary arterial hypertension (PAH) varies greatly depending on the underlying condition for which the splenectomy is performed and whether or not the condition is associated with ongoing intravascular hemolysis (see Table 2). Thromboembolic complications have been most frequently reported after splenectomy in thalassemia intermedia (TI). TI is characterized by chronic intravascular hemolysis and marked ineffective erythropoiesis. In a recent large survey of 8860 patients with thalassemia, the prevalence of thromboembolic events among those with TI was 4%.⁷ Remarkably, 94% of these complications occurred after splenectomy. The rate of thromboembolic events in other hematologic disorders after splenectomy has not been thoroughly investigated. The following discussion summarizes these conditions in which vascular complications have been reported and reviews purported pathophysiologic mechanisms.

Table 2. Hypothetical risk of vascular complications based on the presence or absence of intravascular hemolysis and splenic function

Pathophysiologic state	Example	Risk of vascular complications
Neither hemolysis nor splenectomy	Normal person	Baseline*
Hemolysis with intact spleen	Hereditary spherocytosis Glucose-6-phosphate dehydrogenase deficiency β-thalassemia trait Other chronic hemolytic anemia	Baseline* or decreased
Splenectomy without hemolysis	Hereditary spherocytosis† Immune thrombocytopenic purpura Trauma	Increased or baseline*
Splenectomy and ongoing hemolysis‡	Thalassemia intermedia Hgb E/β-thalassemia Sickle cell anemia§ Hereditary stomatocytosis	Greatly increased

*Dependent on the person's unique genetic and environmental risk factors.
 †Very mild hemolysis may be present in some persons.
 ‡Especially intravascular hemolysis.
 §Nearly all patients lack functional spleens.

Table 3. Summary of strength of evidence of the cited studies of vascular complications following splenectomy in various hematologic disorders

Medical condition	Reported vascular complications	Strength of evidence
Thalassemia	Arteriothrombosis	Large multi-institutional survey ^{7,20} Individual case reports or small series ²¹⁻²⁵
	DVT or PE	Large multi-institutional survey ⁷ Institutional retrospective review ^{36,37} Individual case reports or small series ³⁸⁻⁴⁰
	PAH	Cross-sectional studies ^{56,67}
Sickle cell disease*	Arteriothrombosis	Large prospective cohort (stroke) ²⁸
	DVT or PE	National database review ^{41,42}
	PAH	Prospective cohort ⁶⁸
Hereditary spherocytosis	Arteriothrombosis	Cross-sectional studies ^{9,10} Individual case reports ¹¹
	DVT or PE	Cross-sectional study ¹⁰ Individual case reports ^{46,60}
	PAH	Case control studies ^{55,56} Individual case reports ^{60,62}
Hereditary stomatocytosis	Arteriothrombosis	Individual case reports ⁵⁰
	DVT or PE	Small case series, but includes majority of known cases ¹⁷ Individual case reports ⁵⁰
	PAH	Small case series, but includes majority of known cases ¹⁷ Individual case reports ^{50,51,53}
Other asplenic states	Arteriothrombosis	Case control study ⁸ Individual case reports ^{18,19,26,27}
	DVT or PE	Case control study (autopsies) ⁴³ Population-based prevalence study ⁴⁹ Individual case reports ^{44,45,47,48}
	PAH	Case control studies ⁵⁵⁻⁵⁷ Individual case reports ^{61,63-65}

*None of the reported complications specifically mention their occurrence after splenectomy, yet most patients with sickle cell disease are functionally asplenic.

Table 3 summarizes the cited reports of vascular complications in various disorders along with a description of the strength of the evidence.

Arteriothrombosis

In 1977, a case-control study of 745 World War II servicemen who had splenectomy resulting from trauma demonstrated that they were 1.9 times more likely than controls to die of ischemic heart disease.⁸ Twenty years later, Schilling reported a 5.6-fold increased rate of arteriosclerotic events (defined as stroke, myocardial infarction, and coronary or carotid artery surgery) in persons older than age 40 years with HS who had undergone splenectomy compared with persons with HS who had not had splenectomy.⁹ This finding was recently confirmed in a follow-up study of arterial events in the same groups of HS patients, which reported a hazard ratio of 7.2 for those with previous splenectomy.¹⁰ Only 7 cases of stroke have otherwise been reported in HS,¹¹⁻¹⁶ and only one of

these cases clearly followed splenectomy.¹¹ Otherwise, no documented case reports of myocardial infarction and other forms of atherothrombosis have been described in HS.

Individual reports have described arterial thrombotic events (primarily involving peripheral extremity, pulmonary, and cerebral arteries) after splenectomy in hemolytic disorders other than HS.¹⁷⁻¹⁹ For example, patients with thalassemia have been reported to be at risk of arteriothrombosis after splenectomy, especially ischemic stroke,²⁰⁻²⁵ but the incidence appears to be less than with venous events. In the largest series of thrombotic complications in thalassemia reviewing 8860 patients, arteriothrombosis (primarily stroke) was more common in the thalassemia major population (n = 28) compared with TI (n = 9, *P* = .005) with all but one patient in each group having previously had splenectomy.⁷ This difference in prevalence is unclear. In addition, rapidly progressive multi-infarct dementia and acute coronary syndrome have been observed in a series of patients with ITP and previous splenectomy.^{26,27} Although stroke is a frequent complication of sickle cell disease,²⁸ these patients also do not appear to be at increased risk of other forms of atherothrombosis.²⁹ Interestingly, myocardial infarction and acute coronary syndrome have rarely been reported in thalassemia or any other hemolytic disorders after splenectomy beyond the limited literature in HS.

Venous thrombosis

Local. Acute portal vein thrombosis has been reported after splenectomy for a wide variety of conditions. Indeed, prospective cohort studies reveal that the incidence of thrombosis involving the portal venous system after splenectomy ranges from 5% to 37%, all occurring within 2 months, and the majority within 2 weeks of the surgery.³⁰⁻³⁵ This is probably the result of local surgical factors (given that the splenic vein remnant is attached to the portal vein) rather than to the absence of the spleen^{30,35} per se. A higher incidence of this complication appears with laparoscopic than with open splenectomy³⁰ and with greater splenic weight.^{30,32,33,35} Higher platelet count shortly after splenectomy does not consistently correlate with this increased risk of thrombosis.^{30,33-35}

Systemic. Patients with thalassemia and previous splenectomy appear to have an increased incidence of venous thromboembolism beyond the portal venous system.^{7,36-40} Although less commonly reported than in thalassemia, patients with sickle cell disease have also been found to be at increased risk of venous thrombosis.^{41,42} In the largest series to date of 8860 patients, 32% of the 146 reported thromboembolic events were deep venous thrombosis, 16% portal venous thrombosis, and 13% pulmonary embolism.⁷ In a large review of more than 37 000 autopsies, the odds of fatal pulmonary embolism were 5-fold higher in persons with previous splenectomy (n = 202, performed for any indication) compared with matched controls (n = 403) not having had splenectomy but with comparable trauma or similar surgery.⁴³ Deep vein thrombosis and pulmonary embolism have also been reported in other conditions after splenectomy.^{17,44-49} Patients with hereditary stomatocytosis (whose hemolytic anemia results from abnormalities in red cell cation permeability) are at such a high risk of thrombotic complications after splenectomy that persons with this condition are advised not to undergo the procedure.^{17,50-53} The largest series of 7 kindreds with hereditary stomatocytosis (which represents > 50% of the reported cases) includes 13 persons with prior splenectomy, 11 of whom suffered severe, recurrent thromboses (the 2 without thrombosis were still children). Nine of 10 affected persons with intact spleens did not have thromboembolic disease. The reported vascular

events in these patients were all venous, for example, deep vein thrombosis, pulmonary embolism, or portal vein thrombosis.¹⁷

PAH

Splenectomy appears to be a risk factor for the development of pulmonary hypertension. This is perhaps the most intriguing and frequently reported vascular complication of splenectomy. The classification of pulmonary hypertension has undergone many changes in the past decade and most recently includes a subcategory of PAH in patients with hemoglobinopathies and/or splenectomy (Table 4).⁵⁴ The pulmonary hypertension in asplenic conditions, including thalassemia and sickle cell disease, is generally classified as PAH (previously known as primary pulmonary hypertension) or chronic thromboembolic pulmonary hypertension, usually occurring in distal pulmonary arteries and having characteristic histopathology.⁵⁵ However, it is possible that the process instead represents “in situ” thrombosis with medial hypertrophy, intimal fibrosis, and plexiform lesions such as seen in idiopathic PAH.⁵⁶ In this current review, we will simply refer to all reported types as PAH.

Splenectomy has also been reported as an independent risk factor in patients with PAH referred for lung transplantation^{55,56} and in persons with chronic thromboembolic pulmonary hypertension.⁵⁷⁻⁵⁹ The reported prevalence of splenectomy in patients with PAH ranged from 8.6% to 11.5% compared with 0% to 0.6% in the control groups (patients with other forms of pulmonary disease). It seems probable that the absence of a spleen, not the underlying condition for which the splenectomy was performed, is the primary cause of PAH because the indications for splenectomy in the aforementioned studies were varied, including trauma, ITP, and HS, states where ongoing hemolysis was generally absent, in contrast to disorders where intravascular hemolysis is prominent.⁵⁵⁻⁵⁷

In addition to the case-control studies described in the previous paragraph, PAH has rarely been described in individual case reports after splenectomy in HS, hereditary stomatocytosis, myeloid metaplasia, paroxysmal nocturnal hemoglobinuria, and unstable hemoglobinopathies.^{17,46,51,53,60-65} In thalassemia patients, after splenectomy PAH occurs at greatly increased frequency (prevalence as high as 70%) over the general population.^{66,67} The prevalence of PAH in sickle cell anemia (where the spleen is usually nonfunctional) appears to be approximately 30%.⁶⁸ It is important to note that in most studies of sickle cell disease and thalassemia PAH is defined by echocardiographic techniques using a tricuspid regurgitant jet velocity of more than 2.5 m/s as the cutoff for diagnosing PAH. Although this is generally considered to correlate fairly well with right heart catheterization,⁶⁹ few studies have included this “gold standard” measurement of PAH in their design, so prevalence data must be interpreted cautiously. Although lack of splenic function has never been directly implicated as its cause, PAH also occurs in various conditions treated with splenectomy, such as

Table 4. Abbreviated World Health Organization Clinical Classification of Pulmonary Hypertension

1. Pulmonary arterial hypertension (PAH)
1.1. Idiopathic (IPAH)
1.2. Familial (FPAH)
1.3. Associated with other underlying disease (including Gaucher disease, hemoglobinopathies, myeloproliferative disorders, splenectomy)
2. Pulmonary hypertension with left heart disease
3. Pulmonary hypertension associated with lung disease and/or hypoxia
4. Pulmonary hypertension due to chronic thrombotic and/or embolic disease
5. Miscellaneous

Adapted from Simmoneau et al⁵⁴ with permission.

Gaucher disease,^{70,71} or associated with functional hyposplenism, such as sarcoidosis, rheumatoid arthritis, systemic lupus erythematosus, HIV infection, and other autoimmune and immunologic disorders.⁶⁹ One case of PAH in isolated congenital asplenia has also been reported.⁷² Unfortunately, no published studies have prospectively evaluated the prevalence of PAH in HS or other patients after splenectomy for conditions other than thalassemia or sickle cell disease.

Miscellaneous vascular complications

Leg ulcers and priapism are additional vascular complications that commonly occur in sickle cell disease,⁷³ and the frequency of these reported events has been correlated with severity of hemolysis.⁷⁴ Priapism also occurs in thalassemia and in certain rare hemoglobinopathies, and its prevalence appears to be higher in splenectomized patients.^{19,75-78} However, only a limited number of cases of priapism after splenectomy for other indications have been reported.⁷⁹

Animal models of vascular complications after splenectomy

Very few studies describing postsplenectomy vascular complications in animal models have been published. A rabbit model of hemolysis showed evidence of pulmonary platelet thromboemboli occurring only after ligation of the splenic artery.⁸⁰ In mouse models of spherocytosis, multiorgan thrombosis and infarction occur at high frequency, although these events occur even with the spleen intact and more probably reflect the extremely high rate of hemolysis.^{81,82} To our knowledge, the effect of splenectomy in these murine models has not been evaluated.

Potential pathophysiologic mechanisms of vascular complications after splenectomy

Vascular events after splenectomy are likely multifactorial, probably resulting from some combination of hypercoagulability, platelet activation, disturbance and activation of the endothelium, and altered lipid profiles. The spleen's primary phagocytic function is to remove infectious organisms, other insoluble cellular debris, and senescent or abnormal red cells. This filtration function results from the blood moving slowly through the splenic sinusoids in the red pulp lined with macrophages actively ingesting that which does not easily pass around them. Absence of this extremely sensitive filter may permit particulate matter and damaged cells to persist in the bloodstream, therefore perturbing and activating the vascular endothelium leading to a shift in vascular homeostasis toward enhanced coagulation. There is limited evidence that splenectomy, irrespective of the indication, increases platelet count,⁸³⁻⁸⁷ hemoglobin concentration,^{84,87,88} plasma cholesterol,⁸⁹⁻⁹¹ leukocyte count,⁸⁴ and C-reactive protein levels.⁹² Each of these elevations independently is associated with increased risk of arteriothrombosis⁹³⁻⁹⁷ so in combination might be expected to foster a highly unfavorable prothrombotic state.

Lipid alterations

In animal models, altered plasma lipid profiles (combinations of increased total cholesterol, low-density lipoprotein cholesterol and triglycerides, and decreased high-density lipoprotein) after splenectomy have been suggested to contribute to accelerated arteriosclerosis even in the absence of hemolysis.^{98,99} These changes were most pronounced in animals after being fed high-fat diets. Similarly,

Caligiuri et al demonstrated that splenectomy in hypercholesterolemic apoE receptor knockout mice accelerated atherosclerosis.¹⁰⁰ The authors proposed that this phenomenon was the result of novel mechanisms in which the splenectomized mice were not able to develop immunity against atherosclerosis. Changes in lipid profiles were not considered to play a large role because cholesterol levels remained constant after splenectomy. Although convincing evidence exists for the presence of hypocholesterolemia in humans with hemolytic anemia,¹⁰¹⁻¹⁰⁴ it is not clear that splenectomy in these persons results in hypercholesterolemia.¹⁰⁵

Hypercoagulability

Evidence in thalassemia supports the presence of a hypercoagulable state greatly exacerbated by splenectomy, which is the result of platelet activation,^{36,66,106-109} enhanced red blood cell adherence to the endothelium,^{110,111} reduced levels of the natural anticoagulants protein C and protein S,^{36,112} and increased thrombin generation.^{36,86,112} Procoagulant cell-derived microparticles have also been shown to be elevated after splenectomy for hematologic disorders, suggesting that the spleen may play a role in clearing them from the circulation.¹¹³⁻¹¹⁶ Several studies have reported testing for inherited or acquired thrombophilias at the time of the thrombotic insults, but no consistent risk factor has been shown in a large cohort of splenectomized persons so the contribution of thrombophilia is yet to be determined.^{7,30,33,117-119}

It has been hypothesized that the hypercoagulable state after splenectomy may usually be secondary to persistence of abnormal erythrocytes in the circulation, which have been rendered "procoagulant" resulting from increased exposure of phosphatidylserine on the outer membrane surface.^{120,121} These abnormal red cells have been shown to be increased in thalassemia and sickle cell disease,^{118,122} but not in HS patients.¹²³ Thrombin generation induced by red cells was increased only in thalassemia patients with prior splenectomy but not in healthy splenectomized persons, suggesting that splenectomy alone was not sufficient to produce these procoagulant erythrocytes.³⁶ Similarly, erythrocyte fragmentation during hemolysis may lead to formation of microparticles, which also have similar procoagulant properties.^{124,125} Therefore, the observed vascular complications after splenectomy in conditions, such as thalassemia and sickle cell disease, may be related to this mechanism, yet it does not explain such complications in other postsplenectomy states without ongoing intravascular hemolysis.

Whole blood viscosity is increased after splenectomy irrespective of indication,^{126,127} and in experimental animal models the spleen appears to play a role in maintaining normal hemorheology.¹²⁸ This hyperviscosity after splenectomy may be the result in part of the persistence of aged and damaged red cells in the circulation as well as intracellular inclusions, such as Howell-Jolly bodies, siderotic granules, and Heinz bodies, all of which promote decreased erythrocyte deformability.

Pulmonary vascular remodeling

Putative pathophysiologic mechanisms of PAH in patients with hemolytic anemia include platelet and/or endothelial activation induced by asymmetric red blood cell membrane phospholipids^{36,122,129} and nitric oxide scavenging by free hemoglobin and subsequent endothelial dysfunction and platelet activation resulting from nitric oxide depletion.¹³⁰ Either mechanism may explain PAH in disorders, such as sickle cell disease and thalassemia, where intravascular hemolysis continues despite splenectomy, but not when hemolysis is absent (eg, after splenectomy for trauma) or

when it virtually resolves after splenectomy (eg, as in HS). The consistent factor in PAH affecting such persons with hemolysis is the lack of a spleen. Asplenia may therefore be an important, if not the primary, determinant. Future studies should investigate whether PAH commonly occurs in patients with intravascular hemolysis who have not undergone splenectomy.

Potential protective role of hemolysis with intact spleen

Recently, Schilling et al reported that adult HS patients who had not undergone splenectomy had only 20% the lifetime risk of arteriosclerotic events as first-degree relatives unaffected with HS.¹³¹ This remarkable observation was confirmed in a recent follow-up study in which subjects with HS and prior splenectomy experienced a greater risk of cumulative arteriosclerotic events than HS subjects without prior splenectomy (hazard ratio = 7.15, $P < .001$).¹⁰ However, there was no difference in the frequency of such events between the group with prior splenectomy and their siblings who did not have HS (hazard ratio = 1.56, $P = .11$). Hemolysis in HS before splenectomy occurs almost exclusively in the spleen (ie, extravascularly), so it is possible that their anemia may protect them from thrombotic events in addition to or rather than splenectomy increasing their risk.¹³¹ Most of the literature regarding vascular complications describes conditions with variable degrees of intravascular hemolysis and reduced or absent splenic function, so these several confounders render interpretation of the existing data quite difficult.

Several mechanisms may explain this apparent protection from cardiovascular events in HS and several other hemolytic anemias described below. Patients with hemolysis generally have low plasma cholesterol levels before splenectomy.¹⁰¹⁻¹⁰⁴ This feature, plus reduced blood viscosity resulting from anemia, may reduce the risk of thrombosis, especially atherothrombosis. Even persons with the β -thalassemia trait appear to have fewer cardiovascular events, possibly resulting from their slightly lower hemoglobin and cholesterol levels.¹³²⁻¹³⁴ Several studies of persons with glucose-6-phosphate dehydrogenase deficiency (a disorder characterized by life-long, intermittent hemolysis that is primarily extravascular) suggest a reduced incidence of cardiovascular events,¹³⁵ possibly also the result in part of decreased cholesterol levels.^{136,137} In addition, hemolysis leads to increased production of bilirubin, which has antioxidant properties and may protect from cardiovascular disease.¹³⁸ To add even more complexity to the subject, persons with sickle cell disease (although clearly at increased risk of PAH, stroke, and peripheral vascular disease) appear to have reduced incidence of atherothrombosis, perhaps resulting from lower cholesterol levels^{29,102,139} (Table 2).

Conclusions

There appears to be compelling evidence for a hypercoagulable state after splenectomy performed for various indications. This finding is probably compounded by other underlying conditions,

especially intravascular hemolysis. Although we think there is substantial evidence that PAH and venous thrombosis occur at increased rates in asplenic hosts, the strength of evidence is less convincing that atherothrombosis (including myocardial infarction and stroke) occurs more frequently after splenectomy.

Clinical significance and implications

It is important to better characterize the thrombotic risks of splenectomy in persons without ongoing hemolysis (eg, HS, ITP, trauma) to better define the role of the spleen in vascular homeostasis. PAH in persons with reduced or absent splenic function has a high mortality rate when untreated¹⁴⁰ but may be amenable to novel therapies under development.¹⁴¹ Greater knowledge about the prevalence, age, and risk factors regarding PAH and other vascular events after splenectomy should assist with the often difficult decision of whether and when splenectomy should be performed in patients with HS, ITP, autoimmune hemolytic anemia, or other conditions for which splenectomy may be beneficial. If validated biomarkers of thrombosis and PAH or actual arterial or venous thrombotic events are indeed problematic after splenectomy, the current practice of recommending splenectomy in many children with HS, for example, may require reassessment. In addition, children and adults who have undergone splenectomy for any reason might be advised to seek monitoring or thromboprophylaxis long after they seem to have been "cured" of their primary condition. Several authors have advocated for the use of short-term and/or long-term thromboprophylaxis after splenectomy, including anticoagulation or antiplatelet agents, especially in thalassemia patients,^{36,47,119,142} although no prospective studies have yet evaluated the utility of this approach. Future studies should therefore be directed at better defining the risks of subsequent thrombotic and vascular events accompanying hemolysis, splenectomy, or the combination thereof.

Acknowledgments

The authors thank Dr Cindy E. Neunert for her thoughtful review of the manuscript.

This work was supported in part by the National Institutes of Health (grants 5 KL2 RR024983-02 and 5 UL1 RR024982-02; North and Central Texas Clinical and Translational Science Initiative).

Authorship

Contribution: S.E.C. performed extensive literature review and wrote the manuscript; and G.R.B. assisted in writing and revising the manuscript.

Conflict-of-interest disclosure: The authors declare no competing financial interests.

Correspondence: Shelley E. Crary, University of Texas Southwestern Medical Center at Dallas, Department of Pediatrics, 5323 Harry Hines Blvd, Dallas, TX 75390; e-mail: Shelley.crary@utsouthwestern.edu.

References

- Rose V, Izukawa T, Moes CAF. Syndrome of asplenia and polysplenia: a review of cardiac and non-cardiac malformations in 60 cases with special references to diagnosis and prognosis. *Br Heart J*. 1975;37(8):840-852.
- DeFrances CJ, Cullen KA, Kozak LJ. National Hospital Discharge Survey: 2005 annual summary with detailed diagnosis and procedure data. National Center for Health Statistics. *Vital Health Stat*. 2007;13(165):178.
- Kraus MD, Fleming MD, Vonderheide RH. The spleen as a diagnostic specimen. *Cancer*. 2001; 91(11):2001-2009.
- Rose AT, Newman MI, Debelak J, et al. The incidence of splenectomy is decreasing: lessons learned from trauma experience. *Am Surg*. 2000; 66(5):481-486.

5. Eraklis AJ, Keyv SV, Diamond LK, Gross RE. Hazard of overwhelming infection after splenectomy in childhood. *N Engl J Med*. 1967;276(22):1225-1229.
6. King H, Shumacker HB Jr. Splenic studies: I. Susceptibility to infection after splenectomy performed in infancy. *Ann Surg*. 1952;136(2):239-242.
7. Taher A, Isma'eel H, Mehio G, et al. Prevalence of thromboembolic events among 8860 patients with thalassaemia major and intermedia in the Mediterranean area and Iran. *Thromb Haemost*. 2006;96(4):488-491.
8. Robinette CD, Fraumeni JF Jr. Splenectomy and subsequent mortality in veterans of the 1939-45 war. *Lancet*. 1977;2(8029):127-129.
9. Schilling RF. Spherocytosis, splenectomy, strokes, and heart attacks. *Lancet*. 1997;350(9092):1677-1678.
10. Schilling RF, Gangnon RE, Traver MI. Delayed adverse vascular events after splenectomy in hereditary spherocytosis. *J Thromb Haemost*. 2008;6(8):1289-1295.
11. Bruguier A, Clement MC, Texier P, Ponsot G. Cerebral ischemic accident and hereditary spherocytosis [in French]. *Arch Fr Pediatr*. 1983;40(8):653-654.
12. Holz A, Woldenberg R, Miller D, Kalina P, Black K, Lane E. Moyamoya disease in a patient with hereditary spherocytosis. *Pediatr Radiol*. 1998;28(2):95-97.
13. Monge-Argiles JA, Lopez-Garcia MD, Ortega-Ortega MD, et al. Ischemic cerebral infarction due to factor V Leiden, hereditary spherocytosis and smoking [in Spanish]. *Rev Neurol*. 2005;40(2):125-126.
14. Ramu CS, Raju GB, Rao KS, Venkateswarlu K. Uncommon neurological manifestations of hemolytic anemia: a report of two cases. *Neurol India*. 2008;56(2):201-203.
15. Tokunaga Y, Ohga S, Suita S, Matsushima T, Hara T. Moyamoya syndrome with spherocytosis: effect of splenectomy on strokes. *Pediatr Neurol*. 2001;25(1):75-77.
16. van Hilten JJ, Haan J, Wintzen AR, et al. Cerebral infarction in hereditary spherocytosis. *Stroke*. 1989;20(12):1755-1756.
17. Stewart GW, Amess JA, Eber SW, et al. Thrombo-embolic disease after splenectomy for hereditary stomatocytosis. *Br J Haematol*. 1996;93(2):303-310.
18. Kim BJ, Park KW, Koh SB, et al. Stroke induced by splenectomy in hemoglobin Madrid: autopsy clues to the underlying mechanism. *Blood Coagul Fibrinolysis*. 2005;16(2):141-144.
19. Gyan E, Darre S, Jude B, et al. Acute priapism in a patient with unstable hemoglobin Perth and Factor V Leiden under effective oral anticoagulant therapy. *Hematol J*. 2001;2(3):210-211.
20. Borgna Pignatti C, Carnelli V, Caruso V, et al. Thromboembolic events in beta thalassaemia major: an Italian multicenter study. *Acta Haematol*. 1998;99(2):76-79.
21. Incorpora G, Di Gregorio F, Romeo MA, Pavone P, Triffietti RR, Parano E. Focal neurological deficits in children with beta-thalassaemia major. *Neuropediatrics*. 1999;30(1):45-48.
22. Sanefuji M, Ohga S, Kira R, Yoshiura T, Torisu H, Hara T. Moyamoya syndrome in a splenectomized patient with beta-thalassaemia intermedia. *J Child Neurol*. 2006;21(1):75-77.
23. Singh R, Venketasubramanian N. Recurrent cerebral infarction in beta thalassaemia major. *Cerebrovasc Dis*. 2004;17(4):344-345.
24. Wong Y, Yu YL, Liang RH, Tso WK, Li AM, Chan TK. Cerebral thrombosis in beta-thalassaemia/hemoglobin E disease. *Stroke*. 1990;21(5):812-816.
25. Brankovic-Sreckovic V, Milic Rasic V, Djordjevic V, Kuzmanovic M, Pavlovic S. Arterial ischemic stroke in a child with β -thalassaemia trait and methylentetrahydrofolate reductase mutation. *J Child Neurol*. 2007;22(2):208-210.
26. Ahn YS, Horstman LL, Jy W, Jimenez JJ, Bowen B. Vascular dementia in patients with immune thrombocytopenic purpura. *Thromb Res*. 2002;107(6):337-344.
27. Ozner MD, Ahn YS, Horstman LL, Jy W, Kolodny L, Myerburg RJ. Chronic platelet activation and acute coronary syndrome in 13 middle-aged patients. *Clin Appl Thromb Hemost*. 1997;3(1):46-53.
28. Ohene-Frempong K, Weiner SJ, Sleeper LA, et al. Cerebrovascular accidents in sickle cell disease: rates and risk factors. *Blood*. 1998;91(1):288-294.
29. Gerry JL, Bulkley BH, Hutchins GM. Clinicopathologic analysis of cardiac dysfunction in 52 patients with sickle cell anemia. *Am J Cardiol*. 1978;42(2):211-216.
30. Ikeda M, Sekimoto M, Takiguchi S, et al. High incidence of thrombosis of the portal venous system after laparoscopic splenectomy: a prospective study with contrast-enhanced CT scan. *Ann Surg*. 2005;241(2):208-216.
31. Hassn AM, Al-Fallouji MA, Ouf TI, Saad R. Portal vein thrombosis following splenectomy. *Br J Surg*. 2000;87(3):362-373.
32. Chaffanjon PC, Brichon PY, Ranchoup Y, Gressin R, Sotto JJ. Portal vein thrombosis following splenectomy for hematologic disease: prospective study with Doppler color flow imaging. *World J Surg*. 1998;22(10):1082-1086.
33. Stamou KM, Toutouzias KG, Kekis PB, et al. Prospective study of the incidence and risk factors of postsplenectomy thrombosis of the portal, mesenteric, and splenic veins. *Arch Surg*. 2006;141(7):663-669.
34. Skarsgard E, Doski J, Jaksic T, et al. Thrombosis of the portal venous system after splenectomy for pediatric hematologic disease. *J Pediatr Surg*. 1993;28(9):1109-1112.
35. Romano F, Caprotti R, Scaini A, et al. Elective laparoscopic splenectomy and thrombosis of the spleno-portal axis: a prospective study with ecocolordoppler ultrasound. *Surg Laparosc Endosc Percutan Tech*. 2006;16(1):4-7.
36. Cappellini MD, Robbiolo L, Bottasso BM, Coppola R, Fiorelli G, Mannucci AP. Venous thromboembolism and hypercoagulability in splenectomized patients with thalassaemia intermedia. *Br J Haematol*. 2000;111(2):467-473.
37. Moratelli S, De Sanctis V, Gemmati D, et al. Thrombotic risk in thalassaemic patients. *J Pediatr Endocrinol Metab*. 1998;11[suppl 3]:915-921.
38. Tso SC, Chan TK, Todd D. Venous thrombosis in haemoglobin H disease after splenectomy. *Aust N Z J Med*. 1982;12(6):635-638.
39. van Teunenbroek A, Wijburg FA, ten Cate JW, van den Berg W, Weening RS. Thromboembolic complications in an asplenic HbE-beta-thalassaemia patient. *Neth J Med*. 1989;35(3):123-127.
40. Wijburg FA, van den Berg W, van Teunenbroek A, Weening RS. Thrombo-embolic complications after splenectomy in HbE-beta-thalassaemia. *Eur J Pediatr*. 1988;147(4):444-445.
41. James AH, Jamison MG, Brancazio LR, Myers ER. Venous thromboembolism during pregnancy and the postpartum period: incidence, risk factors, and mortality. *Am J Obstet Gynecol*. 2006;194(5):1311-1315.
42. Stein PD, Beemath A, Meyers FA, Skaf E, Olson RE. Deep venous thrombosis and pulmonary embolism in hospitalized patients with sickle cell disease. *Am J Med*. 2006;119(10):897-911.
43. Pimpl W, Dapunt O, Kaindl H, Thalhamer J. Incidence of septic and thromboembolic-related deaths after splenectomy in adults. *Br J Surg*. 1989;76(5):517-521.
44. Tiede MP, Ahn ER, Jy W, et al. Life-threatening hypercoagulable state following splenectomy in ITP: successful management with aggressive antithrombotic therapy and danazol. *Clin Appl Thromb Hemost*. 2005;11(3):347-352.
45. Chou R, DeLoughery TG. Recurrent thromboembolic disease following splenectomy for pyruvate kinase deficiency. *Am J Hematol*. 2001;67(3):197-199.
46. Hayag-Barin JE, Smith RE, Tucker FC Jr. Hereditary spherocytosis, thrombocytosis, and chronic pulmonary emboli: a case report and review of the literature. *Am J Hematol*. 1998;57(1):82-84.
47. Mohren M, Markmann I, Dworschak U, et al. Thromboembolic complications after splenectomy for hematologic diseases. *Am J Hematol*. 2004;76(2):143-147.
48. Delaitre B, Blezel E, Samama G, et al. Laparoscopic splenectomy for idiopathic thrombocytopenic purpura. *Surg Laparosc Endosc Percutan Tech*. 2002;12(6):412-419.
49. Linet MS, Nyren O, Gridley G, et al. Causes of death among patients surviving at least one year following splenectomy. *Am J Surg*. 1996;172(4):320-323.
50. Gallagher PG, Chang SH, Rettig MP, et al. Altered erythrocyte endothelial adherence and membrane phospholipid asymmetry in hereditary hydrocytosis. *Blood*. 2003;101(11):4625-4627.
51. Yoshimoto A, Fujimura M, Nakao S. Pulmonary hypertension after splenectomy in hereditary stomatocytosis. *Am J Med Sci*. 2005;330(4):195-197.
52. Perel Y, Dhermy D, Carrere A, et al. Portal vein thrombosis after splenectomy for hereditary stomatocytosis in childhood. *Eur J Pediatr*. 1999;158(8):628-630.
53. Murali B, Drain A, Sellar D, Dunning J, Vuylsteke A. Pulmonary thromboendarterectomy in a case of hereditary stomatocytosis. *Br J Anaesth*. 2003;91(5):739-741.
54. Simonneau G, Galie N, Rubin LJ, et al. Clinical classification of pulmonary hypertension. *J Am Coll Cardiol*. 2004;43[suppl 12]:S5-S12S.
55. Jais X, loos V, Jardim C, et al. Splenectomy and chronic thromboembolic pulmonary hypertension. *Thorax*. 2005;60(12):1031-1034.
56. Hoepfer MM, Niedermeier J, Hoffmeyer F, Flemming P, Fabel H. Pulmonary hypertension after splenectomy? *Ann Intern Med*. 1999;130(6):506-509.
57. Bonderman D, Jakowitsch J, Adlbrecht C, et al. Medical conditions increasing the risk of chronic thromboembolic pulmonary hypertension. *Thromb Haemost*. 2005;93(3):512-516.
58. Lang I, Kerr K. Risk factors for chronic thromboembolic pulmonary hypertension. *Proc Am Thorac Soc*. 2006;3(7):568-570.
59. Suntharalingam J, Machado RD, Sharples LD, et al. Demographic features, BMPR2 status and outcomes in distal chronic thromboembolic pulmonary hypertension. *Thorax*. 2007;62(7):617-622.
60. Jardine DL, Laing AD. Delayed pulmonary hypertension following splenectomy for congenital spherocytosis. *Intern Med J*. 2004;34(4):214-216.
61. Lode HN, Krings G, Schulze-Neick I, et al. Pulmonary hypertension in a case of Hb-Mainz hemolytic anemia. *J Pediatr Hematol Oncol*. 2007;29(3):173-177.
62. Smedema JP, Louw VJ. Pulmonary arterial hypertension after splenectomy for hereditary spherocytosis. *Cardiovasc J Afr*. 2007;18(2):84-89.
63. Heller PG, Grinberg AR, Lencioni M, Molina MM, Roncoroni AJ. Pulmonary hypertension in paroxysmal nocturnal hemoglobinuria. *Chest*. 1992;102(2):642-643.
64. Rich S, Hart K. Familial pulmonary hypertension

- in association with an abnormal hemoglobin: insights into the pathogenesis of primary pulmonary hypertension. *Chest*. 1991;99(5):1208-1210.
65. Marvin KS, Spellberg RD. Pulmonary hypertension secondary to thrombocytosis in a patient with myeloid metaplasia. *Chest*. 1993;103(2):642-644.
 66. Singer ST, Kuypers FA, Styles L, Vichinsky EP, Foote D, Rosenfeld H. Pulmonary hypertension in thalassemia: association with platelet activation and hypercoagulable state. *Am J Hematol*. 2006;81(9):670-675.
 67. Phrommintikul A, Sukonthasarn A, Kanjanavanit R, Nawarawong W. Splenectomy: a strong risk factor for pulmonary hypertension in patients with thalassaemia. *Heart*. 2006;92(10):1467-1472.
 68. Gladwin MT, Sachdev V, Jison ML, et al. Pulmonary hypertension as a risk factor for death in patients with sickle cell disease. *N Engl J Med*. 2004;350(9):886-895.
 69. McGoon M, Guterman D, Steen V, et al. Screening, early detection, and diagnosis of pulmonary arterial hypertension: ACCP evidence-based clinical practice guidelines. *Chest*. 2004;126[suppl]:14S-34S.
 70. Mistry PK, Sirrs S, Chan A, et al. Pulmonary hypertension in type 1 Gaucher's disease: genetic and epigenetic determinants of phenotype and response to therapy. *Mol Genet Metab*. 2002;77(1):91-98.
 71. Elstein D, Klutstein MW, Lahad A, Abrahamov A, Hadas-Halpern I, Zimran A. Echocardiographic assessment of pulmonary hypertension in Gaucher's disease. *Lancet*. 1998;351(9115):1544-1546.
 72. Takahashi F, Uchida K, Nagaoka T, et al. Isolated congenital spleen agenesis: a rare cause of chronic thromboembolic pulmonary hypertension in an adult. *Respirology*. 2008;13(6):913-915.
 73. Powars DR, Chan LS, Hiti A, Ramicone E, Johnson C. Outcome of sickle cell anemia: a 4-decade observational study of 1056 patients. *Medicine (Baltimore)*. 2005;84(6):363-376.
 74. Taylor JG, Nolan VG, Mendelsohn L, Kato GJ, Gladwin MT, Steinberg MH. Chronic hyperhemolysis in sickle cell anemia: association of vascular complications and mortality with less frequent vasoocclusive pain. *PLoS ONE*. 2008;3(5):e2095.
 75. Jackson N, Franklin IM, Hughes MA. Recurrent priapism following splenectomy for thalassaemia intermedia. *Br J Surg*. 1986;73(8):678.
 76. Macchia P, Massei F, Nardi M, Favre C, Brunori E, Barba V. Thalassaemia intermedia and recurrent priapism following splenectomy. *Haematologica*. 1990;75(5):486-487.
 77. Sharma R, Prakash R, Kamboj S, Jain V. Hemoglobin-E beta thalassaemia presenting with recurrent priapism: a rare complication. *Am J Hematol*. 2008;83(3):257.
 78. Thuret I, Bardakdjian J, Badens C, et al. Priapism following splenectomy in an unstable hemoglobin: hemoglobin Olmsted beta 141 (H19) Leu→Arg. *Am J Hematol*. 1996;51(2):133-136.
 79. Atala A, Amin M, Harty JI, Liu YK, Keeling MM. Priapism associated with asplenic state. *Urology*. 1992;40(4):371-373.
 80. Kisanuki A, Kietthubthew S, Asada Y, Marutsuka K, Funahara Y, Sumiyoshi A. Intravenous injection of sonicated blood induces pulmonary microthromboembolism in rabbits with ligation of the splenic artery. *Thromb Res*. 1997;85(2):95-103.
 81. Kaysser TM, Wandersee NJ, Bronson RT, Barker JE. Thrombosis and secondary hemochromatosis play major roles in the pathogenesis of jaundiced and spherocytic mice, murine models for hereditary spherocytosis. *Blood*. 1997;90(11):4610-4619.
 82. Wandersee NJ, Olson SC, Holzhauser SL, Hoffmann RG, Barker JE, Hillery CA. Increased erythrocyte adhesion in mice and humans with hereditary spherocytosis and hereditary elliptocytosis. *Blood*. 2004;103(2):710-716.
 83. Hathirat P, Mahaphan W, Chuansumrit A, Pintadit P, Sasanakul W, Isarangkul P. Platelet counts in thalassemic children before and after splenectomy. *Southeast Asian J Trop Med Public Health*. 1993;24[suppl 1]:213-215.
 84. Schilling RF. Hereditary spherocytosis: a study of splenectomized persons. *Semin Hematol*. 1976;13(3):169-176.
 85. Boxer MA, Braun J, Ellman L. Thromboembolic risk of postsplenectomy thrombocytosis. *Arch Surg*. 1978;113(7):808-809.
 86. Visudhiphan S, Ketsa-Ard K, Pankijagum A, Tumliang S. Blood coagulation and platelet profiles in persistent post-splenectomy thrombocytosis: the relationship to thromboembolism. *Biomed Pharmacother*. 1985;39(6):264-271.
 87. Troendle SB, Adix L, Cray SE, Buchanan GR. Laboratory markers of thrombosis risk in children with hereditary spherocytosis. *Pediatr Blood Cancer*. 2007;49(6):781-785.
 88. Cynober T, Mohandas N, Tchernia G. Red cell abnormalities in hereditary spherocytosis: relevance to diagnosis and understanding of the variable expression of clinical severity. *J Lab Clin Med*. 1996;128(3):259-269.
 89. Sugihara T, Yawata Y. Observations on plasma and red cell lipids in hereditary spherocytosis. *Clin Chim Acta*. 1984;137(2):227-232.
 90. Goldfarb AW, Rachmilewitz EA, Eisenberg S. Abnormal low and high density lipoproteins in homozygous beta-thalassaemia. *Br J Haematol*. 1991;79(3):481-486.
 91. Aviram M, Brook JG, Tatarsky I, Levy Y, Carter A. Increased low-density lipoprotein levels after splenectomy: a role for the spleen in cholesterol metabolism in myeloproliferative disorders. *Am J Med Sci*. 1986;291(1):25-28.
 92. Archarit N, Chuncharunee S, Pornvoranunt A, Atamasirikul K, Rachakom B, Atichartakam V. Serum C-reactive protein level in postsplenectomized thalassemic patients. *J Med Assoc Thai*. 2000;83[suppl 1]:S63-S69.
 93. Gagnon DR, Zhang TJ, Brand FN, Kannel WB. Hematocrit and the risk of cardiovascular disease—the Framingham study: a 34-year follow-up. *Am Heart J*. 1994;127(3):674-682.
 94. Wilson PW, D'Agostino RB, Levy D, Belanger AM, Silbershatz H, Kannel WB. Prediction of coronary heart disease using risk factor categories. *Circulation*. 1998;97(18):1837-1847.
 95. Bassuk SS, Rifai N, Ridker PM. High-sensitivity C-reactive protein: clinical importance. *Curr Probl Cardiol*. 2004;29(8):439-493.
 96. Vannucchi AM, Barbui T. Thrombocytosis and thrombosis. *Hematology Am Soc Hematol Educ Program*. 2007:363-370.
 97. Collier BS. Leukocytosis and ischemic vascular disease morbidity and mortality: is it time to intervene? *Arterioscler Thromb Vasc Biol*. 2005;25(4):658-670.
 98. Asai K, Kuzuya M, Naito M, Funaki C, Kuzuya F. Effects of splenectomy on serum lipids and experimental atherosclerosis. *Angiology*. 1988;39(6):497-504.
 99. Fatouros M, Bourantas K, Bairaktari E, Elisaf M, Tsolas O, Cassiouis D. Role of the spleen in lipid metabolism. *Br J Surg*. 1995;82(12):1675-1677.
 100. Caligiuri G, Nicoletti A, Poirier B, Hansson GK. Protective immunity against atherosclerosis carried by B cells of hypercholesterolemic mice. *J Clin Invest*. 2002;109(6):745-753.
 101. Johnsson R, Saris NE. Plasma and erythrocyte lipids in hereditary spherocytosis. *Clin Chim Acta*. 1981;114(2-3):263-268.
 102. Sasaki J, Waterman MR, Buchanan GR, Cottam GL. Plasma and erythrocyte lipids in sickle cell anaemia. *Clin Lab Haematol*. 1983;5(1):35-44.
 103. Atac B, Brahad J, Frishman WH, Lerner R. Anemia and hypocholesterolemia. *Heart Dis*. 2003;5(1):65-71.
 104. Shalev H, Kapelushnik J, Moser A, Knobler H, Tamary H. Hypocholesterolemia in chronic anemias with increased erythropoietic activity. *Am J Hematol*. 2007;82(3):199-202.
 105. Cray SE, Troendle SB, Torres R, Ahmad N, Buchanan GR. Prospective assessment of laboratory markers of cardiovascular risk in hereditary spherocytosis. http://www3.interscience.wiley.com/homepages/106561790/ASPHO_Abstracts_-_2008.pdf. 2008;Poster 199:35. Accessed July 2009.
 106. Eldor A, Maclouf J, Lellouche F, et al. A chronic hypercoagulable state and life-long platelet activation in beta thalassaemia major. *Southeast Asian J Trop Med Public Health*. 1993;24[suppl 1]:92-95.
 107. Atichartakam V, Angchaisuksiri P, Aryurachai K, Chuncharunee S, Thakkinian A. In vivo platelet activation and hyperaggregation in hemoglobin E/beta-thalassaemia: a consequence of splenectomy. *Int J Hematol*. 2003;77(3):299-303.
 108. Oparattikul N, Funahara Y, Fucharoen S, Talalak P. Increase in spontaneous platelet aggregation in beta-thalassaemia/hemoglobin E disease: a consequence of splenectomy. *Southeast Asian J Trop Med Public Health*. 1992;23[suppl 2]:36-41.
 109. Winichagoon P, Fucharoen S, Wasi P. Increased circulating platelet aggregates in thalassaemia. *Southeast Asian J Trop Med Public Health*. 1981;12(4):556-560.
 110. Hovav T, Goldfarb A, Artmann G, Yedgar S, Barshtein G. Enhanced adherence of beta-thalassaemic erythrocytes to endothelial cells. *Br J Haematol*. 1999;106(1):178-181.
 111. Butthep P, Bunyaratvej A, Funahara Y, et al. Possible evidence of endothelial cell activation and disturbance in thalassaemia: an in vitro study. *Southeast Asian J Trop Med Public Health*. 1997;28[suppl 3]:141A-148A.
 112. Eldor A, Durst R, Hy-Am E, et al. A chronic hypercoagulable state in patients with beta-thalassaemia major is already present in childhood. *Br J Haematol*. 1999;107(4):739-746.
 113. Fontana V, Jy W, Dudkiewicz P, Horstman LL, Bidot CJ, Ahn YS. Cell-derived microparticles (C-MP) are elevated in splenectomized patients and depressed in those with hypersplenism, suggesting spleen as scavenger of C-MP. *Blood*. 2006;108(11):1482.
 114. Fontana V, Jy W, Ahn ER, et al. Increased procoagulant cell-derived microparticles (C-MP) in splenectomized patients with ITP. *Thromb Res*. 2008;122(5):599-603.
 115. Westerman M, Pizzey A, Hirschman J, et al. Microvesicles in haemoglobinopathies offer insights into mechanisms of hypercoagulability, haemolysis and the effects of therapy. *Br J Haematol*. 2008;142(1):126-135.
 116. Habib A, Kunzelmann C, Shamseddeen W, Zobairi F, Freyssinet JM, Taher A. Elevated levels of circulating procoagulant microparticles in patients with B-thalassaemia intermedia. *Haematologica*. 2008;93(6):941-942.
 117. Rahimi Z, Ghaderi M, Nagel RL, Muniz A. Prevalence of thrombotic risk factors among beta-thalassaemia patients from Western Iran. *J Thromb Thrombolysis*. 2008;26(3):229-233.
 118. Ataga KI, Cappellini MD, Rachmilewitz EA. Beta-thalassaemia and sickle cell anaemia as paradigms of hypercoagulability. *Br J Haematol*. 2007;139(1):3-13.
 119. Eldor A, Rachmilewitz EA. The hypercoagulable state in thalassaemia. *Blood*. 2002;99(1):36-43.
 120. Ataga KI, Key NS. Hypercoagulability in sickle cell disease: new approaches to an old problem. *Hematology Am Soc Hematol Educ Program*. 2007:91-96.

121. Cappellini MD. Coagulation in the pathophysiology of hemolytic anemias. *Hematology Am Soc Hematol Educ Program*. 2007;74-78.
122. Borenstein-Ben Yashar V, Barenholz Y, Hy-Am E, Rachmilewitz EA, Eldor A. Phosphatidylserine in the outer leaflet of red blood cells from beta-thalassemia patients may explain the chronic hypercoagulable state and thrombotic episodes. *Am J Hematol*. 1993;44(1):63-65.
123. de Jong K, Larkin SK, Eber S, Franck PF, Roelofsen B, Kuypers FA. Hereditary spherocytosis and elliptocytosis erythrocytes show a normal transbilayer phospholipid distribution. *Blood*. 1999;94(1):319-325.
124. Horne MK 3rd, Cullinane AM, Merryman PK, Hoddeson EK. The effect of red blood cells on thrombin generation. *Br J Haematol*. 2006;133(4):403-408.
125. Stief TW. Thrombin generation by hemolysis. *Blood Coagul Fibrinolysis*. 2007;18(1):61-66.
126. Robertson DA, Simpson FG, Losowsky MS. Blood viscosity after splenectomy. *Br Med J (Clin Res Ed)*. 1981;283(6291):573-575.
127. Murphy JR. The influence of pH and temperature on some physical properties of normal erythrocytes and erythrocytes from patients with hereditary spherocytosis. *J Lab Clin Med*. 1967;69(5):758-775.
128. Baskurt OK. The role of spleen in suppressing the rheological alterations in circulating blood. *Clin Hemorheol Microcirc*. 1999;20(3):181-188.
129. Peacock AJ. Pulmonary hypertension after splenectomy: a consequence of loss of the splenic filter or is there something more? *Thorax*. 2005;60(12):983-984.
130. Rother RP, Bell L, Hillmen P, Gladwin MT. The clinical sequelae of intravascular hemolysis and extracellular plasma hemoglobin: a novel mechanism of human disease. *JAMA*. 2005;293(13):1653-1662.
131. Schilling RF, Gangnon RE, Traver M. Arteriosclerotic events are less frequent in persons with chronic anemia: evidence from families with hereditary spherocytosis. *Am J Hematol*. 2006;81(5):315-317.
132. Wang CH, Schilling RF. Myocardial infarction and thalassemia trait: an example of heterozygote advantage. *Am J Hematol*. 1995;49(1):73-75.
133. Tassiopoulos T, Stamatelos G, Zakopoulos N, Fessas P, Eliopoulos GD. Low incidence of acute myocardial infarction in beta-thalassaemia trait carriers. *Haematologia (Budap)*. 1995;26(4):199-203.
134. Tassiopoulos S, Deftereos S, Konstantopoulos K, et al. Does heterozygous beta-thalassemia confer a protection against coronary artery disease? *Ann N Y Acad Sci*. 2005;1054:467-470.
135. Cocco P, Todde P, Fornera S, Manca MB, Manca P, Sias AR. Mortality in a cohort of men expressing the glucose-6-phosphate dehydrogenase deficiency. *Blood*. 1998;91(2):706-709.
136. Dessi S, Batetta B, Spano O, et al. Serum lipoprotein pattern as modified in G6PD-deficient children during haemolytic anaemia induced by fava bean ingestion. *Int J Exp Pathol*. 1992;73(2):157-160.
137. Muntoni S, Batetta B, Dessi S, Pani P. Serum lipoprotein profile in the Mediterranean variant of glucose-6-phosphate dehydrogenase deficiency. *Eur J Epidemiol*. 1992;8[suppl 1]:48-53.
138. Morita T. Heme oxygenase and atherosclerosis. *Arterioscler Thromb Vasc Biol*. 2005;25(9):1786-1795.
139. Martin CR, Johnson CS, Cobb C, Tatter D, Haywood LJ. Myocardial infarction in sickle cell disease. *J Natl Med Assoc*. 1996;88(7):428-432.
140. Bonderman D, Skoro-Sajer N, Jakowitsch J, et al. Predictors of outcome in chronic thromboembolic pulmonary hypertension. *Circulation*. 2007;115(16):2153-2158.
141. Raiesdana A, Loscalzo J. Pulmonary arterial hypertension. *Ann Med*. 2006;38(2):95-110.
142. Cohen AR, Galanello R, Pennell DJ, Cunningham MJ, Vichinsky E. Thalassemia. *Hematology Am Soc Hematol Educ Program*. 2004:14-34.