

Chronic myeloid leukemia: a prospective comparison of interphase fluorescence in situ hybridization and chromosome banding analysis for the definition of complete cytogenetic response: a study of the GIMEMA CML WP

Nicoletta Testoni,¹ Giulia Marzocchi,¹ Simona Luatti,¹ Marilina Amabile,¹ Carmen Baldazzi,¹ Monica Stacchini,¹ Mauro Nanni,² Giovanna Rege-Cambrin,³ Emilia Giugliano,³ Ursula Giussani,⁴ Elisabetta Abruzzese,⁵ Simonetta Kerim,⁶ Maria Grazia Grimoldi,⁷ Alessandro Gozzetti,⁸ Barbara Crescenzi,⁹ Carlo Carcassi,¹⁰ Paolo Bernasconi,¹¹ Antonio Cuneo,¹² Francesco Albano,¹³ Giuseppina Fugazza,¹⁴ Alfonso Zaccaria,¹⁵ Giovanni Martinelli,¹ Fabrizio Pane,¹⁶ Gianantonio Rosti,¹ and Michele Baccarani¹

¹Department of Hematology-Oncology, L and A Seràgnoli, University of Bologna and S Orsola-Malpighi University Hospital, Bologna; ²Department of Cellular Biotechnologies and Hematology, La Sapienza University, Roma; ³Department of Clinical and Biological Sciences, University of Torino, San Luigi Hospital, Orbassano; ⁴Medical Genetic Laboratory, Ospedali Riuniti, Bergamo; ⁵Department of Hematology, S Eugenio, Tor Vergata University Hospital, Roma; ⁶Department of Biomedical Sciences and Human Oncology, San Giovanni Hospital, University of Torino, Torino; ⁷Department of Medicine, Pathology Unit, University of Milano, Milano; ⁸Department of Medicine and Immunological Sciences, Division of Hematology and Transplants, University of Siena, Siena; ⁹Institute of Haematology, University of Perugia, Perugia; ¹⁰Department of Medical Sciences, Medical Genetics, University of Cagliari, Cagliari; ¹¹Institute of Hematology, Fondazione Policlinico San Matteo Istituto Di Ricovero e Cura a Carattere Scientifico, University of Pavia, Pavia; ¹²Department of Biomedical Sciences and Advanced Therapies, University of Ferrara, Ferrara; ¹³Division of Hematology, University of Bari, Bari; ¹⁴Department of Internal Medicine, University of Genova, Genova; ¹⁵Department of Oncology and Hematology, Santa Maria delle Croci Hospital, Ravenna; and ¹⁶Department University Federico II, Napoli, Italy

In chronic myeloid leukemia, different methods are available to monitor the response to therapy: chromosome banding analysis (CBA), interphase fluorescence in situ hybridization (I-FISH), and real-time quantitative polymerase chain reaction (RT-Q-PCR). The GIMEMA CML WP (Gruppo Italiano Malattie Ematologiche Adulto Chronic Myeloid Leukemia Working Party) has performed a prospective study to compare CBA and I-FISH for the definition of complete cytogenetic re-

sponse (CCgR). Samples (n = 664) were evaluated simultaneously by CBA and I-FISH. Of 537 cases in CCgR, the number of positive nuclei by I-FISH was less than 1% in 444 cases (82.7%). Of 451 cases with less than 1% positive nuclei by I-FISH, 444 (98.4%) were classified as CCgR by CBA. The major molecular response rate was significantly greater in cases with I-FISH less than 1% than in those with I-FISH 1% to 5% (66.8% vs 51.6%, $P < .001$) and in cases with CCgR

and I-FISH less than 1% than in cases with CCgR and I-FISH 1% to 5% (66.1% vs 49.4%, $P = .004$). I-FISH is more sensitive than CBA and can be used to monitor CCgR. With appropriate probes, the cut-off value of I-FISH may be established at 1%. These trials are registered at <http://www.clinicaltrials.gov> as NCT00514488 and NCT00510926. (Blood. 2009;114:4939-4943)

Introduction

Chronic myeloid leukemia (CML) is a clonal hematopoietic stem cell disorder induced by a chimeric gene that results from the fusion of the *ABL* gene on chromosome 9 with the *BCR* gene on chromosome 22, leading to the formation of a new leukemia-specific fusion gene that codes for constitutionally activated protein tyrosine kinases (PTK) of different molecular weight (p210, rarely p230 or p185/190). The oncogenic PTK, which is located in the cytoplasm, is responsible of the leukemic phenotype through the constitutive activation of several downstream pathways.¹⁻³ In more than 95% of cases, the translocation between chromosome 9 and 22 is balanced and results in the formation of a small chromosome 22 that was identified originally as Philadelphia (Ph). Rare variant translocations may be masked and detected by fluorescence in situ hybridization of interphase nuclei (I-FISH).²⁻⁵ The *BCR-ABL* mRNA is detected by reverse transcriptase polymerase chain reaction (PCR) and may be quantified by real-time quantitative PCR (RT-Q-PCR).⁶⁻⁸

After the introduction of interferon- α and the PTK inhibitor imatinib mesylate (IM), it has become more and more important to monitor cytogenetically the response to treatment and the course of the disease.⁹⁻¹³ On the basis of chromosome banding analysis (CBA) of marrow cell metaphases, the cytogenetic response (CgR) is classified as none, minimal, minor, partial, or complete according to the percentage of Ph⁺ metaphases (95%, 95%-66%, 65%-36%, 35%-1%, and none).¹⁴ The achievement and the maintenance of a complete cytogenetic response (CCgR) are of particular importance because a CCgR is the most solid, confirmed, early surrogate marker of progression-free survival and overall survival.¹¹⁻¹⁶ The definition of CgR by CBA requires marrow cells, which cannot be always sampled, and an adequate number of banded metaphases, which cannot be always obtained. For these reasons, I-FISH is sometimes used, with increasing frequency, as a substitute for CBA, but although there is a fairly good relationship between I-FISH and CBA data,¹⁷⁻²⁸ there are no controlled and shared

Submitted July 13, 2009; accepted September 6, 2009. Prepublished online as *Blood* First Edition paper, October 1, 2009; DOI 10.1182/blood-2009-07-229864.

The online version of this article contains a data supplement.

The publication costs of this article were defrayed in part by page charge payment. Therefore, and solely to indicate this fact, this article is hereby marked "advertisement" in accordance with 18 USC section 1734.

Table 1. Distribution of I-FISH data according to CBA data

Cytogenetic response by CBA	Cytogenetic response by I-FISH, n (%)		
	Less than 1% <i>BCR-ABL</i> ⁺ nuclei	1%-5% <i>BCR-ABL</i> ⁺ nuclei	More than 5% <i>BCR-ABL</i> ⁺ nuclei
CCgR (n = 537), no Ph ⁺ metaphases	444 (82.7)	71 (13.2)	22 (4.1)
PCgR (n = 77), 1%-35% Ph ⁺ metaphases	7 (9.1)	32 (41.6)	38 (49.3)
<i>P</i>	< .001	< .001	< .001

CBA indicates chromosome banding analysis; CCgR, complete cytogenetic response; I-FISH, interphase fluorescence in situ hybridization; and PCgR, partial cytogenetic response.

definitions of CgR by I-FISH. This study was designed with the purpose of comparing CBA and I-FISH data for the definition of CCgR.

Methods

Patients

All the patients, at least 18 years of age, were required to have a Ph⁺ and *BCR-ABL*⁺ CML in early chronic phase (CP). They were enrolled in 3 prospective and concurrent studies: CML/021 (ClinTrials.gov no. NCT00514488), a phase 2 trial exploring IM 800 mg daily in intermediate Sokal risk patients with CP CML; CML/022 (ClinTrials.gov no. NCT00510926), a phase 3 trial comparing IM 400 versus 800 mg daily in high Sokal risk patients with CP CML; and CML/023, an observational study of IM 400 mg daily in patients with CP CML. These studies were promoted, sponsored, and operated by the CML Working Party of GIMEMA (previously the Italian Cooperative Study Group on CML). These studies included a prospective comparison of CBA and I-FISH data on marrow cells after 6 and 12 months of therapy. The studies were approved by the independent ethics committee of S. Orsola-Malpighi Bologna University Hospital and the ethic committees of all participating institutions and were operated according to Good Clinical Practices and the Declaration of Helsinki. Written informed consent was required and provided by all patients. The bone-marrow samples of 68 non-CML patients, who had given informed consent to cytogenetic studies for diagnostic purposes, also were analyzed by the use of I-FISH in the Bologna laboratory.

Cytogenetics: CBA

The GIMEMA CML Working Party has established a network of cytogenetic laboratories throughout Italy (supplemental Table 1, available on the *Blood* website; see the Supplemental Materials link at the top of the online article). In 14 of these laboratories, researchers performed CBA and FISH studies for more than one clinical center (reference laboratories), and researchers in 24 laboratories performed CBA and FISH studies only for their respective clinical center. In most laboratories, CBA and I-FISH were performed by different technicians. All studies were performed on unfractionated bone marrow cells, which were referred to the laboratories within 24 hours from sampling. CBA was performed after short-term culture (24 and/or 48 hours). The cells were treated with colchicine and hypotonic solution, then they were centrifuged, and the resulting pellet was fixed and washed in methanol/acetic acid (3:1). The cells were resuspended in fixative and dropped on slides. Karyotypes were examined after the G banding technique and described according to International System for Human Cytogenetic Nomenclature (ISCN 1995).²⁹ At least 20 metaphases per sample were analyzed in 87% of samples and 10 to 19 metaphases in 13% of samples. A central review was not performed.

I-FISH analysis

I-FISH was performed on marrow cells prepared according to the technique described previously and by the use of DNA probes that hybridize to *BCR* and *ABL* regions. The probes were commercially available. All labs used *BCR-ABL* Extra-Signal (ES; Vysis-Abbott), Dual-Color Dual-Fusion (DCDF; Vysis-Abbott), or Dual-Fusion FISH (D-FISH; Q-Biogene-Onco)

probes. Only one laboratory used a "home-brew" PAC/BAC system: a pool of PAC, RP5-1132H12, and RP5-835J22, for *ABL* gene and BAC, RP11-164n13 for *BCR* gene.⁵ Dual-color single-fusion probes were never used in this study because they generate only one fusion signal in Ph⁺ cells, and therefore cutoff values may be remarkably high. The number of nuclei analyzed by I-FISH ranged between 166 and 2677 (median, 300; mean, 298).

Molecular studies

Qualitative reverse transcription polymerase chain reaction (PCR) for *BCR-ABL* transcript was routinely performed at enrollment for determining the type of transcript. Peripheral blood samples for real-time quantitative PCR (RT-Q-PCR) were collected before therapy; after 3, 6, and 12 months; and every 6 months thereafter. All samples and tests were centralized in Bologna. Whole buffy-coat cells were used. RNA extraction, reverse transcription, RT-Q-PCR were performed according to European recommendations, as previously described.³⁰⁻³³ RT-Q-PCR was performed on the ABI PRISM 7700 Sequence Detector (Perkin Elmer).³⁴ *ABL* was used as a housekeeping gene to correct differences in RNA quality and/or reverse transcription efficacy. *BCR-ABL* and *ABL* plasmid dilutions (Ipsogen Inc) were used as standards, and the final results were calculated as the ratios *BCR-ABL* to *ABL* and expressed in percentages. All experiments were performed in duplicate, and the results were expressed as percent ratio to *ABL*. The *BCR-ABL/ABL* ratios were further multiplied by the conversion factor of the Bologna laboratory to set the results on an international scale.^{8,11,15,34} Samples yielding an *ABL* threshold cycle greater than 30, corresponding to less than 1000 *ABL* transcript copies, were considered as having degraded RNA and discarded. We defined major molecular response (MMoR) as a *BCR-ABL/ABL* ratio equal less than 0.1%.^{11,13,14}

Statistical analysis

The statistical significance of differences was assessed with the Student *t* test and the Mann-Whitney *U* test of continuous variables and with the Fisher exact test for categorical variables. The GraphPad Prism 4 (GraphPad Software Inc) was used throughout.

Results

Five hundred sixty-seven patients were studied at baseline; 515 of them (90.8%) were evaluable cytogenetically. Thirty had a variant translocation; 665 marrow samples were evaluable for CBA and I-FISH at the same time; and 614 were in complete or partial CgR (PCgR) by standard CBA-based definition and were analyzed for comparison of CBA and I-FISH data. In addition, 512 of these 614 samples (83.4%) were evaluable for *BCR-ABL* transcripts level at the same time and were analyzed for comparison of CBA, I-FISH, and RT-Q-PCR data. In these samples, the number of metaphases analyzed by CBA ranged between 10 and 100 (median, 20; mean, 24), and the number of nuclei analyzed for I-FISH ranged between 166 and 2677 (median, 300; mean, 298).

First, we analyzed I-FISH data according to CBA data (Table 1). There were 537 cases of CCgR by CBA, of which 444 (82.7%) had

Table 2. Distribution of CBA data according to I-FISH data

Cytogenetic response by I-FISH	Cytogenetic response by CBA, n (%)	
	No Ph ⁺ metaphases, CCgR	1%-35% Ph ⁺ metaphases, PCgR
Less than 1% <i>BCR-ABL</i> ⁺ nuclei (n = 451)	444 (98.4) <i>P</i> < .001	7 (1.6) <i>P</i> < .001
1%-5% <i>BCR-ABL</i> ⁺ nuclei (n = 103)	71 (68.9) <i>P</i> < .001	32 (31.5) <i>P</i> < .001
More than 5% <i>BCR-ABL</i> ⁺ nuclei (n = 60)	22 (36.7)	38 (63.3)

CBA indicates chromosome banding analysis; CCgR, complete cytogenetic response; I-FISH, interphase fluorescence in situ hybridization; and PCgR, partial cytogenetic response.

less than 1% positive nuclei by I-FISH, but 71 (13.2%) had 1% to 5% positive nuclei, and 22 (4.1%) even more than 5% positive nuclei. In contrast, there were 77 cases of PCgR by CBA, of which only 7 (9.1%) had less than 1% positive nuclei by I-FISH, whereas 32 (41.6%) had 1% to 5% positive nuclei, and 38 (49.3%) had more than 5% positive nuclei. The data show that a substantial minority (17.3% overall) of cases who were classified as CCgR by CBA had 1% or greater positive nuclei by I-FISH, whereas only few cases who were classified as PCgR by CBA had less than 1% positive nuclei by I-FISH.

Second, we analyzed CBA data according to I-FISH data (Table 2). There were 451 cases with less than 1% positive nuclei by I-FISH, of whom almost all (98.4%) had no Ph⁺ metaphases. In contrast, 68.9% and 36.7% of cases with 1% to 5% or more than 5% positive nuclei, respectively had no Ph⁺ metaphases.

Third, we compared the MoIR of the patients classified as CCgR by CBA (100% Ph⁻) with the MoIR of the cases with less than 1% positive nuclei by I-FISH. The proportion of MMoIR was 63.3% in the former and 66.8% (*P* = .24) in the latter, showing no significant differences between the group defined by CBA and the group defined by I-FISH. However, the percentage of MMoIR was significantly greater in the 425 cases with no Ph⁺ metaphases and less than 1% I-FISH-positive nuclei, than the 87 cases with no Ph⁺ metaphases CCgR and more than 1% I-FISH-positive nuclei (66.1% vs 49.4%, *P* = .004; Table 3).

Next, we examined the distribution of MMoIR according to the number of positive nuclei by I-FISH (Table 3) There were no differences between the cases with no positive nuclei and those with 0.1% to 0.9% positive nuclei, whereas in the cases with 1% to 5% positive nuclei the percentage of MMoIR was significantly lower and the *BCR-ABL* transcript level was significantly greater.

Table 3. Distribution of MoIR according to CBA and I-FISH data

Cytogenetic response by I-FISH and by CBA	Molecular response			
	Major MoIR, n (%)	<i>P</i>	Median <i>BCR-ABL</i> transcripts level*	<i>P</i>
No Ph ⁺ metaphases and < 1% <i>BCR-ABL</i> ⁺ nuclei (n = 425)	281 (66.1)	.004	0.035	.06
	43 (49.4)		0.079	
No <i>BCR-ABL</i> ⁺ nuclei (n = 396)	263 (66.4)	.68	0.031	.43
	21 (72.4)		0.043	
	49 (51.6)		0.082	
0.1%-0.9% <i>BCR-ABL</i> ⁺ nuclei (n = 29)		< .001		< .001
1%-5% <i>BCR-ABL</i> ⁺ nuclei (n = 95)				

CBA indicates chromosome banding analysis; I-FISH, interphase fluorescence in situ hybridization; and MoIR, molecular response. **BCR-ABL*/*ABL*% on the international scale.

Table 4. Control non-CML samples

<i>BCR-ABL</i> ⁺ nuclei by I-FISH	Samples, n (%)	Positive nuclei, n (%)
0 (none)	63/68 (92.6)	0/18 900
0.1%-0.9%	2/68 (2.9)	3/600 (0.5)
1%-5%	3/68 (4.4)	13/900 (1.4)
More than 5%	0/68	0/0

Positive nuclei were recorded in 5 of 68 samples, but the positivity was greater than 1% in only 3 of 68 samples.

I-FISH indicates interphase fluorescence in situ hybridization.

The total number of cases classified as MMoIR was 333, of which 263 (79%) had no positive nuclei at I-FISH, 21 (6.3%) had 0.1% to 0.9% positive nuclei, and 49 (14.7%) had 1% to 5% positive nuclei. No cases of MMoIR were found among the cases with more than 5% positive nuclei.

I-FISH was performed at one laboratory (Bologna) also on 68 selected *BCR-ABL*⁻ bone-marrow samples, scoring 300 nuclei in each sample (Table 4). The percentage of positive nuclei was 1% or greater (1%, 1.3%, and 2%) in 3 samples, was less than 1% (0.3% and 0.7%) in 2 samples, and was 0 in 63 samples. The mean percentage of *BCR-ABL*⁺ nuclei was 0.08 (SD = 0.3). Thus a positivity cutoff value was fixed at 1% (mean + 3 SD).

Discussion

Progress in treatment of Ph⁺ CML has been always measured by the degree of CgR. The degree of CgR is assessed by the use of CBA of marrow cell metaphases, based on the percentage of Ph⁺ metaphases.⁹⁻¹³ Although the number of banded metaphases that are conventionally required is small (n = 20),^{13,14} the value of the different degrees of CgR, which were established during interferon-α studies, also has been validated with IM, and achieving a CCgR is still the most robust early surrogate of the outcome of therapy.¹¹⁻¹⁶ The authors of several reports¹⁷⁻²⁸ pointed out that the relationship between CBA and I-FISH data was significant and excellent but they did not allow one to translate the figures that define CgR with CBA data into those that should define CgR with I-FISH data. A percentage of Ph⁺ metaphases ranging between 1% and 35% defines a CgR as partial, but the same figures cannot be applied by use of the percentage of *BCR-ABL*⁺ nuclei. However, the 2 techniques can be concordant in the most important definition of CgR, that is, CCgR, corresponding to the absence of Ph⁺ metaphases of a total of at least 20 banded metaphases.

Table 5. Cutoff values for “false *BCR-ABL*⁺ cells,” as reported in previous studies, by the use of I-FISH

References	Probe used	Cutoff	Samples
Dewald et al, 1998 ¹⁹	D-FISH	Not identified*	35 normal subjects
Buño et al, 1998 ²⁰	D-FISH	0.8%	PBL from 10 patients without any malignant hematologic disorder and BM from 10 normal bone marrow transplant donors
Chomel et al, 2000 ⁹	D-FISH	1%†	PBL or BM from 5 patients with hematologic disorders other than CML
Le Gouill et al, 2000 ²²	ES	1.2%†	14 BM from AML or <i>BCR/ABL</i> ⁻ myeloproliferative syndrome patients and 10 PBL from healthy donors
Chase et al, 2000 ³⁵	DCDF	1%‡	Not specified
Primo et al, 2003 ³⁶	ES	0.9%§	20 <i>BCR/ABL</i> ⁻ BM samples
Raanani et al, 2004 ²⁶	ES	1%	Not specified
El-Zimaity et al, 2004 ³⁷	ES	1.2%†	20 normal control samples

Landstrom et al,²⁷ in a study of Ph⁺ CML, used a cutoff of 0.6%, citing Dewald et al.¹⁹

AML indicates acute myeloid leukemia; D-FISH, dual FISH (from Q-Biogene); DCDF, dual-color dual-fusion (from Vysis-Abbott); ES, extra-signal FISH (from Vysis-Abbott); BM, bone marrow; PBL, peripheral blood.

*Based on 200 nuclei, the mean percentage of false-positive cells was 0.25 ± 0.39 . The normal cutoff for 6000 nuclei (for each of 3 normal specimens) was 0.079%.”

†Mean + 3 SD.

‡Although in experienced hands with good quality preparations the technique can give a very low false positive rate of 0.1%-0.2%, in practice we prefer to use cutoff rate of 1% because accuracy is seriously compromised by several factors, including slide quality and operator experience.”

§Mean + 2SD.

||Because the FISH (ES) method has false positive rate of 1%, results were considered negative when < 1% of interphases nuclei were positive for *BCR-ABL* double-labeled probes.”

Historically, the definition of the number of *BCR-ABL*⁺ cells by I-FISH was disturbed by the use of single fusion probes, which may lead to false-positive data, and tends to overestimate the number of positive nuclei, compelling one to establish a cutoff value of 5% or more. In this study, as in all other recent studies,^{9,19,20,22,24-27,35-37} only Dual Fusion (DF) and ES FISH strategies were used. The DF strategy uses probes that span the common breakpoints in the *ABL* and *BCR* gene regions, generating 2 novel fusion signals caused by reciprocal t(9;22): 1 on the derivative chromosome 9 and 1 on the derivative of chromosome 22. Thus, cells with a classical t(9;22) will display a FISH pattern with 1 red (normal 9 chromosome), 1 green (normal 22 chromosome), and 2 “yellow” fusion signals (derivative 9 and 22 chromosomes with *BCR-ABL* and *ABL-BCR* fusions). The detection of 2 fusion signals can virtually exclude false-positive nuclei. With the ES FISH strategy, the Ph⁺ cells display 2 red (normal and derivative 9 chromosome), 1 green (normal 22 chromosome), and 1 yellow fusion signals (derivative 22 chromosome).

Using these strategies, we examined 20 400 interphase nuclei in 68 non-CML marrow samples. We found only 16 positive nuclei (0.078%) and only 3 samples (4.4%) with more than 1% positive nuclei. Table 5 lists the results of 8 independent studies, reporting the *BCR/ABL* positivity detected by DF or ES FISH in non-CML samples.^{9,19,20,22,26,35-37} The cutoff values that were proposed by these studies ranged between 0.8% and 1.2% (median, 1%). Therefore, we conclude that a cutoff value of 1% is fairly reasonable, even conservative, and may prevent any overestimate of the completeness of the response.

In this prospective comparative study of CBA versus I-FISH, almost all the cases (98.4%) below the cutoff value of 1% were CCgR by CBA. On the opposite, 13.2% and 4.1% of the samples that were defined as CCgR by CBA had 1% to 5% and more than 5% positive nuclei, respectively. These data suggest that I-FISH may be more sensitive than CBA for the assessment of minimal residual disease, which is not surprising, considering that more cells are examined with I-FISH than with CBA. As a matter of fact, the rate of MMolR was greater and the *BCR/ABL* transcripts level was lower in the patients who were negative by I-FISH and CBA than in those who were negative only by CBA (Table 3).

I-FISH cannot be used to assess all the different degrees of the response, from minimal to partial; although the relationship between the percentage of Ph⁺ metaphases and the percentage of *BCR-ABL*⁺ nuclei may be significant, there are no data showing that the percentages are the same. However, I-FISH can be used to substitute for metaphase CBA, once all the metaphases are Ph⁻, that is to say once the CgR is defined as complete by CBA. The cost of the reagents is greater for I-FISH, but CBA is technically more demanding and requires specifically and well-trained technicians. Moreover, the use of I-FISH will allow monitoring the completeness of the response better than CBA if the number of metaphases is small. In this study, I-FISH was performed on marrow cells, but the authors of several studies^{20-22,24,26,27} have already shown that there are no differences in I-FISH results between marrow and blood samples.

Taking into account that I-FISH may be more sensitive than CBA because it also correlates better with MolR, the detection of *BCR-ABL*⁺ cells by I-FISH always requires a confirmatory test with CBA before one can conclude that a CCgR has been lost. CBA also is required to identify additional chromosome abnormalities in Ph⁺, in case of response loss, and to identify other chromosome abnormalities in Ph⁻ cells, in case of hematologic abnormalities, suggesting the development of a myelodysplastic condition.

Whether I-FISH may be the preferred technique for monitoring CML patients who are in CCgR is still a matter of debate and also depends on the availability and the reproducibility of the techniques for the quantification of *BCR-ABL* transcripts level.

Acknowledgments

We thank Katia Vecchi for skillful assistance.

This work was supported by grants from the Italian Association Against Leukemia–Lymphoma and Myeloma (AIL), the Fondazione del Monte di Bologna e Ravenna, the Italian Ministry of Education (PRIN 2005 and PRIN 2007), the

University of Bologna, the European Union (European LeukemiaNet), and the IRST (Istituto Romagnolo per lo Studio e la Cura dei Tumori).

Authorship

Contribution: N.T. coordinated all cytogenetics studies with the contribution of G. Marzocchi, S.L., C.B., M.S., M.N., G.R.-C., E.G., U.G., E.A., S.K., M.G.G., A.G., B.C., C.C., P.B., A.C., F.A., G.F., and A.Z; M.A. coordinated the molecular studies;

G.R. coordinated the clinical studies with the contribution of G. Martinelli and F.P.; and N.T. and M.B. analyzed the data and wrote the report, which was reviewed by all centers.

Conflict-of-interest disclosure: The authors declare no competing financial interests.

For a complete list of GIMEMA CML WP participants, see the supplemental Appendix, available on the *Blood* website.

Correspondence: Nicoletta Testoni, BS, Department of Hematology-Oncology, L and A Seràgnoli, S Orsola-Malpighi Hospital, Via Massarenti 9, 40138 Bologna, Italy; e-mail: nicoletta.testoni@unibo.it.

References

- Rowley JD. A new consistent chromosomal abnormality in chronic myelogenous leukaemia identified by quinacrine fluorescence and Giemsa staining. *Nature*. 1973;243(5405):290-293.
- Jabbour E, Kantarjian H. Chronic myeloid leukemia. *Semin Hematol*. 2007;44(suppl 1):S1-S150.
- Quintás-Cardama A, Cortes J. Molecular biology of BCR-ABL1-positive chronic myeloid leukemia. *Blood*. 2009;113(8):1619-1630.
- Chase A, Huntly BJP, Cross NCP. Cytogenetics of chronic myeloid leukemia. *Best Pract Res Clin Haematol*. 2001;14(3):553-571.
- Albano F, Anelli L, Zagaria A, et al. "Home-brew" FISH assay shows higher efficiency than BCR-ABL dual-color, dual fusion probe in detecting microdeletion and complex rearrangements associated with t(9;22) in chronic myeloid leukemia. *Cancer Genet Cytogenet*. 2007;174(2):121-126.
- Rosti G, Martinelli G, Bassi S, et al. Study Committee, Italian Cooperative Study Group for Chronic Myeloid Leukemia; Writing Committee, Italian Cooperative Study Group for Chronic Myeloid Leukemia. Molecular response to imatinib in late chronic-phase chronic myeloid leukemia. *Blood*. 2004;103(6):2284-2290.
- Cortes J, Talpaz M, O'Brien S, et al. Molecular responses in patients with chronic myelogenous leukemia in chronic phase treated with imatinib mesylate. *Clin Cancer Res*. 2005;11(9):3425-3432.
- Hughes T, Deininger M, Hochhaus A, et al. Monitoring CML patients responding to treatment with tyrosine kinase inhibitors: review and recommendations for harmonizing current methodology for detecting BCR-ABL transcripts and kinase domain mutations and for expressing results. *Blood*. 2006;108(1):28-37.
- Chomel JC, Briard F, Einstein A, et al. Persistence of BCR-ABL genomic rearrangement in chronic myeloid leukemia patients in complete and sustained cytogenetic remission after interferon- α therapy or allogeneic bone marrow transplantation. *Blood*. 2000;95(2):404-408.
- O'Brien SG, Guilhot F, Larson RA, et al. Imatinib compared with interferon and low-dose cytarabine for newly diagnosed chronic-phase chronic myeloid leukemia. *N Engl J Med*. 2003;348(11):994-1004.
- Hughes TP, Kaeda J, Branford S, et al. Frequency of major molecular responses to imatinib or interferon alpha plus cytarabine in newly diagnosed chronic myeloid leukemia. *N Engl J Med*. 2003;349(15):1423-1432.
- Druker BJ, Guilhot F, O'Brien SG, et al. Five-year follow-up of patients receiving imatinib for chronic myeloid leukemia. *N Engl J Med*. 2006;355(23):2408-2417.
- Baccarani M, Pane F, Saglio G. Monitoring treatment of chronic myeloid leukemia. *Haematologica*. 2008;93(2):161-166.
- Baccarani M, Saglio G, Goldman J, et al. Evolving concepts in the management of chronic myeloid leukemia: recommendations from an expert panel on behalf of the European LeukemiaNet. *Blood*. 2006;108(6):1809-1820.
- Branford S, Fletcher L, Cross NC, et al. Desirable performance characteristics for BCR-ABL measurement on an international reporting scale to allow consistent interpretation of individual patient response and comparison of response rates between clinical trials. *Blood*. 2008;112(8):3330-3338.
- Hochhaus A, O'Brien SG, Guilhot F, et al. Six-year follow-up of patients receiving imatinib for the first-line treatment of chronic myeloid leukemia. *Leukemia*. 2009;23(6):1954-1061.
- Cuneo A, Bigoni R, Emmanuel B, et al. Fluorescence in situ hybridization for the detection and monitoring of the Ph-positive clone in chronic myelogenous leukemia: comparison with metaphase banding analysis. *Leukemia*. 1998;12(11):1718-1723.
- Mühlmann J, Thaler J, Hilbe W, et al. Fluorescence in situ hybridization (FISH) on peripheral blood smears for monitoring Philadelphia chromosome-positive chronic myeloid leukemia (CML) during interferon treatment: a new strategy for remission assessment. *Genes Chromosomes Cancer*. 1998;21(2):90-100.
- Dewald GW, Wyatt WA, Juneau AL, et al. Highly sensitive fluorescence in situ hybridization method to detect double BCR/ABL fusion and monitor response to therapy in chronic myeloid leukemia. *Blood*. 1998;91(9):3357-3365.
- Buño I, Wyatt WA, Zinsmeister AR, Dietz Band J, Silver RT, Dewald GW. A special fluorescent in situ hybridization technique to study peripheral blood and assess the effectiveness of interferon therapy in chronic myeloid leukemia. *Blood*. 1998;92(7):2315-2321.
- Yanagi M, Shinjo K, Takeshita A, et al. Simple and reliably sensitive diagnosis and monitoring of Philadelphia chromosome-positive cells in chronic myeloid leukemia by interphase fluorescence in situ hybridization of peripheral blood cells. *Leukemia*. 1999;13(4):542-552.
- Le Gouill S, Talmant P, Milpied N, et al. Fluorescence in situ hybridization on peripheral-blood specimens is a reliable method to evaluate cytogenetic response in chronic myeloid leukemia. *J Clin Oncol*. 2000;18(7):1533-1538.
- Schoch C, Schnitter C, Bursch S, et al. Comparison of chromosome banding analysis, interphase- and hypermetaphase-FISH, qualitative and quantitative PCR for diagnosis and for follow-up in chronic myeloid leukemia: a study on 350 cases. *Leukemia*. 2002;16(1):53-55.
- Lesser M, Dewald GW, Sison CP, Silver RT. Correlation of three methods of measuring cytogenetic response in chronic myelocytic leukemia. *Cancer Genet Cytogenet*. 2002;137(2):79-84.
- Reinhold U, Henning E, Leiblein S, Niederwieser D, Deininger MWN. FISH for BCR-ABL on interphase of peripheral blood neutrophils but not of unselected white bone marrow cytogenetics in CML patients treated with imatinib. *Leukemia*. 2003;17(10):1925-1929.
- Raanani P, Ben-Bassat I, Gan S, et al. Assessment of the response to imatinib in chronic myeloid leukemia patients: comparison between the FISH, multiplex and RT-PCR methods. *Eur J Haematol*. 2004;73(4):243-2550.
- Landstrom AP, Ketterling RP, Knudson RA, Tefferi A. Utility of peripheral blood dual color dual fusion fluorescence in situ hybridization for BCR/ABL fusion to assess cytogenetic remission status in chronic myeloid leukemia. *Leuk Lymphoma*. 2006;47(10):2055-2061.
- Lundán T, Juvonen V, Müller MC, et al. Comparison of bone marrow high mitotic index metaphase fluorescence in situ hybridization to peripheral blood and bone marrow real time quantitative PCR on the International Scale for detecting residual disease in chronic myeloid leukemia. *Haematologica*. 2008;93(2):178-186.
- ISCN 1995. In: Mitelman F, ed. Guidelines for Cancer Cytogenetics, Supplement to: An International System for Human Cytogenetic Nomenclature. Basel: S Karger; 1995.
- van Dongen JJ, Macintyre EA, Gabert JA, et al. Standardized RT-PCR analysis of fusion gene transcripts from chromosome aberrations in acute leukemia for detection of minimal residual disease. Report of the BIOMED-1 Concerted Action: investigation of minimal residual disease in acute leukemia. *Leukemia*. 1999;13(12):1901-1928.
- van der Velden VH, Hochhaus A, Cazzaniga G, et al. Detection of minimal residual disease in hematologic malignancies by real-time quantitative PCR: principles, approaches, and laboratory aspects. *Leukemia*. 2003;17(6):1013-1034.
- Gabert J, Beillard E, van der Velden VH, et al. Standardization and quality control studies of 'real-time' quantitative reverse transcriptase polymerase chain reaction of fusion gene transcripts for residual disease detection in leukemia—a Europe Against Cancer program. *Leukemia*. 2003;17(12):2318-2357.
- Iacobucci I, Saglio G, Rosti G, et al. GIMEMA Working Party on Chronic Myeloid Leukemia. Achieving a major molecular response at the time of a complete cytogenetic response (CCgR) predicts a better duration of CCgR in imatinib-treated chronic myeloid leukemia patients. *Clin Cancer Res*. 2006;12(10):3037-3042.
- Müller MC, Erben P, Saglio G, et al. Harmonization of BCR-ABL mRNA quantification using a uniform multifunctional control plasmid in 37 international laboratories. *Leukemia*. 2008;22(1):96-102.
- Chase A, Parker S, Kaeda J, Sivalingam R, Cross NCP, Goldman JM. Absence of host-derived cells in the blood of patients in remission after allografting for chronic myeloid leukemia. *Blood*. 2000;96(2):777-780.
- Primo D, Tabernero A, Rasillo A, et al. Patterns of BCR/ABL gene rearrangements by interphase fluorescence in situ hybridization (FISH) in BCR/ABL⁺ leukemias: incidence and underlying genetic abnormalities. *Leukemia*. 2003;17(6):1124-1129.
- El-Zimaity MM, Kantarjian H, Talpaz M, et al. Results of imatinib mesylate therapy in chronic myeloid leukaemia with variant Philadelphia chromosome. *Br J Haematol*. 2004;125(2):187-195.