

## Brief report

Prognostic impact of *IDH2* mutations in cytogenetically normal acute myeloid leukemia

Felicita Thol,<sup>1</sup> Frederik Damm,<sup>1</sup> Katharina Wagner,<sup>1</sup> Gudrun Göhring,<sup>2</sup> Brigitte Schlegelberger,<sup>2</sup> Dieter Hoelzer,<sup>3</sup> Michael Lübbert,<sup>4</sup> Wolfgang Heit,<sup>5</sup> Lothar Kanz,<sup>6</sup> Günter Schlimok,<sup>7</sup> Aruna Raghavachar,<sup>8</sup> Walter Fiedler,<sup>9</sup> Hartmut Kirchner,<sup>10</sup> Gerhard Heil,<sup>1,11</sup> Michael Heuser,<sup>1</sup> \*Jürgen Krauter,<sup>1</sup> and \*Arnold Ganser<sup>1</sup>

<sup>1</sup>Department of Hematology, Hemostasis, Oncology, and Stem Cell Transplantation, and <sup>2</sup>Institute of Cell and Molecular Pathology, Hannover Medical School, Hannover; <sup>3</sup>Department of Internal Medicine III, University of Frankfurt, Frankfurt; <sup>4</sup>Department of Hematology-Oncology, University of Freiburg Medical Center, Freiburg; <sup>5</sup>Department of Hematology/Oncology, Ev. Krankenhaus Essen-Werden, Essen; <sup>6</sup>Department of Internal Medicine II, University of Tübingen, Tübingen; <sup>7</sup>Department of Hematology and Oncology, Klinikum Augsburg, Augsburg; <sup>8</sup>Department of Hematology, Oncology and Nephrology, HELIOS Klinikum Wuppertal, Wuppertal; <sup>9</sup>Department of Medicine II, Oncological Center, Hubertus Wald University Cancer Center, University Hospital Hamburg-Eppendorf, Hamburg; <sup>10</sup>Department of Internal Medicine III, Krankenhaus Siloah, Hannover; and <sup>11</sup>Department of Internal Medicine V, Klinikum Lüdenscheld, Germany

Mutations in the nicotinamide adenine dinucleotide phosphate<sup>+</sup>-dependent isocitrate dehydrogenase gene 2 (*IDH2*) have recently been found in patients with acute myeloid leukemia (AML) as well as in patients with leukemic transformation of myeloproliferative neoplasms. We analyzed 272 adult patients with cytogenetically normal AML (CN-AML) for the presence of *IDH2* mutations in codons R140

and R172. *IDH2* mutations of amino acid 140 or 172 could be identified in 12.1% of CN-AML patients, with the majority of mutations (90%) occurring at position R140. The incidence of *IDH2* mutations in AML patients with aberrant karyotypes (n = 130) was significantly lower (3.8%, *P* = .006). *IDH2* mutations were mutually exclusive with mutations in *IDH1*. *IDH2* mutation status alone or in combination

with *IDH1* mutations had no impact on response to therapy, overall survival, and relapse-free survival in patients with CN-AML. In conclusion, *IDH2* mutations are frequently found in CN-AML, but in our analysis these mutations did not influence treatment outcome. This study was registered at [www.clinicaltrials.gov](http://www.clinicaltrials.gov) as #NCT00209833. (*Blood*. 2010;116(4):614-616)

## Introduction

In an attempt to discover unknown molecular alterations in patients with acute myeloid leukemia (AML), whole genome sequencing was performed on AML patients.<sup>1</sup> With this approach, a novel mutation in nicotinamide adenine dinucleotide phosphate<sup>+</sup>-dependent isocitrate dehydrogenase gene (*IDH1*) at codon R132 has been identified<sup>2</sup> previously described in gliomas.<sup>3</sup> Since this first description of *IDH1* mutations in AML, several reports have confirmed that *IDH1* mutations occur in patients with cytogenetically normal AML (CN-AML), with a frequency of 5.5% to 11%.<sup>4-7</sup> A strong association between *IDH1* mutations and the intermediate-risk karyotypes and concurrent *NPM1* mutations has been described.<sup>4-7</sup> The prognostic impact of *IDH1* mutations in AML is currently being investigated.<sup>4-7</sup> A recent study also found mutations in *IDH2* in codon R172 at a low frequency in AML patients.<sup>6</sup> In addition, a novel mutation in *IDH2* in codon R140 could be identified in 2 patients with leukemic transformation of myeloproliferative neoplasm as well as in patients with AML.<sup>8,9</sup> *IDH2* has the same enzymatic activity as *IDH1* but is located in the mitochondrial matrix.

In the present study, we performed a comprehensive analysis of mutations occurring in exon 4 of *IDH2* (including both codons R140 and R172) in 272 patients with CN-AML in the context of other known prognostic markers. These patients were intensively treated with a uniform protocol in 2 consecutive multicenter trials.

Overall, our data indicate that *IDH2* mutations in codons R140 and R172 are frequent but have no prognostic implications in patients with CN-AML when considered alone or in combination with *IDH1* mutations and treated with these intensive protocols.

## Methods

## Patients

We examined 272 patients with CN-AML and 130 AML patients with aberrant karyotypes for mutations in *IDH2*. Of the latter, 45 had a core-binding factor leukemia, 15 had aberrations of chromosome band 11q23, 13 had a complex karyotype, 12 had an isolated trisomy 8, 7 had a monosomy 7, 5 had aberrations of chromosome 3q, 4 had a del(9q), 3 had a t(6;9), and 26 had various other aberrations. All patients were treated within the multicenter treatment trials AML SHG 0295 or AML SHG 0199. Details of the treatment protocols have been previously reported.<sup>10,11</sup> Written informed consent was obtained in accordance with the Declaration of Helsinki, and the study was approved by the Institutional Review Board of Hannover Medical School.

Mutational analysis of *IDH2*

Preparation of mononuclear cells and analysis of prognostic markers were performed as previously reported.<sup>5,12</sup> The genomic region of the *IDH2* gene

Submitted March 2, 2010; accepted April 16, 2010. Prepublished online as *Blood* First Edition paper, April 26, 2010; DOI 10.1182/blood-2010-03-272146.

\*J.K. and A.G. contributed equally to this study.

The online version of this article contains a data supplement.


The publication costs of this article were defrayed in part by page charge payment. Therefore, and solely to indicate this fact, this article is hereby marked "advertisement" in accordance with 18 USC section 1734.

© 2010 by The American Society of Hematology

(exon 4), containing the mutational hotspot codons R140 and R172, was amplified using polymerase chain reaction. Sequences of the primers were as follows: forward 5'-GGGGTCAAATTCTGGTTGA-3', and reverse 5'-CTAGGCGAGGAGCTCCAGT-3'. Purified polymerase chain reaction fragments were directly sequenced as previously described.<sup>12</sup> Point mutations were confirmed in an independent second experiment.

## Results and discussion

*IDH2* mutations were found in 33 of 272 patients (12.1%) with CN-AML (30 patients [11%] in codon R140 and 3 patients [1.1%] in codon R172) and 3.8% of 130 AML patients with aberrant karyotypes. As already described for *IDH1*,<sup>7</sup> *IDH2* mutations were associated with a normal karyotype ( $P = .006$ ), whereas no association with a single chromosomal aberration could be identified for the group of AML patients with aberrant karyotypes (supplemental Table 1, available on the *Blood* Web site; see the Supplemental Materials link at the top of the online article). *IDH2* R140 mutations in CN-AML seem to be similarly common as *IDH1* mutations,<sup>4,5</sup> but *IDH2* mutations in AML appear to be different from those in gliomas where *IDH2* R140 mutations have not been described yet.<sup>3</sup> The majority (31 of 33 patients, 93.9%) of *IDH2* R140 mutations showed a heterozygous conversion of CGG → CAG leading to an arginine to glutamine substitution, whereas one patient had a homozygous CGG → CAG conversion and one patient showed a heterozygous conversion of CGG → CTG leading to an arginine to leucine substitution. Four of 5 patients with mutations in codon 172 had a heterozygous conversion of AGG → AAG leading to an arginine to lysine substitution, whereas one patient showed a heterozygous conversion of AGG → AGT leading to an arginine to serine substitution. By comparing the clinical and hematologic characteristics of *IDH2*-mutated versus unmutated CN-AML patients, apart from a higher platelet count in patients with *IDH2* mutations, no significant difference in relevant markers was found between the 2 groups (supplemental Table 2). However, *IDH2* mutations were mutually exclusive with mutations in *IDH1*. This has also been described in gliomas suggesting overlapping effects of both mutations.<sup>3,9</sup> To determine whether patients lose their *IDH2* mutation in remission, we examined 5 patients with an *IDH2* mutation at the time of diagnosis during the postinduction phase. All 4 patients in continuous remission lost the *IDH2* mutation during remission. However, 1 relapsing patient, who had lost the *IDH2* mutation in remission, was again positive at relapse. An additional patient with mutant *IDH2* at the time of diagnosis was again positive at the time of relapse. These data suggest that *IDH2* mutations are stable during disease progression and might be a suitable marker for minimal residual disease monitoring. To analyze the prognostic impact of *IDH2* mutations in CN-AML, *IDH2* R140 and R172 mutations were combined because of the low incidence of R172 mutations. In univariate and multivariate analysis with adjustment for *NPM1*, *CEBPA*, *WT1* SNP rs16754, and *FLT3-ITD*, *IDH2* mutations had no influence on achieving complete remission. The complete remission rate was 81.8% versus 78.2% in the *IDH2* mutated versus in the *IDH2* WT group, respectively ( $P = .82$ ). Of the 272 patients, 145 died, resulting in a median overall survival (OS) of 46 months. *IDH2* mutations had no impact on OS in univariate or multivariate analysis (Figure 1A, Table 1, supplemental Table 3). In addition, no influence on OS was found for *IDH2* mutations in an exploratory



**Figure 1. Kaplan-Meier curves for OS.** (A) OS in CN-AML patients according to *IDH2* mutations. Dotted line represents patients with mutated *IDH2* (n = 33); and solid line, patients with unmutated *IDH2* (n = 239; log-rank test,  $P = .792$ ). (B) RFS in CN-AML patients according to *IDH2* mutations. Dotted line represents patients with mutated *IDH2* (n = 27); and solid line, patients with unmutated *IDH2* (n = 182) (log-rank test,  $P = .524$ ).

subgroup analysis for the *NPM1*<sup>mut</sup>/*FLT3-ITD*<sup>-</sup> patients (supplemental Figure 1A). Of the 214 patients who achieved complete remission, 98 relapsed and 30 died in complete remission, resulting in a median relapse-free survival (RFS) of 25 months and a 6-year RFS of 40% (95% confidence interval, 37%-43%). *IDH2* also had no impact on RFS in univariate or multivariate analysis (Figure 1B, Table 1, supplemental Table 3). Again, no influence on RFS was found for *IDH2* mutations in an exploratory subgroup analysis for the *NPM1*<sup>mut</sup>/*FLT3-ITD*<sup>-</sup> patients (supplemental Figure 1B). The lack of a prognostic impact of *IDH2* mutations on treatment response and survival is in accordance with results for *IDH1* R132 mutations for which we and others did not find any prognostic implications in CN-AML.<sup>4,5</sup> When analyzing *IDH1*/*IDH2* mutations together, *IDH* mutations remained without prognostic effect. The high frequency of *IDH2* mutations in CN-AML but the missing prognostic implications for this marker proposes that the potential oncogenic effect of the mutation might be overcome by therapy. As already demonstrated for markers, such as *RAS* and *WT1* mutations, the therapy administered can have a significant influence on prognostic markers.<sup>13,14</sup>

In conclusion, we have identified *IDH2* mutations of amino acids 140 and 172 in 12.1% of patients with CN-AML, with the majority of the mutations (90%) occurring in position R140.

**Table 1. Significant variables in multivariate analysis for OS and RFS**

Endpoint/variable	HR	95% CI	P
<b>OS</b>			
Age: above vs below median	2.04	1.41-2.99	< .001
<i>NPM1/FLT3</i> mutation status: low risk* vs high risk	0.66	0.44-0.97	.035
Platelets: above vs below median	0.61	0.42-0.88	.008
<i>WT1</i> SNP rs16754	0.43	0.26-0.71	.001
<b>RFS</b>			
<i>WT1</i> SNP rs16754	0.60	0.39-0.92	.019
<i>NPM1</i> mutation	0.51	0.35-0.74	< .001
<i>CEBPA</i> mutation	0.36	0.19-0.67	.001

Variables with *P* values of less than or equal to .1 in univariate analysis were included for the multivariate analysis. For OS, these included age, *FLT3*-ITD, *WT1* expression, extramedullary disease, platelets, *NPM1/FLT3* status low risk vs others, *CEBPA* mutation, and *WT1* SNP rs16754. For RFS, these included *CEBPA* mutation, *WT1* SNP rs16754, *NPM1* mutation, and *NPM1/FLT3* status low risk vs others. Variables evaluated in univariate analysis (see also supplemental Table 3) included *FLT3*-ITD, age, *WT1* expression, extramedullary disease, platelets high vs low, *NPM1/FLT3* status low risk vs others, *CEBPA* mutation, *WT1* SNP rs16754, *NPM1* mutation, *WT1* mutation, sibling vs no sibling donor, white blood cells high vs low, peripheral blasts high vs low, hemoglobin high vs low, de novo vs secondary AML, sex, *IDH2* mutation, and *IDH1* or *IDH2* mutation combined.

HR indicates hazard ratio; and CI, confidence interval.

\**NPM1/FLT3* low risk is defined as the presence of an *NPM1* mutation in the absence of an *FLT3*-ITD.

Although in our study these mutations had no prognostic impact, further analyses are necessary to define their role in leukemogenesis and their use as a marker for minimal residual disease monitoring.

## References

- Ley TJ, Mardis ER, Ding L, et al. DNA sequencing of a cytogenetically normal acute myeloid leukemia genome. *Nature*. 2008;456(7218):66-72.
- Mardis ER, Ding L, Dooling DJ, et al. Recurring mutations found by sequencing an acute myeloid leukemia genome. *N Engl J Med*. 2009;361(11):1058-1066.
- Yan H, Parsons DW, Jin G, et al. *IDH1* and *IDH2* mutations in gliomas. *N Engl J Med*. 2009;360(8):765-773.
- Chou WC, Hou HA, Chen CY, et al. Distinct clinical and biological characteristics in adult acute myeloid leukemia bearing isocitrate dehydrogenase 1 (*IDH1*) mutation. *Blood*. 2010;115(14):2749-2754.
- Wagner K, Damm F, Göhring G. Impact of *IDH1* R132 mutations and an *IDH1* single nucleotide polymorphism in cytogenetically normal acute myeloid leukemia: SNP rs11554137 is an adverse prognostic factor. *J Clin Oncol*. 2010;28(14):2356-2364.
- Gross S, Cairns RA, Minden MD, et al. Cancer-associated metabolite 2-hydroxyglutarate accumulates in acute myelogenous leukemia with isocitrate dehydrogenase 1 and 2 mutations. *J Exp Med*. 2010;207(2):339-344.
- Schnittger S, Haferlach C, Ulke U, et al. *IDH1* mutations are detected in 9.3% of all AML and are strongly associated with intermediate risk karyotype and unfavourable prognosis: a study of 999 patients [abstract]. *Blood (ASH Annual Meeting Abstracts)*. 2009;114:LBA-3.
- Green A, Beer P. Somatic mutations of *IDH1* and *IDH2* in the leukemic transformation of myeloproliferative neoplasms. *N Engl J Med*. 2010;362(4):369-370.
- Ward PS, Patel J, Wise DR, et al. The common feature of leukemia-associated *IDH1* and *IDH2* mutations is a neomorphic enzyme activity converting alpha-ketoglutarate to 2-hydroxyglutarate. *Cancer Cell*. 2010;17(3):225-234.
- Heil G, Krauter J, Raghavachar A, et al. Risk-adapted induction and consolidation therapy in adults with de novo AML aged  $\leq 60$  years: results of a prospective multicenter trial. *Ann Hematol*. 2004;83(6):336-344.
- Krauter J, Wagner K, Schafer I, et al. Prognostic factors in adult patients up to 60 years old with acute myeloid leukemia and translocations of chromosome band 11q23: individual patient data-based meta-analysis of the German Acute Myeloid Leukemia Intergroup. *J Clin Oncol*. 2009;27(18):3000-3006.
- Damm F, Heuser M, Morgan M, et al. Single nucleotide polymorphism in the mutational hotspot of *WT1* predicts a favorable outcome in patients with cytogenetically normal acute myeloid leukemia. *J Clin Oncol*. 2010;28(4):578-585.
- Neubauer A, Maharry K, Mrozek K, et al. Patients with acute myeloid leukemia and RAS mutations benefit most from postremission high-dose cytarabine: a Cancer and Leukemia Group B study. *J Clin Oncol*. 2008;26(28):4603-4609.
- Gaidzik VI, Schlenk RF, Moschny S, et al. Prognostic impact of *WT1* mutations in cytogenetically normal acute myeloid leukemia: a study of the German-Austrian AML Study Group. *Blood*. 2009;113(19):4505-4511.

## Acknowledgments

The authors thank the patients and the members of the AML-Study Group as well as Kerstin Görlich, Martin Wichmann, Elvira Lux, Diana Dudacy, and Sylvia Horter for their excellent technical support.

This work was supported by the Deutsche-José-Carreras Leukämie-Stiftung e.V. (grant DJCLS H 06/04v), Kompetenznetz "Akute und chronische Leukämien" (grant 01GI0378), and the Bundesministerium für Bildung und Forschung and the Dieter-Schlag-Stiftung (grant 01KG0605).

## Authorship

Contribution: F.T., M.H., J.K., and A.G. designed the research; F.T., F.D., K.W., and M.H. performed the research; D.H., M.L., W.H., L.K., G.S., A.R., W.F., H.K., and G.H. contributed patient samples and clinical data; G.G. and B.S. performed cytogenetic studies; F.T., J.K., and A.G. analyzed the data and wrote the paper; and all authors read and agreed to the final version of the manuscript.

Conflict-of-interest disclosure: The authors declare no competing financial interests.

Correspondence: Felicitas Thol, Department of Hematology, Hemostasis, Oncology, and Stem Cell Transplantation, Hannover Medical School, Carl-Neuberg Str 1, 30625 Hannover, Germany; e-mail: thol.felicitas@mh-hannover.de.