

Regenerating islet-derived 3-alpha is a biomarker of gastrointestinal graft-versus-host disease

*James L. M. Ferrara,¹ *Andrew C. Harris,¹ Joel K. Greenson,² Thomas M. Braun,³ Ernst Holler,⁴ Takanori Teshima,⁵ John E. Levine,¹ Sung W. J. Choi,¹ Elisabeth Huber,⁶ Karin Landfried,⁴ Koichi Akashi,⁵ Mark Vander Lugt,¹ Pavan Reddy,⁷ Alice Chin,⁸ Qing Zhang,⁸ Samir Hanash,⁸ and Sophie Paczesny¹

Departments of ¹Pediatrics, ²Pathology, and ³Biostatistics, University of Michigan, Ann Arbor, MI; ⁴Department of Hematology/Oncology, University Medical Center Regensburg, Regensburg, Germany; ⁵Center for Cellular and Molecular Medicine, Kyushu University Graduate School of Science, Fukuoka, Japan; ⁶Institute of Pathology, University Medical Center Regensburg, Regensburg, Germany; ⁷Department of Internal Medicine, University of Michigan, Ann Arbor, MI; and ⁸Molecular Diagnostics Program, Fred Hutchinson Cancer Research Center, Seattle, WA

There are no plasma biomarkers specific for GVHD of the gastrointestinal (GI) tract, the GVHD target organ most associated with nonrelapse mortality (NRM) following hematopoietic cell transplantation (HCT). Using an unbiased, large-scale, quantitative proteomic discovery approach to identify candidate biomarkers that were increased in plasma from HCT patients with GI GVHD, 74 proteins were increased at least 2-fold; 5 were of GI origin. We validated the lead candidate,

REG3 α , by ELISA in samples from 1014 HCT patients from 3 transplantation centers. Plasma REG3 α concentrations were 3-fold higher in patients at GI GVHD onset than in all other patients and correlated most closely with lower GI GVHD. REG3 α concentrations at GVHD onset predicted response to therapy at 4 weeks, 1-year NRM, and 1-year survival ($P \leq .001$). In a multivariate analysis, advanced clinical stage, severe histologic damage, and high REG3 α concentrations at GVHD diag-

nosis independently predicted 1-year NRM, which progressively increased with higher numbers of onset risk factors present: 25% for patients with 0 risk factors to 86% with 3 risk factors present ($P < .001$). REG3 α is a plasma biomarker of GI GVHD that can be combined with clinical stage and histologic grade to improve risk stratification of patients. (*Blood*. 2011;118(25): 6702-6708)

Introduction

Acute GVHD, a leading cause of nonrelapse mortality (NRM) after allogeneic hematopoietic cell transplantation (HCT), is measured by dysfunction in 3 organ systems: the skin, liver, and gastrointestinal (GI) tract.¹⁻⁴ Acute GVHD of the GI tract affects up to 60% of patients receiving allogeneic HCT,^{5,6} causing nausea, vomiting, anorexia, secretory diarrhea, and, in more severe cases, abdominal pain and/or hemorrhage.⁷ Acute GVHD typically occurs between 2 and 8 weeks after transplantation, but may occur later,⁴ and is often clinically indistinguishable from other causes of GI dysfunction such as conditioning regimen toxicity, infection, or medication. Endoscopic biopsy is often used to confirm the diagnosis,^{1,8} but histologic severity on biopsy has not consistently correlated with clinical outcome.^{3,8-10} Clinical stage II or greater (> 1 L of diarrhea/d) is associated with reduced survival,^{5,6} but daily stool volume can vary considerably. Lower GI GVHD responds poorly to treatment compared with other target organs,⁶ and treatment with high-dose systemic steroid therapy carries significant risks, especially infectious complications in profoundly immunosuppressed patients.^{11,12} A noninvasive, reliable blood biomarker specific for GVHD of the GI tract would thus significantly aid in the management of patients with this disorder.

Here, we report the discovery and validation of a plasma biomarker of acute GI GVHD: regenerating islet-derived 3-alpha (REG3 α), a C-type lectin secreted by Paneth cells.^{13,14}

Methods

Proteomic analysis

Methods for sample preparation, protein fractionation, mass spectrometry (MS) analysis, protein identification, and quantitative analysis of protein concentrations during the intact protein analysis system (IPAS) have been previously reported.¹⁵⁻¹⁷

Patients and samples

Heparinized blood samples were collected weekly for 4 weeks after allogeneic HCT, then monthly for 2 months, and also at the time of key clinical events, including the development of symptoms consistent with GVHD (eg, the onset of diarrhea). Plasma samples were collected prospectively under protocols approved by the University of Michigan Institutional Review Board and stored at the University of Michigan. GVHD assessments, sample processing, and storage were performed as previously described.^{7,17} In Regensburg, Germany, and Kyushu, Japan, serum samples were collected weekly and at the onset of GVHD symptoms, prepared, frozen, and stored per institutional guidelines. Samples were shipped and received frozen on dry ice and no sample was thawed more than twice before analysis. REG3 α concentrations were stable in samples frozen for at least 5 years. REG3 α concentrations of 12 paired healthy donors plasma and serum were similar (mean \pm SEM: 20 \pm 3 vs 24 \pm 3 ng/mL, respectively).

Submitted August 20, 2011; accepted September 27, 2011. Prepublished online as *Blood* First Edition paper, October 6, 2011; DOI 10.1182/blood-2011-08-375006.

*J.L.M.F. and A.C.H. contributed equally to this work.

The online version of this article contains a data supplement.

The publication costs of this article were defrayed in part by page charge payment. Therefore, and solely to indicate this fact, this article is hereby marked "advertisement" in accordance with 18 USC section 1734.

© 2011 by The American Society of Hematology

Table 1. Patient characteristics of the University of Michigan validation set

Total, N = 871	GI GVHD*†, N = 167	No GVHD, N = 362	Non-GVHD enteritis‡, N = 52	Skin GVHD, N = 290	P
Median age, y (range)	50 (0-67)	46 (0-68)	48 (3-66)	49 (0-70)	.003
Disease, %					.002
Malignant	99 (n = 165)	92 (n = 334)	96 (n = 50)	97 (n = 282)	
Other	1 (n = 2)	8 (n = 28)	4 (n = 2)	3 (n = 8)	
Disease status at transplantation, %§					.63
Other/low/intermediate risk	64 (n = 105)	69 (n = 232)	68 (n = 34)	68 (n = 192)	
High risk	36 (n = 60)	31 (n = 102)	32 (n = 16)	32 (n = 90)	
Donor type, %					< .001
Related donor	45 (n = 75)	64 (n = 233)	54 (n = 28)	40 (n = 115)	
Unrelated donor	55 (n = 92)	36 (n = 129)	46 (n = 24)	60 (n = 175)	
Donor match, %					< .001
Matched donor	70 (n = 117)	90 (n = 325)	92 (n = 48)	73 (n = 212)	
Mismatched donor	30 (n = 50)	10 (n = 37)	8 (n = 4)	27 (n = 78)	
Conditioning regimen intensity, %					.06
High intensity	57 (n = 95)	67 (n = 243)	63 (n = 33)	57 (n = 165)	
Moderate intensity	43 (n = 72)	33 (n = 119)	37 (n = 19)	43 (n = 125)	
Grade of GVHD at onset, %					
0	0 (n = 0)	100 (n = 362)	100 (n = 52)	0 (n = 0)	
I	0 (n = 0)	0 (n = 0)	0 (n = 0)	69 (n = 201)	
Skin stage 1	0 (n = 0)	0 (n = 0)	0 (n = 0)	41 (n = 118)	
Skin stage 2	0 (n = 0)	0 (n = 0)	0 (n = 0)	29 (n = 83)	
II	57 (n = 96)	0 (n = 0)	0 (n = 0)	30 (n = 88)	
Isolated skin stage 3	0 (n = 0)	0 (n = 0)	0 (n = 0)	30 (n = 88)	
Isolated upper GI stage 1†	17 (n = 29)	0 (n = 0)	0 (n = 0)	0 (n = 0)	
Lower GI stage 1†	40 (n = 67)	0 (n = 0)	0 (n = 0)	0 (n = 0)	
III-IV	43 (n = 71)	0 (n = 0)	0 (n = 0)	1 (n = 1)	
Isolated skin stage 4	0 (n = 0)	0 (n = 0)	0 (n = 0)	1 (n = 1)	
GI stage 2†	13 (n = 22)	0 (n = 0)	0 (n = 0)	0 (n = 0)	
GI stage 3†	16 (n = 27)	0 (n = 0)	0 (n = 0)	0 (n = 0)	
GI stage 4†	13 (n = 22)	0 (n = 0)	0 (n = 0)	0 (n = 0)	
Median d after HCT (range)	33 (11-216)	31 (7-185)	24 (7-93)	28 (5-175)	< .001

GI indicates gastrointestinal; HCT, hematopoietic cell transplantation; and CIBMTR, Center for International Blood and Marrow Transplant Research.

*Including 29 patients with isolated upper GI GVHD and 138 with lower ± upper GI GVHD.

†With or without other GVHD target organ involvement.

‡Including 13 patients with isolated upper GI non-GVHD enteritis and 39 patients with lower ± upper GI non-GVHD enteritis.

§High risk of disease status at HCT is according to CIBMTR guidelines.

All patients received pharmacologic GVHD prophylaxis with at least 2 agents, including a calcineurin inhibitor. No donor grafts were depleted of T cells. All patients with available samples were analyzed, including patients who developed other complications of HCT, such as sinusoidal obstruction syndrome (SOS), idiopathic pneumonia syndrome (IPS), and sepsis/bacteremia. Patients were excluded from analysis only if a plasma sample at the time of GVHD onset was not available, or if methylprednisolone > 1 mg/kg (or equivalent) had been administered for > 48 hours at the time of sample acquisition. One sample was analyzed per patient; patients who developed GVHD had samples selected at the time of initial GVHD diagnosis.

The discovery set consisted of plasma samples from 10 HCT patients at the onset of biopsy-proven GI GVHD (clinical stage 1-3) and 10 HCT patients who never developed GVHD and who were matched for key transplantation characteristics (supplemental Table 1, available on the Blood Web site; see the Supplemental Materials link at the top of the online article). Patient samples in the discovery set were not included in the validation set.

The University of Michigan validation set consisted of 4 groups: patients with newly diagnosed GVHD involving the GI tract (with or without other organ involvement; GI GVHD); patients at similar time points who never developed GVHD symptoms (no GVHD); patients with GI distress that was inconsistent with GVHD either by clinical or histologic criteria (non-GVHD enteritis); and patients who presented with isolated skin GVHD (skin GVHD). Patient numbers and characteristics are shown in Table 1. Enteritis was determined to be inconsistent with GVHD on clinical grounds by documentation of infected stool and by resolution of symptoms without steroid treatment. The etiologies of non-GVHD enteritis are listed in Table 2.

Patients from the Regensburg/Kyushu validation set were divided into the same 4 groups; patient characteristics are detailed in supplemental Table 2, with causes of non-GVHD enteritis listed in supplemental Table 3.

Histopathology

GI biopsies were obtained and prepared per institutional guidelines. GVHD was histologically confirmed by duodenal/colonic biopsy in 183 of 197 GI GVHD patients and by skin biopsy in an additional 5 patients with both rash

Table 2. Causes of non-GVHD enteritis in the University of Michigan validation set

Causes of non-GVHD enteritis	% (n)
Non-GVHD lower GI enteritis ± upper GI symptoms: N = 39	
<i>Clostridium difficile</i> infection	54 (21)
Diarrhea with negative biopsy	15 (6)
Nausea/vomiting and diarrhea with negative biopsies	28 (11)
Ulcerative esophagitis and diarrhea (negative biopsies)	3 (1)
Non-GVHD upper GI enteritis without diarrhea (all biopsy negative): N = 13	
Nausea/vomiting	54 (7)
Anorexia	15 (2)
Chemical gastropathy	23 (3)
<i>Helicobacter pylori</i> gastritis	8 (1)

GI indicates gastrointestinal.

and GI symptoms.⁹ Skin GVHD was confirmed by biopsy in 272 of 341 patients with rashes and by biopsy of another target organ later affected by GVHD in an additional 8 patients. One hundred sixty-two of 197 patients with GI GVHD had diarrhea. One hundred forty of those 162 patients had biopsies (duodenal = 87, colonic = 53) available for formal grading as described by Lerner.¹⁸ If both duodenal and colonic biopsies were available, colonic biopsies were graded only if duodenal biopsies were negative. We did not impute values for unavailable biopsies.

ELISAs

REG3 α ELISA kits were purchased from MBL International (Ab-Match Assembly Human PAPI kit and Ab-Match Universal kit), and measurements were performed according to the manufacturer's protocol. Samples (diluted 1:10) and standards were run in duplicate, absorbance was measured with a SpectraMax M2 (Molecular Devices), and results were calculated with SoftMax Pro Version 5.4 (Molecular Devices). Elafin, IL2R α , HGF, TNFR1, and IL-8 ELISAs were performed in duplicate as previously reported.^{17,19} Measurements of samples from 66 patients (6.5% of the total population) were repeated in a second ELISA at random intervals and were comparable; correlation coefficient $r = 0.82$, $P < .0001$. Details of the assay parameters are provided in supplemental Table 4.

Statistical analysis

The statistical methods used for the IPAS are as previously described.¹⁵⁻¹⁷ REG3 α and albumin concentrations from individual samples in the discovery and validation sets were compared using 2-sample t tests applied to log-transformed concentrations. Differences in characteristics between patient groups were assessed with a Kruskal-Wallis test for continuous values and χ^2 tests of association for categorical values. Receiver operating characteristic (ROC) area under the curves (AUC) were estimated nonparametrically. NRM and relapse mortality were modeled with cumulative incidence regression methods as described by Fine and Gray.²⁰ One-year overall survival (OS) was modeled with Cox regression methods and probability of response was modeled with logistic regression.

Results

Discovery study

We used a proteomics approach to identify candidate biomarkers in a discovery set of pooled plasma samples taken at similar times after HCT from 10 patients with biopsy-proven GI GVHD and 10 patients without GVHD as previously described (supplemental Table 1).¹⁵⁻¹⁷ We identified and quantified 562 proteins of which 74 were increased at least 2-fold in patients with GVHD (supplemental Table 5). Five proteins (carboxypeptidase N catalytic chain precursor, pancreatic secretory trypsin inhibitor precursor, palladin, lithostathine 1- α precursor, and regenerating islet-derived 3- α) were preferentially expressed in the GI tract based on the relevant literature²¹⁻²⁵ and the Human Protein Atlas (<http://www.proteinatlas.org/>). Commercially available Abs suitable for quantification of plasma concentrations by ELISA were available for only 1 of these 5 proteins, regenerating islet-derived 3- α (REG3 α ; supplemental Table 5). The MS characteristics of the identified REG3 α peptides are shown in supplemental Figure 1 and supplemental Table 6. The plasma concentrations of REG3 α in the individual plasma samples in the discovery set were 4 times higher in the patients with GI GVHD than in asymptomatic controls (supplemental Figure 2, $P = .01$).

Validation study

We next evaluated REG3 α plasma concentration as a biomarker of GI GVHD in samples from a validation set of 871 allogeneic HCT

recipients from the University of Michigan (Table 1). Older transplant recipients, an underlying diagnosis of malignant disease, graft sources from unrelated and HLA-mismatched donors were overrepresented in the groups with GVHD. The median day of sample acquisition for patients with non-GVHD enteritis was closer to the day of transplantation than for all other groups.


Plasma REG3 α concentrations were 3 times higher in patients at the onset of GI GVHD than in all other patients, including those with non-GVHD enteritis (Figure 1A). There was no specific cause of non-GVHD diarrhea associated with higher REG3 α concentrations. Serum REG3 α concentrations were also higher in GI GVHD in an independent validation set of 143 HCT patients from Regensburg, Germany, and Kyushu, Japan, although the absolute values were lower (Figure 1B). This difference may be because of a center effect that depends on several factors, including variations in transplantation conditioning regimens and supportive care; patients receiving high-intensity conditioning regimens had REG3 α concentrations that were twice as high as those receiving moderate intensity conditioning, but this difference did not reach statistical significance (Figure 1C). In addition, all patients in Regensburg and Kyushu received oral antibiotics as GVHD prophylaxis, whereas Michigan patients did not and thus increased GI flora might account for greater REG3 α secretion.²⁶ Neither total body irradiation (TBI)-based conditioning nor GVHD prophylaxis regimen significantly impacted REG3 α concentrations (data not shown).

We next analyzed REG3 α concentrations according to diagnosis and type of GI symptom. In patients with diarrhea caused by GVHD, REG3 α concentrations at the onset of GVHD were 5 times higher than in patients with diarrhea from other causes (Figure 1D). In patients without diarrhea, REG3 α concentrations were 25% higher when attributable to GVHD compared with other causes, a difference that was not statistically significant.

We measured concentrations of 4 previously reported diagnostic markers of systemic acute GVHD (IL2R α , TNFR1, IL-8, and HGF),¹⁹ and of elafin, a biomarker for GVHD of the skin,¹⁷ in all patients with diarrhea (Figure 1C, $N = 204$). ROC curves for these biomarkers distinguished GVHD from non-GVHD with an AUC of 0.80 for REG3 α alone and an AUC of 0.81 for a composite panel of all 6 biomarkers (Figure 2). In this analysis, 52% of patients with lower GI GVHD also had skin involvement at onset, and thus the AUC for elafin, which is specific for GVHD of the skin,¹⁷ was greater than expected (supplemental Table 7). ROC curves of REG3 α concentrations in patients with diarrhea had similar AUCs in both validation sets (supplemental Figure 3). REG3 α was therefore the best single diagnostic biomarker at the onset of symptoms of lower GI GVHD, and additional biomarkers provided no further increased sensitivity or specificity. Using REG3 α at the median concentration provided a positive predictive value (PPV) of 95% and a negative predictive value (NPV) of 32% for GVHD as the etiology of diarrhea. Additional predictive values at other REG3 α concentrations are provided in supplemental Table 8.

When we categorized patients by the volume of diarrhea, REG3 α concentrations at the onset of symptoms continued to distinguish between GVHD and non-GVHD etiologies (Figure 3A, $P < .001$) but did not correlate with the clinical stage of GVHD. Twenty-three of 26 patients with clinical stage IV GI GVHD at onset received full-intensity conditioning, and these patients showed a trend toward higher REG3 α concentrations than those with stage 1-3 GI GVHD ($P = .07$; data not shown). Comparing patients who had < 1 L of stool per day because of GVHD versus other causes, the AUC for REG3 α was 0.81 (supplemental Figure 4). Plasma REG3 α concentrations at the onset of GVHD were significantly

Figure 1. REG3 α concentrations in plasma samples from HCT patients of 2 independent validation sets. (A) University of Michigan patients (n = 871). (B) Regensburg, Germany, and Kyushu, Japan (n = 143). (C) Plasma REG3 α concentrations in patients classified by GI symptoms and histologic diagnosis and categorized by conditioning regimen intensity. High-intensity regimens included: cyclophosphamide \pm cytarabine, thiopeta, fludarabine and/or TBI; cyclophosphamide/VP-16/BCNU; busulfan + cytarabine, clofarabine, melphalan, cyclophosphamide/anasacrin, or cytarabine/cyclophosphamide; BCNU/VP-16/cytarabine/melphalan; TBI \pm VP-16; melphalan. Moderate-intensity regimens included: fludarabine + busulfan or treosulfan \pm TBI, melphalan, zevalin, or anasacrin/cytarabine; fludarabine \pm TBI, melphalan, or cyclophosphamide; fludarabine/BCNU/melphalan; TBI. (D) Patients classified by symptoms and etiology (n = 675).



higher in patients whose GI biopsies showed evidence of severe GVHD with mucosal denudation (histologic grade 4) compared with less severe GVHD (Figure 3B; $P = .03$). Hypoalbuminemia is associated with the protein-losing enteropathy in GI GVHD,²⁷ and we analyzed the serum albumin level as a potential marker for loss of intravascular proteins into the intestinal lumen. Albumin levels at the onset of GI GVHD also correlated with both the clinical GI GVHD severity (supplemental Figure 5A) and histopathologic severity (supplemental Figure 5B).

Prognostic value of REG3 α concentrations in patients with lower GI GVHD

The clinical use of any biomarker is greatly enhanced when it provides prognostic information regarding the future status of a disease and/or patient, for example, the likelihood of response to treatment. We therefore evaluated the prognostic significance of REG3 α plasma levels in 162 patients taken at the time of diagnosis of lower GI GVHD. REG3 α concentrations were 3-fold higher at the time of GVHD diagnosis in patients who had no response to therapy at 4 weeks^{28,29} than in patients who experienced a complete or partial response (Figure 4A; $P < .001$)^{28,29}; patients responding to therapy still exhibited REG3 α concentrations more than twice that of non-GVHD controls. REG3 α concentrations at diagnosis also correlated with eventual maximal clinical stage of GI GVHD (supplemental Figure 6); patients presenting with isolated skin GVHD who later developed GI GVHD had concentrations comparable with those with skin GVHD who never developed GI GVHD ($P = .2$; data not shown). Because maximal GVHD grade correlates with NRM,¹¹ we hypothesized that the REG3 α concentration at GVHD diagnosis would also correlate with NRM. We therefore divided the 162 patients into 2 equal groups based on the median REG3 α concentration: high (> 151ng/mL, n = 81) and low (\leq 151 ng/mL, N = 81). NRM was twice as high in patients with high REG3 α concentrations, and this difference remained significant after adjusting for known risk factors of donor type, degree of HLA match, conditioning intensity, age, and baseline disease severity (59% [95% confidence interval [CI], 48%-69%] vs 34%[95% CI, 24%-46%], $P < .001$, Figure 4B). The incidence of relapse mortality was comparable for both groups (14% [95% CI, 8-24] vs 17% [95% CI, 8-24], $P = .5$; Figure 4C), and thus patients with high REG3 α concentrations at the time of GVHD diagnosis experienced significantly inferior 1-year OS (27% [95% CI, 19%-39%] vs 48% [95% CI, 38%-61%], $P = .001$; Figure 4D).




Figure 2. ROC curves for patients with post-HCT diarrhea. ROC curves comparing REG3 α concentrations for patients with diarrhea caused by GVHD (n = 162) and not caused by GVHD (N = 42). REG α alone (thick blue): AUC = 0.80; IL2R α (thick brown): AUC = 0.69; Elafin (thick red): AUC = 0.68; IL-8 (thin blue): AUC = 0.61; HGF (thin brown): AUC = 0.61; TNFR1 (thin red): AUC = 0.60; composite of all 6 biomarkers (solid black): AUC = 0.81.




Figure 3. REG3 α expression according to severity of GVHD at diagnosis. Patients were classified by volume of diarrhea (A) and histologic grade (B).

Causes of 1-year mortality for these patients are listed in supplemental Table 9.

Of the 162 patients with diarrhea at the onset of GVHD, we possessed all 4 data points of clinical stage, histologic grade, REG3 α concentration, and serum albumin level in 140 patients. As shown in Table 3, the plasma concentration of REG3 α , the clinical severity of GVHD, the histologic severity, and serum albumin level at GVHD diagnosis independently predicted lack of response to

GVHD therapy 4 weeks following treatment after adjustment for the aforementioned risk factors (odds ratios: 4.8, 3.9, 18.9, and 2.5, respectively). When lack of response to therapy and NRM were modeled simultaneously on all 4 parameters, all but albumin remained statistically significant. When only advanced clinical stage and severe histologic grade were considered, NRM was 71% (Figure 4E). The inclusion of high REG3 α concentration further risk-stratified patients who had either advanced clinical




Figure 4. Prognostic value of REG3 α concentrations at onset of GVHD. (A) Patients were classified by response to GVHD therapy after 4 weeks (N = 160). (B-D) Patients were classified by REG3 α concentration: low (≤ 151 ng/mL, n = 81; thin line) and high (> 151 ng/mL, n = 81; thick line). (B) NRM (34% vs 59%, $P < .001$) (C) Relapse mortality (17% vs 14%, $P = .59$). (D) One-year survival (48% vs 27%, $P = .001$). All P values are adjusted for donor source, HLA match, conditioning intensity, recipient age, and baseline disease severity according to the Center for International Blood and Marrow Transplant Research (CIBMTR) guidelines. (E) One-year NRM for patients classified by number of risk factors at GVHD onset, using clinical stage (high risk = stage 2-4) and histologic grade (high risk = grade 4); 0 (purple, NRM = 26%); 1 (red, NRM = 60%); 2 (blue, NRM = 71%); 0 vs 1, $P < .001$; 1 vs 2, $P = .006$. (F) One-year NRM for patients classified by number of risk factors at the time of GVHD diagnosis as in panel E and including REG3 α concentration (high risk > 151 ng/mL); 0 (purple, NRM = 25%); 1 (red, NRM = 34%); 2 (purple, NRM = 66%); 3 (brown, NRM = 86%); 0 vs 1, $P = .2$; 1 vs 2, $P < .001$; 2 vs 3, $P < .001$.

Table 3. REG3 α concentrations and characteristics at onset of GVHD diarrhea predict 4-week response to GVHD therapy and 1-year NRM

	Independent		Simultaneous	
	Ratio	P*	Ratio	P*
No response to treatment (at 4 wk)	Odds		Odds	
REG3 α (high vs low)	4.8	< .001	5.7	.001
GVHD GI onset stage (2-4 vs 1)	3.9	.001	3.0	.027
Histologic grade (4 vs 1-3)	18.9	< .001	16.7	< .001
Albumin (low vs high)	2.5	.02	1.4	.5
1-y NRM	Hazard		Hazard	
REG3 α (high vs low)	2.2	.003	2.4	.002
GVHD GI onset stage (2-4 vs 1)	3.0	< .001	3.1	< .001
Histologic grade (4 vs 1-3)	3.6	< .001	2.9	< .001
Albumin (low vs high)	2.3	.004	1.6	.2

NRM indicates nonrelapse mortality; and GI, gastrointestinal.

*Adjusted for age, donor type, HLA match, conditioning intensity, and disease status at transplantation.

stage or histologic severity (Figure 4F; 34% vs 66% for 1 or 2 risk factors, respectively, $P < .001$), and patients who had all 3 risk factors experienced significantly greater NRM than those with any 2 of the risk factors (86% vs 66%, $P < .001$). Details of patient risk factors are listed in supplemental Table 10; NRM by all other risk factor combinations are shown in supplemental Figure 7.

Discussion

The etiology of diarrhea following HCT presents a common diagnostic dilemma.^{30,31} We identified REG3 α as a candidate biomarker specific for lower GI GVHD through an unbiased, in-depth tandem MS-based discovery approach that can quantify proteins at low concentrations and that we previously used successfully to identify elafin as a plasma biomarker specific for GVHD of the skin.¹⁷ Our discovery approach identified 74 proteins that were increased at least 2-fold in the plasma from patients with GI GVHD. Of note, the list did not include cytokeratin-18 (KRT18), which has been reported to be specific for both liver and intestinal GVHD.³² This discrepancy may be explained by limitations in proteomics technology and the significantly later acquisition times of samples in the earlier report.

REG proteins act downstream of IL-22 to protect the epithelial barrier function of the intestinal mucosa^{33,34} through the binding of bacterial peptidoglycans.¹³ Intestinal stem cells (ISCs) are principal cellular targets of GVHD in the GI tract,^{3,35} where intestinal flora are critical for amplification of GVHD damage.^{36,37} A leading hypothesis is that ISCs are protected by antibacterial proteins such as REG3 α secreted by neighboring Paneth cells into the crypt microenvironment.³⁸ If death of an ISC eventually manifests itself as denudation of the mucosa, the patchy nature of GVHD histologic damage may be explained as the lack of mucosal regeneration following the dropout of individual ISCs.^{3,35} REG3 α reduces the inflammation of human intestinal crypts in vitro,^{14,39} and its administration protects ISCs and prevents GI epithelial damage in vivo,³⁴ raising interesting therapeutic possibilities for this molecule.

REG3 α plasma concentrations correlate with disease activity in inflammatory bowel disease, and can distinguish infectious and autoimmune causes of diarrhea.¹⁴ The correlation of mucosal denudation (histologic grade 4) with high REG3 α concentrations suggests that microscopic breaches in the mucosal epithelial barrier

caused by severe GVHD permit REG3 α to traverse into the systemic circulation. The tight proximity of Paneth cells with ISCs concentrates their secretory contents in that vicinity, so that mucosal barrier disruption caused by stem cell dropout may preferentially allow Paneth cell secretions, including REG3 α , to traverse into the bloodstream. We hypothesize that plasma levels of REG3 α may therefore serve as a surrogate marker for the cumulative area of these breaches to GI mucosal barrier integrity, a parameter impossible to measure by individual tissue biopsies. Such an estimate of total damage to the mucosal barrier may also help explain the prognostic value of REG3 α with respect to therapy responsiveness and NRM.

In this study, 3 high-risk parameters each independently correlated with lack of response to treatment and to higher NRM: elevated plasma REG3 α concentration, higher clinical stage of GVHD at diagnosis and grade 4 histologic severity. All 3 of these values thus provided important prognostic information before the initiation of therapy rather than at the time of maximum grade of GVHD, which by definition includes responsiveness to therapy.^{5,6,11} This study confirms earlier reports where higher clinical stage of GI GVHD^{5,6} and more severe histology correlated with worse survival.¹⁰ In our study the 1-year NRM was 33% (22 of 67 patients) in patients with clinical stage I lower GI GVHD when considering clinical severity alone. Seven of 8 patients (88%) who had the 2 other high risk factors present experienced 1-year NRM while 25% (15 of 59) of patients with 1 or no risk factors experienced 1-year NRM. In this regard it should be noted that REG3 α levels did not obviate the need for biopsy. If the prognostic value of REG3 α is confirmed in additional patients, we believe the integration of clinical stage, histologic grade and REG3 α plasma concentrations into a single grading system will permit better risk stratification and rapid identification of those patients with severe GI damage in whom standard treatment is likely to be insufficient.

Acknowledgments

The authors thank Dr David Beer for critical reading of the manuscript, the clinicians of the University of Michigan Blood and Marrow Transplantation division: Daniel Couriel, Greg Yanik, Carrie Kitko, James Connelly, Craig Byersdorfer, Shin Mineishi, John Magenau, Ed Peres, and Attaphol Pawarode; David Hanauer for assistance with data abstraction using the EMERSE tool; the University of Michigan clinical research support staff: Joel Whitfield, Dawn Jones, and Connie Varner; the University of Michigan BMT division data managers: Pamela Jones, Rachel Young, Charlotte Zinkus, Katherine Archangeli, Tamara Cummings, Sean Kelley, and Jennifer Lay-Luskin; the members of the Paczesny laboratory: Aurelie Gomez, Megan Conlon, and Jeffrey Crawford; the members of the Hanash laboratory: Hong Wang, Vitor Faca, and Sharon Pitteri; and the Seattle Proteomics Core, particularly Phillip Gafken and Jason Hogan.

This work was supported by National Institutes of Health (NIH) grants RC1-HL-101102, P01-CA039542, T32-HL007622, the Hartwell Foundation, and the Doris Duke Charitable Foundation. Informatics assistance with data abstraction and the EMERSE tool was provided by the University of Michigan Cancer Center's (UMCC) Biomedical Informatics Core with partial support from the NIH through the UMCC Support grant (CA46592). Partial funding for the LTQ-FT mass spectrometer used in this work was generously provided by the M. J. Murdock Charitable Trust.

J.L.M.F. is a clinical research professor of the American Cancer Society and a visiting fellow of the Oxford All Souls College. S.P. is an investigator of the Eric Hartwell fund and the Amy Strelzer Manasevit Research Program.

Authorship

Contribution: J.L.M.F. planned the study, interpreted the data, and wrote the manuscript; A.C.H. designed and planned the experiments, performed research, performed data collection and quality assurance, analyzed data, and wrote the manuscript; J.K.G. and E. Huber performed pathology evaluations and wrote the manuscript; T.M.B. was the study statistician and wrote the manuscript;

E. Holler, T.T., J.E.L., S.W.J.C., K. L., K.A., and P.R. contributed to patient accrual, clinical data collection and quality assurance, research discussion, and wrote the manuscript; M.V.L. performed experiments and wrote the manuscript; A.C., Q.Z., and S.H. performed the proteomics experiments, interpreted data, and wrote the manuscript; and S.P. conceived and planned the study design, performed experiments, interpreted data, and wrote the manuscript.

Conflict-of-interest disclosure: The authors declare no competing financial interests.

Correspondence: Dr Sophie Paczesny, Blood and Marrow Transplant Program, University of Michigan Comprehensive Cancer Center, Rm 6410, 1500 E Medical Center Dr, Ann Arbor, MI, 48109-5942; e-mail: sophiep@med.umich.edu.

References

- Cutler C, Antin JH. Manifestation and treatment of acute graft-versus-host-disease. In: Appelbaum F, Forman SJ, Negrin RS, Blume KG, eds. *Thomas' Hematopoietic Cell Transplantation*. Oxford, United Kingdom: Blackwell Publishing Ltd; 2009: 1287-1303.
- Welniak LA, Blazar BR, Murphy WJ. Immunobiology of allogeneic hematopoietic stem cell transplantation. *Annu Rev Immunol*. 2007;25:139-170.
- Mowat A, Socie G. Intestinal graft-vs.-host disease. In: Ferrara JLM, Cooke KR, Deeg HJ, eds. *Graft-vs-Host Disease*. New York, NY: Marcel Dekker; 2004:279-327.
- Ferrara JL, Levine JE, Reddy P, Holler E. Graft-versus-host disease. *Lancet*. 2009;373(9674):1550-1561.
- Martin PJ, McDonald GB, Sanders JE, et al. Increasingly frequent diagnosis of acute gastrointestinal graft-versus-host disease after allogeneic hematopoietic cell transplantation. *Biol Blood Marrow Transpl*. 2004;10(5):320-327.
- MacMillan ML, Weisdorf DJ, Wagner JE, et al. Response of 443 patients to steroids as primary therapy for acute graft-versus-host disease: comparison of grading systems. *Biol Blood Marrow Transpl*. 2002;8(7):387-394.
- Przepiorka D, Weisdorf D, Martin P, et al. 1994 Consensus Conference on Acute GVHD Grading. *Bone Marrow Transpl*. 1995;15(6):825-828.
- Shulman HM, Kleiner D, Lee SJ, et al. Histopathologic diagnosis of chronic graft-versus-host disease: National Institutes of Health Consensus Development Project on Criteria for Clinical Trials in Chronic Graft-versus-Host Disease: II. Pathology Working Group Report. *Biol Blood Marrow Transpl*. 2006;12(1):31-47.
- Washington K, Jagasia M. Pathology of graft-versus-host disease in the gastrointestinal tract. *Hum Pathol*. 2009;40(7):909-917.
- Ertault-Daneshpouy M, Leboeuf C, Lemann M, et al. Pericapillary hemorrhage as criterion of severe human digestive graft-versus-host disease. *Blood*. 2004;103(12):4681-4684.
- Weisdorf D, Haake R, Blazar B, et al. Treatment of moderate/severe acute graft-versus-host disease after allogeneic bone marrow transplantation: an analysis of clinical risk features and outcome. *Blood*. 1990;75(4):1024-1030.
- Deeg HJ. How I treat refractory acute GVHD. *Blood*. 2007;109(10):4119-4126.
- Cash HL, Whitham CV, Behrendt CL, Hooper LV. Symbiotic bacteria direct expression of an intestinal bactericidal lectin. *Science*. 2006;313(5790):1126-1130.
- Gironella M, Iovanna JL, Sans M, et al. Anti-inflammatory effects of pancreatitis associated protein in inflammatory bowel disease. *Gut*. 2005;54(9):1244-1253.
- Faca V, Coram M, Phanstiel D, et al. Quantitative analysis of acrylamide labeled serum proteins by LC-MS/MS. *J Proteome Res*. 2006;5(8):2009-2018.
- Faca V, Pitteri SJ, Newcomb L, et al. Contribution of protein fractionation to depth of analysis of the serum and plasma proteomes. *J Proteome Res*. 2007;6(9):3558-3565.
- Paczesny S, Braun T, Levine JE, et al. Elafin is a biomarker of graft-versus-host disease of the skin. *Sci Transl Medicine*. 2010;2(13):50-57.
- Lerner KG, Kao GF, Storb R, Buckner CD, Clift RA, Thomas ED. Histopathology of graft-vs.-host reaction (GvHR) in human recipients of marrow from HL-A-matched sibling donors. *Transplant Proc*. 1974;6(4):367-371.
- Paczesny S, Krijanovski OI, Braun TM, et al. A biomarker panel for acute graft-versus-host disease. *Blood*. 2009;113(2):273-278.
- Fine JP, Gray RJ. A proportional hazards model for the redistribution of a competing risk. *J Am Stat Assoc*. 1999;94:496-509.
- Skidgel RA, Erdos EG. Structure and function of human plasma carboxypeptidase N, the anaphylatoxin inactivator. *Int Immunopharmacol*. 2007;7(14):1888-1899.
- Marchbank T, Freeman TC, Playford RJ. Human pancreatic secretory trypsin inhibitor. Distribution, actions and possible role in mucosal integrity and repair. *Digestion*. 1998;59(3):167-174.
- Mykkanen OM, Gronholm M, Ronty M, et al. Characterization of human palladin, a microfilament-associated protein. *Mol Biol Cell*. 2001;12(10):3060-3073.
- Watanabe T, Yonekura H, Terazono K, Yamamoto H, Okamoto H. Complete nucleotide sequence of human reg gene and its expression in normal and tumoral tissues. The reg protein, pancreatic stone protein, and pancreatic thread protein are one and the same product of the gene. *J Biol Chem*. 1990;265(13):7432-7439.
- Christa L, Carnot F, Simon MT, et al. HIP/PAP is an adhesive protein expressed in hepatocarcinoma, normal Paneth, and pancreatic cells. *Am J Physiol*. 1996;271(6 Pt 1):G993-G1002.
- Ayabe T, Satchell DP, Wilson CL, Parks WC, Selsted ME, Ouellette AJ. Secretion of microbicidal alpha-defensins by intestinal Paneth cells in response to bacteria. *Nat Immunol*. 2000;1(2):113-118.
- Weisdorf SA, Salati LM, Longsdorf JA, Ramsay NK, Sharp HL. Graft-versus-host disease of the intestine: a protein losing enteropathy characterized by fecal alpha 1-antitrypsin. *Gastroenterology*. 1983;85(5):1076-1081.
- MacMillan ML, DeFor TE, Weisdorf DJ. The best endpoint for acute GVHD treatment trials. *Blood*. 2010;115(26):5412-5417.
- Levine JE, Logan B, Wu J, et al. Graft-versus-host disease treatment: predictors of survival. *Biol Blood Marrow Transpl*. 2010;16(12):1693-1699.
- Cox GJ, Matsui SM, Lo RS, et al. Etiology and outcome of diarrhea after marrow transplantation: a prospective study. *Gastroenterology*. 1994;107(5):1398-1407.
- Barker CC, Anderson RA, Sauve RS, Butzner JD. GI complications in pediatric patients post-BMT. *Bone Marrow Transpl*. 2005;36(1):51-58.
- Luft T, Conzelmann M, Benner A, et al. Serum cytokeratin-18 fragments as quantitative markers of epithelial apoptosis in liver and intestinal graft-versus-host disease. *Blood*. 2007;110(13):4535-4542.
- Sanos SL, Vonarbourg C, Mortha A, Diefenbach A. Control of epithelial cell function by interleukin-22-producing RORgamma(+) innate lymphoid cells. *Immunology*. 2011;132(4):453-465.
- Zheng Y, Valdez PA, Danilenko DM, et al. Interleukin-22 mediates early host defense against attaching and effacing bacterial pathogens. *Nat Med*. 2008;14(3):282-289.
- Takashima S, Kadowaki M, Aoyama K, et al. The Wnt agonist R-spondin1 regulates systemic graft-versus-host disease by protecting intestinal stem cells. *J Exp Med*. 2011;208(2):285-294.
- van Bekkum DW, Roodenburg J, Heidt PJ, van der Waaij D. Mitigation of secondary disease of allogeneic mouse radiation chimeras by modification of the intestinal microflora. *J Natl Cancer Inst*. 1974;52(2):401-404.
- Gerbitz A, Schultz M, Wilke A, et al. Probiotic effects on experimental graft-versus-host disease: let them eat yogurt. *Blood*. 2004;103(11):4365-4367.
- Elphick DA, Mahida YR. Paneth cells: their role in innate immunity and inflammatory disease. *Gut*. 2005;54(12):1802-1809.
- Closa D, Motoo Y, Iovanna JL. Pancreatitis-associated protein: from a lectin to an anti-inflammatory cytokine. *World J Gastroenterol*. 2007;13(2):170-174.