Brief report

Simple genetic diagnosis of hairy cell leukemia by sensitive detection of the *BRAF*-V600E mutation

*Enrico Tiacci,¹ *Gianluca Schiavoni,¹ Francesco Forconi,² Alessia Santi,¹ Livio Trentin,³ Achille Ambrosetti,⁴ Debora Cecchini,¹ Elisa Sozzi,² Paola Francia di Celle,⁵ Cristiana Di Bello,⁵ Alessandro Pulsoni,⁶ Robin Foà,⁶ Giorgio Inghirami,⁵ and Brunangelo Falini¹

¹Institute of Hematology, University of Perugia, Perugia, Italy; ²Hematology, University of Siena and Azienda Ospedaliera Universitaria Senese, Siena, Italy; ³Department of Clinical and Experimental Medicine, Hematology and Clinical Immunology Section, University of Padua, Padua, Italy; ⁴Department of Medicine, Hematology Section, University of Verona, Italy; ⁵Department of Pathology, Center for Experimental Research and Medical Studies, University of Torino, Turin, Italy; and ⁶Division of Hematology, Department of Cellular Biotechnologies and Hematology, Sapienza University, Rome, Italy

Hairy cell leukemia (HCL) is a distinct clinicopathologic entity that responds well to purine analogs but is sometimes difficult to differentiate from HCL-like disorders (eg, splenic marginal zone lymphoma and HCL variant). We recently identified the *BRAF*-V600E mutation as the disease-defining genetic event in HCL. In this study, we describe a new, simple, and inexpensive test for genetics-based diagnosis of HCL in whole-blood samples that detects *BRAF*-V600E through a sensitive allele-specific PCR qualitative assay followed by agarose-gel electrophoresis. This approach detected *BRAF*-V600E in all 123 leukemic HCL samples investigated containing as few as 0.1% leukemic cells. *BRAF*-V600E was detected at different time points during the disease course, even after therapy, pointing to its pivotal role in HCL pathogenesis and maintenance of the leukemic clone. Conversely, 115 non-HCL chronic B-cell neoplasms, including 79 HCL-like disorders, were invariably negative for *BRAF*-V600E. This molecular assay is a powerful tool for improving the diagnostic accuracy in HCL. (*Blood.* 2012;119(1):192-195)

Introduction

Hairy cell leukemia (HCL) is a distinct entity usually characterized by splenomegaly (without lymphadenopathy), pancytopenia, and infiltration of BM, spleen, and liver by leukemic B cells with "hairy" appearance. In contrast to other chronic B-cell leukemias, HCL cells circulate at low percentages in the blood,¹ and exhibit distinct functional features and gene expression profile.^{2,3}

HCL diagnosis relies on morphologic and immunophenotypic criteria¹⁻⁴ that usually allow its distinction from HCL-like disorders of the 2008 World Health Organization classification, that is, splenic marginal zone lymphoma (SMZL) and splenic lymphoma/ leukemia unclassifiable (SLLU; including HCL variant [HCL-v]).⁵ A correct diagnosis is critical because purine analogs (pentostatin and cladribine) are highly effective only in HCL.⁴

The most problematic cases can be diagnosed using annexin-A1 immunostaining,⁶⁻⁸ which we previously reported to be highly sensitive and specific for HCL among B-cell lymphomas.⁶ However, because annexin-A1 is also expressed by myeloid and T cells,⁶ this immunohistochemical staining may be difficult to interpret in BM biopsies with low percentages of HCL cells. Moreover, immunocytochemistry for annexin-A1 is not readily applicable to routine hematologic samples, such as peripheral blood or diluted BM aspirate (because of dry tap), that are also usually poor in HCL cells and rich in neutrophils and T cells.

An ideal solution would be a sensitive and specific test for a genetics-based diagnosis of HCL. We recently identified *BRAF*-

The online version of this article contains a data supplement.

V600E as the HCL-defining genetic lesion (present in all HCL cases, absent in other B-cell neoplasms)⁹ by Sanger sequencing of *BRAF* exon 15. However, this technique required $\geq 30\%$ leukemic cells for reliably detecting a clonal heterozygous mutation. Thus, the rare HCL cells typically present in the blood of most patients had to be purified through cell sorting,⁹ a laborious procedure not amenable to a routine diagnostic setting.

Aims of this study were: (1) to develop a sensitive, easy, and inexpensive test for the routine clinical diagnosis of HCL in blood samples, based on *BRAF*-V600E detection by allele-specific PCR (AS-PCR) followed by conventional agarose-gel electrophoresis; and (2) to assess the diagnostic accuracy of this test in a large cohort of HCL and HCL-like disorders.

Methods

Tumor samples

We studied 117 HCL patients: 96 before therapy and 21 after therapy, all with detectable disease (at least 0.1% leukemic cells). We also investigated 16 HCL patients after therapy in complete flow cytometric remission (< 0.1% leukemic cells), 112 patients with other B-cell neoplasms (61 SMZL; 18 SLLU, including 11 HCL-v; 31 chronic lymphocytic leukemias [CLL]; 5 unclassifiable CD5-negative mature B-cell neoplasms), and 9 healthy blood donors (see supplemental Methods for details, available on the *Blood* Web site; see the Supplemental Materials link at the top of the

The publication costs of this article were defrayed in part by page charge payment. Therefore, and solely to indicate this fact, this article is hereby marked "advertisement" in accordance with 18 USC section 1734.

© 2012 by The American Society of Hematology

Submitted August 3, 2011; accepted October 5, 2011. Prepublished online as *Blood* First Edition paper, October 25, 2011; DOI 10.1182/blood-2011-08-371179.

^{*}E.T. and G.S. equally contributed to the work.



Figure 1. Flow cytometry and AS-PCR assays in HCL and HCL-like disorders. (Top panel) Flow cytometry dot plots of a whole-blood sample subjected to RBC lysis from a representative HCL patient (top graphs) and of purified peripheral blood leukemic cells from a representative patient with HCL-v (bottom graphs). HCL cells (CD19⁺/CD25⁺ red events in the top right panel) represent 2% of all nucleated cells (CD45⁺ black events in the top left panel). HCL-v cells (CD19⁺/CD25⁻ and CD19⁺/CD103⁺ red events in the bottom left and right panels, respectively) represent 92% of all cells. (Bottom panel) Conventional agarose-gel electrophoresis of samples from 13 HCL patients (top panels; 12 pretreatment, 1 with residual disease after treatment) and 16 HCL-like patients (bottom gels; 6 SLLU, 10 SMZL), after AS-PCR for the mutant allele (1st and 3rd gel from the top) and for the wild-type allele (2nd and 4th gel from the top). In the top-most gels, serial dilutions of mutated and wild-type alleles (from 3.1% to 0% muted alleles) are also included to show the analytical sensitivity of the mutant AS-PCR ($\ge 0.1\%$ mutated alleles). All HCL samples gave rise to a mutant *BRAF*-V600E band as opposed to one of the HCL-like samples. One of the latter (SMZL case 30), which did not give rise to the wild-type band either, was not evaluable in this particular experiment (shown on purpose), but on repetition turned out to be evaluable (ie, strong wild-type band) and negative for *BRAF*-V600E (ie, mutant band not visible). To facilitate the visualization of the results, the gel lane of HCL case 13 was repositioned in the 2 top gels and the gel lane of the 50-bp DNA ladder was repositioned in the 2 bottom gels.

online article). Samples from 23 HCL patients and 38 non-HCL patients were previously reported.⁹ Diagnosis of HCL and non-HCL tumors conformed to the WHO 2008 classification.^{1,5} Patients gave verbal or

written consent for the analysis of their sample material in accordance with the Declaration of Helsinki, and the study was approved by the local ethics committee at Perugia University.

Table 1. Results of AS-PCR in HCL cases and other B-cell	
neoplasms	

Sample type	Leukemic cells	No. of cases	Mutated cases, no. (%)
Hairy cell leukemia			
Blood, pretreatment	1%-90%	81	81 (100)
Blood, MRD	0.1%-13%		10 (100)
Blood, complete remission	Not detectable	16	0 (0)
BM biopsies	30%-80%	14	14 (100)
Purified leukemic cells	$\ge 90\%$	17	17 (100)
Splenic marginal zone lymphoma	2%-97%	61	0 (0)
Splenic lymphoma/leukemia unclassifiable	15%-97%	18*	0 (0)
Chronic lymphocytic leukemia	18%-98%	31	0 (0)
CD5-negative mature B-cell neoplasm unclassifiable	15%-97%	5	0 (0)
Healthy blood donors	Not present	9	0 (0)

AS-PCR indicates allele-specific PCR; HCL, hairy cell leukemia; and MRD, minimal residual disease.

*Including 11 HCL variants.

Qualitative AS-PCR for BRAF-V600E

Briefly, the assay consists of 2 PCRs sharing the same reverse primer (5'-GTAACTCAGCAGCATCTCAGGG-3') but differing for the forward primer, which is complementary to either the wild-type (T) or the mutated (A) base causing the V600E replacement. Forward primers were: 5'-AGGTGATTTTGGTCTAGCTACAGA-3' (mutated base in bold) and 5'-AGGTGATTTTGGTCTAGCTACAGT-3' (wild-type base in bold). After mutant AS-PCR, detection of a band on agarose-gel electrophoresis indicates the presence of the mutation. Conversely, absence of the band indicates lack of the mutation as long as the wild-type AS-PCR (positive control) gives a readily visible product (see supplemental Methods for details).

Results and discussion

We first assessed the analytical sensitivity of our AS-PCR in serial dilutions of DNA from a *BRAF*-V600E homozygous sample with DNA from a *BRAF* wild-type sample, and established the lower detection limit to be 0.1% of mutant alleles (Figure 1), corresponding to 0.2% of diploid tumor cells harboring a clonal heterozygous *BRAF*-V600E mutation (or 0.1% if the mutation is homozygous).

We then analyzed samples from 123 HCL patients (101 before treatment; 22 with residual or relapsing disease) and all tested positive (100% diagnostic sensitivity; Table 1, Figure 1), including 23 samples previously known to harbor *BRAF*-V600E by Sanger sequencing.⁹ Notably, among the newly reported 94 HCL cases, 24 (17 before treatment, 7 after treatment) had only 0.1% to 5% leukemic cells and 11 were analyzed after the onset of the disease (range 1-26 years).

These findings demonstrate the excellent analytical and diagnostic sensitivity of our test. They also confirm and extend our previous report that *BRAF*-V600E occurs and persists over the disease course in virtually all HCL cases,⁹ further supporting the view that *BRAF*-V600E represents the key pathogenetic event in HCL and therefore a new therapeutic target. Indeed, persistence of *BRAF*-V600E at partial remission or relapse after conventional therapy establishes the rationale for using active BRAF inhibitors¹⁰ in this setting. Our test may also serve as a new tool (in addition to IHC, flow cytometry, and Ig gene rearrangement analysis^{11,12}) to assess minimal residual disease (MRD) after therapy, although the clinical relevance of MRD in HCL remains unclear.¹²

We next evaluated the diagnostic specificity of our test by analyzing blood samples from 9 healthy donors and 16 HCL patients in complete flow cytometric remission (< 0.1% leukemic cells) after therapy, and all tested negative (Table 1, Figure 1). Specificity was also assessed in 115 patients with non-HCL chronic B-cell neoplasms. Because absence9 or very rare occurrence13-17 of BRAF-V600E has been already reported in several B-cell tumors, we focused on HCL-like disorders that, being rare, have been so far poorly investigated. Therefore, we included, among the 115 cases, 79 patients with SMZL and SLLU, of which 65 previously unreported.9 Notably, all 112 cases tested negative (Table 1, Figure 1), showing a 100% diagnostic specificity of our assay and further confirming in a larger patient series the absence of BRAF-V600E in HCL-like disorders. Considering that 37 HCL-like cases had $\geq 40\%$ neoplastic cells and that our test can detect 0.1% mutant alleles, these data also argue against the presence of small BRAF-V600E-mutated subclones (down to 0.5% of a whole leukemic population representing $\geq 40\%$ of the sample) in HCLlike disorders. This finding further supports the concept that, among B-cell lymphomas and leukemias, BRAF-V600E is the genetic lesion defining HCL.9 Although we have collectively analyzed 99 HCL-like disorders (SMZL and SLLU, the latter including 19 HCL-v) without finding BRAF-V600E in any of them (this article and Tiacci et al⁹), we cannot exclude that this mutation may be rarely found in these and other B-cell neoplasms if a larger number of cases is investigated.

Our diagnostic test is especially useful for patients with a low tumor burden in the blood (as typically occurs in HCL) or BM. In this setting, it appears superior to annexin-A1 immunostaining, which may be difficult to interpret (because of annexin-A1 expression by myeloid and T cells) unless a technically demanding double staining for annexin-1 and a B-cell marker (eg, PAX5) is performed. Our gel-based AS-PCR is considerably more sensitive than a recently described high-resolution-melting analysis (HRMA)– based PCR, which was applied to fewer HCL samples (n = 48) containing more (\geq 10%) leukemic cells.¹⁸ Notably, our test detected *BRAF*-V600E in all blood samples having < 10% HCL cells (37 of 88 samples; 27 before therapy; 10 after therapy) and does not require the expensive instrumentation needed for HRMA.

In conclusion, our sensitive, simple, and reliable assay confirms the constant presence of *BRAF*-V600E in HCL (as also reported in Boyd et al¹⁸ and Arcaini et al¹⁹) and its absence in HCL-like disorders, and adds to the already available armamentarium for improving the diagnostic accuracy in HCL and HCL-like disorders.

Acknowledgments

The authors are grateful to Barbara Bigerna, Alessandra Pucciarini, Roberta Pacini, Alessia Tabarrini, Ariele Spanhol-Rosseto, Valentina Pettirossi, and Ilaria Gionfriddo for technical help; Maria Paola Martelli and Alain Sylvin Bell for helpful discussions; Monia Capponi and Franca Falzetti for providing clinical and flow cytometry data on some patients; Stefano Ascani for help in hematopathologic diagnosis; Pellegrino Musto, Angelo Michele Carella, Francesco Di Raimondo, Michael Bennet, Mauro Di Ianni, Michele Cimminiello, Massimo Catarini, and Adriano Venditti for providing blood samples of 1 to 4 patients each; and Claudia Tibidò for secretarial assistance. This work was supported by the Associazione Italiana Ricerca Cancro (AIRC), the HCL Research Foundation, the Ministero dell'Istruzione, dell'Università e della Ricerca (MIUR-Programma Futuro in Ricerca 2010), and the Associazione Umbra contro le Leucemie e i Linfomi (AULL).

Authorship

Contribution: E.T. designed the study, decided the allele-specific PCR strategy, analyzed and interpreted the data, and wrote the manuscript; G.S. developed the allele-specific PCR assay and performed all related experiments, analyzed and interpreted the data, and contributed to the writing of the manuscript; F.F., L.T.,

References

- Foucar K, Falini B, Catovsky D, Stein H. Hairy cell leukaemia. In: Swerdlow S, Campo E, Harris NL, et al, eds. WHO Classification of Tumours of Haematopoietic and Lymphoid Tissues. 4th ed. Lyon, France: International Agency for Research on Cancer (IARC); 2008:188-190.
- Basso K, Liso A, Tiacci E, et al. Gene expression profiling of hairy cell leukemia reveals a phenotype related to memory B cells with altered expression of chemokine and adhesion receptors. *J Exp Med.* 2004;199(1):59-68.
- Tiacci E, Liso A, Piris M, Falini B. Evolving concepts in the pathogenesis of hairy-cell leukaemia. *Nat Rev Cancer*. 2006;6(6):437-448.
- 4. Grever MR. How I treat hairy cell leukemia. *Blood.* 2010;115(1):21-28.
- Piris M, Foucar K, Mollejo M, Campo E, Falini B. Splenic B-cell lymphoma/leukaemia, unclassificable. In: Swerdlow S, Campo E, Harris NL, et al, eds. WHO Classification of Tumours of Haematopoietic and Lymphoid Tissues. 4th ed. Lyon, France: International Agency for Research on Cancer (IARC); 2008:191-193.
- Falini B, Tiacci E, Liso A, et al. Simple diagnostic assay for hairy cell leukaemia by immunocytochemical detection of annexin A1 (ANXA1). *Lancet*. 2004;363(9424):1869-1870.

- Dong HY, Weisberger J, Liu Z, Tugulea S. Immunophenotypic analysis of CD103+ B-lymphoproliferative disorders: hairy cell leukemia and its mimics. *Am J Clin Pathol.* 2009;131(4): 586-595.
- Sadik W, Coupland S, Thachil J. Hairy cell leukaemia, negative for conventional cell markers, diagnosed using antibodies to annexin A1 and T-bet. *Br J Haematol.* 2010;151(3):207.
- Tiacci E, Trifonov V, Schiavoni G, et al. BRAF mutations in hairy-cell leukemia. N Engl J Med. 2011;364(24):2305-2315.
- Flaherty KT, Puzanov I, Kim KB, et al. Inhibition of mutated, activated BRAF in metastatic melanoma. N Engl J Med. 2010;363(9):809-819.
- Noel P. Definition of remission, minimal residual disease, and relapse in hairy cell leukemia bone marrow biopsy histology and immunohistology specimens. *Leuk Lymphoma*. 2011;52(suppl 2): 62-64.
- Tallman MS. Implications of minimal residual disease in hairy cell leukemia after cladribine using immunohistochemistry and immunophenotyping. *Leuk Lymphoma*. 2011;52(suppl 2):65-68.
- Case M, Matheson E, Minto L, et al. Mutation of genes affecting the RAS pathway is common in

A.A., D.C., E.S., P.F.d.C., C.D.B, A.P., R.F., and G.I. provided patients' samples and clinicopathologic data, and commented on the manuscript; A.S. purified leukemic cells from patients' peripheral blood samples; and B.F. led the project, supervised the study, and wrote the manuscript.

Conflict-of-interest disclosure: E.T. and B.F. applied for a patent on the clinical use of BRAF mutants in HCL. The remaining authors declare no competing financial interests.

Correspondence: Prof Brunangelo Falini, Institute of Hematology, University of Perugia, Ospedale S. Maria della Misericordia, S. Andrea delle Fratte, 06132 Perugia, Italy; e-mail: faliniem@ unipg.it; or Dr Enrico Tiacci, Institute of Hematology, University of Perugia, Ospedale S. Maria della Misericordia, S. Andrea delle Fratte, 06132 Perugia, Italy; e-mail: etiacci@ solido.umbria.it.

childhood acute lymphoblastic leukemia. *Cancer Res.* 2008;68(16):6803-6809.

- Gustafsson B, Angelini S, Sander B, et al. Mutations in the BRAF and N-ras genes in childhood acute lymphoblastic leukaemia. *Leukemia*. 2005; 19(2):310-312.
- Davidsson J, Lilljebjorn H, Panagopoulos I, et al. BRAF mutations are very rare in B- and T-cell pediatric acute lymphoblastic leukemias. *Leukemia*. 2008;22(8):1619-1621.
- Lee JW, Yoo NJ, Soung YH, et al. BRAF mutations in non-Hodgkin's lymphoma. *Br J Cancer.* 2003;89(10):1958-1960.
- Chapman MA, Lawrence MS, Keats JJ, et al. Initial genome sequencing and analysis of multiple myeloma. *Nature*. 2011;471(7339):467-472.
- Boyd EM, Bench AJ, van 't Veer MB, et al. High resolution melting analysis for detection of BRAF exon 15 mutations in hairy cell leukaemia and other lymphoid malignancies. *Br J Haematol.* 2011;155(5):609-612.
- Arcaini L, Zibellini S, Boveri E, et al. The BRAF V600E mutation in hairy cell leukemia and other mature B-cell neoplasms [published online ahead of print November 9, 2011]. *Blood*. doi: 10.1182/ blood-2011-08-368209.