Brief report

Dicer is selectively important for the earliest stages of erythroid development

*Natalija Buza-Vidas,^{1,2} *Valeriu B. Cismasiu,¹ Susan Moore,² Adam J. Mead,¹ Petter S. Woll,¹ Michael Lutteropp,¹ Luca Melchiori,¹ Sidinh Luc,¹ Tiphaine Bouriez-Jones,¹ Deborah Atkinson,¹ Donal O'Carroll,³ †Sten Eirik W. Jacobsen,¹ and †Claus Nerlov^{2,3}

¹Haematopoietic Stem Cell Laboratory, MRC Molecular Haematology Unit, Weatherall Institute of Molecular Medicine, University of Oxford, John Radcliffe Hospital, Oxford, United Kingdom; ²Institute for Stem Cell Research, MRC Centre for Regenerative Medicine, University of Edinburgh, Edinburgh, United Kingdom; and ³EMBL Mouse Biology Program, Adriano Buzzati-Traverso Campus, Monterotondo, Italy

MicroRNAs (miRs) are involved in many aspects of normal and malignant hematopoiesis, including hematopoietic stem cell (HSC) self-renewal, proliferation, and terminal differentiation. However, a role for miRs in the generation of the earliest stages of lineage committed progenitors from HSCs has not been identified. Using *Dicer* inactivation, we show that the miR complex is not only essential for HSC

maintenance but is specifically required for their erythroid programming and subsequent generation of committed erythroid progenitors. In bipotent pre-MegEs, loss of *Dicer* up-regulated transcription factors preferentially expressed in megakaryocyte progenitors (*Gata2* and *Zfpm1*) and decreased expression of the erythroid-specific *Klf1* transcription factor. These results show a specific requirement for *Dicer* in acquisition of erythroid lineage programming and potential in HSCs and their subsequent erythroid lineage differentiation, and in particular indicate a role for the miR complex in achieving proper balance of lineage-specific transcriptional regulators necessary for HSC multilineage potential to be maintained. (*Blood.* 2012;120(12):2412-2416)

Introduction

MicroRNAs (miRs) are an abundant class of endogenous small noncoding RNAs that regulate gene expression, such as gene silencing and degradation, uncovering a new layer of regulatory control of gene expression in many organisms.¹ To date, almost 1000 different miRs have been identified, playing important roles in proliferation, differentiation, and apoptosis.²

miR biogenesis requires successive processing events carried out by the RNAses Drosha and Dicer. Thus, in the absence of Dicer, essentially all miRs are depleted, making Dicer deletion a method by which their role in biological processes may be assessed. Recently, it was suggested that Dicer is involved in regulation of the hematopoietic stem cell (HSC) niche³ as well as in intrinsic regulation of HSCs themselves.⁴ Individual miRs have been implicated in differentiation of mature blood cell lineages, such as B cells,⁵ T cells,⁶ and granulocytes.⁷ Recently, miR-141/451⁸ and miR-191⁹ were implicated in terminal erythroid differentiation, and in vitro studies have suggested that miR-150 is involved in the development of megakaryocytes from megakaryocyte/erythroid progenitors.¹⁰ However, a requirement for Dicer in regulating the early stages of myelo/erythroid lineage commitment remains to be demonstrated.

By deleting Dicer from the hematopoietic system, we show that Dicer is not only crucial for maintenance of HSCs but is selectively required for generation of the most primitive erythroid committed progenitors, whereas megakaryocyte and myeloid lineage commitment remains intact.

Methods

Mice

Dicer1^{fl/fl,11} *Mx1-Cre*,¹² and *R26CreERT2*¹³ mice have been described. C57BI/6SJLCD45.1 mice were from The Jackson Laboratory. Experiments were performed with United Kingdom Home Office approval.

Gene expression analysis

Microarray analysis was performed as previously described.¹⁴ Microarray data have been deposited in the ArrayExpress database (www.ebi.ac.uk/arrayexpress; accession no. E-MTAB-674). Quantitative PCR analysis was performed using the BioMark Dynamic Array platform (Fluidigm) and TaqMan Gene Expression Assays (Applied Biosystems; supplemental Methods, available on the *Blood* Web site; see the Supplemental Materials link at the top of the online article).

In vitro progenitor assays

Evaluation of erythroid (E) and myeloid (GM) potential was performed as described,¹⁵ and megakaryocyte (Mk) potential with the Megacult assay (StemCell Technologies) according to the manufacturer's instructions (supplemental Methods).

FACS

FACS analysis of peripheral blood and BM stem and progenitor cells was done on a LSRII analyzer and sorts on a FACSAriaII (BD Biosciences). For further details, see supplemental Methods.

Statistical analysis

Statistical significance was calculated using Mann-Whitney t test.

Submitted October 2, 2011; accepted July 26, 2012. Prepublished online as *Blood* First Edition paper, August 6, 2012; DOI 10.1182/blood-2011-10-383653.

*N.B.-V. and V.B.C. contributed equally to this study.

†S.E.W.J. and C.N. contributed equally to this study.

The online version of this article contains a data supplement.

The publication costs of this article were defrayed in part by page charge payment. Therefore, and solely to indicate this fact, this article is hereby marked "advertisement" in accordance with 18 USC section 1734.

© 2012 by The American Society of Hematology



Figure 1. Dicer regulates erythroid intege transcriptional priming in HSCs. Leftaily irradiated (9 Gy) wild-type CD45.1 mice were transplanted noncompetitively with 6 million BM cells from control (*Dicer*^(M) or *Dicer*^(M), *Mx1Cre*) or *Dicer*^(M), *Mx1Cre*) mice. At least 4 weeks after transplantation, Cre-mediated recombination was induced with polyriboinosinic acid/polyribocytidylic acid (polyl:C), by intraperitoneal injections of 200 µg polyl:C every second day, 3 injections in total. (A) Representative FACS profiles of HSCs (LSKCD105⁺CD150⁺) in *Dicer*^(M) and *Dicer*^(M) mice, respectively. Numbers in FACS profiles indicate percentage within indicated gate, relative to total BM cells. (B-C) Mean (± SEM) frequency relative to total BM cells (B; 10 or 11 mice/group, analyzed individually) and absolute number per 2 tibiae (C) of LSKCD105⁺CD150⁺ HSCs in *Dicer*^(M) and *Dicer*^(M) mice. respectively. The context of t

Results and discussion

Loss of erythroid transcriptional priming in Dicer-deficient HSCs

Previous reports suggested that miRs control HSC numbers.⁴ Through competitive transplantation experiments, using 2 different Cre lines, we found Dicer to be essential for maintenance of HSCs

(supplemental Figure 1). To assess whether loss of *Dicer* from HSCs affects their lineage transcriptional priming we noncompetitively transplanted *Dicer*^{fl/fl};*Mx1Cre* and control *Dicer*^{fl/fl} BM cells. BM cellularity was reduced > 2-fold in *Dicer*^{Δ/Δ} mice 13 days after deletion (supplemental Figure 2A). However, at this time point, phenotypic LSKCD105⁺CD150⁺ HSCs were present at normal levels (Figure 1A-C), even though *Dicer* mRNA was efficiently



Figure 2. Dicer is critical for the earliest stages of the ervthroid progenitor compartment. (A-C) Mean (\pm SEM) CFU-GM (A), CFU-Mk (B), and BFU-E (C) colonies at 8 to 9 days of culture from unfractionated BM cells isolated 12 days after intraperitoneal injections of 200 µg polyI:C every second day, 3 injections in total, of 11-day-old $Dicer^{fl/fl}$ (n = 4) and $Dicer^{fl/fl}$; Mx1Cre mice (n = 8). Each mouse was analyzed in quadruplicate. (D) Representative FACS profiles of myelo/erythroid progenitor cells from lethally irradiated CD45.1 wild-type mice transplanted noncompetitively with Dicer#/# and Dicer Δ/Δ (CD45.2) BM cells, analyzed 12 days after polyI:C treatment. Numbers indicate percentage within indicated gate, of total BM cells, (E-F) Mean (± SEM) frequencies of test cell-derived myelo/erythroid progenitors in transplanted mice. n = 10 or 11 mice/group, analyzed individually. (G) Mean (± SEM) numbers of test cell-derived myelo/erythroid progenitors in transplanted mice. n = 4 or 5 mice/group, analyzed individually. (H) Dicer deletion efficiency in pre-GMs, GMPs, pre-MegEs, and MkPs isolated from mice transplanted with $Dicer^{\Delta/\Delta}$ BM cells and injected with polyI:C 12 days prior to analysis. Data are mean (\pm SEM) values from cells purified from 2 different pools of mice/genotype, 2 or 3 replicate wells/pool. (I) Megakarvocvte/ervthroid gene expression in 100-200 pre-MegEs isolated from Dicer#/# and Dicer $\Delta \Delta$ mice. Mean (\pm SEM) expression in Dicer $\Delta \Delta$ relative to control *Dicer^{fl/fl}* pre-MegEs, calculated using the $2^{-\Delta\Delta CT}$ method.¹⁷ Data generated from sorts performed on 6 cohorts of mice/genotype (2 or 3 mice/ cohort), 13 or 14 technical replicates, 3 experiments in total. *P < .05. **P < .01. ***P < .001. ns indicates not significant. Statistics refer to comparison between Dicer mice and Dicer^{fl/fl} control mice.

deleted (Figure 1D). Using global gene profiling of $Dicer^{\Delta/+}$ and $Dicer^{\Delta/-}$ LSKFlt3⁻ HSCs (Figure 1E) and gene set enrichment analysis,¹⁶ we observed that while lymphoid (pre-B), myeloid (GM), and megakaryocyte (Mk) lineage programming was not significantly affected by *Dicer* deletion (Figure 1F-H), there was a marked overall down-regulation of erythroid (E) genes in $Dicer^{\Delta/-}$ HSCs (Figure 1I). Importantly, because Dicer-deleted LSKCD105⁺CD150⁺ HSCs were present at normal levels at this time point (Figure 1A-B), any observed changes in transcriptional

lineage priming in HSCs, or output of distinct progenitors, are unlikely in this setting to be because of depletion of HSCs.

In vitro colony assays revealed normal Mk colony numbers and mildly reduced CFU-GM, in striking contrast to the complete loss of BFU-Es in *Dicer*^{Δ/Δ} BM, both upon direct deletion (Figure 2A-C) and deletion after noncompetitive transplantation (supplemental Figure 2B-E). To establish at which stage of the erythroid differentiation pathway *Dicer* is required, detailed FACS staging of myelo/erythroid progenitors in *Dicer*^{Δ/Δ} BM was performed.¹⁸ This analysis demonstrated a specific and complete loss of the earliest erythroid committed progenitors (pre-CFU-Es, CFU-Es) and a reduction in the preceding pre-MegE progenitors in *Dicer*^{Δ/Δ} BM (Figure 2D-G), whereas MkPs and pre-GMs were present at normal levels, indicating unimpaired megakaryocytic and myeloid lineage commitment. Consistent with the lower number of CFU-GM, granulocyte-monocyte progenitor (GMPs) were reduced (Figure 2D-G). Specific deletion of Dicer in myeloid progenitors, using *Cebpa-Cre*, does not result in impaired GMP formation.¹⁹ The observed reduction of GMPs may therefore reflect that earlier, and possibly more efficient, Dicer depletion is able to reveal a role for miRs in myeloid lineage progression.

Blood analysis 12 days after deletion revealed a significant reduction in red blood cells, hemoglobin, and hematocrit, but not platelet numbers (supplemental Figure 2F), consistent with Dicer being required for erythroid, but not megakaryocyte lineage commitment. White blood cells were also reduced in Dicer-deficient mice (supplemental Figure 2F), most probably reflecting the reported role of Dicer in maturation of granulocytes and lymphocytes.⁷⁻⁹

Importantly, analysis of myelo-erythroid progenitors in $Dicer^{\Delta/\Delta}$ BM confirmed the absence of Dicer mRNA (Figure 2H). Consistent with this, the Dicer-dependent miR-148a, but not the Dicerindependent miR-451, was depleted in both HSCs and pre-MegEs (supplemental Figure 2G). Expression of common Mk/E transcription factors Gata1, Gfilb, Runx1, and Fli1 was unaffected in $Dicer^{\Delta/\Delta}$ pre-MegEs (Figure 2I), whereas expression of the Mkrelated Gata2, Zfpm1, and Mpl was increased, and the E genes Epor and Klf1 decreased in pre-MegEs. Notably, at the normal Mk/E bifurcation, Gata2 and Zfpm1 are up-regulated in MkPs, whereas Klf1 is up-regulated in pre-CFU-Es.¹⁸ Likewise, Mpl, the key cytokine receptor for early Mk development, was up-regulated in *Dicer* Δ/Δ pre-MegEs. These data provide support for miRs acting as developmental switches during cell fate determination by finetuning the lineage-specific transcriptomes. The changes in lineagespecific transcription factor and cytokine receptor expression are probably the result of a wide-ranging deregulation of miRcontrolled genes. Using gene set enrichment analysis, we observed that 203 of 219 gene sets defined by specific miR seeds being present in the 3'-untranslated region showed up-regulation of gene expression in $Dicer^{\Delta/\Delta}$ HSCs, with 122 of 219 of these being significantly up-regulated (P < .05; false discovery rate < 0.25; no gene sets significantly down-regulated; supplemental Table 1). These data confirm a general role for miRs in suppression of mRNA expression but also indicate that a large number of miRs may contribute to the observed phenotype, although it needs to be considered that pre-miR accumulation may also play a role. Furthermore, recently it was suggested that Dicer may also regulate gene expression by binding chromatin directly, suggesting that Dicer might also affect hematopoiesis through siRNA-independent mechanisms.20

So far, miRs have not been implicated in the early stages of erythroid progenitors, but rather in maturation of erythroid cells, such as miR-144/451⁸ and miR-191.⁹ Furthermore, miR-150 has been suggested to drive the differentiation of megakaryocyte/ erythroid progenitors into the Mk lineage at the expense of the erythroid lineage.¹⁰ The results obtained here show that *Dicer*-deficient hematopoiesis already at the stem cell level is depleted of erythroid gene expression; and in line with this, *Dicer*^{Δ/Δ} HSCs subsequently fail to generate the earliest stages of erythroid lineage commitment seems unaffected. This both identifies the miR complex as selectively required for the generation of the earliest stages of erythroid commitment and supports the proposed role of HSC lineage priming in subsequent commitment events.²¹⁻²⁴

Acknowledgments

The authors thank the Biomedical Services at Oxford University and Theresa Furey and Helen Ferry for technical assistance, Paul Collier and Vladimir Benes (EMBL, Heidelberg) for help with miR quantitative RT-PCR analysis, and Elizabeth Zuo (Stanford University) for help with affymetrix analysis.

This work was supported by the United Kingdom MRC (Strategic Awards and Program Grants, S.E.W.J. and C.N.) and the Swedish Research Council (HematoLinne Strategic Award). A.J.M. was supported by the Leukemia and Lymphoma Research Senior Bennett Fellowship. D.A. was supported by the Leukemia and Lymphoma Research Foundation.

Authorship

Contribution: S.E.W.J., C.N., V.B.C., and N.B.-V. designed and conceptualized the overall research, analyzed the data, and wrote the manuscript; V.B.C., N.B.-V., P.S.W., M.L., A.J.M., L.M., S.L., T.B.-J., and D.A. performed experiments; V.B.C., S.M., and C.N. performed Affymetrix analysis; and D.O. provided Dicer conditional knockout mice.

Conflict-of-interest disclosure: The authors declare no competing financial interests.

Correspondence: Claus Nerlov, Institute for Stem Cell Research, University of Edinburgh, Edinburgh EH9 16UU, United Kingdom; e-mail: cnerlov@staffmail.ed.ac.uk; and Sten Eirik W. Jacobsen, Weatherall Institute of Molecular Medicine, University of Oxford, Oxford OX3 9DS, United Kingdom; e-mail: sten.jacobsen@imm.ox.ac.uk.

- 1. Bushati N, Cohen SM. microRNA functions. Annu Rev Cell Dev Biol. 2007;23:175-205.
- Havelange V, Garzon R. MicroRNAs: emerging key regulators of hematopoiesis. *Am J Hematol.* 2010;85(12):935-942.
- Raaijmakers MH, Mukherjee S, Guo S, et al. Bone progenitor dysfunction induces myelodysplasia and secondary leukaemia. *Nature*. 2010; 464(7290):852-857.
- Guo S, Lu J, Schlanger R, et al. MicroRNA miR-125a controls hematopoietic stem cell number. Proc Natl Acad Sci U S A. 2010;107(32):14229-14234.
- 5. Xiao C, Calado DP, Galler G, et al. MiR-150 con-

trols B cell differentiation by targeting the transcription factor c-Myb. *Cell.* 2007;131(1):146-159.

- Muljo SA, Ansel KM, Kanellopoulou C, Livingston DM, Rao A, Rajewsky K. Aberrant T cell differentiation in the absence of Dicer. J Exp Med. 2005;202(2):261-269.
- Johnnidis JB, Harris MH, Wheeler RT, et al. Regulation of progenitor cell proliferation and granulocyte function by microRNA-223. *Nature*. 2008;451(7182):1125-1129.
- Rasmussen KD, Simmini S, Abreu-Goodger C, et al. The miR-144/451 locus is required for erythroid homeostasis. J Exp Med. 2010;207(7):1351-1358.
- 9. Zhang L, Flygare J, Wong P, Lim B, Lodish HF.

miR-191 regulates mouse erythroblast enucleation by down-regulating Riok3 and Mxi1. *Genes Dev.* 2011;25(2):119-124.

- Lu J, Guo S, Ebert BL, et al. MicroRNA-mediated control of cell fate in megakaryocyte-erythrocyte progenitors. *Dev Cell*. 2008;14(6):843-853.
- Yi R, O'Carroll D, Pasolli HA, et al. Morphogenesis in skin is governed by discrete sets of differentially expressed microRNAs. *Nat Genet.* 2006; 38(3):356-362.
- Kuhn R, Schwenk F, Aguet M, Rajewsky K. Inducible gene targeting in mice. *Science*. 1995; 269(5229):1427-1429.
- 13. Soriano P. Generalized lacZ expression with the

ROSA26 Cre reporter strain. *Nat Genet.* 1999; 21(1):70-71.

- Kirstetter P, Schuster MB, Bereshchenko O, et al. Modeling of C/EBPalpha mutant acute myeloid leukemia reveals a common expression signature of committed myeloid leukemia-initiating cells. *Cancer Cell*. 2008;13(4):229-310.
- Luc S, Anderson K, Kharazi S, et al. Downregulation of Mpl marks the transition to lymphoid-primed multipotent progenitors with gradual loss of granulocyte-monocyte potential. *Blood.* 2008;111(7):3424-3434.
- Subramanian A, Tamayo P, Mootha VK, et al. Gene set enrichment analysis: a knowledgebased approach for interpreting genome-wide expression profiles. *Proc Natl Acad Sci U S A*. 2005;102(43):15545-15550.
- 17. Livak KJ, Schmittgen TD. Analysis of relative

gene expression data using real-time quantitative PCR and the 2(-delta delta C(T)) Method. *Methods.* 2001;25(4):402-408.

- Pronk CJ, Rossi DJ, Mansson R, et al. Elucidation of the phenotypic, functional, and molecular topography of a myeloerythroid progenitor cell hierarchy. *Cell Stem Cell*. 2007;1(4):428-442.
- Alemdehy MF, van Boxtel NG, de Looper HW, et al. Dicer1 deletion in myeloid-committed progenitors causes neutrophil dysplasia and blocks macrophage/dendritic cell development in mice. *Blood*. 2012;119(20):4723-4730.
- Cernilogar FM, Onorati MC, Kothe GO, et al. Chromatin-associated RNA interference components contribute to transcriptional regulation in Drosophila. *Nature.* 2011;480(7377):391-395.
- 21. Hu M, Krause D, Greaves M, et al. Multilineage gene expression precedes commitment in the

hemopoietic system. *Genes Dev.* 1997;11(6):774-785.

- Adolfsson J, Mansson R, Buza-Vidas N, et al. Identification of Flt3+ lympho-myeloid stem cells lacking erythro-megakaryocytic potential a revised road map for adult blood lineage commitment. *Cell.* 2005;121(2):295-306.
- Bereshchenko O, Mancini E, Moore S, et al. Hematopoietic stem cell expansion precedes the generation of committed myeloid leukemiainitiating cells in C/EBPalpha mutant AML. *Cancer Cell*. 2009;16(5):390-400.
- Mansson R, Hultquist A, Luc S, et al. Molecular evidence for hierarchical transcriptional lineage priming in fetal and adult stem cells and multipotent progenitors. *Immunity.* 2007;26(4):407-419.