

Brief report

Targeted genomic sequencing of pediatric Burkitt lymphoma identifies recurrent alterations in antiapoptotic and chromatin-remodeling genes

Lisa Giulino-Roth,¹⁻³ Kai Wang,⁴ Theresa Y. MacDonald,³ Susan Mathew,³ Yifang Tam,³ Maureen T. Cronin,⁴ Gary Palmer,⁴ Norma Lucena-Silva,⁵ Francisco Pedrosa,⁵ Marcia Pedrosa,⁵ Julie Teruya-Feldstein,⁶ Govind Bhagat,⁷ Bachir Alobeid,⁷ Lorenzo Leoncini,⁸ Cristiana Bellan,⁸ Emily Rogena,⁹ Kerice A. Pinkney,¹⁰ Mark A. Rubin,³ Raul C. Ribeiro,¹¹ Roman Yelensky,⁴ Wayne Tam,³ Philip J. Stephens,⁴ and Ethel Cesarman³

¹Department of Pediatrics, Memorial Sloan-Kettering Cancer Center, New York, NY; ²Department of Pediatrics, Division of Pediatric Hematology and Oncology, Weill Cornell Medical College, New York, NY; ³Department of Pathology and Laboratory Medicine, Weill Cornell Medical College, New York, NY; ⁴Foundation Medicine, Cambridge, MA; ⁵Department of Pediatric Oncology, Institute of Integrative Medicine Prof Fernando Figueira (IMIP), Recife, Brazil; ⁶Department of Pathology, Memorial Sloan-Kettering Cancer Center, New York, NY; ⁷Department of Pathology and Cell Biology, Division of Hematopathology, Columbia University Medical Center, New York Presbyterian Hospital, New York, NY; ⁸Department of Human Pathology and Oncology, University of Siena, Siena, Italy; ⁹University of Nairobi, Nairobi, Kenya; ¹⁰Department of Pediatrics, Division of Pediatric Hematology/Oncology, Columbia University Medical Center, New York, NY; and ¹¹International Outreach Program and Department of Oncology, St Jude Children's Research Hospital, Memphis, TN

To ascertain the genetic basis of pediatric Burkitt lymphoma (pBL), we performed clinical-grade next-generation sequencing of 182 cancer-related genes on 29 formalin-fixed, paraffin embedded primary pBL samples. Ninety percent of cases had at least one mutation or genetic alteration, most commonly involving *MYC* and *TP53*.

EBV(–) cases were more likely than EBV(+) cases to have multiple mutations ($P < .0001$). Alterations in tumor-related genes not previously described in BL were identified. Truncating mutations in *ARID1A*, a member of the SWI/SNF nucleosome remodeling complex, were seen in 17% of cases. *MCL1* pathway alterations were found in 22% of

cases and confirmed in an expanded panel. Other clinically relevant genomic alterations were found in 20% of cases. Our data suggest the roles of *MCL1* and *ARID1A* in BL pathogenesis and demonstrate that comprehensive genomic profiling may identify additional treatment options in refractory disease. (*Blood*. 2012;120(26):5181-5184)

Introduction

Burkitt lymphoma (BL) is an aggressive B-cell malignancy that predominantly affects the pediatric population. Although most children are cured with intensive chemotherapy, up to 20% die of relapsed or refractory disease.¹⁻³ Cure rates are significantly lower in developing countries that have a greater incidence of BL, making BL a global health concern. The molecular hallmark of BL is the translocation of the *MYC* proto-oncogene to the immunoglobulin-heavy or one of the light chain genes, leading to constitutive *MYC* activation. Additional molecular alterations that may counteract *MYC*-induced proapoptotic signals are likely relevant in the pathogenesis of BL. RNA sequencing recently has been performed to investigate the genetic landscape of BL via the use of a cohort that combined pediatric and adult cases.⁴ In contrast to adult cases, which typically have a simple karyotype, 60%-90% of pediatric tumors have secondary chromosomal abnormalities, the consequences of which are less well characterized.⁵⁻⁸ Therefore, we focused on pediatric BL (pBL) to better understand the driving genomic alterations in this disease and to aid the development of rational therapeutics.

Genomic studies in rare tumors previously have been limited by the availability of frozen tissue to obtain DNA. In this report, we demonstrate comprehensive next-generation sequencing on formalin-fixed, paraffin embedded (FFPE) tissue, which allowed the use of archived specimens. We identified mutations in a

significant proportion of pBL, including recurrent alterations in *ARID1A* and *MCL1* not previously reported.

Methods

Patient samples

Eighty-two FFPE samples of pBL were collected from sites in the United States, Kenya, and Brazil. Cases were included if the patient age was < 21 years and the diagnosis of BL was confirmed. Cases included endemic (n = 20), sporadic (n = 60), and HIV-associated (n = 2) pBL. All samples were obtained with institutional research board and biospecimen-use approval. From this panel, a cohort was selected for next-generation sequencing on the basis of tumor representation > 80%, sufficient tissue availability, and adequate DNA yield (supplemental Methods, available on the *Blood* Web site; see the Supplemental Materials link at the top of the online article). This study was conducted in accordance with the Declaration of Helsinki.

Tumor characterization, FISH, and immunohistochemistry

See supplemental Methods.

Targeted genomic sequencing

DNA was extracted from FFPE tissue (supplemental Methods). Molecular barcode-indexed ligation-based sequencing libraries were constructed

Submitted June 15, 2012; accepted October 2, 2012. Prepublished online as *Blood* First Edition paper, October 22, 2012; DOI 10.1182/blood-2012-06-437624.

The online version of this article contains a data supplement.

The publication costs of this article were defrayed in part by page charge payment. Therefore, and solely to indicate this fact, this article is hereby marked "advertisement" in accordance with 18 USC section 1734.

© 2012 by The American Society of Hematology




Figure 1. Genomic alterations in pBL. (A) Diagram of mutations identified by targeted genomic sequencing. Columns represent individual samples. Alterations listed include only those likely to be somatic as referenced in COSMIC,²² as well as *MYC* point mutations and indels of unknown significance. We detected copy number alterations by comparing targeted genomic DNA sequence coverage with a process-matched normal control sample. We detected genomic rearrangements by clustering chimeric reads mapping to targeted introns. (B) Distribution of frequency of alterations by EBV status. EBV-negative cases (EBV Neg) are significantly more likely to have > 1 alteration compared with EBV-positive cases (EBV Pos; $P < .0001$).

using 50-200 ng of total DNA. Libraries were hybridization captured with custom biotinylated RNA oligo pools (custom SureSelect kit; Agilent) representing 3230 exons of 182 cancer-related genes plus 37 introns from 14 genes often rearranged in cancer (supplemental Table 1). Paired end sequencing (49 × 49 cycles) was performed with the HiSeq2000 (Illumina). Sequence analysis is detailed in the supplemental Methods.

Results and discussion

Sequence analysis of 29 cases of pBL at an average coverage of 653-fold identified 99 somatic genetic alterations in 19 genes, including 68 base substitutions, 10 insertions/deletions, 12 copy number alterations, and 9 structural rearrangements (Figure 1A, supplemental Table 2). Twenty-six cases were sporadic (14 from the United States; 12 from Brazil) and 3 endemic (Kenya); 2 were HIV+ (both sporadic). The most frequent genomic alterations were point mutations/indels in *MYC* (58.6%) and *TP53* mutations (41.4%). *MYC* mutations spanned the coding region and included

hot spots previously documented in lymphomas (supplemental Table 3, supplemental Figure 1).⁹ Cases with *MYC* mutations also demonstrated *MYC* translocation, confirming that mutations may functionally cooperate with translocation to promote *MYC*-mediated oncogenesis. Twenty of 29 cases (69%) had genetic alterations in addition to those involving *MYC*.

Although mutations did not group with epidemiologic subtypes, EBV(-) cases were more likely to have multiple genetic alterations than EBV(+) cases. When we excluded *MYC* alterations, 12 of 14 EBV(-) cases had > 1 alteration and 0 of 11 EBV(+) cases had > 1 alteration ($P < .0001$; Figure 1B). This is consistent with the tumorigenic role of EBV in a subset of BL. *TP53* was altered in 4 EBV(+) and 8 EBV(-) cases.

Recurrent alterations were identified in cancer-related genes not previously reported in BL, including truncating mutations in *ARID1A* and amplification of *MCL1* (Figure 2, supplemental Table 4). *ARID1A* is a member of the SWI/SNF family of complexes, which function as chromatin remodelers, and has been implicated




Figure 2. Primary BL cases demonstrate recurrent alterations in *ARID1A* and *MCL1*. (A) Representation of *ARID1A* gene and mutations identified by sequencing. The 20 exons of *ARID1A* are represented in the green boxes. Diamonds represent deletion; triangles represent point mutation. In all 5 cases, the mutation was truncating. ARID indicates AT-rich interactive domain; LXXLL, leucine-rich motifs; nt, nucleotide; and UTR, untranslated region. (B) *ARID1A* protein expression in mutated and wild-type cases as determined by IHC. Cases B-6 and BL-17 have truncating mutations upstream to the antibody epitope and demonstrate decreased staining compared with cases lacking mutations. Case BL-27 has a mutation downstream of the antibody probe. (C) *MCL1* pathway-altered cases as identified by sequencing. Five cases had a 1.8 to 3.1 copy gain of *MCL1* relative to a diploid control, corresponding to a predicted 4-6 copies of *MCL1* per tumor cells. One case had a mutation in *FBXW7*, a gene in the *MCL1* pathway. (D) FISH for *MCL1* using a *MCL1* probe (red) and centromeric probe for chromosome 1 (green). A representative wild-type case (WT; left) and an amplified case (right) are shown. Amplified cases had 3-4 copies, but in some a signal was stronger, suggesting tandem duplication. (E) *MCL1* protein expression was evaluated by immunohistochemistry. Quantitative analysis of *MCL1* staining intensity in tonsil ($n = 3$), *MCL1* WT ($n = 24$), and *MCL1* amplified ($n = 5$) cases is shown. Error bars represent standard error of the mean. Unpaired *t* test was performed to evaluate statistical significance between expression in *MCL1* WT and *MCL1* amplified cases. (F) Representative normal and amplified cases are shown, as well as the pattern in a tonsil control showing stronger cytoplasmic expression in the follicles. Panels B and F, original magnification $\times 400$ with $40\times$ objective lens. Microscope: Olympus BX 41; camera: Olympus Q-COLOR3; software: QCapture Version 2.9.8.0 (Quantitative Imaging). Panel D, original magnification $\times 1000$; Apochromatic $100\times$ lens with 1.4 aperture; microscope: Olympus BX51; camera: Jai CV-A10CL; software: Cytovision Imaging Version 3.6 (Genetix).

as a tumor suppressor.¹⁰ Inactivating *ARID1A* mutations have been described in solid malignancies, including ovarian and gastric cancer,¹¹⁻¹³ and a tumor suppressor role of *ARID1A* is supported by functional studies.¹²

Mutations in *ARID1A* were found in 5 of 29 (17.2%; 95% confidence interval 5.8%-35.8%) of pBL cases, one of which was EBV(+). Mutations were distributed throughout the gene and all resulted protein truncation (Figure 2A), consistent with tumor suppressor role. In addition, one case with *ARID1A* mutation had a secondary mutation in *SNF5*, also a member of the SWI/SNF family. *ARID1A* protein expression was evaluated by immunohistochemistry in an expanded cohort ($n = 50$) that included 17 cases with known *ARID1A* mutation status. Cases with *ARID1A* alterations leading to truncated protein lacking the antibody epitope showed decreased expression (Figure 2B). In contrast, 10 of 12 cases with WT *ARID1A* demonstrated increased expression (Figure 2B, supplemental Figure 2). Overall 15 of 50 cases (30%) demonstrated *ARID1A* expression that was equal to or lower than that seen in cases with known mutation. There was no evidence of loss of heterozygosity in mutated cases because the mutation was present in close to 50% of the reads. As has been proposed for other malignancies, haploinsufficiency of *ARID1A* may be enough for cellular transformation.

Recurrent amplification was found in *MCL1*, a member of the *BCL2* family. *MCL1* and related proteins inhibit apoptosis by blocking the cell death mediators BAK and BAX.¹⁴ The impor-

tance of *MCL1* as an oncogene has been implicated in transgenic mice that develop aggressive B-cell lymphomas.^{15,16} *MCL1* is located on 1q21.2, a genomic region amplified in approximately 25% of pBL cases.¹⁷ Amplification of *MCL1* has been described in a BL subline¹⁸ but has not been reported in primary BL samples. *MCL1* overexpression may be clinically relevant because it has been linked to chemotherapy resistance,^{18,19} and several inhibitors that may target *MCL1* are in clinical development (supplemental Table 5).²⁰ We identified *MCL1* amplification in 5 of 29 (17.2%, 95% confidence interval 5.8%-35.8%) cases ranging from 1.8X to 3.1X copy gain relative to a diploid control (Figure 2C). In addition, one case had a point mutation in *FBXW7*, which encodes an ubiquitin ligase that targets *MCL1* for degradation.²¹ FISH for *MCL1* confirmed the sequencing results, and in an independent cohort, 5 of 17 (29%) cases demonstrated *MCL1* amplification by FISH (Figure 2D). In total 10 of 46 (21.7%) pBL cases demonstrated *MCL1* amplification. Evaluation of *MCL1* protein expression by immunohistochemistry and densitometry analysis revealed increased *MCL1* protein expression in amplified cases ($P = .002$; Figure 2E-F).

Alterations also were found in other cancer-related genes, including point mutations in *LRP6*; truncating alterations in *LRP1B*, *PTPRD*, *PTEN*, *NOTCH*, and *ATM*; amplifications of *RAF1*, *MDM4*, *MDM2*, *KRAS*, *IKBKE*, and *CDK6*; and a deletion of *CDKN2A*, many of which are targetable by therapies in clinical trials (supplemental Table 5).

This is the first report of next-generation sequencing focusing specifically on pediatric BL. Our work demonstrates the feasibility of genomic sequencing using FFPE specimens. We identified novel recurrent alterations in members of the SWI/SNF family of chromatin remodeling genes, the antiapoptotic gene *MCL1*, as well as other therapeutically actionable alterations. As the spectrum of mutations in pBL becomes further defined and genotype-phenotype and clinical correlations are established, inclusion of mutation profiling should become part of routine diagnostic testing, prognostic evaluation, and treatment of pBL. In particular, as we move into the era of precision medicine, the specific genomic information reported here should be useful in the molecular subclassification of pBL and use of therapies that target key biologic pathways such as chromatin remodeling and suppression of apoptosis.

Acknowledgments

The authors acknowledge Sharon Barouk for organizing the materials. They thank the Translational Research Program of the Department of Pathology and Laboratory Medicine of Weill Cornell Medical College for support with the construction of tissue microarrays and immunohistochemical characterization of the samples, and Robert Kim for help with image analysis.

This work was supported by funding from the Rally Foundation for Childhood Cancer Research to L.G.-R., the American Syrian

Lebanese Associated Charities (ALSAC) to R.C.R., and National Institutes of Health grant R01CA068939 to E.C.

Authorship

Contribution: L.G.-R. designed the study, performed research, analyzed data, and wrote the paper; K.W., M.T.C., G.P., R.Y., and P.J.S. performed sequencing studies and analyzed data; N.L.-S., F.P., M.P., J.T.F., G.B., B.A., L.L., C.B., E.R., and R.C.R. provided precious patient samples, characterized some of the samples, collected data, reviewed and approved the manuscript; K.A.P. collected data; T.Y.M., S.M., Y.T., and W.T. performed research and analyzed data; M.A.R. designed and supervised part of the study; and E.C. designed the study, supervised research, analyzed the data, and assisted in writing the paper.

Conflict-of-interest disclosure: K.W., M.T.C., G.P., R.Y., and P.J.S. are stockholders in Foundation Medicine and employees of Foundation Medicine. The remaining authors declare no competing financial interests.

Correspondence: Ethel Cesaran, MD, Department of Pathology and Laboratory Medicine, Weill Cornell Medical College, 1300 York Ave, New York, NY 10065; e-mail: ecesarn@med.cornell.edu; or Lisa Giulino-Roth, MD, Weill Cornell Medical College, 525 E 68th St, Payson 695, New York, NY 10065; e-mail: giulino@mskcc.org.

References

- Cairo MS, Gerrard M, Sposto R, et al. Results of a randomized international study of high-risk central nervous system B non-Hodgkin lymphoma and B acute lymphoblastic leukemia in children and adolescents. *Blood*. 2007;109(7):2736-2743.
- Patte C, Auferin A, Gerrard M, et al. Results of the randomized international FAB/LMB96 trial for intermediate risk B-cell non-Hodgkin lymphoma in children and adolescents: it is possible to reduce treatment for the early responding patients. *Blood*. 2007;109(7):2773-2780.
- Cairo MS, Krailo MD, Morse M, et al. Long-term follow-up of short intensive multiagent chemotherapy without high-dose methotrexate ('Orange') in children with advanced non-lymphoblastic non-Hodgkin's lymphoma: a children's cancer group report. *Leukemia*. 2002;16(4):594-600.
- Schmitz R, Young RM, Ceribelli M, et al. Burkitt lymphoma pathogenesis and therapeutic targets from structural and functional genomics. *Nature*. 2012;490(7418):116-120.
- Lai JL, Fenaux P, Zandacki M, Nelken B, Huart JJ, Deminatti M. Cytogenetic studies in 30 patients with Burkitt's lymphoma or L3 acute lymphoblastic leukemia with special reference to additional chromosome abnormalities. *Ann Genet*. 1989;32(1):26-32.
- Lones MA, Sanger WG, Le Beau MM, et al. Chromosome abnormalities may correlate with prognosis in Burkitt/Burkitt-like lymphomas of children and adolescents: a report from Children's Cancer Group Study CCG-E08. *J Pediatr Hematol Oncol*. 2004;26(3):169-178.
- Nelson M, Perkins SL, Dave BJ, et al. An increased frequency of 13q deletions detected by fluorescence in situ hybridization and its impact on survival in children and adolescents with Burkitt lymphoma: results from the Children's Oncology Group study CCG-5961. *Br J Haematol*. 2010;148(4):600-610.
- Onciu M, Schlette E, Zhou Y, et al. Secondary chromosomal abnormalities predict outcome in pediatric and adult high-stage Burkitt lymphoma. *Cancer*. 2006;107(5):1084-1092.
- Bahram F, von der Lehr N, Cetinkaya C, Larsson LG. c-Myc hot spot mutations in lymphomas result in inefficient ubiquitination and decreased proteasome-mediated turnover. *Blood*. 2000;95(6):2104-2110.
- Wilson BG, Roberts CW. SWI/SNF nucleosome remodellers and cancer. *Nat Rev Cancer*. 2011;11(7):481-492.
- Wiegand KC, Shah SP, Al-Agha OM, et al. ARID1A mutations in endometriosis-associated ovarian carcinomas. *N Engl J Med*. 2010;363(16):1532-1543.
- Zang ZJ, Cutcutache I, Poon SL, et al. Exome sequencing of gastric adenocarcinoma identifies recurrent somatic mutations in cell adhesion and chromatin remodeling genes. *Nat Genet*. 2012;44(5):570-574.
- Wang K, Kan J, Yuen ST, et al. Exome sequencing identifies frequent mutation of ARID1A in molecular subtypes of gastric cancer. *Nat Genet*. 2011;43(12):1219-1223.
- Youle RJ, Strasser A. The BCL-2 protein family: opposing activities that mediate cell death. *Nat Rev Mol Cell Biol*. 2008;9(1):47-59.
- Zhou P, Levy NB, Xie H, et al. MCL1 transgenic mice exhibit a high incidence of B-cell lymphoma manifested as a spectrum of histologic subtypes. *Blood*. 2001;97(12):3902-3909.
- Zhou P, Qian L, Bieszczyk CK, et al. Mcl-1 in transgenic mice promotes survival in a spectrum of hematopoietic cell types and immortalization in the myeloid lineage. *Blood*. 1998;92(9):3226-3239.
- Schiffman JD, Lorimer PD, Rodic V, et al. Genome wide copy number analysis of paediatric Burkitt lymphoma using formalin-fixed tissues reveals a subset with gain of chromosome 13q and corresponding miRNA over expression. *Br J Haematol*. 2011;155(4):477-486.
- Vrana JA, Bieszczyk CK, Cleveland ES, et al. An MCL1-overexpressing Burkitt lymphoma subline exhibits enhanced survival on exposure to serum deprivation, topoisomerase inhibitors, or staurosporine but remains sensitive to 1-beta-D-arabinofuranosylcytosine. *Cancer Res*. 2002;62(3):892-900.
- Wertz IE, Kusam S, Lam C, et al. Sensitivity to antitubulin chemotherapeutics is regulated by MCL1 and FBW7. *Nature*. 2011;471(7336):110-114.
- Azmi AS, Wang Z, Philip PA, Mohammad RM, Sarkar FH. Emerging Bcl-2 inhibitors for the treatment of cancer. *Expert Opin Emerg Drugs*. 2011;16(1):59-70.
- Inuzuka H, Shaik S, Onoyama I, et al. SCF(FBW7) regulates cellular apoptosis by targeting MCL1 for ubiquitylation and destruction. *Nature*. 2011;471(7336):104-109.
- Forbes SA, Bindal N, Bamford S, et al. COSMIC: mining complete cancer genomes in the Catalogue of Somatic Mutations in Cancer. *Nucleic Acids Res*. 2011;39(Database issue):D945-950.