

Multiple SNP testing improves risk prediction of first venous thrombosis

Hugoline G. de Haan,¹ Irene D. Bezemer,¹ Carine J. M. Doggen,^{1,2} Saskia Le Cessie,^{1,3} Pieter H. Reitsma,^{4,5} Andre R. Arellano,⁶ Carmen H. Tong,⁶ James J. Devlin,⁶ Lance A. Bare,⁶ Frits R. Rosendaal,^{1,4,5} and Carla Y. Vossen^{1,7}

¹Department of Clinical Epidemiology, Leiden University Medical Center, Leiden, The Netherlands; ²Department of Health Technology & Services Research, MIRA Institute for Biomedical Technology and Technical Medicine, Enschede, The Netherlands; ³Department of Medical Statistics, Leiden University Medical Center, Leiden, The Netherlands; ⁴Eindhoven Laboratory for Experimental Vascular Medicine, Leiden University Medical Center, Leiden, The Netherlands; ⁵Department of Thrombosis and Hemostasis, Leiden University Medical Center, Leiden, The Netherlands; ⁶Celera, Alameda, CA; and ⁷Department of Medical Genetics, University Medical Center Utrecht, Utrecht, The Netherlands

There are no risk models available yet that accurately predict a person's risk for developing venous thrombosis. Our aim was therefore to explore whether inclusion of established thrombosis-associated single nucleotide polymorphisms (SNPs) in a venous thrombosis risk model improves the risk prediction. We calculated genetic risk scores by counting risk-increasing alleles from 31 venous thrombosis-associated SNPs for subjects of a large case-control study,

including 2712 patients and 4634 controls (Multiple Environmental and Genetic Assessment). Genetic risk scores based on all 31 SNPs or on the 5 most strongly associated SNPs performed similarly (areas under receiver-operating characteristic curves [AUCs] of 0.70 and 0.69, respectively). For the 5-SNP risk score, the odds ratios for venous thrombosis ranged from 0.37 (95% confidence interval [CI], 0.25-0.53) for persons with 0 risk alleles to 7.48 (95% CI, 4.49-12.46) for persons

with more than or equal to 6 risk alleles. The AUC of a risk model based on known nongenetic risk factors was 0.77 (95% CI, 0.76-0.78). Combining the nongenetic and genetic risk models improved the AUC to 0.82 (95% CI, 0.81-0.83), indicating good diagnostic accuracy. To become clinically useful, subgroups of high-risk persons must be identified in whom genetic profiling will also be cost-effective. (*Blood*. 2012;120(3):656-663)

Introduction

Venous thrombosis is the result of innate thrombotic tendency and nongenetic triggers. Many common genetic variants, mainly single nucleotide polymorphisms (SNPs), with modest effects on risk of venous thrombosis have been reported.¹ Individual SNPs have little predictive value because of their modest effect on risk, but combinations of gene variants may improve the predictive ability and could be used to model susceptibility to venous thrombosis.

Simulation studies have shown that so-called genetic profiling may be useful to discriminate between persons with high risk of disease and those with low risk. The discriminative accuracy of genetic profiling depends on the heritability and incidence of the disease and on the frequencies of risk alleles.^{2,3}

Genetic profiling has become a popular aim in epidemiologic studies of many common diseases because a large amount of data from genome-wide association studies has become available.²⁻⁸ For recurrent venous thrombosis, we previously investigated the potential clinical utility of multiple SNP testing for recurrent events.⁹ In that study, individual SNPs were not significantly associated with recurrent venous thrombosis. However, when the risk alleles of the individual SNPs were combined, the risk estimates as well as the significance of the association increased. The predictive ability of multiple SNP analysis has not been studied for first events of venous thrombosis. Genetic profiling may guide decisions on prophylactic measures in high-risk groups, such as cancer patients, persons undergoing surgery, persons requiring a plaster cast, or those subject to prolonged immobilization.

To explore to what extent venous-thrombosis associated SNPs can be used as predictors for a first venous thrombosis in the general population and in high-risk groups, we investigated 31 SNPs in 2 large population-based case-control studies, of which one was used as a validation set. We created genetic risk scores based on these SNPs and a risk score based on nongenetic risk factors. We also compared and combined our genetic risk score with the nongenetic risk score to determine whether genetic profiling with the currently known SNPs will improve the assessment of venous thrombosis risk.

Methods

Study populations

The Multiple Environmental and Genetic Assessment of risk factors for venous thrombosis (MEGA study) is a population-based case-control study of venous thrombosis. Collection and ascertainment of events have been described in detail previously.^{10,11} The MEGA analysis included 2712 consecutive patients with a diagnosis of a first deep vein thrombosis of the leg or arm (with or without pulmonary embolism) and 4634 control subjects (partners of patients and random population controls).

The Leiden Thrombophilia Study (LETS), another population-based case-control study of venous thrombosis, was used to validate the risk scores and included 443 consecutive patients with a diagnosis of a first deep vein thrombosis of the leg (with or without pulmonary embolism) and 453 control subjects (acquaintances or partners of patients), all without a

Submitted December 9, 2011; accepted April 27, 2012. Prepublished online as *Blood* First Edition paper, May 14, 2012; DOI 10.1182/blood-2011-12-397752.

There is an Inside *Blood* commentary on this article in this issue.

The online version of this article contains a data supplement.

The publication costs of this article were defrayed in part by page charge payment. Therefore, and solely to indicate this fact, this article is hereby marked "advertisement" in accordance with 18 USC section 1734.

© 2012 by The American Society of Hematology

Table 1. The 31 SNP associations with venous thrombosis in MEGA and in the literature^{13-17,19-26}

Gene	SNP	Chromosome	Position	MEGA				Literature average OR
				Risk allele frequency, %		OR	95% CI	
				Cases	Controls			
<i>F5</i>	rs6025	1	167.785.673	10	3	4.30	3.70-4.99	3.79
<i>F2</i>	rs1799963	11	46.717.631	6	2	3.01	2.36-3.85	2.78
<i>ABO</i>	rs8176719	9	136.132.908	47	34	1.74	1.63-1.87	1.85
<i>FGG</i>	rs2066865	4	155.744.726	34	27	1.41	1.32-1.51	1.56
<i>F11</i>	rs2036914	4	187.429.475	59	52	1.35	1.26-1.44	1.32
<i>PROCR</i>	rs2069951	20	33.227.425	7	5	1.32	1.16-1.51	1.30
<i>F11</i>	rs2289252	4	187.444.375	48	41	1.36	1.28-1.45	1.26
<i>F9</i>	rs4149755	X	138.451.778	7	6	1.11	0.99-1.24	1.24
<i>PROCR</i>	rs2069952	20	33.227.612	64	60	1.21	1.13-1.29	1.21
<i>SERPINC1</i>	rs2227589	1	172.152.839	11	9	1.27	1.15-1.41	1.20
<i>HIVEP1</i>	rs169713	6	11.920.517	22	20	1.10	1.01-1.19	1.20
<i>F2</i>	rs3136516	11	46.717.332	52	49	1.12	1.06-1.20	1.19
<i>F5</i>	rs1800595	1	167.776.972	6	5	1.18	1.03-1.36	1.18
<i>PROC</i>	rs1799809	2	127.892.345	47	43	1.17	1.10-1.25	1.17
<i>PROCR</i>	rs867186	20	33.228.215	14	12	1.18	1.07-1.29	1.17
<i>VWF</i>	rs1063856	12	6.153.534	37	33	1.18	1.10-1.26	1.16
<i>GP6</i>	rs1613662	19	60.228.407	84	82	1.18	1.09-1.29	1.15
<i>F2</i>	rs3136520	11	46.699.808	3	2	1.09	0.89-1.32	1.13
<i>F8</i>	rs1800291	X	153.811.479	85	83	1.12	1.05-1.20	1.13
<i>STXBP5</i>	rs1039084	6	147.635.413	42	45	0.90	0.84-0.96	0.90
<i>NAT8B</i>	rs2001490	2	73.781.606	40	37	1.13	1.06-1.20	1.10
<i>F13B</i>	rs6003	1	195.297.644	9	10	1.11	1.00-1.24	1.09
<i>RGS7</i>	rs670659	1	239.228.398	67	64	1.14	1.06-1.22	1.09
<i>F9</i>	rs6048	X	138.460.946	72	70	1.09	1.03-1.16	1.08
<i>F5</i>	rs4524	1	167.778.379	79	74	1.31	1.22-1.42	0.92
<i>F13A1</i>	rs5985	6	6.263.794	76	76	1.03	0.95-1.10	0.93
<i>F3</i>	1208 indel	1	94.780.000	46	46	1.02	0.96-1.09	1.06
<i>TFPI</i>	rs8176592	2	188.040.937	69	68	1.04	0.97-1.11	1.06
<i>F11</i>	rs3822057	4	187.425.146	55	49	1.31	1.23-1.39	1.06
<i>NR1I2</i>	rs1523127	3	120.983.729	41	38	1.15	1.08-1.23	1.05
<i>CPB2</i>	rs3742264	13	45.546.095	69	68	1.04	0.97-1.11	1.01

known malignancy. Collection and ascertainment of events have been described in detail previously.¹² Both studies were approved by the Medical Ethics Committee of the Leiden University Medical Center, Leiden, The Netherlands.

SNP selection

Initially, we selected 40 SNPs for the genetic risk score, based on the literature and our previous work. Eighteen SNPs had been reported and repeatedly confirmed to be associated with venous thrombosis.^{1,13} Twelve SNPs were added from the Group Health study^{13,14}; these SNPs were associated with venous thrombosis in the original study and replicated in the MEGA study. Nine SNPs were added from a large SNP association analysis, including subsequent fine mapping that we performed recently in LETS and MEGA.^{15,16} Another added SNP was recently identified in a follow-up study of a genome-wide association study and replicated in the FARIVE study and the MEGA study.¹⁷ Among the 40 SNPs in the initial selection, we studied linkage disequilibrium and mutually adjusted SNPs within genes. Four SNPs in *PROC* (rs1799808, rs1799810, rs2069915, and rs5937) were explained by rs1799809 in *PROC*; 4 SNPs in the fibrinogen genes (rs6050 and rs2070006 in *FGA*, rs1800788 in *FGB* and rs2066854 in *FGG*) were explained by rs2066865 in *FGG*; and rs3753305 in *F5* was explained by rs6025 (factor V Leiden). Consequently, we excluded 9 SNP associations that were explained by other SNPs. The remaining 31 SNPs (Table 1) were included in the genetic risk score.

Genetic risk score

We defined a genetic risk score that counts the total number of risk-increasing alleles in persons. To take into account the stronger association of some SNPs with venous thrombosis, we also constructed

a weighted risk score assigning weights to the risk alleles of each SNP corresponding to the logarithm of the average risk estimates found in the literature. In addition to the full genetic model including 31 SNPs, we constructed a parsimonious model with fewer SNPs. To determine which SNPs should be included in this model, we added SNPs one-by-one to create the genetic risk score. We started with the SNP with the highest odds ratios (ORs) in the literature and assessed whether adding SNPs to the risk score improved the area under the curve (AUC) after each SNP addition. The addition of SNPs was stopped when the AUC of the risk score, including the newly added SNP, did not differ from the AUC of the full genetic model.

Nongenetic risk factors

We constructed a nongenetic risk score, which included the following risk factors: recent (within 3 months before the index date) leg injury, surgery, pregnancy or postpartum, immobilization (ie, plaster cast, bedridden at home, hospitalization), travel for more than 4 hours in 2 months before the index date, oral contraceptives (OC) use or hormone replacement therapy (HRT) at the index date, obesity (body mass index > 30 kg/m²), and a cancer diagnosis between 5 years before and 6 months after the index date. The index date was defined as date of diagnosis for patients and their partner controls and the date of completing the questionnaire for random controls. We also included family history in the nongenetic risk score. Family history was defined as positive when a parent or sibling had experienced venous thrombosis and negative when none of these relatives had experienced venous thrombosis, or when the participant was not aware of venous thrombosis in the family. We assigned weights to each nongenetic risk factor




Figure 1. The 31-SNP risk allele distribution in patients with venous thrombosis and control subjects and corresponding ORs. The number of risk alleles was counted for cases and control subjects (top panel). ORs (95% CI) for venous thrombosis were calculated relative to the median number of risk alleles among control subjects (24 risk alleles; bottom panel). Persons with 15 or less and 36 or more risk alleles were combined for the calculation of the OR because of the low numbers of persons with that few or many risk alleles (bottom panel).

corresponding to the logarithm of the risk estimates in MEGA (supplemental Table 1, available on the *Blood* Web site; see the Supplemental Materials link at the top of the online article) and constructed a simple risk scoring system counting the weighted risk factors. We also constructed a combined risk score, including both the genetic risk score and the nongenetic risk score using a logistic regression model.

Application of genetic profiling may be most useful in high-risk groups (ie, persons exposed to known nongenetic risk factors). We therefore studied the discriminative accuracy of our genetic risk score as well as the combined score in high-risk situations of surgery, plaster cast, hospitalization, young women (younger than 50 years) using OCs, women using HRT, pregnancy or postpartum, middle-aged persons (older than 50 years) and travel. We also studied persons with a positive family history and persons with malignant disorders.

Statistical analyses

Crude and sex-adjusted (in case SNPs were located on the X chromosome) ORs and 95% CIs were calculated by logistic regression for individual SNPs and the genetic, nongenetic, and combined risk scores. When assessing the magnitude of risk associated with number of risk alleles, we used the median number of risk alleles among control subjects as the reference group.

To assess how well a score classifies venous thrombosis patients and control subjects, we calculated the area under the receiver-operating characteristic (ROC) curve (AUC). The AUC ranges from 0.5 (no discrimination between patients and control subjects) to 1.0 (perfect discrimination). We compared the AUCs of the different genetic and nongenetic risk models according to the method of Hanley and McNeil.¹⁸ Nagelkerke pseudo- r^2 statistic was used to approximate the proportion of variability explained by the different risk models. All analyses, including ROC curves and AUC calculation, were performed in SPSS Version 17.0.2 for Windows (SPSS Inc).

Results

SNPs associated with venous thrombosis

Table 1 lists all associations between SNPs and venous thrombosis in the MEGA population and the average estimated effect size in the literature.^{13-17,19-26} Not all SNPs were associated with venous thrombosis in our study populations; nevertheless, we included all 31 SNPs in the genetic risk score because these SNPs had been associated with venous thrombosis in other studies.

Genetic risk score

We first included all 31 SNPs in the genetic risk score. For each person, we counted the number of risk-increasing alleles. The number of risk alleles ranged from 13 to 38 with a median of 24 among control subjects and 26 among cases (Figure 1). The risk for venous thrombosis was estimated for each number of risk alleles, relative to the median number of risk alleles of 24, and ranged from an OR of 0.27 (95% confidence interval [CI], 0.13-0.56) for 16 risk alleles to an OR of 3.23 (95% CI, 1.96-5.30) for 33 risk alleles. At the more extreme ends of the risk distribution, CIs around risk estimates became very wide because of small numbers. The average relative risk increase per risk allele, when treated as an ordinal variable, however, could be estimated with a high level of precision, and was 1.14 (95% CI, 1.12-1.16). This corresponds to an about 100-fold difference in risk between the lowest and the highest number of risk alleles in our population.

We also constructed a weighted risk score thereby assigning weight to the risk alleles according to their risk estimates found

Table 2. Venous thrombosis prediction using genetic, nongenetic, and combined risk scores

	MEGA (N = 7092)		LETS (N = 881)	
	AUC (95% CI)	Nagelkerke pseudo r^2	AUC (95% CI)	Nagelkerke pseudo r^2
31-SNP risk score	0.71 (0.69-0.72)	0.161	0.69 (0.65-0.72)	0.149
5-SNP risk score	0.69 (0.67-0.70)	0.135	0.67 (0.64-0.71)	0.138
Nongenetic risk score	0.77 (0.76-0.78)	0.288	0.71 (0.68-0.74)	0.200
Combined risk score	0.82 (0.81-0.83)	0.378	0.77 (0.74-0.80)	0.292

The LETS study was used as a validation set.

in the literature (Table 1). A few SNPs have only been studied in the MEGA population; in that case, we used the risk estimate in MEGA as weight. The ROC curve for the weighted 31-SNP risk score had an AUC of 0.71 (Table 2; 95% CI, 0.69-0.72; ie, there is a 71% probability that a randomly chosen patient will have a higher score than a randomly chosen control subject). The weighted 31-SNP risk score was a better predictor than the nonweighted 31-SNP risk score (AUC = 0.64, 95% CI, 0.63-0.65). The average relative risk increase per unit in the risk score, when treated as an ordinal variable, was 7.89 (95% CI, 6.76-9.21). The proportion of variability explained by the 31-SNP risk score was 16.1% (Nagelkerke pseudo- r^2 ; Table 2).

To construct a genetic risk score using the most parsimonious model, we added SNPs one-by-one to the genetic risk score, starting with the SNP with the highest OR in literature (factor V Leiden, rs6025), and calculated the AUC after the addition of each SNP (Figure 2). The AUC for each single SNP ranged from 0.50 (95% CI, 0.49-0.52) for rs3136520 in *F2* to 0.60 (95% CI, 0.59-0.61) for rs8176719 in *ABO*. The discriminative accuracy of the model improved rapidly with the addition of each SNP, until 5 SNPs were included in the model (Figure 2). These SNPs were rs6025 (*F5*, factor V Leiden), rs1799963 (*F2*, 20210 G > A), rs8176719 (*ABO*), rs2066865 (*FGG*, 10034 C > T), and rs2036914 (*F11*). The AUC for this 5-SNP risk score was 0.69 (Table 2, 95% CI, 0.67-0.70). Moreover, a model based on the 3 most well-known prothrombotic polymorphisms (ie, rs6025, rs1799963, and rs8176719; AUC = 0.65, 95% CI, 0.64-0.66) performed significantly worse than the 5-SNP risk score. The average relative risk increase per unit in the risk score, when treated as an ordinal, was 9.50 (95% CI, 7.92-11.39). The 5-SNP risk score explained 13.5% of the total variability (Nagelkerke pseudo- r^2 ; Table 2).

The number of risk alleles in the 5-SNP risk score ranged from 0 (OR = 0.37, 95% CI, 0.26-0.53) to 8 (OR = 7.48, 95% CI, 4.49-12.46 for ≥ 6 risk alleles), with a median number of risk alleles of 2 among control subjects (Figure 3). The relative increase in risk per increase in number of risk alleles was 1.61 (95% CI,

1.54-1.68), again corresponding to an over 100-fold difference in risk between the lowest and the highest number of risk alleles. The weighted 5-SNP risk score was a better predictor than a non-weighted model based on number of risk alleles (AUC = 0.66, 95% CI, 0.64-0.67).


No difference between the discriminative accuracy of the 5-SNP risk score in men (AUC = 0.69, 95% CI, 0.67-0.71) and women (AUC = 0.67, 95% CI, 0.65-0.69) was found. However, differences were found when we constructed and compared the 5-SNP genetic risk score in patients with DVT in the arm, patients with DVT in the leg, and patients with DVT in the leg combined with pulmonary embolism. The AUC of the 5-SNP risk score in patients with DVT in the arm (AUC = 0.62, 95% CI, 0.57-0.67) was significantly lower than in patients with DVT in the leg (AUC = 0.68, 95% CI, 0.67-0.70) or for DVT combined with pulmonary embolism (AUC = 0.68, 95% CI, 0.67-0.70).

High-risk groups and SNP testing

To explore clinical applications of genetic profiling, we studied groups exposed to known nongenetic factors in more detail. The discriminative accuracy of the genetic risk scores in these subgroups was similar to the discriminative accuracy in the overall study population, except among cancer patients (Table 3). Subanalysis in cancer patients according to therapy (chemotherapy, surgery, radiation) or tumor class (solid vs other) did not improve the discriminative accuracy of the weighted 5-SNP risk score (data not shown).

To assess whether the genetic risk score performs better than the current clinical practice of assessing family history, we compared the discriminative accuracy of the genetic risk score with a risk score with family history alone. The AUC of the 5-SNP risk score (0.68, 95% CI, 0.67-0.70) was significantly higher than the AUC of family history (0.58, 95% CI, 0.57-0.60), with a similar trend observed among all subgroups of high-risk persons (Table 3).

Figure 2. Area under the ROC of genetic risk scores based on increasing numbers of SNPs. SNPs were added in order of the OR as found in the literature, starting with rs6025 in the score based on 1 SNP, and ending with *CPB2* included in the score of 31 SNPs (Table 1).






Figure 3. The 5-SNP risk allele distribution in patients with venous thrombosis and control subjects and corresponding ORs. The number of risk alleles was counted for cases and control subjects (top panel). ORs (95% CI) for venous thrombosis were calculated relative to the median number of risk alleles among control subjects (score 2; bottom panel). Persons with 6 or more risk alleles were combined for the calculation of the OR because of the low numbers of persons with that few or many risk alleles (bottom panel).

Combining nongenetic and genetic risk scores

We assessed the discriminative accuracy of a nongenetic risk score based on known nongenetic risk factors for venous thrombosis (leg injury, surgery, pregnancy, plaster cast, bedridden at home, hospitalization, travel, OC use, HRT, obesity, and malignancy) and family history. For the individual components, the AUC ranged from 0.50 (95% CI, 0.48-0.51) for recent travel to 0.67 (95% CI, 0.65-0.69) for OC use by women. The AUC for the nongenetic risk score including family history was 0.77 (95% CI, 0.76-0.78). When

we added the genetic risk score to the nongenetic score, the AUC significantly increased to 0.82 (Figure 4: 95% CI, 0.81-0.83) compared with the nongenetic risk score alone ($P < .0001$) using either the 31-SNP or the 5-SNP risk score. In addition, 28.8% of the total variability in venous disease risk was explained by the nongenetic risk score, which significantly improved to 37.8% (Nagelkerke pseudo r^2 ; Table 2) when combining the nongenetic and genetic risk scores. Both the nongenetic and the combined risk score models performed better in women than in men (nongenetic risk score: AUC = 0.81, 95% CI, 0.80-0.83 for women and AUC = 0.74, 95% CI, 0.72-0.75 for men; combined risk score: AUC = 0.85, 95% CI, 0.83-0.86 for women and AUC = 0.80, 95% CI, 0.78-0.81 for men).

We also studied the discriminative accuracy of the combined risk score model in the high-risk groups. For all subgroups, the AUC improved when using the combined risk score compared with the nongenetic risk score, which was significant for persons using OCs, persons with a positive family history of venous thrombosis, and persons older than 50 years old (Table 3).

Validation of the risk scores

To validate the genetic, nongenetic, and combined risk scores, we studied their discriminative accuracy in subjects from another population, the LETS population. As described in "Study populations," LETS and MEGA are both population-based case-control studies and are similar with respect to mean age at index of patients (45 years in LETS, 47 years in MEGA) or control subjects (45 years in LETS, 48 years in MEGA) and sex distribution (43% men in LETS, 47% men in MEGA). Associations between the 31 SNPs and venous thrombosis in LETS can be found in supplemental Table 2. The discriminative accuracy of the weighted 31-SNP and 5-SNP risk scores in LETS were 0.69 (95% CI, 0.65-0.72) and 0.67 (95% CI, 0.64-0.71), respectively, which are similar to those found in MEGA (Table 2).

We also constructed the nongenetic risk score weighted according to the risk estimates of each risk factor from MEGA, except for malignancies as having cancer was an exclusion criterion in LETS. In addition, information of some nongenetic risk factors (ie, HRT, recent travel, leg injury and plaster cast) was not assessed in LETS or not in such detail as in MEGA. Therefore, these risk factors were excluded from the nongenetic risk score. The discriminative accuracy of the nongenetic risk score in LETS was 0.71 (95% CI, 0.68-0.74) and improved to 0.77 (95% CI, 0.74-0.80) when combined with the genetic risk score. Both risk scores performed slightly better in MEGA than in LETS (Table 2).


Table 3. Risk score prediction in subgroups of persons exposed to known nongenetic risk factors

High-risk group	Patients, N	Control subjects, N	Family history risk score, AUC (95% CI)	5-SNP risk score, AUC (95% CI)	Nongenetic risk score, AUC (95% CI)	Combined risk score, AUC (95% CI)
Surgery	292	111	0.60 (0.55-0.66)	0.66 (0.60-0.72)	0.67 (0.61-0.72)	0.73 (0.67-0.78)
Plaster cast	111	18	0.61 (0.48-0.73)	0.73 (0.59-0.87)	0.70 (0.56-0.84)	0.78 (0.64-0.91)
Hospitalization	278	93	0.57 (0.50-0.63)	0.66 (0.59-0.72)	0.60 (0.53-0.66)	0.66 (0.59-0.72)
Oral contraceptives*	513	327	0.58 (0.54-0.62)	0.71 (0.68-0.75)	0.73 (0.69-0.76)	0.81 (0.78-0.84)
HRT	58	90	0.59 (0.49-0.68)	0.71 (0.63-0.80)	0.74 (0.66-0.82)	0.80 (0.72-0.87)
Pregnancy/postpartum*	67	46	0.54 (0.44-0.65)	0.70 (0.60-0.79)	0.68 (0.57-0.79)	0.76 (0.66-0.85)
Age > 50 y	944	1534	0.57 (0.55-0.60)	0.68 (0.66-0.70)	0.73 (0.71-0.75)	0.79 (0.77-0.81)
Travel	379	610	0.58 (0.54-0.62)	0.70 (0.67-0.73)	0.77 (0.73-0.80)	0.82 (0.80-0.85)
Family history	659	551	NA	0.68 (0.65-0.71)	0.74 (0.71-0.76)	0.81 (0.78-0.83)
Malignancies	156	65	0.57 (0.49-0.65)	0.60 (0.52-0.68)	0.71 (0.64-0.78)	0.72 (0.65-0.80)

NA indicates not applicable.

*Women younger than 50 years.

Figure 4. ROC (AUC) curves of the weighted 5-SNP risk score (light gray line), the nongenetic risk score (dotted gray line), and the combined risk score (black line). The striped black line represents the reference line (no discrimination).



Discussion

We calculated a genetic risk score based on SNPs consistently associated with venous thrombosis and observed a “dose-response” relationship between this score and the risk of venous thrombosis. The more risk alleles or genotypes present, the higher the risk of venous thrombosis. A score constructed of the 5 most strongly associated SNPs appeared to differentiate between patients and control subjects equally as well as the initial genetic risk score based on 31 SNPs. The discriminative accuracy of both the 5-SNP and 31-SNP risk score was replicated in another study (LETS) suggesting robustness of the genetic models.

When preventive measures after a positive test are invasive or can have harmful side effects, strict discrimination is required between those at high risk and low risk of developing a specific disease. In the case of venous thrombosis, indiscriminate may lead to an increased risk of thrombosis in high-risk persons receiving insufficient prophylactic anticoagulant treatment, whereas persons at low risk receiving treatment are at an increased risk of major bleeding. We investigated the extent to which genetic risk scores can improve the accuracy of thrombosis risk assessment by ROC curves. The 5-SNP genetic risk score performed better than family history assessment, which is the current clinical practice of risk assessment in persons exposed to known nongenetic risk factors. However, the 5-SNP genetic risk score performed worse than a risk score of nongenetic risk factors. A recent study by Hippisley-Cox and Coupland²⁷ showed that an algorithm of nongenetic risk factors is able to discriminate between patients and control subjects with an AUC of 0.75. This is similar to the AUC observed with our nongenetic risk score (0.77). However, the AUC may be an overestimation because we used (the logarithm of) the risk estimates from MEGA as weights.

Here, we showed that addition of the 5-SNP genetic risk score to the nongenetic risk score model significantly improved the AUC to 0.82, indicating good diagnostic accuracy. In our validation study, information on the nongenetic risk factors was less complete, which explains the lower discriminative accuracy of both the nongenetic risk score (0.71) and the combined risk score (0.77).

Identification of persons at risk of developing venous thrombosis is most useful in high-risk populations. This is because the incidence of venous thrombosis in the general population is too low

(1 per 1000 persons a year²⁸) to justify genotyping of all persons. In all subgroups of high-risk persons, the combined risk score performed better than the nongenetic score alone, which may indicate the potential clinical value of genetic profiling in these high-risk persons.

We defined a basic genetic risk score that counts the total number of risk-increasing alleles in persons. To take into account the stronger association of some SNPs with venous thrombosis, we assigned literature-based weights to each SNP, which discriminated patients better from controls than a non-weighted genetic risk score. Although the proportion of variability explained by the 5-SNP risk score is smaller than by the 31-SNP risk score, we showed that the discriminative accuracy of the 5-SNP and 31-SNP risk scores was similar. The genetic risk score is still limited, though, by its assumption that all SNPs act independently and in an additive manner in venous thrombosis susceptibility. An additive effect was assumed for the different genotypes, whereas we cannot exclude a multiplicative effect. Gene-gene interaction and gene-environment interaction are not taken into account, although in reality many interactions exist. Examples for venous thrombosis are the synergistic effects between factor V Leiden (rs6025) and OC use²⁹ and between the *F13A1* Val34Leu variant (rs5985) and fibrinogen levels.³⁰ We chose to include SNPs on their contribution to risk (effect size) and gave weights corresponding to the logarithm of this effect size. This is the most relevant for a person who has a certain genotype. One could argue that, on a population level, the prevalence of risk alleles is of relevance. However, this would not be expected to improve the performance of the risk prediction model, and indeed a genetic risk model based on the 5 SNPs with the highest risk allele frequency in MEGA performed worse than the nonweighted 5-SNP risk score, which is based on the 5 SNPs with the highest effect size (AUC = 0.54, 95% CI, 0.53-0.56; and AUC = 0.66, 95% CI, 0.64-0.67, respectively).

In the future, adding newly discovered predictive SNPs to the model may further improve discrimination. In a simulation study, Janssens et al showed that the AUC depends on the number of SNPs included, and their OR and risk allele frequency.² The heritability of a disease determines the maximum obtainable AUC. For venous thrombosis, the heritability is estimated to be approximately 60%.^{31,32} The simulation study indicated that at this level

high AUCs (>0.90) can be obtained, given that all genetic contributors are in the prediction model. Identification of new genetic predictors and validation of the genetic risk score in other study populations will reveal whether genetic profiling is useful in venous thrombosis.

In conclusion, we demonstrated that addition of a 5-SNP risk score to a risk scoring system based on nongenetic risk factors significantly improved the risk prediction of venous thrombosis. Although additional predictive markers may be required for a risk score to be clinically useful in the general population, the 5-SNP risk score may aid the management of subgroups of high-risk persons.

Acknowledgments

The Leiden Thrombophilia Study was supported by The Netherlands Heart Foundation (grant 89.063). The Multiple Environmental and Genetic Assessment of Risk Factors for Venous Thrombosis was supported by The Netherlands Heart Foundation (grant NHS 98.113), the Dutch Cancer Foundation (RUL99/1992), and The Netherlands Organization for Scientific Research (grant 912-03-0331 2003).

Authorship

Contribution: C.Y.V. had full access to all of the data in the study, takes full responsibility for the integrity of the data and the

accuracy of the data analysis, performed statistical analysis, and supervised the study; F.R.R. had full access to all of the data in the study, takes full responsibility for the integrity of the data and the accuracy of the data analysis, conceived and designed the study, acquired the data, performed statistical analysis, obtained funding, and supervised the study; I.D.B. conceived and designed the study, acquired the data, drafted the manuscript, and performed statistical analysis; J.J.D. conceived, designed, provided administrative, technical, or material support, and supervised the study; P.H.R. conceived, designed, and supervised the study; A.R.A. acquired the data and provided administrative, technical, or material support; H.G.d.H. drafted the manuscript, performed statistical analysis; S.L.C. performed statistical analysis; C.J.M.D. conceived and designed the study, acquired the data, provided administrative, technical, or material support and supervised the study; C.H.T. provided administrative, technical, or material support; L.A.B. provided administrative, technical, or material support and supervised the study; and all authors analyzed and interpreted the data and critically revised the manuscript for important intellectual content.

Conflict-of-interest disclosure: L.A.B., J.J.D., P.H.R., I.D.B., and F.R.R. hold or have applied for patents related to SNPs in this manuscript (notably rs6025, rs2066865, and rs2036914). A.R.A., C.H.T., J.J.D., and L.A.B. are employees of Celera USA. The remaining authors declare no competing financial interests.

Correspondence: Frits R. Rosendaal, Department of Clinical Epidemiology, C7-P, Leiden University Medical Center, PO Box 9600, 2300 RC Leiden, The Netherlands; e-mail: f.r.rosendaal@lumc.nl.

References

- Bezemer ID, Rosendaal FR. Predictive genetic variants for venous thrombosis: what's new? *Semin Hematol*. 2007;44(2):85-92.
- Janssens AC, Aulchenko YS, Elefante S, Borsboom GJ, Steyerberg EW, van Duijn CM. Predictive testing for complex diseases using multiple genes: fact or fiction? *Genet Med*. 2006;8(7):395-400.
- Janssens AC, Moonesinghe R, Yang Q, Steyerberg EW, van Duijn CM, Khoury MJ. The impact of genotype frequencies on the clinical validity of genomic profiling for predicting common chronic diseases. *Genet Med*. 2007;9(8):528-535.
- Brautbar A, Ballantyne CM, Lawson K, et al. Impact of adding a single allele in the 9p21 locus to traditional risk factors on reclassification of coronary heart disease risk and implications for lipid-modifying therapy in the Atherosclerosis Risk in Communities Study. *Circ Cardiovasc Genet*. 2009;2(3):279-285.
- Paynter NP, Chasman DI, Pare G, et al. Association between a literature-based genetic risk score and cardiovascular events in 19 313 women. *JAMA*. 2010;303(7):631-637.
- Davies RW, Dandona S, Stewart AFR, et al. Improved prediction of cardiovascular disease based on a panel of single nucleotide polymorphisms identified through genome-wide association studies. *Circ Cardiovasc Genet*. 2010;3(5):468-474.
- Chen H, Poon A, Yeung C, et al. A genetic risk score combining ten psoriasis risk loci improves disease prediction. *PLoS One*. 2011;6(4):e19454.
- Ripatti S, Tikkanen E, Orho-Melander M, et al. A multilocus genetic risk score for coronary heart disease: case-control and prospective cohort analyses. *Lancet*. 2010;376(9750):1393-1400.
- van Hylckama Vlieg A, Baglin CA, Bare LA, Rosendaal FR, Baglin TP. Proof of principle of potential clinical utility of multiple SNP analysis for prediction of recurrent venous thrombosis. *J Thromb Haemost*. 2008;6(5):751-754.
- Blom JW, Doggen CJM, Osanto S, Rosendaal FR. Malignancies, prothrombotic mutations, and the risk of venous thrombosis. *JAMA*. 2005;293(6):715-722.
- van Stralen KJ, Rosendaal FR, Doggen CJM. Minor injuries as a risk factor for venous thrombosis. *Arch Intern Med*. 2008;168(1):21-26.
- Koster T, Rosendaal FR, De Ronde H, Briët E, Vandenbroucke JP, Bertina RM. Venous thrombosis due to poor anticoagulant response to activated protein C: Leiden Thrombophilia Study. *Lancet*. 1993;342(8886):1503-1506.
- Smith NL, Hindorf LA, Heckbert SR, et al. Association of genetic variations with nonfatal venous thrombosis in postmenopausal women. *JAMA*. 2007;297(5):489-498.
- Smith NL, Rice KM, Bovill EG, et al. Genetic variation associated with plasma von Willebrand factor levels and the risk of incident venous thrombosis. *Blood*. 2011;117(22):6007-6011.
- Bezemer ID, Bare LA, Doggen CJM, et al. Gene variants associated with deep vein thrombosis. *JAMA*. 2008;299(11):1306-1314.
- Li Y, Bezemer I, Rowland CM, et al. Genetic variants associated with deep vein thrombosis: the F11 locus. *J Thromb Haemost*. 2009;7(11):1802-1808.
- Morange PE, Bezemer I, Saut N, et al. A follow-up study of a genome-wide association scan identifies a susceptibility locus for venous thrombosis on chromosome 6p24.1. *Am J Hum Genet*. 2010;86(4):592-595.
- Hanley JA, McNeil BJ. A method of comparing the areas under receiver operating characteristic curves derived from the same cases. *Radiology*. 1983;148(3):839-843.
- Delluc A, Gourhant L, Lacut K, et al. Association of common genetic variations and idiopathic venous thromboembolism: results from EDITH, a hospital-based case-control study. *Thromb Haemost*. 2010;103(6):1161-1169.
- Emmerich J, Rosendaal FR, Cattaneo M, et al. Combined effect of factor V Leiden and prothrombin 20210A on the risk of venous thromboembolism: pooled analysis of 8 case-control studies including 2310 cases and 3204 controls. Study Group for Pooled-Analysis in Venous Thromboembolism. *Thromb Haemost*. 2001;86(3):809-816.
- Jick H, Slone D, Westerholm B, et al. Venous thromboembolic disease and ABO blood type: a cooperative study. *Lancet*. 1969;15:1(7594):539-542.
- Grünbacher G, Weger W, Marx-Neuhold E, et al. The fibrinogen gamma (Fgg) 10034C>T polymorphism is associated with venous thrombosis. *Thromb Res*. 2007;121(1):33-36.
- Uitte de Willige S, Pyle ME, Vos HL, et al. Fibrinogen gamma gene 3'-end polymorphisms and risk of venous thromboembolism in the African-American and Caucasian population. *Thromb Haemost*. 2009;101(6):1078-1084.
- Smith NL, Wiggins KL, Reiner AP, et al. Replication of findings on the association of genetic variation in 24 hemostasis genes and risk of incident venous thrombosis. *J Thromb Haemost*. 2009;7(10):1743-1746.
- Austin H, De Staercke C, Lally C, et al. New gene variants associated with venous thrombosis: a replication study in white and black Americans. *J Thromb Haemost*. 2011;9(3):489-495.
- Wells PS, Anderson JL, Scarvelis DK, et al. Factor XIII Val34Leu variant is protective against venous thromboembolism: a HuGe review and meta-analysis. *Am J Epidemiol*. 2006;164(2):101-109.

27. Hippisley-Cox J, Coupland C. Development and validation of risk prediction algorithm (QThrombosis) to estimate future risk of venous thromboembolism: prospective cohort study. *BMJ*. 2011;343:d4656.
28. Naess IA, Christiansen SC, Romundstad P, Cannegieter SC, Rosendaal FR, Hammerström J. Incidence and mortality of venous thrombosis: a population-based study. *J Thromb Haemost*. 2007;5(4):692-699.
29. Vandenbroucke JP, Koster T, Briët E, Reitsma PH, Bertina RM, Rosendaal FR. Increased risk of venous thrombosis in oral-contraceptive users who are carriers of factor V Leiden mutation. *Lancet*. 1994;344(8935):1453-1457.
30. Vossen CY, Rosendaal FR. The protective effect of the factor XIII Val34Leu mutation on the risk of deep venous thrombosis is dependent on the fibrinogen level. *J Thromb Haemost*. 2005;3(5):1102-1103.
31. Souto JC, Almasy L, Borrell M, et al. Genetic susceptibility to thrombosis and its relationship to physiological risk factors. The GAIT study: Genetic Analysis of Idiopathic Thrombophilia. *Am J Hum Genet*. 2000;67(6):1452-1459.
32. Larsen TB, Sorensen HT, Skytthe A, Johnsen SP, Vaupel JW, Christensen K. Major genetic susceptibility for venous thromboembolism in men: a study of Danish twins. *Epidemiology*. 2003;14(3):328-332.