


To the editor:

Somatic mutations in calreticulin can be found in pedigrees with familial predisposition to myeloproliferative neoplasms

JAK2-V617F is the most frequently observed somatic mutation in patients with myeloproliferative neoplasms (MPNs).^{1,2} Recently, mutations in *CALR* were described in patients with

essential thrombocythemia (ET) and primary myelofibrosis (PMF) that were negative for *JAK2-V617F*.^{3,4} All *CALR* mutations associated with MPNs described to date are either insertions or

A Pedigrees of MPN families with mutations in *CALR*



B Summary of *CALR* and *JAK2-V617F* mutations in MPN and HT families

	family members analyzed	affected			unaffected		
		<i>V617F</i>	<i>CALR</i>	double negatives	<i>V617F</i>	<i>CALR</i>	double negatives
MPN family 1	44	6	1	0	0	0	37
MPN family 2	10	1	1	0	0	0	8
MPN family 3	3	1	1	0	0	0	1
MPN family 4	26	5	0	0	0	0	21
MPN family 5	29	4	0	0	1	0	24
MPN family 6	7	3	0	0	0	0	4
MPN family 7	3	3	0	0	0	0	0
MPN family 8	2	2	0	0	0	0	0
MPN family 9	11	2	0	0	0	0	9
MPN family 10	3	2	0	0	0	0	1
MPN family 11	2	2	0	0	0	0	0
MPN family 12	8	2	0	0	0	0	6
Total	148	33	3	0	1	0	111
HT family 1	9	0	0	3	0	0	6
HT family 2	13	0	0	3	0	0	10
HT family 3	2	0	0	2	0	0	0
HT family 4	4	0	0	3	0	0	1
HT family 5	5	0	0	2	0	0	3
HT family 6	2	0	0	2	0	0	0
HT family 7	3	0	0	2	0	0	1
HT family 8	9	0	0	3	0	0	6
HT family 9	10	0	0	5	0	0	5
HT family 10	12	0	0	4	0	0	8
HT family 11	23	0	0	5	0	0	18
Total	92	0	0	34	0	0	58

Figure 1. Somatic *CALR* mutations in familial myeloproliferative diseases. (A) The 3 pedigrees with somatic *CALR* mutations are shown in detail. Screening for *CALR* mutations was performed by using the sizing polymerase chain reaction (PCR) assay described by Klampfl et al.³ The *JAK2-V617F* mutation was screened by allele-specific PCR.⁵ Percentages in blue represent the *CALR* mutant allele burden. %T (red), percentage of G>T *JAK2-V617F* mutant allele burden; *CALR*+, presence of a 52-base deletion in exon 9 of the *CALR* gene; DN, double-negative (ie, absence of detectable mutations in *CALR* exon 9 and *JAK2-V617F*); V617F, presence of the *JAK2-V617F* mutation. (B) Summary of families analyzed. Affected and unaffected family members from families with familial predisposition to MPN (upper part) or HT (lower part) were screened for the presence or absence of mutations in *CALR*.

deletions and result in a frameshift into the alternative reading frame 1. Familial predisposition to MPNs also frequently leads to somatic *JAK2-V617F* mutations that are found in the majority of the affected family members.⁵ Because polycythemia vera (PV), ET, and PMF phenotypes can be observed in these pedigrees, we tested whether mutations in *CALR* also occur in familial MPNs.

We studied 12 pedigrees with familial MPNs in which at least 1 affected family member carried a somatic *JAK2-V617F* mutation, and we also examined 11 pedigrees with hereditary thrombocytosis (HT), in which we excluded mutations in the *THPO*, *MPL*, and *JAK2* genes (data not shown). We found a *CALR* mutation in 3 of the 12 familial MPN pedigrees (Figure 1A). In each of these 3 pedigrees, 1 affected family member carried a 52-base deletion in exon 9 of the *CALR* gene (not shown), which represents the most frequent form of the *CALR* mutations (type 1). One of these patients was initially diagnosed with ET⁷ and later progressed to PMF (MPN family 1), and the two remaining patients had ET (MPN families 2 and 3). All *CALR* mutations occurred in patients with an MPN diagnosis, whereas a somatic *JAK2-V617F* mutation was found in 1 healthy family member with platelets in the upper normal range but otherwise normal blood counts (MPN family 5; data not shown). A similar finding was reported previously.⁸ We did not detect *CALR* mutations in any of the 11 families with HT and a total of 44 affected and family members (Figure 1B).

In the 12 pedigrees with familial MPNs, all affected family members carried either a somatic *JAK2-V617F* mutation or a mutation in *CALR*. In MPN family 1, the underlying predisposition is inherited as an autosomal dominant trait with low penetrance (Figure 1A), whereas in other pedigrees, the mode of transmission is more difficult to determine because too few family members in too few generations were available (eg, MPN families 2 and 3). Two mechanistic models have been proposed regarding how such a germline predisposition may result in MPNs with clonal hematopoiesis.^{9,10} First, the germline mutation could increase the mutation rate in *JAK2* and *CALR* genes (hypermutability hypothesis), or second, it could synergize with *JAK2-V617F* or *CALR* in MPN disease initiation (fertile ground hypothesis). Because the G>T transversion in *JAK2-V617F* and the 52-base deletion in *CALR* are mechanistically very different, it seems unlikely that they could be promoted by the same germline mutation through a hypermutability mechanism. Therefore, our findings favor the fertile ground hypothesis of germline predisposition to MPN.

Pontus Lundberg

*Experimental Hematology, Department of Biomedicine,
University Hospital Basel,
Basel, Switzerland*

Ronny Nienhold

*Experimental Hematology, Department of Biomedicine,
University Hospital Basel,
Basel, Switzerland*

Achille Ambrosetti
*Section of Hematology, Department of Medicine,
University of Verona,
Verona, Italy*

Francisco Cervantes
*Hospital Clínic,
Institut d'Investigacions Biomèdiques August Pi i Sunyer,
Barcelona, Spain*

Manuel M. Pérez-Encinas
*Servicio de Hematología,
Hospital Clínico Universitario de Santiago de Compostela,
Santiago de Compostela, Spain*

Radek C. Skoda
*Experimental Hematology, Department of Biomedicine,
University Hospital Basel,
Basel, Switzerland*

Acknowledgments: This work was supported by grants 310000-120724/1 and 32003BB_135712/1 from the Swiss National Science Foundation and KLS-02398-02-2009 from the Swiss Cancer League (R.C.S.).

Contribution: P.L. designed and performed research, analyzed data, and wrote the paper; R.N. performed research and analyzed data; A.A., F.C., and M.M.P.-E. provided clinical data and patient samples and analyzed results; and R.C.S. designed research, analyzed data, and wrote the paper.

Conflict-of-interest disclosure: The authors declare no competing financial interests.

Correspondence: Radek C. Skoda, Department of Biomedicine, Experimental Hematology, University Hospital Basel, Hebelstrasse 20, 4031 Basel, Switzerland; e-mail: radek.skoda@unibas.ch.

References

- Levine RL, Gilliland DG. Myeloproliferative disorders. *Blood*. 2008;112(6):2190-2198.
- Vainchenker W, Delhommeau F, Constantinescu SN, Bernard OA. New mutations and pathogenesis of myeloproliferative neoplasms. *Blood*. 2011;118(7):1723-1735.
- Klampfl T, Gisslinger H, Harutyunyan AS, et al. Somatic mutations of calreticulin in myeloproliferative neoplasms. *N Engl J Med*. 2013;369(25):2379-2390.
- Nangalia J, Massie CE, Baxter EJ, et al. Somatic *CALR* mutations in myeloproliferative neoplasms with nonmutated *JAK2*. *N Engl J Med*. 2013;369(25):2391-2405.
- Bellanné-Chantelot C, Chaumarel I, Labopin M, et al. Genetic and clinical implications of the Val617Phe *JAK2* mutation in 72 families with myeloproliferative disorders. *Blood*. 2006;108(1):346-352.
- Kralovics R, Teo SS, Li S, et al. Acquisition of the V617F mutation of *JAK2* is a late genetic event in a subset of patients with myeloproliferative disorders. *Blood*. 2006;108(4):1377-1380.
- Pérez-Encinas M, Bello JL, Pérez-Crespo S, De Miguel R, Tome S. Familial myeloproliferative syndrome. *Am J Hematol*. 1994;46(3):225-229.
- Bellanné-Chantelot C, Jego P, Lionne-Huyghe P, Tulliez M, Najman A; French group on myeloproliferative disorders. The *JAK2(V617F)* mutation may be present several years before the occurrence of overt myeloproliferative disorders. *Leukemia*. 2008;22(2):450-451.
- Skoda RC. Hereditary myeloproliferative disorders. *Haematologica*. 2010;95(1):6-8.
- Jones AV, Cross NC. Inherited predisposition to myeloproliferative neoplasms. *Ther Adv Hematol*. 2013;4(4):237-253.

© 2014 by The American Society of Hematology

To the editor:

Physical activity limits pulmonary endothelial activation in sickle cell SAD mice

Vaso-occlusion (VOC) in sickle cell disease (SCD) results from many pathophysiological mechanisms including sickling of red blood cells, hemolysis, inflammation, vascular adhesion, and reduced nitric oxide (NO) bioavailability.¹ All these mechanisms interact

together to trap blood cells and to accentuate the local blood flow decrease and subsequent local hypoxia.

In healthy and pathological conditions, physical training results in physiological and molecular adaptations including anti-inflammatory