How does lenalidomide target the chronic lymphocytic leukemia microenvironment?

Arnon P. Kater,^{1,2} Sanne H. Tonino,^{1,2} Alexander Egle,³ and Alan G. Ramsay⁴

¹Department of Hematology and ²Lymphoma and Myeloma Centre Amsterdam (LYMMCARE), Academic Medical Center, Amsterdam, The Netherlands; ³Department of Internal Medicine III, Salzburg Cancer Research Institute, Paracelsus Medical University Salzburg, Salzburg, Austria; and ⁴Department of Haemato-Oncology, Division of Cancer Studies, Faculty of Life Sciences & Medicine, King's College London, London, United Kingdom

Immunotherapy has emerged as a viable clinical strategy to harness endogenous antitumor T-cell immunity. Lenalidomide is an oral immunomodulatory drug that repairs antitumor T-cell function and is showing efficacy in ongoing chronic lymphocytic leukemia (CLL) and lymphoma clinical trials. This article focuses on advances in our understanding of its mechanism of action in the tumor microenvironment and provides a clinical update in CLL. Challenges associated with this drug and its potential use in the targeted drug treatment era are discussed. (*Blood.* 2014; 124(14):2184-2189)

Introduction

Chronic lymphocytic leukemia (CLL) is one of the most common B-cell malignancies in adults, characterized by an accumulation of monoclonal CD5⁺ mature B cells in lymphoid tissues and the peripheral blood. Clonal expansion and invasive migration typically causes the lymph nodes, spleen, and the bone marrow to become infiltrated with tumor. Current standard therapy combines chemotherapy with an anti-CD20 monoclonal antibody (mAb) (chemoimmunotherapy [CIT]). Although highly potent, CIT induces substantial toxicity and is not curative, with nearly all patients eventually relapsing. Recent advances using kinase inhibitors (eg, ibrutinib and idelalisib) that target B-cell receptor (BCR) signaling indicate an exciting shift toward a nonchemotherapy treatment era (reviewed by Jones and Byrd¹). Present indications suggest these drugs are not producing many complete responses and should be taken continuously to avoid relapse. Unanswered questions include whether long-term persistent disease and prolonged therapy promote drug-resistant variants through clonal evolution and/or activation of compensatory oncogenic signaling in CLL. Ibrutinib resistance has already been detected in genetically high-risk patients,² highlighting the necessity to identify combinatorial therapy using agents with distinct mechanism of action (MOA).

In addition to genome alterations, CLL exhibits another dimension of complexity: leukemic cells are nurtured and protected from anticancer therapies by a variety of resident and recruited ostensibly normal cells that constitute the tumor microenvironment (TME) in lymphoid organs. Nonmalignant components of the TME include: mesenchymal stromal/stem cells (MSCs), endothelial cells, tumorassociated macrophages (TAMs or "nurse-like cells" [NLCs]), dendritic cells (DCs) and T cells. A recent breakthrough in cancer therapeutics has been the use of immunotherapies (immune checkpoint blockade) that target mechanisms of T-cell evasion by tumors.³ In this review, we focus on the immunomodulatory drug (IMiD) lenalidomide which activates antitumor T-cell activity and is showing clinical activity in ongoing CLL clinical trials. Lenalidomide (Revlimid) is a derivative of thalidomide that is US Food and Drug Administration (FDA) approved for the treatment of multiple myeloma, myelodysplastic syndromes, and mantle cell lymphoma. Remarkably, in contrast to the BCR inhibitor drugs, IMiDs exemplify successful bedside-to-bench research, in that their clinical effectiveness was known before recent MOA data emerged that help explain their pleotropic effects in the TME.

The TME in CLL

Active crosstalk between leukemic cells and nonmalignant cells in the TME plays a critical role in activating tumor migration, survival, proliferation, and fostering immune privilege (reviewed by Burger and Gribben⁴).

Briefly, CLL cells migrate into their niches via prosurvival CXC chemokine ligand 12 (CXCL12)- and CXCL13-chemokine gradients released by MSCs, NLCs, follicular helper CD4⁺ T cells (T_{FH}), and follicular DCs (FDCs). In the TME, MSCs and endothelial cells induce signal transducer and activator of transcription 3 (STAT3) and nuclear factor–κB (NF-κB)–mediated prosurvival signaling in CLL cells. Monocyte-derived NLCs supply pro-CLL survival signals and resemble proangiogenic and immuno-suppressive TAMs.⁴ Notably, CLL-stroma crosstalk is bidirectional and stromal cells in turn also become activated by leukemic cells. CLL-secreted soluble factors also activate receptors on MSCs leading to AKT activation, proliferation, and secretion of proangiogenic vascular endothelial growth factor (VEGF) and fibroblast growth factor (FGF).⁵

BCR activation in the TME has emerged as a central oncogenic pathway⁶ essential for CLL survival and proliferation. BCR engagement likely involves extrinsic autoantigens and/or microbial antigens, as well as autoreactive autonomous activation.⁶ Signals from the BCR and the tissue TME (cellular and molecular interactions) converge on several key intracellular signaling pathways including the phosphatidylinositol 3-kinase (PI3K)–AKT axis.⁴ FDCs and T_{FH} cells are specialized reservoirs of intact antigen and cognate B-cell help, respectively, and likely contribute to BCR activation.

Intriguingly, subverted CD4⁺ T cells support CLL cells in the TME, whereas antileukemic CD8⁺ T cells are suppressed. Activated

Submitted May 28, 2014; accepted August 14, 2014. Prepublished online as *Blood* First Edition paper, August 26, 2014; DOI 10.1182/blood-2014-05-578286.

The online version of this article contains a data supplement.

^{© 2014} by The American Society of Hematology

CD4⁺ T cells and T_{FH} cells can provide CD40L costimulation and prosurvival T helper 2 (Th2) cytokines (interleukin-4 [IL-4], IL-6, IL-21) that trigger extracellular signal-regulated kinase (ERK), STAT-3/-6, and NF-κB signaling and leukemic proliferation. However, CLL cells also subvert T cells to avoid immune destruction by inducing defective immune synapsis.⁷ Immune synapses (formed at T-cell–APC interfaces during antigen recognition) have a master role in T-cell activation and polarized delivery of effector molecules. CLL cells express low levels of costimulatory molecules while coopting multiple immune checkpoint coinhibitory pathways including PD-L1 that deliver an inhibitory signal into PD1⁺ T cells to actively suppress synapses and secretion of IL-2 and lytic granules.⁸ Additional immune evasion mechanisms include leukemic-derived factors that suppress NK-cell lytic activity and induction of regulatory T cells (T_{regs}).⁴

How does lenalidomide target malignant B cells?

Seminal breakthrough studies have recently identified that IMiDs (thalidomide, lenalidomide, and pomalidomide) bind to the protein target cereblon (CRBN), altering an E3 ligase complex that modulates expression of critical transcription factors. In the absence of IMiDs, CRBN-directed E3 ubiquitin ligase activity is directed to CRBN itself and possibly other specific proteins that are tagged for proteasome-mediated degradation. IMiDs bind to CRBN and modulate its substrate recognition, preventing autoubiquitylation. Three independent studies have simultaneously revealed that IMiDs retarget CRBN-dependent ligase activity toward the transcription factors Ikaros (IKZF1) and Aiolos (IKZF3) and induce their proteasomal degradation.9-11 Their decreased abundance leads to loss of viability in myeloma B cells. Ikaros and Aiolos are highly expressed in CLL cells and there is evidence supporting their role in regulating survival signaling.^{12,13} Testing whether the ability of lenalidomide to downregulate these transcription factors in lymphocytes contributes to both direct and indirect anti-CLL activity will be of great interest. Lenalidomide is not directly cytotoxic to CLL cells in vitro, but does alter CLL-TME protumoral signals. Direct anti-CLL effects include inhibition of TME-induced proliferation in a CRBN/p21-dependent manner that is associated with reduced expression of IKZF1 and IKZF3 in common with myeloma (Figure 1i).¹⁴

How does lenalidomide simultaneously modulate immune/stromal cells in the TME?

Importantly, Ikaros and Aiolos are repressors of the IL-2 promoter, and their degradation in response to IMiDs explains enhanced costimulation and IL-2 secretion in T cells.¹¹ Thus, the ability of IMiDs to target CRBN and the ubiquitin proteasome system modulates the expression of critical transcription factors that costimulate T cells, while degrading B-cell function simultaneously (Figure 1). It should be noted that the CRBN complex is likely to have distinct sets of targets in different healthy and cancer cell types. Moreover, proteins that are degraded (polyubiquitylated), compartmentalized (monoubiquitylated), or sequestered by the CRBN complex, or differentially released and stabilized by IMiDs, are likely to contribute to their biological activity.¹⁵

IMiDs increase the DC cross-priming,¹⁶ expansion, and activation of CD8⁺ T cells while decreasing activated CD4⁺ T-cell–derived

cytokines,¹⁷ promoting Th1 T-cell differentiation, and polarizing Th2 T cells to a Th1 phenotype (interferon- γ [IFN- γ], tumor necrosis factor- α [TNF- α]).¹ This Th2 to Th1 switch is linked to increased expression of the transcription factor T-bet.¹

Lenalidomide repairs the tumor-induced T-cell immune synapse defect in CLL and lymphoma by increasing the assembly and activity of cytoskeletal signaling molecules including protein kinase C- θ (PKC- θ), WASp, and RHO GTPase CDC42 to the synapse, allowing effective T-cell receptor (TCR)/CD28 signal transduction and directional secretion of lytic granules.^{7,8} IMiD-mediated repair of NK-cell lytic synapse formation is also emerging.^{1,4} Lenalidomide also rescues T-cell motility in CLL by normalizing RAC1, RHOA, and CDC42 activity levels. Importantly, knockdown of CRBN blocks lenalidomide repair of T-cell function.¹⁸ Taken together, enhancement of T-cell-mediated responses by lenalidomide is linked to a retargeted CRBN complex and altered transcription and activation of critical RHO GTPases.⁸

Lenalidomide also alters RHO GTPase activation signaling that degrades malignant B-cell function (migration capability) and TME interactions (blocks prosurvival signaling interactions with NLCs).^{19,20} Upregulation of costimulatory molecules such as CD80 and CD86 on CLL cells by lenalidomide contributes to the repair of T-cell synapses.⁷ Enhanced expression of CD40L on CLL cells by lenalidomide promotes immunoglobulin production by normal B cells²¹ that has been detected in long-term treatment responders. Thus, repair of the humoral defect may contribute to enhanced antitumor immunity. Lenalidomide downregulates the PD-L1:PD-1 immunosuppressive axis⁸ in CLL, lymphoma and myeloma, allowing T cells and NK cells to form lytic synapses with target tumor cells. Paradoxically, PD-1 positively regulates the suppressive activity of T_{regs}. Decreased PD-1⁸ and transcription factor FOXP3 expression may explain how IMiDs reduce T_{regs} while activating CD8⁺ T cells¹⁷ (Figure 1ii).

CLL cells position themselves in close contact with endothelial cells in the TME.⁴ Lenalidomide inhibits bidirectional prosurvival crosstalk between endothelial cells and tumor cells including proangiogenic signals.²² Notably, plasma levels of proangiogenic VEGF and FGF were reduced in CLL patients who responded to therapy.²² IMiD inhibition of MSC-derived CXCL12¹ may interfere with the CXC chemokine receptor 4 (CXCR4)-CXCL12 migratory axis in the CLL-TME (Figure 1iii).

Clinical experience with lenalidomide

The initial dose of lenalidomide in CLL clinical trials was chosen based on experience in myeloma but this induced serious side effects including rapid tumor cell death (tumor lysis syndrome [TLS]) and acute inflammation (tumor flare reaction [TFR], that may be associated with immune-mediated clinical response). Therefore, a low initial dose with dose escalation has been applied in subsequent trials and has improved tolerability.

Published results discussed here are summarized and referenced in Table 1 (ongoing studies: supplemental Table 1, see supplemental Data available at the *Blood* Web site).²³⁻⁴¹ When used as monotherapy, overall response rates (ORRs) have been up to 72% (mostly partial responses) for first-line therapy and 25% to 30% for fludarabine-refractory CLL. Lenalidomide has shown clinical activity in patients with high-risk features such as del(17p) (31%-38% ORR). Lenalidomide was shown to antagonize NK-cell killing of rituximabtreated CLL cells if both agents were used simultaneously.⁴² However, this has not translated to the clinic (likely due to drug scheduling) as



Figure 1. How lenalidomide targets the CLL TME. In contrast to conventional targeted drugs, lenalidomide seems to exert most of its anti-CLL activity by interfering with protumoral TME interactions rather than directly targeting prosurvival signaling in the tumor clone itself. Recent MOA data have shown that the IMiD lenalidomide binds to the protein CRBN, which modulates its substrate recognition and augments the ubiquitylation and degradation of critical transcription factors in lymphocytes. Ikaros (IKZF1) and Aiolos (IKZF3) are the downregulated targets in MM that cause direct malignant B-cell toxicity. Whether these transcription factors are targeted by lenalidomide in the event components of the CLL TME remains to be investigated. This IMiD retargeted CRBN-dependent ubiquitin ligase activity simultaneously alters malignant B-cell and nonmalignant T-cell function. (i) Emerging data have shown that a retargeted CRBN complex directly inhibits TME-mediated proliferation signaling in CLL B cells that is associated with reduced expression of IKZF1 and IKZF3 in common with myeloma. (ii) Importantly, degradation of Ikaros and Aiolos in T cells stimulates IL-2 secretion that enhances immune function. Lenalidomide has been shown to effectively reverse tumor-induced immune suppression/privilege. Immunomodulatory MOA includes: promoting a Th2 to Th1 CD4⁺ T-cell switch, downregulating immunosuppressive signaling axes (eg, PD:L1:PD1) while enhancing costimulatory molecules that positively regulate antitumor CD8⁺ T-cell function and the induction of humoral immunity (residual normal B cells). CRBN-dependent activation of RHO GTPase activation signaling and cytoskeletal signaling repairs T-cell lytic immune synapse and motility function. (iii) Lenalidomide also modulates tumor-educated stromal cells (TAMs/NLCs and MSCs) in the TME that block essential survival signals for the expanding malignant B-cell clone. MM, multiple myeloma.

results suggest improved efficacy and lower occurrence of TLS and TFR when combining rituximab or ofatumumab with lenalidomide.

Experience with lenalidomide in the consolidation/maintenance setting has been limited. Results from a phase 2 trial of CIT followed by lenalidomide showed that 24% of patients improved their quality of response with consolidation and some converted to minimal residual disease–negative status.⁴¹ These results are encouraging for an active placebo-controlled phase 3 trial of lenalidomide maintenance in minimal residual disease–positive patients post-CIT.

How to manage safety and toxicity?

The randomized ORIGIN trial that compared the safety and efficacy of front-line lenalidomide vs chlorambucil in elderly patients was halted by the FDA for safety concerns following an imbalance in the number of deaths in the lenalidomide treatment arm. This experience is in contrast to a trial reporting long-lasting efficacy with lenalidomide monotherapy²⁵ and highlights the safety and toxicity challenges associated with trial management. Clinical experience suggests an individual component to how CLL patients tolerate lenalidomide, and identifying predictive pretreatment factors requires future research. Grade 3/4 neutropenia has been reported in 70% to 83% of treated CLL patients (Table 1). The cause of lenalidomide-induced neutropenia may be related to the downregulation of transcription factor PU.1,⁴³ that can be alleviated with granulocyte colony-stimulating factor (G-CSF) treatment.

Lenalidomide-induced TFR can be effectively managed with anti-inflammatory drugs (eg, dexamethasone). Notably, the combination of lenalidomide with fludarabine and rituximab results in a dramatic reduction in TFR.

TLS has been reported in 0% to 4.5% of patients from phase 2 studies. Slow-dose escalation, close monitoring, and prophylaxis (eg, allopurinol) can effectively prevent TLS.

								1000	فاستعاد	ć		0/11		Tevioit	# /0	
					No. of prior			Lena	aniiine	Ű	sportse,	/011			y, ∕o#	
Reference	Status*	Phase	No. of patients	Age, median (range)	therapies, median (range)	del17p, %†	Scheme‡	Dose, mg/d§	Scheme	ORR	СВ	PR, incl nPR	PFS, mo	Grade 3-4 neutropenia	TLS	TFR
Upfront																
Badoux et al ^{24**}	Completed	0	60	71 (66-85)		10	len mono	5→25	Continuous	65	15	50	NR	83	0	52
Chen et al ²⁶ ††	Completed	0	25	60 (33-78)		32#‡	len mono	2.5→25	21 of 28 d	56	0	56	RN	72	4	88
Chen et al ²⁷ ††	Completed	0	25	60 (33-78)		32#	len mono	2.5→25	21 of 28 d	72	12	60	NR	76	4	88
James et al ²⁸	Completed	N	40	56 (45-64)		10	len+R	2.5→10	21 of 28 d	95	20	75	19	53	N	83
			29	70 (65-80)		7				79	10	69	20	66	N	99
Brown et al ²⁹	Terminated	-	6	59 (37-66)		None	len+FR	2.5→25	21 of 28 d	56	1	45	N	Study halted be	ecause of	
														grade 3-4 he toxicity and 1	matologic 'FS	
Flinn et al ³⁰	Open	1/2	51	62 (44-82)		12 ^a	len+FR	2.5→5	21 of 28 d	43	ŧ	41	NR	47	N	S
Egle et al ^{31b}	Completed	1/2	45	66 (43-79)		٥N°	len+FR	2.5→25	21 of 28 d	87	49	38	NR	88	0	Ŋ
Egle et al ^{32b}	Completed	1/2	40	67 (43-79)		οNο	len+FR + len+R	2.5→25	21 of 28 d	96	67	29	46	72	0	Ŋ
							maint.									
Strati et al ²⁵	Open	c٧	25	68 (44-79)		12	len+R + len maint.	10	Continuous	85	10	75	NR	36	NN	29
Relapsed or refractory																
Chanan-Khan et al ³³	Completed	N	45	64 (42-75)	3 (1-10)	13	len mono	25	21 of 28 d	47	6	38	NR	70	4	58
Ferrajoli et al ³⁴	Completed	N	4	64 (49-86)	5 (1-15)	18	len mono	10→25	Continuous	32	7	25	NR	41 ^d	0	12 ^e
Wendtner et al ³⁵	Completed	-	52	65 (37-80)	4 (1-14)	17	len mono	2.5→20	Continuous	11.5	0	11.5	9	65	4	44
Buhler et al ³⁶	Open	0	104	64 (32-81)	¥	25 ^e	len mono	5→25	Continuous	44	N	N	11	NU	N	S
Ferrajoli et al ³⁷	Open	N	36	62 (34-82)	2 (1-8)	26	len+O	10	Continuous	65	15	50	NR	44	NU	24
Costa et al ³⁸	Completed	N	17	65 (51-80)	2 (1-4)	25	len+O	10	21 of 28 d	43	NU	N	NR	79	NU	57
Sylvan et al ³⁹	Completed	-	12	71 (65-79)	4 (1-6)	42	len+A	2.5 or 5	Continuous	42	0	42	5	75	0	33
Badoux et al ⁴⁰	Open	N	59	62 (42-82)	2 (1-9)	25	len+R	10	Continuous	66	12	54	17.4	73	1.7	27
Maintenance																
Shanafelt et al ⁴¹	Open	5	34	65 (44-78)	After PCR	2.3	len maint.	5→10	Continuous	Improve respc (24%	ed quality onse in 8)	/ of patients	NR	49	N/A	N/A
*As registered in clini †UN, unknown/not rej ‡A, alemtuzumab; C,	caltrials.gov; Ju ported. cyclophosphan	ne 2014. nide; F, flu	darabine; len	, lenalidomide; m	aint, maintenance; n	nono, mono	therapy; MPS, methyl	prednisolone	; O, ofatumum	ab; P, pe	ntostatin	, R, rituxim	de		:	:

Table 1. Published trials with lenalidomide in CLL

sin most studies, dose escalation of lenalidomide was applied; doses before and after the arrow represent the minimum and maximum dose of lenalidomide per protocol. Of note, the highest dose level was not reached in all studies. ICR, complete remission; nPR, nodular partial remission; ORR, overall response rate; PR, partial remission.

INR, median PFS not reached; PFS, progression-free survival; UN, unknown/not reported.

#N/A, not applicable; TFR, tumor flare reaction; TLS, tumor lysis syndrome.

**An update on long-term outcome of these patients was published by Strati et al.25

tTwo reports presenting results of 1 trial at different time points (at median follow-up of 20.7 and 47 mo, respectively).

ttlncluding patients with del11q.

^aIncluding patients with del11q.

^bTwo reports presenting results of 1 trial at different time points. ^cAt least 1 molecular high risk feature (by CD38 expression, fluorescence in situ hybridization analysis, mutation status analysis) was present in 64% of patients. ^dToxicity in percentage of cycles (as opposed to percentage of patients). ^eA TP53 mutation was found in 36/96 (38%) of the patients in this study; of these 20 patients did not have del17p.

Identifying biomarkers of response or resistance to improve patient care

The immune synapse bioassay in combination with the lytic biomarker Granzyme-B⁸ has demonstrated utility as a knowledgebased T-cell-monitoring assay in the phase 2 trial of CIT followed by lenalidomide consolidation.⁴¹ CRBN's central role as a target of IMiD immunomodulation supports its use as a biomarker. However, its multiple splice variants and potential lack of correlation between messenger RNA and protein highlights the challenges when assessing biomarkers. Furthermore, CRBN does not appear to be a good predictive marker in CLL as it exhibits uniform expression regardless of clinical response. In contrast, expression of GSK-3 has shown potential as a response biomarker.^{44,45} Future research on the modulation of CRBN-Ikaros/Aiolos signaling pathways by IMiDs in CLL may reveal effective biomarkers. Proof-of-principle results indicate that Aiolos could act as a biomarker for T-cell activation as in vivo lenalidomide treatment resulted in downregulation of this transcription factor.11

What is the future role of lenalidomide in the evolving treatment era?

Preclinical studies suggest that the PI3Kδ inhibitor idelalisib may antagonize the immune-modulating properties of lenalidomide including repair of the humoral defect.⁴⁵ Whether this

References

- Jones JA, Byrd JC. How will B-cell-receptortargeted therapies change future CLL therapy? *Blood.* 2014;123(10):1455-1460.
- Burger JA, Landau D, Hoellenriegel J, et al. Clonal evolution in patients with chronic lymphocytic leukemia (CLL) developing resistance to BTK inhibition [abstract]. *Blood*. 2013;122(21). Abstract 866.
- Pardoll DM. Immunology beats cancer: a blueprint for successful translation. *Nat Immunol.* 2012; 13(12):1129-1132.
- Burger JA, Gribben JG. The microenvironment in chronic lymphocytic leukemia (CLL) and other B cell malignancies: insight into disease biology and new targeted therapies. *Semin Cancer Biol.* 2014;24:71-81.
- Ding W, Knox TR, Tschumper RC, et al. Plateletderived growth factor (PDGF)-PDGF receptor interaction activates bone marrow-derived mesenchymal stromal cells derived from chronic lymphocytic leukemia: implications for an angiogenic switch. *Blood.* 2010;116(16): 2984-2993.
- Burger JA, Chiorazzi N. B cell receptor signaling in chronic lymphocytic leukemia. *Trends Immunol.* 2013;34(12):592-601.
- Ramsay AG, Johnson AJ, Lee AM, et al. Chronic lymphocytic leukemia T cells show impaired immunological synapse formation that can be reversed with an immunomodulating drug. J Clin Invest. 2008;118(7):2427-2437.
- Ramsay AG, Clear AJ, Fatah R, Gribben JG. Multiple inhibitory ligands induce impaired T-cell immunologic synapse function in chronic lymphocytic leukemia that can be blocked with lenalidomide: establishing a reversible immune evasion mechanism in human cancer. *Blood.* 2012;120(7):1412-1421.

- Krönke J, Udeshi ND, Narla A, et al. Lenalidomide causes selective degradation of IKZF1 and IKZF3 in multiple myeloma cells. *Science*. 2014; 343(6168):301-305.
- Lu G, Middleton RE, Sun H, et al. The myeloma drug lenalidomide promotes the cereblondependent destruction of lkaros proteins. *Science*. 2014;343(6168):305-309.
- Gandhi AK, Kang J, Havens CG, et al. Immunomodulatory agents lenalidomide and pomalidomide co-stimulate T cells by inducing degradation of T cell repressors lkaros and Aiolos via modulation of the E3 ubiquitin ligase complex CRL4(CRBN.). Br J Haematol. 2014;164(6): 811-821.
- Billot K, Soeur J, Chereau F, et al. Deregulation of Aiolos expression in chronic lymphocytic leukemia is associated with epigenetic modifications. *Blood.* 2011;117(6):1917-1927.
- Orozco CA, Acevedo A, Cortina L, et al. The combined expression patterns of Ikaros isoforms characterize different hematological tumor subtypes. *PLoS ONE*. 2013;8(12):e82411.
- Fecteau J-F, Corral LG, Futalan D, et al. Lenalidomide inhibits the proliferation of chronic lymphocytic leukemia cells via a cereblon/ p21WAF1/Cip1-dependent mechanism [abstract]. *Blood.* 2013;122(21). Abstract 4139.
- Fischer ES, Böhm K, Lydeard JR, et al. Structure of the DDB1-CRBN E3 ubiquitin ligase in complex with thalidomide. *Nature*. 2014;512(7512):49-53.
- Henry JY, Labarthe MC, Meyer B, Dasgupta P, Dalgleish AG, Galustian C. Enhanced crosspriming of naive CD8+ T cells by dendritic cells treated by the IMiDs immunomodulatory compounds lenalidomide and pomalidomide. *Immunology.* 2013;139(3):377-385.

translates to the clinic remains to be seen. Combining immune checkpoint blockade (eg, anti-PD-1 mAb) with lenalidomide may enhance antitumor T-cell immunity in CLL and lymphoma as both agents block immunosuppressive signaling.⁸ However, such immunostimulatory trials will need careful monitoring for potential autoimmune reactions. Ibrutinib, via ITK (interleukin-2-inducible T-cell kinase)-mediated inhibition of proleukemic Th2 CD4⁺ T cells,⁴⁶ may also have complementary immunomodulatory potential with IMiDs, enhancing Th1 CD4⁺ and CD8⁺ T-cell immunity. In contrast to lenalidomide, the Bcl-2–specific inhibitor ABT-199/GDC-0199 directly induces tumor lysis. Thus, combining lenalidomide/IMiDs with ibrutinib or other new agents with distinct MOA has strong preclinical rationale and future studies will be of great interest.

Authorship

Contribution: A.G.R., A.P.K., S.H.T., and A.E. wrote the paper and approved the final version.

Conflict-of-interest disclosure: A.P.K. received research funding from Celgene. A.E. received research funding from, and provided consultancy for, Celgene. The remaining authors declare no competing financial interests.

Correspondence: Alan G. Ramsay, Department of Haemato-Oncology, Division of Cancer Studies, The Rayne Institute, King's College London, 123 Coldharbour Lane, London, SE5 9NU, United Kingdom; e-mail: alan.ramsay@kcl.ac.uk.

- Lee BN, Gao H, Cohen EN, et al. Treatment with lenalidomide modulates T-cell immunophenotype and cytokine production in patients with chronic lymphocytic leukemia. *Cancer.* 2011;117(17): 3999-4008.
- Ramsay AG, Evans R, Kiaii S, Svensson L, Hogg N, Gribben JG. Chronic lymphocytic leukemia cells induce defective LFA-1-directed T-cell motility by altering Rho GTPase signaling that is reversible with lenalidomide. *Blood.* 2013;121(14): 2704-2714.
- Troeger A, Johnson AJ, Wood J, et al. RhoH is critical for cell-microenvironment interactions in chronic lymphocytic leukemia in mice and humans. *Blood.* 2012;119(20):4708-4718.
- Schulz A, Dürr C, Zenz T, et al. Lenalidomide reduces survival of chronic lymphocytic leukemia cells in primary cocultures by altering the myeloid microenvironment. *Blood.* 2013;121(13): 2503-2511.
- Lapalombella R, Andritsos L, Liu Q, et al. Lenalidomide treatment promotes CD154 expression on CLL cells and enhances production of antibodies by normal B cells through a PI3kinase-dependent pathway. *Blood.* 2010;115(13): 2619-2629.
- Maffei R, Fiorcari S, Bulgarelli J, et al. Endothelium-mediated survival of leukemic cells and angiogenesis-related factors are affected by lenalidomide treatment in chronic lymphocytic leukemia. *Exp Hematol.* 2014;42(2):126-136.
- Gonzalez-Rodriguez AP, Payer AR, Acebes-Huerta A, et al. Lenalidomide and chronic lymphocytic leukemia. *Biomed Res Int.* 2013; 2013:932010.
- 24. Badoux XC, Keating MJ, Wen S, et al. Lenalidomide as initial therapy of elderly patients

with chronic lymphocytic leukemia. *Blood.* 2011; 118(13):3489-3498.

- Strati P, Keating MJ, Wierda WG, et al. Lenalidomide induces long-lasting responses in elderly patients with chronic lymphocytic leukemia. *Blood*. 2013;122(5):734-737.
- Chen CI, Bergsagel PL, Paul H, et al. Singleagent lenalidomide in the treatment of previously untreated chronic lymphocytic leukemia. J Clin Oncol. 2011;29(9):1175-1181.
- Chen C, Paul H, Wang T, et al. Long-term followup of a phase 2 study of single agent lenalidomide in previously untreated, symptomatic chronic lymphocytic leukemia (CLL) [abstract]. *Blood.* 2012;120(21). Abstract 718.
- James DF, Werner L, Brown JR, et al. Lenalidomide and rituximab for the initial treatment of patients with chronic lymphocytic leukemia: a multicenter clinical-translational study from the chronic lymphocytic leukemia research consortium. J Clin Oncol. 2014;32(19):2067-2073.
- Brown JR, Abramson J, Hochberg E, et al. A phase I study of lenalidomide in combination with fludarabine and rituximab in previously untreated CLL/SLL. *Leukemia*. 2010;24(11):1972-1975.
- Flinn IW, Cooper RS, Thompson DS, et al. Fludarabine, Rituximab, and Lenalidomide in Previously Untreated Patients with Chronic Lymphocytic Leukemia (CLL): A Phase I/II Trial of the Sarah Cannon Research Institute [abstract]. *Blood.* 2012;120(21). Abstract 715.
- Egle A, Steurer M, Gassner F, et al. A combination of fludarabine/rituximab with escalating doses of lenalidomide in previously untreated chronic lymphocytic leukemia (CLL): The REVLIRIT CLL5 AGMT phase I/II study, Clinical and exploratory analyses of induction results [abstract]. *Blood.* 2011;118(21). Abstract 292.
- 32. Egle A, Steurer M, Gassner F, et al. Lenalidomide/rituximab maintenance after

induction with fludarabine/rituximab in combination with escalating doses of lenalidomide in previously untreated chronic lymphocytic leukemia (CLL): The Revlirit CLL5 AGMT phase I/ II study, final results [abstract]. *Blood.* 2013; 122(21). Abstract 4146.

- Chanan-Khan A, Miller KC, Musial L, et al. Clinical efficacy of lenalidomide in patients with relapsed or refractory chronic lymphocytic leukemia: results of a phase II study. *J Clin Oncol.* 2006;24(34): 5343-5349.
- Ferrajoli A, Lee BN, Schlette EJ, et al. Lenalidomide induces complete and partial remissions in patients with relapsed and refractory chronic lymphocytic leukemia. *Blood.* 2008; 111(11):5291-5297.
- Wendtner CM, Hillmen P, Mahadevan D, et al. Final results of a multicenter phase 1 study of lenalidomide in patients with relapsed or refractory chronic lymphocytic leukemia. *Leuk Lymphoma*. 2012;53(3):417-423.
- Buhler A, Wendtner C, Hallek M, et al. TP53 mutation or deletion and efficacy with single-agent lenalidomide in relapsed or refractory chronic lymphocytic leukemia (CLL) (CC-5013-CLL-009 Study) [abstract]. *Blood.* 2013;122(21). Abstract 4146.
- Ferrajoli A, O'Brien S, Wierda WG, et al. Combination therapy with ofatumumab and lenalidomide in patients with relapsed chronic lymphocytic leukemia (CLL): results of a phase II trial [abstract]. *Blood.* 2011;118(21). Abstract 1788.
- Costa LJ, Fanning S, Stephenson J, et al. Phase 2 trial of intracycle sequential ofatumumab and lenalidomide for the treatment of relapsed and refractory chronic lymphocytic leukemia [abstract]. *Blood.* 2012;120(21). Abstract 3933.
- Sylvan SE, Rossmann E, Mozaffari F, et al. Phase I study of lenalidomide and alemtuzumab in refractory chronic lymphocytic leukaemia:

maintaining immune functions during therapyinduced immunosuppression. *Br J Haematol.* 2012;159(5):608-612.

- Badoux XC, Keating MJ, Wen S, et al. Phase II study of lenalidomide and rituximab as salvage therapy for patients with relapsed or refractory chronic lymphocytic leukemia. *J Clin Oncol.* 2013; 31(5):584-591.
- 41. Shanafelt TD, Ramsay AG, Zent CS, et al. Longterm repair of T-cell synapse activity in a phase II trial of chemoimmunotherapy followed by lenalidomide consolidation in previously untreated chronic lymphocytic leukemia (CLL). *Blood.* 2013;121(20):4137-4141.
- Lapalombella R, Yu B, Triantafillou G, et al. Lenalidomide down-regulates the CD20 antigen and antagonizes direct and antibody-dependent cellular cytotoxicity of rituximab on primary chronic lymphocytic leukemia cells. *Blood.* 2008; 112(13):5180-5189.
- Pal R, Monaghan SA, Hassett AC, et al. Immunomodulatory derivatives induce PU.1 down-regulation, myeloid maturation arrest, and neutropenia. *Blood.* 2010;115(3): 605-614.
- Trudel S, Scanga S, Mastronardi FG, et al. GSK-3 spatial regulation has an essential role in lenalidomide activity and represents a biomarker of clinical response in CLL. *Blood.* 2012;120(21). Abstract 1793.
- Herman SE, Lapalombella R, Gordon AL, et al. The role of phosphatidylinositol 3-kinase in the immunomodulatory effects of lenalidomide in chronic lymphocytic leukemia. *Blood.* 2011; 117(16):4323-4327.
- Dubovsky JA, Beckwith KA, Natarajan G, et al. Ibrutinib is an irreversible molecular inhibitor of ITK driving a Th1-selective pressure in T lymphocytes. *Blood.* 2013;122(15): 2539-2549.