Check for updates

strand of the human T-cell leukemia virus type 1

10. Macnamara A, Rowan A, Hilburn S, et al.

in HTLV-1 infection. PLoS Pathog. 2010;6(9):

DOI 10.1182/blood-2015-06-652040

HLA class I binding of HBZ determines outcome

© 2015 by The American Society of Hematology

RNA genome encodes a bZIP transcription factor that

down-regulates viral transcription. J Virol. 2002;76(24):

(rVV-HBZ) would elicit specific T-cell responses and whether this vaccine could be used to treat mice inoculated with HBZ lymphoma cells. The HBZ sequence was modified so that the protein could not activate the transforming growth factor  $\beta/Smad$ pathway. First, Sugata et al showed that anti-HBZ cytotoxic T lymphocytes can be obtained in vivo in mice (as well as in rhesus macaques), although HBZ immunogenicity is weak and requires several boosts. Importantly, anti-HBZ CD8<sup>+</sup> produces interferon gamma and tumor necrosis factor  $\alpha$ . These CD8<sup>+</sup> cells recognize and kill mouse T cells pulsed with HBZ peptides and mouse T cells transduced with an HBZ expression vector. More interestingly, CD4<sup>+</sup> T cells isolated from HBZ-transgenic animals were efficiently eliminated when transferred to HBZ-vaccinated mice. Thus, rVV-HBZ induces a cytotoxic response against cells that express HBZ in vivo.

More importantly, Sugata et al<sup>1</sup> showed that adoptive transfer of splenocytes obtained from rVV-HBZ-vaccinated mice significantly improved the survival of animals inoculated with transformed T cells that express only HBZ, as do most ATL cells (see figure). This indicates that rVV-HBZ elicits a T-cell response that is sufficient to eliminate HBZ lymphoma cells in vivo.

Finally, Sugata et al identified an HBZ peptide (amino acids 157-176) that could be used to generate a peptide-based vaccine. Altogether, the results from Sugata et al are important with respect to the generation of an anti-ATL therapeutic vaccine. Now the efficiency of the vaccine needs to be tested against primary human ATL cells.

Conflict-of-interest disclosure: The author declares no competing financial interests.

### REFERENCES

1. Sugata K, Yasunaga J-I, Mitobe Y, et al. Protective effect of cytotoxic T lymphocytes targeting HTLV-1 bZIP factor. Blood. 2015;126(9):1095-1105.

2. Afonso PV, Mekaouche M, Mortreux F, et al. Highly active antiretroviral treatment against STLV-1 infection combining reverse transcriptase and HDAC inhibitors. Blood, 2010;116(19):3802-3808

3. Bazarbachi A, Suarez F, Fields P, Hermine O. How I treat adult T-cell leukemia/lymphoma. Blood. 2011; 118(7):1736-1745.

4. Yamano Y, Sato T. Clinical pathophysiology of human T-lymphotropic virus-type 1-associated myelopathy/tropical spastic paraparesis. Front Microbiol. 2012;3:389.

5. Wattel E, Vartanian JP, Pannetier C, Wain-Hobson S. Clonal expansion of human T-cell leukemia virus type I-infected cells in asymptomatic and symptomatic

carriers without malignancy. 7 Virol. 1995;69(5): 2863-2868

6. de Thé G, Bomford R. An HTLV-I vaccine: why, how, for whom? AIDS Res Hum Retroviruses. 1993;9(5):381-386.

7. Bomford R, Kazanji M, De Thé G. Vaccine against human T cell leukemia-lymphoma virus type I: progress and prospects, AIDS Res Hum Retroviruses, 1996;12(5);403-405.

Rodríguez SM, Florins A, Gillet N, et al. Preventive and therapeutic strategies for bovine leukemia virus: lessons for HTLV. Viruses. 2011;3(7):1210-1248.

9. Gaudray G, Gachon F, Basbous J, Biard-Piechaczyk M, Devaux C, Mesnard JM. The complementary

### • • LYMPHOID NEOPLASIA

Comment on Paggetti et al, page 1106

12813-12822

e1001117.

## Exosomes and CAFs: partners in crime

Benedetta Apollonio and Alan G. Ramsay KING'S COLLEGE LONDON

In this issue of *Blood*, Paggetti et al present novel findings that chronic lymphocytic leukemia (CLL)-derived exosomes and their molecular cargo are actively transferred to stromal cells that reside in the lymphoid tumor microenvironment (TME), promoting the reprogramming of these cells into cancer-associated fibroblasts (CAFs).<sup>1</sup>

LL is characterized by an accumulation of monoclonal CD5<sup>+</sup> mature B cells in lymphoid tissues and the peripheral blood. Clonal expansion and invasive migration typically cause the lymph nodes and bone marrow to become infiltrated with tumor. Recent progress in understanding the genetic landscape of lymphoid malignancies must also be coupled with research on the TME. Accumulating evidence is showing that bidirectional communication between cancer cells and their microenvironment is critical for tumor growth, but also profoundly affects therapeutic response. Malignant cells engage in novel associations/interdependencies with stromal cells including fibroblasts (mesenchymal stem cells [MSCs]), endothelial cells, and immune cells that provide crucial contributions to the licensing of tumor progression (survival, proliferation) and immune evasion.<sup>2</sup> Many of the examples of heterotypic signaling studied to date involve classical paracrine signaling loops of cytokines or growth factors and their receptors. Although these signaling mechanisms are key mediators of cell-cell communication within the TME, more recently, exosome shedding has emerged as another mode of intercellular signaling. Exosomes are nanometer-sized endocytic vesicles manufactured within

multivesicular endosomes and released into the extracellular compartment by many types of cells. Their biogenesis/release is enhanced when cells are stimulated-under stress or in a diseased environment. Transfer of exosomes and their cargo, which includes proteins, messenger RNA, and microRNAs (miRNAs), from cancer cells to other TME cell types has been the subject of intense studies in solid cancers,<sup>3</sup> but understudied in lymphoid neoplasia. In this work, Paggetti et al<sup>1</sup> provide comprehensive molecular analysis of exosomes derived from CLL cells and, in particular, functional data supporting a novel capacity of these extracellular vesicles to modulate the TME by reprogramming previously healthy stromal cells into CAFs.

The study uses a robust protocol to isolate exosomes from the supernatant of human CLL cells (MEC1 and primary samples). Elegant studies show that tumor-derived exosomes are rapidly taken up by stromal cells, including MSCs, endothelial cells, and myeloid cells (but not CLL cells). Interestingly, this later finding may contrast with microvesicles, a different type of vesicle arising from the outward budding of the plasma membrane, with data suggesting integration into CLL cells.<sup>4</sup> Paggetti et al show that active exosome intercellular



CLL-derived exosomes have a paracrine effect on stromal cells residing in the TME. The transfer of exosomal cargoes (miRNA and proteins) to target cells (bone marrow, BM-MSCs, and endothelial cells) induces an inflammatory CAF phenotype in these cells that assume an aberrant stimulatory, protumoral role (increased angiogenesis, release of pro-survival chemokines/cytokines). In parallel, CLL-derived exosomes can act as decoy receptors for rituximab, thus representing a potential drug escape mechanism in the TME. BM, bone marrow; NK-kB, nuclear factor kB.

transfer requires specific functional surface proteins on exosomes (tetraspanins, integrins) and the presence of heparan sulfate proteoglycans on the surface of target cells (that CLL cells lack). The authors perform small RNA sequencing and mass spectrometry analysis of CLL exosomes to reveal predominantly regulatory RNAs, miscellaneous RNAs, and mature miRNAs enriched in CLL exosomes. Notably, some of these miRNAs (miR-150 and miR-155) were previously identified in circulating miRNA<sup>5</sup> and plasma-derived exosomes from CLL patients.<sup>6</sup> In addition, abundant miRNAs (miR-21, miR-146a) have been shown to regulate the induction of aberrant stromal cells. Exosome miRNAs were shown to transfer to MSCs and retain their functional activity. The exosome proteome was enriched in cell membrane, cytoskeletal, antiapoptotic, and angiogenic proteins.

Similar to miRNAs, exosome proteins were also transferred to recipient murine and human stromal cells as exemplified by B-cell marker CD20.

The authors then study the changes induced in the target stromal cells. CLL exosomes induced an inflammatory phenotype in both MSC and endothelial cells linked to activated nuclear factor кB-dependent signaling. Gene expression profiling of exosome-treated stromal cells revealed transcriptional changes enriched for an "inflammatory response" with significant overlap to the hallmark proinflammatory gene signature of activated CAFs.7 Proinflammatory CAFs show increased expression of molecular markers, including  $\alpha$ -smooth muscle actin (α-SMA).<sup>8</sup> Paggetti et al verified the presence of high numbers of  $\alpha$ -SMA-positive stromal cells in the CLL-TME.9 Gene expression changes alter

the CAF secretome, and the authors' study reveals increased levels of tumor necrosis factor family ligands (B-cell activating factor, A proliferation-inducing ligand), proinflammatory cytokines (eg, interleukin-6), chemokines (eg, CCL2, CXCL16), and proangiogenic factors (eg, hepatocyte growth factor) in subverted stromal cells. Distinct from normal tumor-suppressive stromal cells, CAFs promote tumor progression, angiogenesis, and dissemination through paracrine cross-talk and remodeling of the extracellular matrix. Functional data by Paggetti et al show increased collagenase activity, proliferation, and migration of exosome-educated stromal cells that exhibited striking cytoskeletal remodeling, stress fiber formation, and α-SMA expression (absent from healthy B cell-derived exosome-treated stromal cells). In vivo matrigel plug assays showed that CLL exosomes induce the

formation of new blood vessels. Exosomeeducated stromal cells showed an enhanced capacity to support CLL cell survival. Finally, promotion of tumor growth/dissemination was demonstrated using a cotransfer in vivo model (CLL cells with exosomes). These initial data demonstrate a protumorigenic effect of CLL-derived exosomes when tumor cells engage the microenvironment to proliferate and promote angiogenesis. Taken together, the authors conclude that CLL exosomes actively promote disease progression by subverting the function of stromal cells that reside in the TME which acquire features of proinflammatory CAFs (see figure). Clearly, future studies will need to examine the coevolution of tumor cells and the reprogrammed stroma including characterizing the role CAFs play in regulating cancer hallmark capabilities.

This work by Paggetti et al establishes the tumorigenic importance of CLL-derived exosomes, including the transfer of their molecular cargo to other TME cells. This contributes to increasing evidence that the ability of tumor cells to induce CAFs is a universal feature of progression in both solid<sup>8</sup> and blood<sup>7,10</sup> cancers. These data also provide incentive for detailed mechanistic investigation of the role exosomes play in pathogenic signaling processes<sup>6</sup> and regulating the understudied CLL-TME. Examining the contribution of exosomes and their content in predicting response (biomarkers) to different therapeutic agents including newly targeted kinase inhibitors will be of interest. Paggetti et al show that CLL-derived exosomes act as decoys for the therapeutic antibody rituximab (anti-CD20) that highlights a potential drug escape mechanism in the TME. Continued research on TME regulation of tumor progression will provide insight into disease pathogenesis and also the exciting prospect of identifying novel targeted therapies designed to re-educate the TME to have antitumorigenic effects.

Conflict-of-interest disclosure: The authors declare no competing financial interests.

### REFERENCES

1. Paggetti J, Haderk F, Seiffert M, et al. Exosomes released by chronic lymphocytic leukemia cells induce the transition of stromal cells into cancer-associated fibroblasts. *Blood.* 2015;126(9):1106-1117.

2. Scott DW, Gascoyne RD. The tumour microenvironment in B cell lymphomas. *Nat Rev Cancer*. 2014;14(8): 517-534.  Webber J, Yeung V, Clayton A. Extracellular vesicles as modulators of the cancer microenvironment. *Semin Cell Dev Biol.* 2015;40:27-34.

 Ghosh AK, Secreto C, Boysen J, et al. The novel receptor tyrosine kinase Axl is constitutively active in B-cell chronic lymphocytic leukemia and acts as a docking site of nonreceptor kinases: implications for therapy. *Blood*. 2011;117(6):1928-1937.

 Moussay E, Wang K, Cho JH, et al. MicroRNA as biomarkers and regulators in B-cell chronic lymphocytic leukemia. *Proc Natl Acad Sci USA*. 2011;108(16):6573-6578.

 Yeh YY, Ozer HG, Lehman AM, et al. Characterization of CLL exosomes reveals a distinct microRNA signature and enhanced secretion by activation of BCR signaling. *Blood.* 2015;125(21):3297-3305.

7. Lutzny G, Kocher T, Schmidt-Supprian M, et al. Protein kinase c- $\beta$ -dependent activation of NF- $\kappa$ B in stromal cells is indispensable for the survival of chronic

### • • PLATELETS AND THROMBOPOIESIS

Comment on Valet et al, page 1128

# Membrane grease eases platelet maturation

Sang H. Min and Charles S. Abrams UNIVERSITY OF PENNSYLVANIA

In this issue of *Blood*, Valet et al<sup>1</sup> report a novel regulatory role of class II phosphoinositide 3-kinase (PI3K)-C2 $\alpha$  in the morphology and remodeling of platelet membranes and its implications in platelet maturation and arterial thrombosis.

n 1906, James Homer Wright first postulated that platelets are produced in the bone marrow from megakaryocytes.<sup>2</sup> Our megakaryocytes collectively generate and release  $\sim 1$  million platelets per second (10 billion platelets per day) into the bloodstream through a sequence of remodeling events. Megakaryocytes first undergo a series of maturation processes whereby they increase their overall size as they develop highly tortuous membrane invaginations that are called the invaginated membrane system (previously known as the demarcation membrane system).<sup>3</sup> Megakaryocytes use this extensive membrane reservoir to form long and thin cytoplasmic extensions to produce intermediate platelet structures called proplatelets.<sup>4</sup> Once released into the bloodstream, proplatelets undergo further remodeling steps that allow membrane fragmentation and generation of mature platelets. Proplatelet formation and maturation are heavily dependent on the membrane and cytoplasmic skeletal machinery, which include microtubule, actin, and spectrin.5,6 Although

microtubules and actin function as engines to power platelet formation and elongation, spectrin is necessary for the ultimate shape changes that occur when proplatelets mature

lymphocytic leukemia B cells in vivo. Cancer Cell. 2013;

heterogeneity in the cancer wound. J Exp Med. 2014;

9. Ruan J, Hyjek E, Kermani P, et al. Magnitude of

stromal hemangiogenesis correlates with histologic subtype

of non-Hodgkin's lymphoma. Clin Cancer Res. 2006;12(19):

10. Frassanito MA, Rao L, Moschetta M, et al. Bone

© 2015 by The American Society of Hematology

DOI 10.1182/blood-2015-07-655233

marrow fibroblasts parallel multiple myeloma progression

in patients and mice: in vitro and in vivo studies. Leukemia

8. Öhlund D. Elvada E. Tuveson D. Fibroblast

23(1):77-92

5622-5631

211(8):1503-1523.

2014;28(4):904-916.

Megakaryocyte Proplatelet Platelet



Membrane remodeling by class II PI3K-C2 $\alpha$  is essential for platelet maturation. Megakaryocytes initiate platelet production by extending and releasing intermediate platelet structures called proplatelets into the bloodstream. Released proplatelets undergo a series of remodeling steps to generate mature platelets. Valet et al show that PI3K-C2 $\alpha$  generates a distinct pool of PI3P in resting platelets that may modulate membrane elasticity and remodeling by reorganizing membrane skeletal proteins such as spectrin.

BLOOD, 27 AUGUST 2015 · VOLUME 126, NUMBER 9