PLATELETS AND THROMBOPOIESIS

Sex-specific differences in genetic and nongenetic determinants of mean platelet volume: results from the Gutenberg Health Study

Marina Panova-Noeva,¹ Andreas Schulz,² M. Iris Hermanns,¹ Vera Grossmann,² Evgenia Pefani,³ Henri M. H. Spronk,⁴ Dagmar Laubert-Reh,² Harald Binder,⁵ Manfred Beutel,⁶ Norbert Pfeiffer,⁷ Stefan Blankenberg,^{8,9} Tanja Zeller,^{8,9} Thomas Münzel,^{3,10} Karl J. Lackner,¹¹ Hugo ten Cate,⁴ and Philipp S. Wild^{1,2,10}

¹Center of Thrombosis and Hemostasis, ²Preventive Cardiology and Medical Prevention, Department of Medicine II, and ³Department of Medicine II, University Medical Center Mainz, Mainz, Germany; ⁴Laboratory for Clinical Thrombosis and Hemostasis, Departments of Internal Medicine and Biochemistry, Cardiovascular Research Institute Maastricht, Maastricht University Medical Center, Maastricht, The Netherlands; ⁵Institute of Medical Biostatistics, Epidemiology and Informatics, ⁶Department of Psychosomatic Medicine and Psychotherapy, and ⁷Department of Ophthalmology, University Medical Center Mainz, Mainz, Germany; ⁸Department of General and Interventional Cardiology, University Heart Centre, Hamburg-Eppendorf, Germany; ⁹German Center for Cardiovascular Research (DZHK), Partner Site Hamburg/Kiel/Lübeck, Hamburg, Germany; ¹⁰DZHK, Partner Site Rhein-Main, Mainz, Germany; and ¹¹Institute for Clinical Chemistry and Laboratory Medicine, University Medical Center Mainz, Mainz, Germany

Key Points

- Genetic and nongenetic determiners of MPV substantially differ between males and females in a large population-based study.
- MPV in males is significantly determined by the traditional CVRFs, and males with higher MPV are at higher risk of death.

Mean platelet volume (MPV), a measure of platelet size, is a potential biological marker of platelet function. To date, a comprehensive analysis including known genetic and nongenetic factors that determine MPV is still lacking. MPV has been evaluated in 15 010 individuals from the population-based Gutenberg Health Study. Genetic information was available for 4175 individuals. Our results showed that age (β , 0.0346; 95% confidence interval [CI], 0.0255 to 0.0436), cardiovascular risk factors (CVRFs) such as smoking (β , 0.178; 95% CI, 0.128 to 0.229), hypertension (β , 0.05; 95% CI, 0.00289 to .0981), and high glucose level (β , 0.00179; 95% CI, 0.0006 to 0.00299) were linked with higher MPV in males only. Intake of oral contraceptives (β , 0.150; 95% CI, 0.0649 to 0.236) and menstruation (β , 0.123; 95% CI, 0.0231 to 0.224) were strongly associated with higher MPV in females. Seven single nucleotide polymorphisms (SNPs) for females and 4 SNPs for males were associated with higher MPV. The full model, including age, CVRFs, laboratory parameters, medications, and genetic variation, explained 20.4% of the MPV variance in

females and 18.6% in males. The curves of cumulative mortality, stratified for sex, showed worse survival for males only with MPV >9.96 fL vs MPV \leq 9.96 fL (*P* < .0001). This study provides evidence for heterogeneity in the profile of determinants for MPV between sexes. The observed interactions between genetic variability, CVRFs, and MPV and its association with the development of cardiovascular disease or thrombotic risk need to be further investigated. (*Blood.* 2016;127(2):251-259)

Introduction

Platelets play a key role in both hemostasis and thrombosis and, according to recent findings, in inflammation, atherogenesis, and cancer metastasis.¹ Mean platelet volume (MPV), a measure of platelet size, is a potential biological marker of platelet function and is commonly available in the clinical practice at relatively low cost. Several cytokines and growth factors may affect megakaryocytopoiesis and release of larger platelets. In addition, changes in platelet volume may also occur at the sites of activation with changes in platelet shape.² Larger platelets are metabolically and enzymatically more active and have higher thrombotic potential and higher expression of platelet surface activation markers.³⁻⁵ Higher MPV was shown to be associated with traditional cardiovascular risk factors⁶ such as hypertension,⁷ diabetes,⁸ obesity,⁹ hypercholesterolemia,¹⁰ and smoking¹¹ and is correlated with an increased risk for both arterial^{12,13} and venous thrombosis.¹⁴ Patients with preexisting

There is an Inside Blood Commentary on this article in this issue.

coronary artery disease and increased MPV are at higher risk of myocardial infarction (MI).¹⁵ Furthermore, higher MPV implied an increased risk of vascular mortality, particularly of ischemic heart disease, in a large population study.¹⁶

It has been recognized for almost 40 years that sex may influence platelet biology.^{17,18} Therefore, the determinants of MPV may vary between males and females. Recent studies investigating the relation between MPV and vascular risk factors report higher MPV in females except for a Danish population based-study demonstrating higher MPV in males.^{12,14,16,19}

Both genetic and nongenetic factors may influence the variation of MPV. White blood cell (WBC) counts, sex, and age have recently been identified as major determinants of MPV heterogeneity.¹⁹ A genome-wide association study (GWAS) identified eight chromosomal regions with 22 independent single nucleotide polymorphisms (SNPs) that influence

The publication costs of this article were defrayed in part by page charge payment. Therefore, and solely to indicate this fact, this article is hereby marked "advertisement" in accordance with 18 USC section 1734.

© 2016 by The American Society of Hematology

Submitted July 24, 2015; accepted October 19, 2015. Prepublished online as *Blood* First Edition paper, October 30, 2015; DOI 10.1182/blood-2015-07-660308.

The online version of this article contains a data supplement.

MPV across cohorts.²⁰ One of these SNPs, rs342293 (*FLJ36031*; *PIC3CG*) on chromosome 7q22.3 was recently associated with larger MPV and with higher risk for adverse cardiovascular outcome in patients with coronary artery disease who were undergoing percutaneous coronary intervention.²¹

To date, a comprehensive, combined analysis that includes known genetic and nongenetic factors that determine MPV is still lacking. In addition, sex-specific information regarding the relation of MPV and cardiovascular risk factors is limited. The Gutenberg Health Study (GHS) offers the potential to investigate sex-specific disparities in MPV determinants. Because of the observational and representative character of this study, the continuum from healthy individuals to those with cardiovascular disease such as coronary heart disease, peripheral artery disease, history of MI, and heart failure is present in the study sample. The setting offers the advantage of exploring both genetic and nongenetic influences on MPV variation in a comprehensively phenotyped population sample with a standardized setting for MPV measurement.²²

Research design and methods

Population sample

MPV was determined in 15 010 participants enrolled in the GHS between April 2007 and April 2012 at their baseline examination. The cohort was part of a population-based, prospective, observational, single-center study in western mid-Germany conceptualized primarily for studying cardiovascular disease. Residents of Mainz and Mainz-Bingen County age 35 to 74 years were randomly selected from the registries of the local governmental registry offices with stratification for sex, residence (urban vs rural), and decade of age. All participants underwent a 5-hour examination at the study center and followed standard operating procedures provided by certified medical technical assistants. Details of the study design were published earlier.²³ The project was approved by the local ethics committee, and the study was conducted in accordance with the Declaration of Helsinki. Written informed consent was obtained from all participants before they entered the study.

Blood sampling and laboratory measurements

Venous blood sampling was performed by using tubes containing K3-EDTA. Platelet parameters (count and MPV) were automatically determined on an ADVIA 120 Hematology System (Siemens, Erlangen, Germany) between 30 and 90 minutes after blood sampling.

Internal quality control of automated blood counts was performed by using the TESTpoint controls for the ADVIA 120 Hematology System (Siemens). Results had to be within the limits indicated for the respective control material. Coefficients of variation for MPV ranged between 0.8% and 2.2% over time with a mean of 1.4%. The moving average analysis on MPV performed over the duration of the study is presented in supplemental Figure 1 (available on the Blood Web site). Results have been obtained with three ADVIA 120 analyzers (Siemens) at the Institute for Clinical Chemistry and Laboratory Medicine, University Medical Center Mainz, in Mainz, Germany. To continuously validate and align the measuring systems, 10 random blood samples were analyzed on all 3 instruments on a monthly basis, and the results were evaluated. The inter-instrument coefficient of variation of MPV varied between 1.8% and 4.1% during the study period (supplemental Figure 2). Inter-instrument coefficients of variation <5% were considered acceptable.

Biochemical analysis of blood glucose and C-reactive protein (CRP) measured in plasma and cholesterol, high-density lipoprotein, low-density lipoprotein, cholesterol, and triglycerides measured in serum were determined on the day of sampling by routine methods. All laboratory measurements were performed in the central laboratory of the University Medical Center in Mainz.

Selection, genotyping, and imputation of SNPs

MPV-related SNPs were selected from the catalog of published GWASs with the term "mean platelet volume"²⁴ and by a systematic search of PubMed using the keywords "MPV", "mean platelet volume", "SNPs", and "genetic variants". Overall, 36 SNPs were selected (supplemental Table 1) from the available data, which were described by large GWASs for MPV.^{20,25,26}

SNP information was taken from the genetic data available from the GHS. Genotyping was conducted on Affymetrix Genome-Wide Human SNP 6.0 arrays (Affymetrix, Santa Clara, CA) according to the manufacturer's recommendations. After a quality control review, genetic data were available for analysis in 4175 individuals. Before genotype imputation, SNPs showing significant ($P < 10^{-4}$) deviation from the Hardy-Weinberg equilibrium with minor allele frequency <1% or having a genotyping call rate <98% were excluded. A two-step process was performed for imputation. First, prephasing was performed by using MACH v1.0.18c software. Second, genotypes were imputed by using the reference panel 1000G Phase I Integrated Release Version 2 Haplotypes (2010-11 data freeze, 2012-02-14 haplotypes), and minimac (release 2012-03-14) was used. The imputation results were filtered by using a minimum imputation quality score $r^2 > 0.3$ and a minor allele frequency threshold of 2%. Imputation results were summarized as allele dosage, defined as the expected number of copies of the minor allele for each SNP (values of 0, 1, and 2), and genotype. A ratio of observed to expected variance of the dosage statistic for each SNP was calculated for imputation quality.

Data management and statistical analysis

Safe and efficient data management was ensured by acquiring, storing, and analyzing all data electronically. All data underwent quality control with a review for completeness and plausibility by using predefined algorithms and criteria from a central data management unit.

Distribution of MPV is presented as the median with interquartile range and was tested by using the Mann-Whitney U test. Normally distributed continuous variables were described by using mean \pm standard deviation and tested with the *t* test, whereas categorical variables were expressed as absolute numbers and percentages and tested with the χ^2 test if needed. The population sample was categorized according to the 95th percentile of MPV in the reference group. The reference group was a subsample of the population sample and was defined as individuals without known clinically established disease (ie, MI, coronary artery disease, stroke, peripheral arterial disease, venous thromboembolism, atrial fibrillation, chronic heart failure, cancer, kidney disease, liver disease, chronic obstructive pulmonary disease, and anemia) or traditional cardiovascular risk factors (ie, hypertension, obesity, dyslipidemia, diabetes mellitus, smoking, and family history of MI or stroke). For the definition of traditional cardiovascular risk factors and categorization of medications, please see supplemental Data. Relations between continuous variables were explored by using Pearson's correlation coefficients. Quantile regression was used to describe median MPV levels. Multivariable

Table 1. Characteristics of the GHS population sample by stratification for values above and below the reference limits for MPV (N = 14968)

			Females					Males		
	≤95th (n	percentile = 7012)	>95th (n	percentile = 385)		≤95th (n	percentile = 7060)	>95th (n	percentile = 511)	
Characteristic	Median	IQR	Median	IQR	Р	Median	IQR	Median	IQR	Р
Age (y)	55.0	45.0-64.0	56.0	46.0-65.0	.068	55.0	46.0-65.0	58.0	47.0-66.0	.0023
WBC (× 10 ⁹ /L)	6.94	5.90-8.30	7.37	6.04-8.61	.00086	6.84	5.74-8.20	7.10	6.07-8.37	.0025
Platelets (× 10 ⁹ /L)	289.0	249.0-332.0	230.0	197.0-265.3	<.0001	255.0	219.0-296.0	207.5	177.4-241.0	<.0001
CRP (mg/L)	1.60	0.57-3.50	1.80	0.85-3.60	.18	1.50	0.51-2.90	1.70	0.81-3.30	.0018
Fibrinogen (mg/dL)	325.0	283.0-380.0	332.0	293.0-387.0	.071	315.0	272.0-369.0	327.0	285.0-388.0	.00045
Glucose (mg/dL)	90.0	84.0-96.0	89.2	83.7-98.0	.86	93.0	87.0-100.2	94.0	87.0-103.0	.047
LDL (mg/dL)	137.0	114.0-162.0	131.9	108.9-157.0	.013	137.0	115.0-160.0	132.0	109.0-161.9	.0094
HDL (mg/dL)	63.0	53.6-74.0	61.0	51.0-72.0	.012	49.0	42.0-57.0	48.0	40.0-55.0	.0065
Cholesterol (mg/dL)	223	197-252	216	188-247	.0032	215.0	189.0-241.0	210.0	182.0-241.0	.043
Triglycerides (mg/dL)	96.0	72.0-129.0	102.0	75.2-134.0	.068	117.0	85.0-164.0	122.0	89.0-171.0	.0091

Categories are based on the 95th percentile of MVP in the reference group.

HDL, high-density lipoprotein; IQR, interquartile range; LDL, low-density lipoprotein.

linear regression analysis was used to identify determinants of MPV. Because of the explorative character of the analysis, a significance threshold was not defined for *P* values. *P* values should be interpreted as a continuous measure of statistical evidence. The analysis of prospective mortality data with censoring is presented by using plots of cumulative mortality with log-rank test and by Cox proportional hazards models.

Results

Participant characteristics

Table 1 shows continuous biomarkers of cardiovascular risk stratified for the 95th percentile of the reference group, which was determined as 9.9 fL for males, 10.1 fL for females, and 10.1 fL in the overall sample. WBC count, CRP, fibrinogen, and glucose and triglyceride levels were significantly higher, whereas platelet count, low-density lipoprotein, high-density lipoprotein, and total cholesterol were significantly lower in the group of participants with MPV above the reference limit.

Traditional cardiovascular risk factors (CVRFs) (ie, diabetes and smoking) were associated with higher MPV both in males and females (supplemental Figure 3). In addition, obesity was associated with higher MPV in males only. Figure 1 shows the associations between medication and MPV in separate single linear regression models for each drug adjusted for age. Antithrombotic agents, including aspirin and diuretics, were associated with higher MPV in both males and females. Furthermore, antilipemic agents, antihypertensives, angiotensin-converting enzyme inhibitors, and beta blockers were associated with higher MPV in males. Interestingly, oral contraceptives, but not hormone replacement therapy, showed a significant association with higher MPV in females. Sex-specific multivariable linear regression models adjusted for age, classical CVRFs, medication, and laboratory parameters (Table 2) described 11.9% of MPV variation in both males and females. WBC count and fibrinogen were positively associated with MPV, whereas platelet count and CRP were negatively associated. In male participants, hypertension, smoking, and glucose levels were positively associated with MPV (Table 2), whereas in females, a positive association with MPV was observed for menstruation (B, 0.123; 95% CI, 0.0231 to 0.224; P = .016) and oral contraceptives (β , 0.150; 95% CI, 0.0065 to 0.236; P = .00056).

Age- and sex-specific distribution of MPV

Of the 15 010 participants enrolled at baseline in the GHS cohort, MPV was available for 99.7% (n = 14968). The age range of the study sample at the time of MPV measurement was 35 to 74 years with a mean of 55 years, and 49.4% were female. Linear regression analysis presented a higher MPV in females compared with males with an estimated β of 0.357 (95% CI, 0.212 to 0.502; P < .0001; Figure 2); however, there was a significant interaction effect between sex and age. The graphic in Figure 2 displays an increase in MPV in males only, with higher MPV in females until the age of approximately 65 years with an inversion of the relationship at older ages. Females in the reference group had an MPV comparable to that of females in the population sample (8.5 fL [95% CI, 8.0 to 9.1 fL] vs 8.5 fL [95% CI, 8.0 to 9.0 fL]; P = .2). Males from the reference group showed significantly lower MPV compared with males in the population sample (8.3 fL [95% CI, 7.8 to 8.8 fL] vs 8.4 fL [95% CI, 7.9 to 9.1 fL]; P < .0001).

Genetic variation and MPV

Supplemental Table 1 presents the SNPs evaluated in this study. By using genotype linear association analysis, we tested their association with MPV. For 19 of these SNPs, which were not on the Affymetrix array used in this study, an SNP in linkage equilibrium on the array was chosen with $r^2 > .8$ between the lead SNP and tag SNP. By correlation analysis, we identified 14 SNPs that were strongly correlated with each other (Pearson correlation coefficient $R^2 \sim 1$) as shown in supplemental Figure 4. We consequently excluded 7 of these SNPs from multivariable linear regression analysis.

Linear regression analysis adjusted for age revealed 7 SNPs for females and 4 SNPs for males associated with higher MPV (Table 3). Three of these SNPs (rs342293, rs7961894, and rs342251) were associated with MPV in both sexes. A multivariable model that included both genetic and nongenetic factors by using the variable stepwise selection by Akaike information criterion explained 20.4% of the MPV variance in females and 18.6% in males (Table 4). Compared with the sum of all nongenetic factors, genetic factors explained 8.6% of MPV variance in females and 6.8% in males. WBC count and fibrinogen concentrations remained independently positively correlated with MPV in both males and females (P < .001), whereas platelet count persisted as being negatively correlated (P < .001). Oral contraceptives were still associated with higher MPV in females with an estimated β of 0.271 (95% CI, 0.109

	Men	O Wor	nen			
					Beta [95%CI]	p-value
* Oral contraceptives		⊢ ∽			0.11 [0.03; 0.20]	0.0091
HRT	F)]			0.02 [-0.05; 0.09]	0.52
Antithrombotic agents (B01A)					0.15 [0.09; 0.21] 0.10 [0.02; 0.18]	<0.0001 0.0099
Aspirin (B01AC56)					0.12 [0.06; 0.19] 0.08 [0.00; 0.17]	0.00014 0.053
Antilipemic drugs (C10A)	ŀ		1		0.10 [0.04; 0.16] 0.05 [-0.02; 0.12]	0.0012 0.13
Antihypertensives (C02)	F		1		0.11 [0.07; 0.16] 0.02 [-0.03; 0.07]	<0.0001 0.38
ACE inhibitors (C09)	Þ		1		0.11 [0.06; 0.16] 0.01 [-0.04; 0.07]	<0.0001 0.60
Diuretics (C03)	ŀ			-	0.21 [0.12; 0.31] 0.11 [0.02; 0.20]	<0.0001 0.019
ARBs (C09C)					0.02 [-0.07; 0.12] 0.05 [-0.05; 0.15]	0.62 0.31
Calcium channel blockers (C08)	⊢ ⊫		-1		0.04 [-0.03; 0.12] 0.08 [-0.01; 0.16]	0.29 0.077
B-blocker (C07)			•		0.17 [0.11; 0.22] 0.05 [-0.01; 0.11]	<0.0001 0.078
-C bata	.1 0	0.1 (MPV [fl 1)	0.2	0.3	0.4	

Figure 1. Medical treatment and MPV. Estimates of β with 95% CIs for each drug from linear regression modeling for MPV adjusted for age. Results are displayed with stratification for males (black) and females (white). *Self-reported information. ACE, angiotensinconverting enzyme; ARB, angiotensin II receptor blocker; HRT, hormone replacement therapy.

to 0.433; $P = .001$), whereas smoking correlated with higher	i
MPV in males (β , 0.103; 95% CI, 0.0094 to 0.196; $P = .031$). Of	I
7 SNPs initially associated with MPV in females, 6 remained	а

independently associated with higher MPV, and of 4 SNPs in nales, 2 remained as determinants in multivariable linear regression analysis.

Table 2. Sex-specific	β	estimates	of	determinants f	for	MPV	in	the	ро	pulation sa	ample
-----------------------	---	-----------	----	----------------	-----	-----	----	-----	----	-------------	-------

		Females (n = 7081)		Males (n = 7386)			
	β	95% CI	Р	β	95% CI	Р	
Age (per 5 y)	0.00776	-0.00672 to 0.0222	.29	-0.00134	-0.0120 to 0.00930	.80	
Menopause (%)	-0.0595	-0.126 to 0.00731	.081	_		-	
Menstruation (%)	0.123	0.0232 to 0.224	.016	—		_	
HRT (%)	0.0429	-0.0224 to 0.108	.20	_		-	
Oral contraceptives (%)	0.150	0.0649 to 0.236	.00057	_		_	
Hypertension (%)	0.0351	-0.0181 to 0.0883	.20	0.0530	0.00520 to 0.101	.030	
Smoking (%)	0.0476	-0.00686 to 0.102	.087	0.179	0.129 to 0.230	<.0001	
Dyslipidemia (%)	-0.0410	-0.0925 to 0.0105	.12	0.0224	-0.0185 to 0.0632	.28	
Diabetes (%)	0.0591	-0.0428 to 0.161	.26	0.0358	-0.0461 to 0.118	.39	
Obesity (%)	0.0273	-0.0234 to 0.0780	.29	0.0280	-0.0181 to 0.0740	.23	
Family history of MI or stroke (%)	-0.0161	-0.0623 to 0.0300	.49	0.0281	-0.0197 to 0.0760	.25	
Beta blockers (%)	0.00560	-0.0703 to 0.0815	.89	0.0542	-0.0166 to 0.125	.13	
Antiplatelets (%)	0.0150	-0.132 to 0.161	.84	-0.0324	-0.140 to 0.0755	.56	
Antihypertensives (%)	-0.0298	-0.123 to 0.0631	.53	-0.0655	-0.160 to 0.0288	.17	
Aspirin (%)	0.0512	-0.104 to 0.207	.52	0.0819	-0.0294 to 0.193	.15	
Antilipemic drugs (%)	0.0130	-0.0606 to 0.0867	.73	0.00998	-0.0537 to 0.0737	.76	
ACE inhibitors (%)	-0.0282	-0.108 to 0.0518	.49	0.0557	-0.0268 to 0.138	.19	
Diuretics (%)	0.0487	-0.0458 to 0.143	.31	0.0703	-0.0225 to 0.163	.14	
WBC (× 10 ⁹ /L)	0.0685	0.0574 to 0.0796	<.0001	0.0394	0.0308 to 0.0479	<.0001	
Platelets (× 10 ⁹ /L)	-0.00440	-0.00469 to -0.00411	<.0001	-0.00456	-0.00489 to -0.00424	<.0001	
CRP (mg/L)	-0.00470	-0.00910 to -0.000295	.037	-0.00494	-0.00890 to -0.000981	.014	
Fibrinogen (mg/dL)	0.000652	0.000347 to 0.000956	<.0001	0.000812	0.000511 to 0.00111	<.0001	
Glucose (mg/dL)	0.00126	-0.000120 to 0.00263	.074	0.00184	0.000642 to 0.00305	.0027	
Hemoglobin (g/dL)	0.00311	-0.0176 to 0.0239	.77	-0.0100	-0.0293 to 0.00926	.31	

Multivariable linear regression model for MPV with adjustment for all variables included in the table. Males and females: R², 0.119. P values <.05 are shown in bold. ACE, angiotensin-converting enzyme; HRT, hormone replacement therapy.





Figure 2. Distribution of MPV in males and females in the population sample ($N = 14\,968$). Effects of sex and age on MPV by using quantile regression for the median. The graphic shows the median values of MPV (middle lines) with 5th and 95th percentile (bottom and top lines) for males (solid lines) and females (dashed lines).

MPV, SNP, and CVRF interaction

To evaluate effects on MPV of interaction between SNPs and CVRFs, we performed multivariable linear regression analysis that included the associated SNP and an interaction term with each CVRF as a separate variable (supplemental Table 2). Interestingly, the rs649729 variation in females was associated with an MPV increase of 0.145 fL (P = .023), and the presence of this SNP variation together with smoking was associated with a further increase in MPV of 0.251 fL (P = .019). An rs4774471 variation was associated with a significant increase in MPV of 0.197 fL in combination with hypertension in females (P = .021). In males, there was a significant interaction between a family history of MI or stroke and rs10506328 and/or rs2296828 variation leading to MPV increases of 0.194 fL (P = .042) and 0.213 fL (P = .024), respectively.

MPV and total mortality

A total of 365 deaths were registered during the follow-up period up to December 2014 with a median follow-up time of 5.0 years (interquartile range, 3.6-6.3 years). As shown in Figure 3A, the cumulative mortality in individuals with MPV above the reference limit (>10.1 fL) was significantly worse (P = .00035) compared with individuals with MPV below the 95th percentile. Analysis of cumulative mortality, stratified for sex, showed a significant difference for males only (Figure 3B-C). Cox regression analysis adjusted for age and sex revealed that an increase in MPV of 1 fL was associated with an increase of 16% in mortality (hazard ratio, 1.16; 95% CI, 1.04 to 1.29; P = .0063). In addition, adjusting for traditional CVRFs and oral contraceptives did not significantly change the association between higher MPV and increased mortality (supplemental Table 3A). Only after further adjusting the Cox regression analysis for cardiovascular diseases was the significant association between MPV and mortality lost (supplemental Table 3B).

Discussion

This is the first sex-stratified comprehensive analysis of several genetic and nongenetic factors on MPV in a population-based cohort. Interestingly, despite commonalities, the importance of this approach is highlighted by substantial differences in the profiles of genetic and nongenetic determinants for MPV between males and females.

This is the first study of an adult population-based sample that investigates MPV and total mortality in an analysis stratified by sex. In non-sex-specific analyses, higher MPV has already been associated with total mortality in the general population and with vascular mortality in a large population of hospitalized patients.^{12,16} We now show that an MPV outside the reference range is associated with a higher risk of death in males only. Cox regression analysis demonstrated that MPV is a robust determinant of increased mortality after adjusting for age, sex, oral contraceptive use, and traditional CVRFs. The loss of association between MPV and mortality after adjusting for heart failure, peripheral arterial disease, and stroke, which have been associated with an increased MPV,²⁷⁻²⁹ suggests a causal relationship between MPV and the development of cardiovascular disease or vice versa, which needs to be further explored.

Second, analysis displayed another interaction between sex and age without an effect of age on MPV in women. As also observed by others, MPV was significantly higher in females compared with males and it increased with age in the total sample.^{14,16,19,30} However, the study implies that MPV was higher in females until the age of approximately

	able 3. SNPs	positively	associated	with I	MP\
--	--------------	------------	------------	--------	-----

SNP	Alleles	β	95% CI	Р
Females (n = 2042)				
rs342293	C/G vs C/C	0.134	0.0441 to 0.224	.0035
	G/G vs C/C	0.295	0.182 to 0.408	<.0001
rs7961894	C/T vs C/C	0.218	0.121 to 0.316	<.0001
	T/T vs C/C	0.968	0.559 to 1.38	<.0001
rs12485738	G/A vs G/G	0.101	0.0157 to 0.186	.020
	A/A vs G/G	0.320	0.197 to 0.443	<.0001
rs649729	A/T vs A/A	0.152	0.0692 to 0.234	.00032
	T/T vs A/A	0.172	0.0244 to 0.319	.022
rs342251	G/A vs G/G	0.134	0.0454 to 0.223	.0031
	A/A vs G/G	0.250	0.135 to 0.365	<.0001
rs17568628	T/C vs T/T	0.345	0.182 to 0.509	<.0001
	C/C vs T/T	0.625	-1.16 to 2.41	.49
rs4774471	T/A vs T/T	0.0861	0.00249 to 0.170	.044
	A/A vs T/T	0.278	0.146 to 0.411	<.0001
Males (n = 2131)				
rs342293	C/G vs C/C	0.0286	-0.0593 to 0.117	.52
	G/G vs C/C	0.229	0.121 to 0.338	<.0001
rs7961894	C/T vs C/C	0.253	0.155 to 0.351	<.0001
	T/T vs C/C	0.475	0.147 to 0.802	.0045
rs10876550	A/G vs A/A	0.121	0.0326 to 0.209	.0073
	G/G vs A/A	0.205	0.0957 to 0.314	.00024
rs342251	G/A vs G/G	0.0319	-0.0547 to 0.118	.47
	A/A vs G/G	0.243	0.133 to 0.353	<.0001

Linear regression models for females and males adjusted for age. SNPs positively associated with MPV and shared by both males and females are shown in bold. Seventy-two tests were performed per sex (Bonferroni corrected α , .000694).

Table 4. Variable selected model by Akaike information criteria

Variable	Alleles	β	95% CI	Р
Females (n = 1975)				
Age (per 5 y)		-0.0275	-0.0476 to -0.00732	.0076
Hypertension (%)		0.0849	-0.0122 to 0.182	.086
Diabetes (%)		0.206	0.0393 to 0.373	.015
Obesity (%)		-0.0684	-0.164 to 0.0268	.16
WBC (× 10 ⁹ /L)		0.0448	0.0240 to 0.0657	<.001
Platelets (× 10 ⁹ /L)		-0.00421	-0.00476 to -0.00366	<.001
Fibrinogen (mg/dL)		0.00121	0.000692 to 0.00173	<.001
Oral contraceptive (%)		0.271	0.109 to 0.433	.0010
Antithrombotics (%)		0.132	-0.00214 to 0.266	.054
Antihypertensive (%)		-0.114	-0.219 to -0.00860	.034
rs342293	C/G vs C/C	0.0978	0.0145 to 0.181	.021
rs342293	G/G vs C/C	0.236	0.132 to 0.340	<.001
rs10506328	C/A vs C/C	0.0565	-0.0160 to 0.129	.13
rs12485738	A/A vs G/G	0.169	0.0642 to 0.274	.0016
rs2138852	T/C vs T/T	-0.0651	-0.153 to 0.0224	.14
rs2138852	C/C vs T/T	-0.185	-0.287 to -0.0836	<.001
rs10076782	G/A vs G/G	0.0553	-0.0200 to 0.131	.15
rs11734132	G/C vs G/G	-0.0607	-0.141 to 0.0195	.14
rs12969657	T/C vs T/T	-0.128	-0.216 to -0.0396	.0045
rs12969657	C/C vs T/T	-0.105	-0.210 to -0.00119	047
rs13042885	C/T vs C/C	-0.0860	-0 162 to -0 00948	028
rs13042885	T/T vs C/C	-0.150	-0.298 to -0.00252	046
rs2950390	T/T vs C/C	-0.0818	-0 193 to 0 0296	15
rs4812048		-0.0897	-0.172 to -0.00777	032
rs4812048		-0.190	-0.420 to 0.0395	10
re6/0720		0.130	-0.420 to 0.0395	.10
rs649729		0.100	0.0240 to 0.177	.010
re7317038		-0.0624	-0 136 to 0 0116	.040
ro10255794		0.0024	-0.107 to 0.0219	.030
re12355784		-0.164	-0.197 to -0.0218	.014
ro4051194		0.104	-0.207 to 0.0002	.0013
ro4051104		0.0007	-0.145 to 0.0120	.090
154951164		-0.232	$-0.342 \ 10 \ -0.121$	<.001
ro7242300		0.0023	-0.0193 10 0.144	.13
rs10512629		0.147	0.0361 10 0.255	.0001
ro17207120		0.0005	-0.0214 10 0.156	.14
ro17207120		0.125	0.0397 10 0.211	.0041
ro4774471		0.412	0.103 to 0.720	.0009
154774471 ro4774471		0.0011	0.00424 10 0.156	.039
154/74471		0.237	0.115 to 0.300	017
ro17569609		0.0902	0.0103 10 0.104	.017
IS1700020		0.204	0.115 to 0.412	< .001
157901094	1/1 VS C/C	0.222	0.134 10 0.311	<.001
Males (n = 2084)		0.0245	0.0500 to 0.0160	< 001
Age (per 5 y)		-0.0345	-0.052910 - 0.0160	<.001
Obsoits (%)		0.103	0.00940 to 0.196	.031
$\frac{10^9}{10}$		0.0004	0.00368 10 0.109	.040
$\frac{10^{9}}{10^{1}}$		0.0290	0.0165 t0 0.0415	<.001
$\frac{10^{-1}}{CPR} (mc^{/1})$		-0.00468	-0.00528 to -0.00408	<.001
		-0.00743		.007
Appiring (%)		0.105	0.000323 to 0.00144	.0020
Asplinin (%)		0.125	0.00360 10 0.247	.043
Divertice (%)		0.110	-0.000248 to 0.222	.051
		0.149	-0.00325 10 0.302	.055
15042293		0.172	0.0002 10 0.259	<.001
1510506328	U/A VS U/U	0.138		<.001
101000328	A/A VS C/C	0.171	0.0004 10 0.277	.0015
1512465738	AVA VS G/G	0.116	0.0132 10 0.218	.027
152138852	1/C VS 1/1	-0.0741	-0.159 to 0.0103	.085
rs2138852	C/C VS I/I	-0.180	-0.2// to -0.0823	<.001
152180/48	G/TVSG/G	-0.0960	-0.175 to -0.0173	.017
rs2180/48	1/1 VS G/G	-0.110	-0.214 to -0.00552	.039
rs11734132	G/C vs G/G	-0.0665	-0.145 to 0.0116	.095
rs11/34132	C/C vs G/G	-0.188	-0.416 to 0.0405	.11
rs13042885	C/T vs C/C	-0.0758	-0.150 to -0.00184	.045

	lucuj			
Variable	Alleles	β	95% CI	Р
rs13042885	T/T vs C/C	-0.215	-0.352 to -0.0779	.0021
rs2950390	T/T vs C/C	-0.103	-0.214 to 0.00861	.071
rs12355784	A/A vs C/C	-0.0651	-0.146 to 0.0153	.11
rs4951184	T/A vs T/T	-0.0603	-0.136 to 0.0159	.12
rs4951184	A/A vs T/T	-0.213	-0.322 to -0.104	<.001
rs7342306	G/A vs G/G	0.136	0.0560 to 0.215	<.001
rs7342306	A/A vs G/G	0.106	0.00683 to 0.206	.036
rs2296828	C/T vs C/C	0.0962	0.0257 to 0.167	.0075
rs4676428	G/G vs T/T	-0.139	-0.261 to -0.0174	.025
rs2816608	C/T vs C/C	-0.125	-0.197 to -0.0541	<.001
rs10512628	C/C vs T/T	0.0643	-0.0192 to 0.148	.13
rs4774471	T/A vs T/T	0.0700	-0.00400 to 0.144	.064
rs4774471	A/A vs T/T	0.145	0.0219 to 0.267	.021
rs17568628	T/C vs T/T	0.206	0.0662 to 0.346	.0039
rs7961894	T/T vs C/C	0.228	0.140 to 0.316	<.001

Table (continued)

Multivariable linear regression model for MPV in females and males with variables selected by Akaike information criteria. Females: R^2 , 0.204; males: R^2 , 0.186. *P* values <.05 are shown in bold.

65 years, after which MPV was higher in males because of the absent relation with age in women.

Third, this study confirmed the influence of female hormonal status on MPV, particularly in the presence of menstruation or intake of oral contraceptives. Female sex hormones have been suggested to have an effect on the regulation of platelet indices, because the role of endogenous estrogen on proplatelet formation has been demonstrated in animal models.³¹ Estrogen replacement therapy has also been associated with an increase in MPV in postmenopausal women; however, we could not confirm an independent relationship in our study.³² A recent study reported an increased platelet count in menstruating women suggesting that loss of iron during bleeding could stimulate platelet production.³³⁻³⁵ Higher MPV in menstruating women can be explained by the increased platelet production with release of young platelets, which are known to have larger volume and reactivity compared with mature platelets.³⁶

Fourth, MPV was significantly and positively associated with CVRFs in males, in particular with smoking, hypertension, and high glucose levels. Smoking has been previously associated with increased MPV, and studies have demonstrated that duration and intensity of active and/or passive smoking influences MPV.^{12,37} Furthermore, smoking cessation was associated with decreased MPV values.³⁸ Higher MPV in smokers could partly be induced by smoking-mediated inflammation, as demonstrated by the independent correlation between highly sensitive CRP and smoking.³⁹ Our study confirmed this finding and also displays smoking as being associated with higher MPV independently of WBC count, CRP, and fibrinogen. We also confirmed the association between MPV and WBC count and/or fibrinogen, which supports the link between platelets and inflammation in both males and females.^{19,40,41}

Fifth, this is the first study to include all known genetic variants from large GWASs in exploring MPV heterogeneity at the population level. Genetic variants associated with MPV explained 8.6% of the MPV variation in females and 6.8% in males. In a multivariable linear regression analysis, 6 SNPs in females and 2 SNPs in males remained independently positively associated with MPV. In females, we determined an increase in MPV of 0.251 fL for an interaction of the rs649729 polymorphism (which was correlated with MPV only in females) with smoking. Located on chromosome 2p23.1, this



Figure 3. Total mortality and MPV. Plots of cumulative mortality depending on MPV in (A) the overall sample (N = 14576), (B) in males (n = 7,319), and (C) in females (n = 7,257). Individuals were stratified for the 95th percentile of MPV in the reference group as cutoff value (cutoff values for MPV: overall sample, 10.1fL; males, 9.96 fL; females, 10.1 fL).

intronic SNP lies in the Eps15 homology (EH) -domain containing 3 (EHD3) gene which is highly expressed in brain and heart and functionally associated with regulation of intracellular protein transport, membrane trafficking, and endocytosis.⁴² Recently, the EHD3 polymorphisms, including rs649729, have been associated with major depressive disorder in females only.⁴³ This mechanism may explain the recent finding of higher MPV in patients with major depression.⁴⁴ Another interaction in females was found between hypertension and rs4774471. This SNP is associated with the TPM1 gene on chromosome 15q22.2 that encodes tropomyosin I, which regulates the calcium-dependent interaction of actin and myosin, a key step in platelet formation.²⁵ rs4774471 was associated with higher MPV in both sexes. In males, we observed an important interaction between 2 SNPs, rs10506328 (nuclear factor erythroid 2 [NFE2] and coatomer protein complex [1 [COPZ1]) and rs2296828 (dedicator of cytokinesis [DOCK8]), and family history of MI and/or stroke that was associated with higher MPV. rs10506328 (located on chromosome 12q13.13) has been associated with the NFE2 gene which is critical for megakaryocyte development and proplatelet formation, whereas the COPZ1 gene, which encodes a subunit of coatomer protein complex 1 (COPI), is involved in intracellular traffic and autophagy.^{45,46} The DOCK8 gene on chromosome 9p24.3 (rs2296828) regulates interstitial dendritic cell migration and is critical for the survival and function of natural killer T cells.47 Family history of MI and/or stroke is known to be a major cardiovascular risk factor. The presence of these two SNPs identified in our study might represent an additional factor of increased thrombotic risk.

Finally, antithrombotics, aspirin, and diuretics in both males and females and antilipemic and antihypertensive drugs in males were consistent with an altered MPV when using univariate models adjusted for age. However, the effects lost significance after analyzing them in multivariable linear regression. In the literature, statins were described as decreasing MPV irrespective of their cholesterol-lowering effect.^{48,49} Amlodipine-based therapy has been associated with a decrease in MPV, whereas low-dose aspirin showed no effect on MPV in healthy volunteers after 7 days of ingestion of 81 mg aspirin.^{7,50} However, effects of medical treatment on MPV were beyond the scope of this study, and other study designs and/or study samples are necessary for investigating this interesting issue in more detail.

The variability of MPV as a result of changing preanalytics, however, is a major limiting factor in using this parameter as a marker of platelet reactivity in clinical practice. Time to measurement after blood sampling, storage temperature of sample tubes, type of anticoagulant used, and the instrument used for MPV measurements have all been implicated in MPV variation and discordance among studies.⁵¹ Because platelets shrink in citrate and swell in EDTA, several groups proposed different measuring times ranging from within 1 hour regardless of the anticoagulant used to 120 minutes for EDTA and 60 minutes for citrate anticoagulant.^{22,52} A recent study by Shah et al⁵⁰ also observed platelet swelling in EDTA until 180 minutes after blood withdrawal. Interestingly, Diaz et al⁵³ reported a progressive increase in MPV with storage in EDTA measured at different time intervals up to 24 hours. To control for variation in these preanalytical factors, we performed our analysis under standardized conditions measuring MPV between 30 and 90 minutes after blood sampling. Other strengths of this study are the large sample size and the highly standardized setting that supported the comprehensive analysis of MPV with both genetic and nongenetic determinants in a sex-specific approach.

In conclusion, this study provides evidence for substantial heterogeneity in the profile of determinants for MPV between sexes. To the best of our knowledge, this is the first large population-based study showing evidence of interaction between genetic variations, CVRFs, and MPV, suggesting a need for deeper insight into the interplay between environmental factors and genetic variability for MPV as a biomarker that indicates platelet function. Additional studies should investigate the role of MPV in males for increased risk of thrombosis and development of cardiovascular disease, because they are significantly determined by traditional CVRFs and associated with total mortality.

Acknowledgments

This work was supported by contract AZ 961-386261/733 from the government of Rhineland-Palatinate (Stiftung Rheinland-Pfalz für Innovation,), by research programs Wissen schafft Zukunft and Center for Translational Vascular Biology of the Johannes Gutenberg-University of Mainz, and contracts with Boehringer Ingelheim and Philips Medical Systems, including an unrestricted grant for the Gutenberg Health Study. P.S.W., S.B., T.Z., and T.M. are principal investigators of the German Center for Cardiovascular Research. H.t.C. is a Fellow of the Gutenberg Research Foundation.

Authorship

Contribution: M.P.-N. designed and performed research, analyzed and interpreted data, and wrote the paper; A.S. performed the statistical analysis; M.I.H. and V.G. contributed to discussion of the results and to critical review; D.L.-R. collected laboratory and clinical data; H.M.H.S., H.t.C., K.J.L., and T.M. contributed to writing the paper and to critical review; E.P., H.B., M.B., N.P., S.B., and T.Z. performed research; P.S.W. designed and performed research, interpreted data, and contributed to writing the paper; and all authors have read and approved the manuscript in its current form.

Conflict-of-interest disclosure: M.I.H., V.G., and P.S.W. were funded by the Federal Ministry of Education and Research (BMBF 01EO1003). H.t.C. is a Fellow of the Gutenberg Forschungskolleg of the Johannes Gutenberg-University, Mainz, Germany. P.S.W. has

References

- Leslie M. Cell biology. Beyond clotting: the powers of platelets. *Science*. 2010;328(5978):562-564.
- Vizioli L, Muscari S, Muscari A. The relationship of mean platelet volume with the risk and prognosis of cardiovascular diseases. *Int J Clin Pract.* 2009; 63(10):1509-1515.
- Karpatkin S, Khan Q, Freedman M. Heterogeneity of platelet function. Correlation with platelet volume. Am J Med. 1978;64(4):542-546.
- Thompson CB, Jakubowski JA, Quinn PG, Deykin D, Valeri CR. Platelet size as a determinant of platelet function. *J Lab Clin Med.* 1983;101(2): 205-213.
- Bath PM, Butterworth RJ. Platelet size: measurement, physiology and vascular disease. Blood Coagul Fibrinolysis. 1996;7(2):157-161.
- Chu SG, Becker RC, Berger PB, et al. Mean platelet volume as a predictor of cardiovascular risk: a systematic review and meta-analysis. *J Thromb Haemost.* 2010;8(1):148-156.
- Nadar S, Blann AD, Lip GY. Platelet morphology and plasma indices of platelet activation in essential hypertension: effects of amlodipinebased antihypertensive therapy. *Ann Med.* 2004; 36(7):552-557.
- Papanas N, Symeonidis G, Maltezos E, et al. Mean platelet volume in patients with type 2 diabetes mellitus. *Platelets*. 2004;15(8):475-478.
- Coban E, Yilmaz A, Sari R. The effect of weight loss on the mean platelet volume in obese patients. *Platelets*. 2007;18(3):212-216.
- Pathansali R, Smith N, Bath P. Altered megakaryocyte-platelet haemostatic axis in hypercholesterolaemia. *Platelets.* 2001;12(5): 292-297.
- Kario K, Matsuo T, Nakao K. Cigarette smoking increases the mean platelet volume in elderly patients with risk factors for atherosclerosis. *Clin Lab Haematol.* 1992;14(4):281-287.
- Klovaite J, Benn M, Yazdanyar S, Nordestgaard BG. High platelet volume and increased risk of myocardial infarction: 39,531 participants from the general population. J Thromb Haemost. 2011; 9(1):49-56.
- Sansanayudh N, Anothaisintawee T, Muntham D, McEvoy M, Attia J, Thakkinstian A. Mean platelet volume and coronary artery disease: a systematic review and meta-analysis. *Int J Cardiol.* 2014; 175(3):433-440.
- Braekkan SK, Mathiesen EB, Njølstad I, Wilsgaard T, Størmer J, Hansen JB. Mean platelet volume is a risk factor for venous thromboembolism: the Tromsø Study, Tromsø, Norway. J Thromb Haemost. 2010;8(1):157-162.

- Endler G, Klimesch A, Sunder-Plassmann H, et al. Mean platelet volume is an independent risk factor for myocardial infarction but not for coronary artery disease. *Br J Haematol.* 2002; 117(2):399-404.
- Slavka G, Perkmann T, Haslacher H, et al. Mean platelet volume may represent a predictive parameter for overall vascular mortality and ischemic heart disease. *Arterioscler Thromb Vasc Biol.* 2011;31(5):1215-1218.
- Johnson M, Ramey E, Ramwell PW. Sex and age differences in human platelet aggregation. *Nature*. 1975;253(5490):355-357.
- Stevens RF, Alexander MK. A sex difference in the platelet count. *Br J Haematol.* 1977;37(2): 295-300.
- Santimone I, Di Castelnuovo A, De Curtis A, et al; MOLI-SANI Project Investigators. White blood cell count, sex and age are major determinants of heterogeneity of platelet indices in an adult general population: results from the MOLI-SANI project. *Haematologica*. 2011;96(8):1180-1188.
- Shameer K, Denny JC, Ding K, et al. A genomeand phenome-wide association study to identify genetic variants influencing platelet count and volume and their pleiotropic effects. *Hum Genet*. 2014;133(1):95-109.
- Siller-Matula JM, Arbesu I, Jilma B, Maurer G, Lang IM, Mannhalter C. Association between the rs342293 polymorphism and adverse cardiac events in patients undergoing percutaneous coronary intervention. *Thromb Haemost.* 2014; 111(6):1060-1066.
- Lancé MD, van Oerle R, Henskens YM, Marcus MA. Do we need time adjusted mean platelet volume measurements? *Lab Hematol.* 2010; 16(3):28-31.
- Wild PS, Zeller T, Beutel M, et al. [The Gutenberg Health Study]. Bundesgesundheitsblatt Gesundheitsforschung Gesundheitsschutz [in German]. 2012;55(6-7):824-829.
- Hindorff LA, MacArthur J, Morales J, Junkins HA, Hall PN, Klemm AK, Manolio TA. A Catalog of Published Genome-Wide Association Studies. Available at: www.genome.gov/gwastudies. Accessed July 20, 2015.
- Soranzo N, Spector TD, Mangino M, et al. A genome-wide meta-analysis identifies 22 loci associated with eight hematological parameters in the HaemGen consortium. *Nat Genet.* 2009; 41(11):1182-1190.
- Gieger C, Radhakrishnan A, Cvejic A, et al. New gene functions in megakaryopoiesis and platelet formation. *Nature*. 2011;480(7376):201-208.

received research funding from Boehringer Ingelheim; Philips Medical Systems; sanofi-aventis; Bayer Vital; Daiichi Sankyo Europe; Institute for the Modernization of Economic Base and Employment Structures; Portavita; Federal Institute for Occupational Safety and Health; Health Economy Initiative, Ministry of Health, and Ministry of Economics, Rhineland-Palatinate; Federal Ministry of Education and Research; Federal Ministry of Health, Rhineland-Palatinate; and Mainz Heart Foundation, and has received honoraria for lectures or consulting from Boehringer Ingelheim and Public Health, Heinrich-Heine-University Düsseldorf. The remaining authors declare no competing financial interests.

Correspondence: Marina Panova-Noeva, Clinical Epidemiology, Center for Thrombosis and Hemostasis, Universitätsmedizin der Johannes Gutenberg-Universität Mainz, Langenbeckstrasse 1, 55131 Mainz, Germany; e-mail: marina.panova-noeva@unimedizinmainz.de.

- Kandis H, Ozhan H, Ordu S, et al. The prognostic value of mean platelet volume in decompensated heart failure. *Emerg Med J.* 2011;28(7):575-578.
- Berger JS, Eraso LH, Xie D, Sha D, Mohler ER III. Mean platelet volume and prevalence of peripheral artery disease, the National Health and Nutrition Examination Survey, 1999-2004. *Atherosclerosis.* 2010;213(2):586-591.
- Bath P, Algert C, Chapman N, Neal B; PROGRESS Collaborative Group. Association of mean platelet volume with risk of stroke among 3134 individuals with history of cerebrovascular disease. Stroke. 2004;35(3):622-626.
- Lippi G, Meschi T, Borghi L. Mean platelet volume increases with aging in a large population study. *Thromb Res.* 2012;129(4):e159-e160.
- Nagata Y, Yoshikawa J, Hashimoto A, Yamamoto M, Payne AH, Todokoro K. Proplatelet formation of megakaryocytes is triggered by autocrinesynthesized estradiol. *Genes Dev.* 2003;17(23): 2864-2869.
- Ranganath LR, Christofides J, Semple MJ. Increased mean platelet volume after oestrogen replacement therapy. *Ann Clin Biochem.* 1996; 33(Pt 6):555-560.
- 33. Balduini CL, Noris P. Platelet count and aging. *Haematologica*. 2014;99(6):953-955.
- 34. Beguin Y. Erythropoietin and platelet production. *Haematologica*. 1999;84(6):541-547.
- Kadikoylu G, Yavasoglu I, Bolaman Z, Senturk T. Platelet parameters in women with iron deficiency anemia. J Natl Med Assoc. 2006;98(3):398-402.
- Harrison P, Robinson MS, Mackie IJ, Machin SJ. Reticulated platelets. *Platelets*. 1997;8(6): 379-383.
- Yarlioglues M, Ardic I, Dogdu O, et al. The acute effects of passive smoking on mean platelet volume in healthy volunteers. *Angiology*. 2012; 63(5):353-357.
- Cho SY, You E, Lee HJ, Lee WI, Park TS. Smoking cession decreases mean platelet volume in healthy Korean populations. *Clin Lab.* 2014;60(8):1413-1416.
- Alyan O, Kaçmaz F, Ozdemir O, et al. [High levels of high-sensitivity C-reactive protein and impaired autonomic activity in smokers] [in Turkish]. *Turk Kardiyol Dern Ars.* 2008;36(6):368-375.
- Gasparyan AY, Ayvazyan L, Mikhailidis DP, Kitas GD. Mean platelet volume: a link between thrombosis and inflammation? *Curr Pharm Des*. 2011;17(1):47-58.
- 41. Morrell CN, Aggrey AA, Chapman LM, Modjeski KL. Emerging roles for platelets as immune and

inflammatory cells. *Blood.* 2014;123(18): 2759-2767.

- Pohl U, Smith JS, Tachibana I, et al. EHD2, EHD3, and EHD4 encode novel members of a highly conserved family of EH domain-containing proteins. *Genomics*. 2000;63(2):255-262.
- Wang L, Shi C, Zhang K, Xu Q. The genderspecific association of EHD3 polymorphisms with major depressive disorder. *Neurosci Lett.* 2014; 567:11-14.
- Canan F, Dikici S, Kutlucan A, et al. Association of mean platelet volume with DSM-IV major depression in a large community-based population: the MELEN study. *J Psychiatr Res.* 2012;46(3):298-302.
- Avanzi MP, Goldberg F, Davila J, Langhi D, Chiattone C, Mitchell WB. Rho kinase inhibition drives megakaryocyte polyploidization and

proplatelet formation through MYC and NFE2 downregulation. *Br J Haematol.* 2014;164(6): 867-876.

- Shtutman M, Baig M, Levina E, et al. Tumor-specific silencing of COPZ2 gene encoding coatomer protein complex subunit ζ 2 renders tumor cells dependent on its paralogous gene COPZ1. Proc Natl Acad Sci USA. 2011;108(30):12449-12454.
- Nishikimi A, Kukimoto-Niino M, Yokoyama S, Fukui Y. Immune regulatory functions of DOCK family proteins in health and disease. *Exp Cell Res.* 2013;319(15):2343-2349.
- Sivri N, Tekin G, Yalta K, Aksoy Y, Senen K, Yetkin E. Statins decrease mean platelet volume irrespective of cholesterol lowering effect. *Kardiol Pol.* 2013;71(10):1042-1047.
- 49. Akin F, Ayça B, Köse N, et al. Effect of atorvastatin on hematologic parameters in

patients with hypercholesterolemia. *Angiology.* 2013;64(8):621-625.

- Shah B, Valdes V, Nardi MA, Hu L, Schrem E, Berger JS. Mean platelet volume reproducibility and association with platelet activity and antiplatelet therapy. *Platelets*. 2014;25(3):188-192.
- Lippi G, Franchini M. Platelets and immunity: the interplay of mean platelet volume in health and disease. *Expert Rev Hematol.* 2015;8(5): 555-557.
- Dastjerdi MS, Emami T, Najafian A, Amini M. Mean platelet volume measurement, EDTA or citrate? *Hematology*. 2006;11(5):317-319.
- Diaz-Ricart M, Brunso L, Pino M, et al. Preanalytical treatment of EDTA-anticoagulated blood to ensure stabilization of the mean platelet volume and component measured with the ADVIA counters. *Thromb Res.* 2010;126(1):e30-e35.