

LYMPHOID NEOPLASIA

***TNFRSF14* aberrations in follicular lymphoma increase clinically significant allogeneic T-cell responses**

Eleni Kotsiou, Jessica Okosun, Caroline Besley, Sameena Iqbal, Janet Matthews, Jude Fitzgibbon, John G. Gribben, and Jeffrey K. Davies

Centre for Haemato-oncology, Barts Cancer Institute, Queen Mary University, London, United Kingdom

Key Points

- *TNFRSF14* gene aberrations, common in FL, increase the ability of lymphoma cells to stimulate allogeneic T-cell responses.
- *TNFRSF14* lesions were associated with increased acute GVHD supporting stratified transplantation approaches in the allogeneic setting.

Donor T-cell immune responses can eradicate lymphomas after allogeneic hematopoietic stem cell transplantation (AH SCT), but can also damage healthy tissues resulting in harmful graft-versus-host disease (GVHD). Next-generation sequencing has recently identified many new genetic lesions in follicular lymphoma (FL). One such gene, tumor necrosis factor receptor superfamily 14 (*TNFRSF14*), abnormal in 40% of FL patients, encodes the herpes virus entry mediator (HVEM) which limits T-cell activation via ligation of the B- and T-lymphocyte attenuator. As lymphoma B cells can act as antigen-presenting cells, we hypothesized that *TNFRSF14* aberrations that reduce HVEM expression could alter the capacity of FL B cells to stimulate allogeneic T-cell responses and impact the outcome of AH SCT. In an in vitro model of alloreactivity, human lymphoma B cells with *TNFRSF14* aberrations had reduced HVEM expression and greater alloantigen-presenting capacity than wild-type lymphoma B cells. The increased immune-stimulatory capacity of lymphoma B cells with *TNFRSF14* aberrations had clinical relevance, associating with higher incidence of acute GVHD in patients undergoing AH SCT. FL patients with *TNFRSF14* aberrations may benefit from more aggressive immunosuppression to reduce harmful GVHD after transplantation. Importantly, this study is the first to

demonstrate the impact of an acquired genetic lesion on the capacity of tumor cells to stimulate allogeneic T-cell immune responses which may have wider consequences for adoptive immunotherapy strategies. (*Blood*. 2016;128(1):72-81)

Introduction

Follicular lymphoma (FL) is a common germinal center B-cell malignancy characterized by slow progression but inevitable relapse after conventional chemoimmunotherapy.^{1,2} However, some patients can be cured by the graft-versus-lymphoma (GVL) effect provided by donor T cells in the setting of allogeneic hematopoietic stem cell transplantation (AH SCT).³

FL B cells carry the hallmark t(14;18) translocation which results in cytoplasmic overexpression of the Bcl-2 protein. Two recent studies have reported that additional tumor-specific genetic aberrations of the tumor necrosis factor receptor superfamily 14 (*TNFRSF14*) gene, which encodes the herpes virus entry mediator (HVEM), occur at high frequencies (18%-46%) in FL patients. However, these studies reported conflicting impact of *TNFRSF14* aberrations on clinical outcome, suggesting that their functional effects might be influenced by factors such as differing treatment approaches.^{4,5}

HVEM is a type I transmembrane molecule which acts as a molecular switch through interactions with several different ligands including B- and T-lymphocyte attenuator (BTLA), LIGHT, CD160, lymphotoxin A, and glycoprotein D to regulate a range of immune responses.⁶ Interaction between HVEM expressed on antigen-presenting cells and the coinhibitory receptor BTLA on T cells limits T-cell activation and proliferation.⁷ BTLA has intracellular immunoreceptor tyrosine-based inhibition motifs consistent with immune-inhibitory function, and BTLA-deficient animal models display exaggerated immune responses.⁶ Importantly, BTLA is

expressed by naive CD4⁺ and CD8⁺ T cells, the T-cell compartments known to be enriched for alloreactive specificity, and agonistic antibody-mediated BTLA stimulation reduces donor T-cell-mediated acute GVHD in murine transplant models, consistent with a functional role for BTLA in controlling donor T-cell alloresponses in this setting.⁸⁻¹⁰

Activated FL B cells can act as potent alloantigen-presenting cells in vitro¹¹ and patients with FL often undergo AH SCT with significant residual lymphoma. We hypothesized that *TNFRSF14* aberrations would reduce expression of HVEM and increase the ability of FL B cells to stimulate allogeneic T-cell responses. We therefore determined the functional effect of *TNFRSF14* aberrations on the alloantigen-presenting capacity of human FL B cells in vitro. We also determined the impact of *TNFRSF14* aberrations on clinical alloreactivity in FL patients after HLA-matched reduced-intensity conditioning AH SCT.

Materials and methods**Patient samples**

Lymph node biopsies were obtained from FL patients after written consent. The study was approved by the Local Research Ethical Committee (05/Q0605/140) and was conducted in accordance with the Declaration of Helsinki.

Submitted October 30, 2015; accepted April 5, 2016. Prepublished online as *Blood* First Edition paper, April 21, 2016; DOI 10.1182/blood-2015-10-679191.

The publication costs of this article were defrayed in part by page charge payment. Therefore, and solely to indicate this fact, this article is hereby marked "advertisement" in accordance with 18 USC section 1734.

The online version of this article contains a data supplement.

© 2016 by The American Society of Hematology

TNFRSF14 mutation and deletion analysis of FL B cells

Tumor DNA from pre-AHSCT lymph node biopsies from FL patients was screened for *TNFRSF14* mutations by polymerase chain reaction amplification/Sanger sequencing and for deletions by multiplex ligation-probe amplification as previously described.¹² Primers used for Sanger sequencing are summarized in supplemental Table 1 (available on the *Blood* Web site).

FL B-cell sorting, activation, and phenotyping

FL B cells were stained with CD10–fluorescein isothiocyanate (clone 97C5) and CD20–peridinin chlorophyll (PerCP; clone LT20) antibodies (both from Miltenyi Biotec) and purified by fluorescence-activated cell sorting of dual-positive events on a FACS Aria device (Becton Dickinson). Dead cells were excluded using 4',6-diamidino-2-phenylindole (DAPI). Purity of sorted FL B cells was routinely >90% and sorted FL B cells were routinely >95% light chain–restricted assessed with anti-immunoglobulin light chain κ –Alexa Fluor 700 (clone MHK-49) and anti-immunoglobulin light chain λ –allophycocyanin (APC; clone MHL-38) antibodies (supplemental Figure 1). Following sorting, FL B cells were activated for 48 hours with 1 μ g/mL soluble CD40L (InVivoGen), 5 μ g/mL AffiniPure F(ab')₂ fragment goat anti-human immunoglobulin A (IgA) + IgG + IgM (H+L; Jackson ImmunoResearch), 5 μ g/mL CpG (R&D Systems), and 50 ng/mL interleukin-4 (IL-4; R&D Systems) to optimally upregulate expression of molecules involved in antigen presentation as previously described.^{13,14} Immunophenotyping of CD10⁺CD20⁺ FL B cells was performed by flow cytometry using the following antibodies: HVEM-phycoerythrin (PE; clone 122), CD58-PE (clone TS2/9), major histocompatibility complex (MHC) class I–Pacific Blue (clone W6/32) HLA-DR–APC (clone L243), CD80-PE-cyanine 7 (Cy7; 2D10), CD86-APC (clone IT2.2), and their corresponding isotype controls (all from Biolegend).

Measurement of FL-B-cell–stimulated T-cell alloresponses

Untouched CD3⁺ T cells were purified by negative selection from healthy allogeneic donor peripheral blood mononuclear cells using the Pan T-cell isolation kit (Miltenyi Biotec). Postsort purity assessed by flow cytometry was routinely >95%. T cells were stimulated with activated irradiated (30 Gy) FL B cells at a ratio of 3:1 (T:B cell) for primary allogeneic cocultures. For secondary allogeneic cocultures, T cells received a second round of stimulation on day 3 with irradiated FL B cells. Flow cytometry was used to measure expression of CD4–Alexa Fluor 700 (clone RPA-T4), CD8–PerCP (clone SK1), CD25–APC (clone 4E3), CD127–PE (clone MB15-18C9), and FOXP3–PE–Cy7 (clone) on allogeneic T cells. Proliferation of allogeneic T cells was measured by thymidine incorporation as previously described, and T-cell subset-specific proliferation measured by carboxyfluorescein diacetate succinimidyl ester (CFSE) dye dilution assay.¹⁵ Cytokine secretion in allogeneic cocultures was measured by enzyme-linked immunosorbent assay (ELISA) on culture supernatant harvested at day 5 of primary and secondary cocultures using Ready-Set-Go ELISA kits (eBioscience). Intracellular cytokine staining (interferon γ [IFN- γ]–fluorescein isothiocyanate [clone 4S.B3], tumor necrosis factor α [TNF- α]–PerCP Cy5.5 [clone MAb11], IL-2–PE [clone MQ1-17H12]) and CD107a–PE–Cy5 (clone eBioH4A3) assays were performed on T cells in allogeneic cocultures as previously described.¹⁶ Dead cell exclusion was performed with DAPI, V450-50, or R780/60 fixable viability (from eBioscience). For allospecific proliferation and cytokine secretion, values for T cells stimulated with autologous B cells were subtracted from the values for T cells in allogeneic cultures. For some allogeneic coculture experiments, BTLA agonistic (clone MIH26; Biolegend) or antagonistic antibodies (clone 3B1; Genentech) or their respective isotype controls were added at 50 μ g/mL.

Clinical outcome of FL patients undergoing AHSCT

Patients underwent T-replete AHSCT from fully HLA-matched related or unrelated donors using reduced-intensity conditioning with fludarabine and cyclophosphamide. Seventeen of the 23 patients in this cohort were included in our previous series.¹⁷ Patient and donor demographics and details of conditioning are in Table 1. GVHD prophylaxis was methotrexate (5 mg/m² on days +3, +6, and +11), and cyclosporine (3 mg/kg per day) until day +90 to 120. Remission status prior to AHSCT was assessed using Cheson criteria,

modified for computed tomography positron emission tomography status where appropriate.^{18,19} Acute and chronic GVHD was defined using updated National Institutes of Health (NIH) criteria and graded by consensus criteria confirmed when possible by tissue biopsy.^{20,21} All patients with grades 2–4 acute GVHD were treated initially with 1 to 2 mg/kg methylprednisolone. Acute GVHD was defined as steroid-refractory if progression occurred after 3 days of steroid therapy or there was no response after 7 days.²²

Statistical considerations

Data for in vitro experiments were analyzed using Prism version 5.0. (GraphPad Software, Inc). Two-tailed tests are used throughout. Equal variance was not assumed for unpaired Student *t* tests and Welch correction was used where appropriate. Clinical outcome data were analyzed using SPSS Statistics 22 (IBM) and STATA 12 (Statacorp). The statistical significance of differences in frequencies of categorical variables was assessed using the Fisher exact test. Cumulative incidence of death from acute GVHD was calculated using the method of Fine and Gray with competing risks.²³ Patients transplanted in partial remission were deemed to have persistent disease if detectable at day +100 (or earlier if indicated) and were censored at this date. Overall survival curves were constructed using the method of Kaplan and Meier²⁴ and differences between groups were assessed using the log-rank statistic.²⁵

Results

FL B cells with *TNFRSF14* aberrations have reduced HVEM expression but retain expression of other molecules important in alloantigen presentation

We initially sought to determine the impact of *TNFRSF14* aberrations in human FL B cells on the expression of HVEM and other molecules important to stimulation of T-cell alloresponses. We used flow cytometry to compare surface HVEM expression on *TNFRSF14* wild-type (WT) FL B cells and FL B cells with single or dual *TNFRSF14* aberrations. The individual genetic lesions in FL B cells used for in vitro experiments are detailed in supplemental Table 2. Samples with dual *TNFRSF14* aberrations were selected whose genetic lesions were predicted to result in loss of expression of HVEM (eg, homozygous deletions, nonsense mutations plus deletions).

In WT cases, ~50% of FL B cells expressed HVEM at levels above isotype control, whereas HVEM expression was virtually undetectable in FL B cells with dual *TNFRSF14* aberrations. As expected, FL B cells from patients with single *TNFRSF14* aberrations had HVEM expression intermediate between patients with WT and dual aberrations (Figure 1A–B). In contrast, molecules associated with antigen-presenting capacity (including MHC class I, MHC class II, CD80, CD86, and CD58 [LFA-3]) were expressed at similar levels on *TNFRSF14* WT FL B cells and those with dual *TNFRSF14* aberrations both before and after in vitro activation which models the proinflammatory post-transplant in vivo environment.¹¹ Importantly, in vitro activation had no significant effect on HVEM expression levels (Figure 1C).

FL B cells with dual *TNFRSF14* aberrations stimulated greater frequencies of alloreactive effector T cells

After confirming that aberrations in *TNFRSF14* FL B cells resulted in a significant reduction in expression of HVEM, we chose to investigate whether this impacted upon the ability of FL B cells to stimulate allogeneic T-cell responses in vitro. As HVEM expression was more markedly and consistently reduced in FL B cells with dual rather than single *TNFRSF14* aberrations, we compared *TNFRSF14* WT FL B cells to those with dual aberrations in subsequent functional experiments.

Table 1. Patient and donor demographics and transplant outcome

Patient	Age, y	Prior therapies	Prior autograft	Prior rituximab	Donor	Disease status	<i>TNFRSF14</i> mutation	<i>TNFRSF14</i> deletion	Acute GVHD*	Chronic GVHD	Outcome
1	48	8	Yes	Yes	Sibling	Not CR	Yes	Yes	NE	NE	Died PD D+12
2	43	5	No	No	Sibling	Not CR	Yes	Yes	Yes	Yes	Died GVHD D+217
3	66	5	No	Yes	Unrelated	Not CR	Yes	Yes	Yes	Yes	Died GVHD D+378
4	61	4	No	Yes	Unrelated	Not CR	Yes	No	Yes	NE	Died GVHD D+87
5	57	3	Yes	Yes	Unrelated	Not CR	Yes	No	Yes	Yes	Died GVHD D+159
6	54	3	No	Yes	Unrelated	Not CR	No	Yes	Yes	Yes	Died GVHD D+232
7	49	6	Yes	Yes	Sibling	CR	No	Yes	Yes	No	Died PD D+138
8	49	8	No	Yes	Unrelated	CR	No	Yes	Yes	Yes	Alive CR D+2726
9	64	3	No	Yes	Unrelated	CR	No	Yes	Yes	Yes	Died GVHD D+1714
10	52	3	Yes	No	Sibling	CR	Yes	No	No	Yes	Alive CR D+2230
11	60	7	No	Yes	Unrelated	CR	Yes	Yes	No	No	Alive CR D+2292
12	49	4	Yes	Yes	Unrelated	CR	No	Yes	No	No	Alive CR D+1359
13	61	5	No	Yes	Sibling	Not CR	No	No	Yes	No	Alive CR D+3206
14	54	6	Yes	No	Sibling	Not CR	No	No	Yes	No	Died PD D+160
15	51	4	No	Yes	Sibling	CR	No	No	No	Yes	Alive CR D+2313
16	53	5	Yes	Yes	Sibling	CR	No	No	No	Yes	Alive CR D+2988
17	34	8	Yes	Yes	Sibling	Not CR	No	No	No	No	Alive CR† D+4810
18	49	3	No	Yes	Sibling	Not CR	No	No	No	No	Alive CR‡ D+3207
19	50	5	No	Yes	Sibling	Not CR	No	No	No	No	Died CR§ D+582
20	34	3	No	Yes	Sibling	CR	No	No	No	No	Alive CR D+1728
21	46	7	Yes	Yes	Unrelated	Not CR	No	No	No	No	Alive CR D+1205
22	30	3	No	No	Sibling	CR	No	No	No	Yes	Alive CR D+3611
23	46	4	Yes	No	Sibling	CR	No	No	No	No	Alive CR D+4312

CR, complete remission; D, day; DLI, donor lymphocyte infusion; MI, myocardial infarction; NE, not evaluable; PD, progressive disease.

*Spontaneous acute excluding DLI-induced.

†Progressed D+72 given DLI.

‡Progressed D+85 given DLI.

§Died of MI in CR.

Allotransplantation with FL B cells with dual *TNFRSF14* aberrations resulted in significantly higher frequencies of alloreactive responder CD4⁺ T cells identified by upregulation of the activation marker CD25 when compared with *TNFRSF14* WT FL B cells (median 17% vs 8%, $P = .01$ [primary allotransplantation] and 28% vs 13%, $P = .02$ [secondary allotransplantation]). A smaller and less consistent increase of activated

alloreactive CD8⁺ T cells was seen after allotransplantation with FL B cells with dual aberrations when compared with *TNFRSF14* WT FL B cells with a trend toward significance after secondary allotransplantation (median 10% vs 7%, $P = .08$, Figure 2A-B). The increase in the proportion of activated CD4⁺ T cells after allotransplantation with FL B cells with dual *TNFRSF14* aberrations was not due to an increase




Figure 1. Cell surface expression of HVEM is reduced on FL B cells with *TNFRSF14* aberrations but expression of other molecules involved in antigen presentation is maintained. (A) Cell surface expression of HVEM (filled histograms) and isotype controls (open histograms) on FL B cells. Representative data are shown from patients with *TNFRSF14* WT (left panel), single *TNFRSF14* aberrations (middle panel), and dual *TNFRSF14* aberrations (right panel). (B) Mean (\pm standard deviation) frequencies of HVEM⁺ cells, expressed as a percentage of FL B cells, for WT (n = 5), single aberration (n = 3), and dual *TNFRSF14* aberration cases (n = 5). P value is for 2-tailed unpaired Student *t* test. (C) Surface expression of molecules involved in antigen presentation on FL B cells shown pre- and postactivation. The mean values are shown for 5 WT cases and 5 cases with dual *TNFRSF14* aberrations. Error bars show standard deviation. There were no significant differences in the level of expression of CD58, MHC class I and II, CD80 or CD86 in WT FL B cells and FL B cells with dual *TNFRSF14* aberrations either before or after in vitro activation ($P > .05$ in 2-tailed unpaired Student *t* test). MFI, mean fluorescence intensity; ns, not significant.

in CD4⁺ T cells with a regulatory T-cell phenotype identified either by surface CD25⁺CD127^{lo} expression pattern or by expression of intracellular FOXP3 (Figure 2C-D). These data are consistent with an increase in the frequency of activated alloreactive effector T cells after allostimulation with FL B cells with dual *TNFRSF14* aberrations when compared with stimulation with WT FL B cells.

Frequencies of polyfunctional alloreactive T cells are increased after stimulation with FL B cells with dual *TNFRSF14* aberrations

We next measured the generation of the proinflammatory cytokines IL-2, IFN- γ , and TNF- α in allogeneic coculture supernatants. When analyzed individually, there was a trend toward significantly increased IL-2 and IFN- γ levels in culture supernatants after allostimulation with FL B cells with dual aberrations compared with *TNFRSF14* WT FL B cells. However, allostimulation with FL B cells with dual aberrations resulted in significantly higher combined levels of all 3 proinflammatory cytokines when compared with allostimulation with WT FL B cells (Figure 3A).

We next used intracellular cytokine cytometry to measure allospecific cytokine production at a cellular level. Primary allostimulation with FL B cells with dual aberrations resulted in significantly greater frequencies of CD4⁺ T cells accumulating IL-2 and a trend to greater frequencies of CD8⁺ T cells accumulating IL-2 and IFN- γ . In contrast, allostimulation with FL B cells with dual aberrations did not significantly increase frequencies of T cells accumulating TNF- α consistent with our ELISA data (Figure 3B).

As lytic function of alloreactive T cells is associated with clinically significant acute GVHD,²⁶ we also measured surface CD107a expression on alloreactive T cells stimulated with FL B cells as a surrogate marker of degranulation and lytic capacity.²⁷ We observed significantly higher frequencies of CD4⁺ T cells and a trend toward higher frequencies of CD8⁺ T cells expressing CD107a following primary allostimulation with FL B cells with dual aberrations compared with *TNFRSF14* WT FL B cells (Figure 3C).

Polyfunctional T-cell responses in vitro have been shown to correlate with clinical immunity.²⁸ Therefore, we next used multiparameter flow cytometry combining intracellular measurement of IFN- γ , IL-2, and TNF- α accumulation and surface expression of CD107a




Figure 2. FL B cells with dual *TNFRSF14* aberrations stimulate allogeneic T-cell activation more effectively than WT FL B cells. (A-B) Proportion of CD4⁺ and CD8⁺ T cells expressing the activation marker CD25 after primary (A; n = 7) and secondary (B; n = 5) allogeneic cocultures with WT FL B cells or FL B cells with dual *TNFRSF14* aberrations. Horizontal bars and adjacent numbers represent median values and P values are for 2-tailed paired Student t tests. (C) Proportion of CD4⁺ T cells expressing the CD25^{hi}CD127^{lo} regulatory T-cell surface phenotype or (D) FOXP3 after primary (n = 4) and secondary (n = 4) allogeneic cocultures with WT FL B cells or FL B cells with dual *TNFRSF14* aberrations. Mean values are shown and error bars represent standard deviation. P values are for 2-tailed paired Student t tests.

to assess the frequencies of polyfunctional alloreactive T cells after stimulation with FL B cells. We observed significantly higher frequencies of dual-function T cells expressing CD107a and accumulating IFN- γ , IL-2, or TNF- α after allostimulation with FL B cells with dual aberrations compared with *TNFRSF14* WT FL B cells (Figure 3D). Finally, we used SPICE software to compare the proportion of alloreactive T cells with a polyfunctional phenotype (those positive for 2 or more of surface CD107a, and intracellular IFN- γ , IL-2, and TNF- α) after stimulation with allogeneic FL B cells (Figure 3E). We observed significantly higher frequencies of polyfunctional alloreactive T cells after allostimulation with FL B cells with dual aberrations compared with WT *TNFRSF14* FL B cells (9.1% vs 3.7%, $P = .03$).

FL B cells with dual *TNFRSF14* aberrations stimulate greater allogeneic T-cell proliferation than WT FL B cells

As alloreactive effector T cells must expand in vivo to exert clinically effects, we measured allospecific T-cell proliferation after stimulation with FL B cells. Proliferative responses of allogeneic responder T cells measured by thymidine incorporation were significantly

greater after allostimulation with FL B cells with dual aberrations when compared with *TNFRSF14* WT FL B cells after both primary and secondary allostimulation (Figure 4A). We next used CFSE dye dilution to determine whether this difference was due to a selective effect on proliferation of either CD4⁺ or CD8⁺ responder T cells. The proportion of responder CD4⁺ T cells proliferating after allostimulation with FL B cells with dual aberrations was consistently and significantly greater than after allostimulation with *TNFRSF14* WT FL B cells (median 14% vs 7%, $P = .01$ [primary allostimulation] and 37% vs 10%, $P = .007$ [secondary allostimulation]). Consistent with our data for activation marker expression, we also saw a smaller and less consistent increase in proliferation of allogeneic CD8⁺ T cells after allostimulation with FL B cells with dual aberrations compared with *TNFRSF14* WT FL B cells with a trend toward significance after secondary allostimulation (median 50% vs 40%, $P = .10$, Figure 4B-D).

Collectively, these data demonstrate that the capacity of FL B cells with dual *TNFRSF14* aberrations to stimulate T-cell alloresponses in vitro is both quantitatively and qualitatively enhanced when compared with *TNFRSF14* WT FL B cells.




Figure 3. Frequencies of polyfunctional alloreactive T cells are increased after stimulation with FL B cells with dual *TNFRSF14* aberrations. (A) Proinflammatory cytokines generated from allogeneic T cells after primary (n = 9) and secondary (n = 7) coculture with *TNFRSF14* WT or dual *TNFRSF14* aberration FL B cells. (B) Intracellular cytokine accumulation in allogeneic CD4⁺ and CD8⁺ T cells after primary (n = 8) and secondary (n = 7) coculture with FL B cells. Cytokine-positive cell frequencies are expressed as a percentage of T-cell subsets. (C) Surface CD107a⁺ cell frequencies (expressed as a percentage of T-cell subsets) in allogeneic CD4⁺ and CD8⁺ T cells after primary (n = 8) and secondary (n = 7) coculture with FL B cells. (A–C) Mean values \pm standard deviation are shown. (D) Frequencies of allogeneic bifunctional T cells (CD107a⁺ and cytokine⁺) after primary allogeneic coculture with FL B cells. Horizontal lines depict median values. (E) Multifunctional allogeneic T-cell effectors after primary coculture with FL B cells. Cytokine-secreting populations within the alloreactive CD107a⁺ T-cell compartment are shown. Arcs show frequencies of T cells positive for CD107a, IFN- γ , TNF- α , and IL-2, and slices show effector populations with combined expression patterns of CD107a, IFN- γ , TNF- α , and IL-2. Mean frequencies from 7 independent experiments are shown. *P* values are for 2-tailed Student *t* tests throughout.





Figure 4. FL B cells with dual *TNFRSF14* aberrations stimulate greater allogeneic T-cell proliferation than WT FL B cells. (A) Proliferation of allogeneic T cells measured by thymidine incorporation after primary ($n = 7$) and secondary ($n = 6$) coculture with *TNFRSF14* WT or dual *TNFRSF14* aberration FL B cells. Horizontal lines show median values. (B) Representative histograms of CFSE dye dilution in allogeneic T cells in secondary coculture with FL B cells. Numbers are proportions of proliferating cells. (C-D) Proportion of proliferating allogeneic CD4⁺ and CD8⁺ T cells after primary (C; $n = 7$) and secondary (D; $n = 5$) coculture with FL B cells. Horizontal lines show median values. (E) Proportion of proliferating CD4⁺ and CD8⁺ T cells after primary allogeneic coculture with *TNFRSF14* WT FL B cells without and with exogenous BTLA antagonist or isotype control antibody. (F) Proportion of proliferating CD4⁺ and CD8⁺ T cells after primary allogeneic coculture ($n = 3$) with FL B cells with dual *TNFRSF14* aberrations without and with exogenous BTLA agonist or isotype control antibody. Mean values \pm standard deviation for 3 independent experiments are shown in panels E-F. P values are for 2-tailed Student t tests throughout. cpm, counts per minute.

Figure 5. *TNFRSF14* aberrations in FL are associated with severe acute GVHD after AHSCT. (A) Schematic representation denoting *TNFRSF14* status of purified FL B cells from lymph node biopsies from patients prior to allogeneic transplantation and occurrence of clinically significant acute GVHD (grades 2-4). Red text denotes patients whose acute GVHD was refractory to steroid therapy. (B) Cumulative incidence analysis of acute GVHD (grades 2-4) after allogeneic transplantation in FL patients with relapse/progression of lymphoma and death from other causes as competing risks. *P* value is for the Gray test. (C) Cumulative incidence analysis of death from acute GVHD (grades 2-4) after allogeneic transplantation in FL patients with relapse/progression of lymphoma and death from other causes as competing risks. *P* value is for the Gray test. NE, not evaluable.



BTLA ligation reduces T-cell alloresponses stimulated by FL B cells with *TNFRSF14* aberrations

To provide further evidence that the presence of *TNFRSF14* aberrations increases the capacity of FL B cells to stimulate T-cell alloresponses via a reduction in HVEM-BTLA signaling, we next examined the effect of antibody-mediated modulation of BTLA signaling on B-cell-stimulated T-cell alloresponses in vitro. BTLA blockade increased T-cell alloresponses stimulated with *TNFRSF14* WT FL B cells demonstrating that interaction of HVEM and BTLA regulates the magnitude of T-cell alloresponses stimulated by FL B cells, as has previously been shown with dendritic cell-stimulated T-cell alloresponses.²⁹ Importantly, addition of agonistic BTLA antibody to cocultures of FL B cells with *TNFRSF14* dual aberrations and allogeneic T cells reduced the proportion of alloproliferative CD4⁺ and CD8⁺ T cells (Figure 4E-F). Taken together, these results are consistent with BTLA signaling contributing to control of proliferation of alloreactive T cells after stimulation with *TNFRSF14* WT FL B cells, and reduced BTLA signaling contributing to increased proliferation of alloreactive T cells after stimulation with FL B cells with *TNFRSF14* aberrations.

***TNFRSF14* aberrations are associated with increased acute GVHD in FL patients undergoing AHSCT**

Finally, we sought to assess whether the increased alloantigen-presenting capacity we had observed in FL B cells with *TNFRSF14* aberrations had clinical impact after allogeneic transplantation. We assessed the *TNFRSF14* status in lymph node biopsies performed prior to transplantation in a cohort of patients with FL (Table 1). We observed a similar frequency of *TNFRSF14* aberrations to previously published studies^{4,5} with aberrations detected in 12 of 23 patients (52%). Seven patients had mutations (all within the region encoding the extracellular domain, supplemental Figure 2), 9 had deletions (with 4 patients possessing both a mutation and a deletion), and 1 had a homozygous deletion of both *TNFRSF14* alleles.

To assess the impact of *TNFRSF14* aberrations on clinical alloreactivity after transplantation, we determined the incidence of

acute GVHD in evaluable patients who engrafted. Ten of 22 evaluable patients developed significant acute GVHD (grades 2-4). Using a categorical approach, the only pretransplant risk factor associated with significantly increased incidence of acute GVHD was tumor *TNFRSF14* status (8 of 11 patients with *TNFRSF14* aberrations developed acute GVHD compared with only 2 of 11 *TNFRSF14* WT patients (*P* = .03, Figure 5A). There was no significant difference in the incidence of acute GVHD in patients grouped by age, prior histological transformation, prior rituximab therapy, prior lines of therapy, prior autologous transplantation, remission status pretransplant, or donor type, despite there being a higher frequency of unrelated donors in patients with *TNFRSF14* aberrations (Table 2). We also analyzed the impact of pretransplant factors on acute GVHD using the time-dependent approach of cumulative incidence (CI) with competing risks. Confirming our categorical analysis, *TNFRSF14* status was also the only variable associated with significantly increased CI of acute GVHD (Figure 5B; Table 2).

We took a similar approach to determine the impact of pretransplant factors on the incidence of death from acute GVHD. Death from acute GVHD was limited to patients with *TNFRSF14* aberrations, occurring in 5 of 11 patients compared with 0 of 11 *TNFRSF14* WT patients and *TNFRSF14* status was a statistically significant risk factor for death from

Table 2. Effect of pretransplant variables on incidence of acute GVHD

Pretransplant factor	Incidence of acute GVHD	
	Categorical	Cumulative
	<i>P</i> (FET)	<i>P</i> (Gray)
Recipient <i>TNFRSF14</i> status (any aberration vs none)	.03	.05
Recipient age (above vs below median)	.20	.14
Prior history of histological transformation (yes vs no)	.35	.24
Prior rituximab therapy (yes vs no)	1.00	.89
Prior lines of therapy (above 5 vs not)	1.00	.89
Prior autologous transplant (yes vs no)	.42	.51
Remission status pretransplant (CR vs not CR)	.23	.25
Donor (matched sibling vs matched unrelated)	.19	.15

CR, complete remission; FET, Fisher exact test.

acute GVHD in using both categorical and CI approaches (Figure 5C). However, death from acute GVHD was also significantly associated with pretransplant remission status (supplemental Table 3) and all patients who died of acute GVHD had both *TNFRSF14* aberrations and residual nodal lymphoma at the time of transplantation. Although we cannot separate the effects of *TNFRSF14* aberrations and pretransplant remission status on death from acute GVHD in this small cohort, this observation is consistent with the hypothesis that FL B cells with *TNFRSF14* aberrations increase clinically significant alloresponses that result in acute GVHD after AHSCT.

In contrast to acute GVHD, the incidence of chronic GVHD was not significantly associated with *TNFRSF14* status, or any other pretransplant factor using both categorical and CI approaches (supplemental Table 4). The overall incidence of relapse/progression of lymphoma posttransplant was low in our cohort with only 4 patients relapsing/progressing, precluding an assessment of the impact of *TNFRSF14* status on GVL effects, which needs to be addressed prospectively in a larger cohort of patients.

Discussion

We have shown that aberrations in the *TNFRSF14* gene, which occur in up to half of FL patients, increase the capacity of tumor cells to stimulate clinically significant allogeneic donor T-cell immune responses. These findings have implications for AHSCT strategies for FL patients. The study also has wider significance in demonstrating for the first time that genetic alterations in tumor cells may impact the outcome of allogeneic immunotherapy.

Although we had demonstrated that FL B cells with *TNFRSF14* aberrations stimulate allogeneic T-cell responses more effectively than WT FL B cells in vitro, our finding that this tumor-specific genetic lesion was associated with severe acute GVHD after AHSCT was somewhat unexpected. We anticipated *TNFRSF14* lesions would have little effect on allogeneic T-cell responses that initiate GVHD early posttransplant where alloantigen might be presented predominantly by nonmalignant tissue antigen-presenting cells, but would result in increased alloantigen presentation by small numbers of residual tumor cells resulting in increases GVL effects and less lymphoma relapse posttransplant. The increase in acute GVHD in patients with *TNFRSF14* aberrations may have reflected 2 different mechanisms. FL B cells present in the early posttransplant period could contribute to direct alloantigen presentation, with tumor cells with *TNFRSF14* aberrations priming alloresponses more efficiently than their WT counterparts. This is supported both by our observation that steroid-refractoriness and death from acute GVHD was restricted to patients with *TNFRSF14* aberrations with persistent nodal lymphoma pretransplant and by our in vitro experiments. Alternatively, *TNFRSF14* aberrations in lymphoma cells might modulate the recipient microenvironment, establishing proinflammatory conditions which increase alloantigen presentation by nonmalignant host antigen-presenting cells thereby increasing donor T-cell alloresponses. Because reduced BTLA ligation on dendritic cells and tissue macrophages increases proinflammatory cytokine release,^{30,31} this mechanism could further amplify alloresponses in patients with *TNFRSF14* aberrations to increase severe acute GVHD.

It is likely that increased T-cell alloresponses stimulated by lymphoma cells with *TNFRSF14* aberrations occurred predominantly as a result of reduced ligation of BTLA on donor T cells. FL B cells with *TNFRSF14* aberrations induced greater alloproliferation in CD4⁺ T cells than in CD8⁺ T cells in vitro, consistent with retention of high levels of

BTLA expression on both naive and memory CD4⁺ T cells, whereas BTLA expression is progressively lost with CD8⁺ T-cell differentiation.²⁹ Although HVEM also ligates a second coinhibitory T-cell receptor, CD160, it is unlikely that this receptor plays a significant role in this setting, as CD160 is expressed at very low levels on human CD4⁺ T cells in the tumor microenvironment.³²

However, we need to take into account the limitations of our study. The functional in vitro experiments were restricted to FL B cells with dual rather than single *TNFRSF14* aberrations. Although further studies are required to accurately define comparative effects of single *TNFRSF14* lesions, and the differential impact of specific mutations, our clinical data suggest single *TNFRSF14* lesions may also be clinically relevant in this setting as severe acute GVHD was observed in both patients with single and dual *TNFRSF14* aberrations. Additionally, other potentially immune-modulating genetic lesions have been identified in lymphoma, including mutations in genes encoding β 2-microglobulin and CD58 which could result in reduction of antigen-presenting capacity.³³ Although our present study does not directly examine such mutations, it is unlikely they impacted on our results as we demonstrated similar expression of HLA class I and CD58 on FL B cells with and without *TNFRSF14* aberrations. Finally, in the current study, we measured the effect of *TNFRSF14* aberrations on HLA-mismatched alloresponses which are driven predominantly by direct alloantigen presentation by recipient rather than donor antigen-presenting cells. BTLA ligation is known to also limit human minor histocompatibility antigen (mHAg)-specific T-cell responses that mediate acute GVHD in the HLA-matched setting³⁴ and our clinical data are consistent with *TNFRSF14* aberrations impacting on the capacity of FL B cells to present mHags leading to increased acute GVHD after HLA-matched AHSCT. Although we have not extended our functional assays to measure mHAg-specific HLA-matched alloresponses, this will be a focus of future work. The impact of *TNFRSF14* aberrations in FL on alloreactivity in other immunotherapy platforms such as T-depleted approaches or cord blood transplants will need to be further explored.

Given our findings in the allogeneic setting, *TNFRSF14* aberrations which reduce expression of HVEM on FL B cells might also be expected to potentiate autologous T-cell antitumor immunity via reduced BTLA-mediated suppression. However, as previous studies suggest that *TNFRSF14* aberrations may confer a poor prognosis in patients treated with chemoimmunotherapy,⁴ it is likely that additional or alternative HVEM-mediated mechanisms impact on FL B-cell survival in the autologous setting. As BTLA is also expressed on B cells, and negatively regulates B-lymphocyte transformation and malignant B-cell survival,³⁵⁻³⁷ reduction of HVEM ligation of BTLA on FL B cells either in *cis* or *trans* (from neighboring tumor cells) could potentiate tumor growth and survival in this setting.

In summary, these data support the development of risk-adapted allogeneic transplant strategies in FL patients with *TNFRSF14* aberrations to reduce harmful acute GVHD, which could include pretransplant purging of recipient B cells or augmented posttransplant immunosuppression. Genetic lesions in tumor cells that modulate allogeneic immune responses may have wider significance, impacting the outcome of allogeneic cellular immunotherapy not only for lymphoma but also for other hematologic cancers.

Acknowledgments

This work was supported by grants from the Kay Kendall Leukaemia Fund (Junior Clinical Research Fellowship 557 [J.O.]), Cancer Research UK (Programme Grant 15968 [J.F.]) and Centre Grant

C16420/A18066), the Medical Research Council (Clinician Scientist Fellowship G0902269 [J.K.D.]), and Barts Cancer Institute Bridge to the Future Fellowship (E.K.).

Authorship

Contribution: E.K. designed and performed experiments, analyzed data, and wrote the manuscript; J.O. performed genetic analysis;

C.B. performed experiments; S.I. collated patient samples; J.M. provided clinical data; J.F. contributed to the genetic analysis and critically reviewed the manuscript; J.G.G. contributed to experimental design and wrote the manuscript; and J.K.D. conceptualized the study, designed and analyzed experiments and clinical data, and wrote the manuscript.

Conflict-of-interest disclosure: The authors declare no competing financial interests.

Correspondence: Jeffrey K. Davies, Centre for Haemato-oncology, Barts Cancer Institute, Charterhouse Square, London EC1M 6BQ, United Kingdom; e-mail: j.k.davies@qmul.ac.uk.

References

- Czuczman MS, Grillo-López AJ, White CA, et al. Treatment of patients with low-grade B-cell lymphoma with the combination of chimeric anti-CD20 monoclonal antibody and CHOP chemotherapy. *J Clin Oncol*. 1999;17(1):268-276.
- Vose JM, Link BK, Grossbard ML, Czuczman M, Grillo-Lopez A, Fisher RI. Long-term update of a phase II study of rituximab in combination with CHOP chemotherapy in patients with previously untreated, aggressive non-Hodgkin's lymphoma. *Leuk Lymphoma*. 2005;46(11):1569-1573.
- Piñana JL, Martino R, Gayoso J, et al; GELTAMO Group. Reduced intensity conditioning HLA identical sibling donor allogeneic stem cell transplantation for patients with follicular lymphoma: long-term follow-up from two prospective multicenter trials. *Haematologica*. 2010;95(7):1176-1182.
- Cheung KJ, Johnson NA, Affleck JG, et al. Acquired TNFRSF14 mutations in follicular lymphoma are associated with worse prognosis. *Cancer Res*. 2010;70(22):9166-9174.
- Launay E, Pangault C, Bertrand P, et al. High rate of TNFRSF14 gene alterations related to 1p36 region in de novo follicular lymphoma and impact on prognosis. *Leukemia*. 2012;26(3):559-562.
- Shui JW, Steinberg MW, Kronenberg M. Regulation of inflammation, autoimmunity, and infection immunity by HVEM-BTLA signaling. *J Leukoc Biol*. 2011;89(4):517-523.
- Krieg C, Boyman O, Fu YX, Kaye J. B and T lymphocyte attenuator regulates CD8+ T cell-intrinsic homeostasis and memory cell generation. *Nat Immunol*. 2007;8(2):162-171.
- Distler E, Bloetz A, Albrecht J, et al. Alloreactive and leukemia-reactive T cells are preferentially derived from naive precursors in healthy donors: implications for immunotherapy with memory T cells. *Haematologica*. 2011;96(7):1024-1032.
- Chérel M, Choufi B, Trauet J, et al. Naive subset develops the most important alloreactive response among human CD4+ T lymphocytes in human leukocyte antigen-identical related setting. *Eur J Haematol*. 2014;92(6):491-496.
- Albring JC, Sandau MM, Rapaport AS, et al. Targeting of B and T lymphocyte associated (BTLA) prevents graft-versus-host disease without global immunosuppression. *J Exp Med*. 2010;207(12):2551-2559.
- Schultz JL, Cardoso AA, Freeman GJ, et al. Follicular lymphomas can be induced to present alloantigen efficiently: a conceptual model to improve their tumor immunogenicity. *Proc Natl Acad Sci USA*. 1995;92(18):8200-8204.
- Okosun J, Bődör C, Wang J, et al. Integrated genomic analysis identifies recurrent mutations and evolution patterns driving the initiation and progression of follicular lymphoma. *Nat Genet*. 2014;46(2):176-181.
- Jiang W, Lederman MM, Harding CV, Rodriguez B, Mohner RJ, Sieg SF. TLR9 stimulation drives naïve B cells to proliferate and to attain enhanced antigen presenting function. *Eur J Immunol*. 2007;37(8):2205-2213.
- Ruprecht CR, Lanzavecchia A. Toll-like receptor stimulation as a third signal required for activation of human naïve B cells. *Eur J Immunol*. 2006;36(4):810-816.
- Lyons AB. Divided we stand: tracking cell proliferation with carboxyfluorescein diacetate succinimidyl ester. *Immunol Cell Biol*. 1999;77(6):509-515.
- Davies JK, Yuk D, Nadler LM, Guinan EC. Induction of alloantigeny in human donor T cells without loss of pathogen or tumor immunity [published correction appears in *Transplantation*. 2012;93(11):e44]. *Transplantation*. 2008;86(6):854-864.
- Auer RL, MacDougall F, Oakervee HE, et al. T-cell replete fludarabine/cyclophosphamide reduced intensity allogeneic stem cell transplantation for lymphoid malignancies. *Br J Haematol*. 2012;157(5):580-585.
- Cheson BD, Horning SJ, Coiffier B, et al; NCI Sponsored International Working Group. Report of an international workshop to standardize response criteria for non-Hodgkin's lymphomas. *J Clin Oncol*. 1999;17(4):1244.
- Cheson BD, Pfistner B, Juweid ME, et al; International Harmonization Project on Lymphoma. Revised response criteria for malignant lymphoma. *J Clin Oncol*. 2007;25(5):579-586.
- Filipovich AH, Weisdorf D, Pavletic S, et al. National Institutes of Health consensus development project on criteria for clinical trials in chronic graft-versus-host disease: I. Diagnosis and staging working group report. *Biol Blood Marrow Transplant*. 2005;11(12):945-956.
- Przepiorka D, Weisdorf D, Martin P, et al. 1994 Consensus Conference on Acute GVHD Grading. *Bone Marrow Transplant*. 1995;15(6):825-828.
- Deeg HJ. How I treat refractory acute GVHD. *Blood*. 2007;109(10):4119-4126.
- Fine JG, Gray RJ. A proportional hazards model for the subdistribution of a competing risk. *J Am Stat Assoc*. 1999;94:496-509.
- Kaplan E. Non-parametric estimation from incomplete observations. *J Am Stat Assoc*. 1958;53:457-481.
- Peto R, Pike MC, Armitage P, et al. Design and analysis of randomized clinical trials requiring prolonged observation of each patient. II. Analysis and examples. *Br J Cancer*. 1977;35(1):1-39.
- Schmaltz C, Alpdogan O, Horndasch KJ, et al. Differential use of Fas ligand and perforin cytotoxic pathways by donor T cells in graft-versus-host disease and graft-versus-leukemia effect. *Blood*. 2001;97(9):2886-2895.
- Betts MR, Koup RA. Detection of T-cell degranulation: CD107a and b. *Methods Cell Biol*. 2004;75:497-512.
- Yuan J, Gnjatich S, Li H, et al. CTLA-4 blockade enhances polyfunctional NY-ESO-1 specific T cell responses in metastatic melanoma patients with clinical benefit. *Proc Natl Acad Sci USA*. 2008;105(51):20410-20415.
- Serriari NE, Gondois-Rey F, Guillaume Y, et al. B and T lymphocyte attenuator is highly expressed on CMV-specific T cells during infection and regulates their function. *J Immunol*. 2010;185(6):3140-3148.
- Bjordahl RL, Steidl C, Gascoyne RD, Ware CF. Lymphotoxin network pathways shape the tumor microenvironment. *Curr Opin Immunol*. 2013;25(2):222-229.
- Kobayashi Y, Iwata A, Suzuki K, et al. B and T lymphocyte attenuator inhibits LPS-induced endotoxic shock by suppressing Toll-like receptor 4 signaling in innate immune cells. *Proc Natl Acad Sci USA*. 2013;110(13):5121-5126.
- Gertner-Dardenne J, Fauriat C, Orlanducci F, et al. The co-receptor BTLA negatively regulates human Vγ9Vδ2 T-cell proliferation: a potential way of immune escape for lymphoma cells. *Blood*. 2013;122(6):922-931.
- Challa-Malladi M, Lieu YK, Califano O, et al. Combined genetic inactivation of β2-microglobulin and CD58 reveals frequent escape from immune recognition in diffuse large B cell lymphoma. *Cancer Cell*. 2011;20(6):728-740.
- Hobo W, Norde WJ, Schaap N, et al. B and T lymphocyte attenuator mediates inhibition of tumor-reactive CD8+ T cells in patients after allogeneic stem cell transplantation. *J Immunol*. 2012;189(1):39-49.
- Vendel AC, Calemine-Fenaux J, Izrael-Tomasevic A, Chauhan V, Arnott D, Eaton DL. B and T lymphocyte attenuator regulates B cell receptor signaling by targeting Syk and BLNK. *J Immunol*. 2009;182(3):1509-1517.
- Gururajan M, Jennings CD, Bondada S. Cutting edge: constitutive B cell receptor signaling is critical for basal growth of B lymphoma. *J Immunol*. 2006;176(10):5715-5719.
- Gupta N, Delrow J, Drawid A, Sengupta AM, Fan G, Gélinas C. Repression of B-cell linker (BLNK) and B-cell adaptor for phosphoinositide 3-kinase (BCAP) is important for lymphocyte transformation by rel proteins. *Cancer Res*. 2008;68(3):808-814.