

*Conflict-of-interest disclosure:* The author declares no competing financial interests. ■

## REFERENCES

1. Payne H, Ponomaryov T, Watson SP, Brill A. Mice with a deficiency in CLEC-2 are protected against deep vein thrombosis. *Blood*. 2017;129(14):2013-2020.
2. Suzuki-Inoue K, Osada M, Ozaki Y. Physiological and pathophysiological roles of interaction between CLEC-2 and podoplanin: partners from in utero to adulthood. *J Thromb Haemost*. 2017;15(2):219-229.
3. Suzuki-Inoue K, Inoue O, Ding G, et al. Essential in vivo roles of the C-type lectin receptor CLEC-2: embryonic/neonatal lethality of CLEC-2-deficient mice by blood/lymphatic misconnections and impaired thrombus formation of CLEC-2-deficient platelets. *J Biol Chem*. 2010;285(32):24494-24507.
4. Hughes CE, Navarro-Núñez L, Finney BA, Mourão-Sá D, Pollitt AY, Watson SP. CLEC-2 is not required for platelet aggregation at arteriolar shear. *J Thromb Haemost*. 2010;8(10):2328-2332.
5. Bender M, May F, Lorenz V, et al. Combined in vivo depletion of glycoprotein VI and C-type lectin-like receptor 2 severely compromises hemostasis and abrogates

arterial thrombosis in mice. *Arterioscler Thromb Vasc Biol*. 2013;33(5):926-934.

6. Inoue O, Hokamura K, Shirai T, et al. Vascular smooth muscle cells stimulate platelets and facilitate thrombus formation through platelet CLEC-2: implications in atherothrombosis. *PLoS One*. 2015;10(9):e0139357.
7. Boulaftali Y, Hess PR, Getz TM, et al. Platelet ITAM signaling is critical for vascular integrity in inflammation. *J Clin Invest*. 2013;123(2):908-916.
8. Herzog BH, Fu J, Wilson SJ, et al. Podoplanin maintains high endothelial venule integrity by interacting with platelet CLEC-2. *Nature*. 2013;502(7469):105-109.
9. Hitchcock JR, Cook CN, Bobat S, et al. Inflammation drives thrombosis after Salmonella infection via CLEC-2 on platelets. *J Clin Invest*. 2015;125(12):4429-4446.
10. Shirai T, Inoue O, Tamura S, et al. CLEC-2 promotes hematogenous tumor metastasis and prothrombotic state in tumor-bearing mice. *J Thromb Haemost*. 2017;15(3):513-525.

DOI 10.1182/blood-2017-02-764670

© 2017 by The American Society of Hematology

## ● ● ● TRANSPLANTATION

Comment on Riesner et al, page 2021

# Vessels, T cells, and GVHD: time matters

Sebastian P. Haen EBERHARD KARLS UNIVERSITÄT TÜBINGEN

In this issue of *Blood*, Riesner et al describe that neovascularization precedes infiltration of inflammatory cells in graft-versus-host disease (GVHD) and colitis models, revealing a new pathophysiological sequence of disease initiation.<sup>1</sup>


In Greek philosophy, the term *kairos* (καῖρός) defines the perfect time point. With regard to therapeutic strategies and decisions in medicine, the determination of the *kairos* is equally important as the identification of major mechanisms and pathways leading to substantiated understanding of (patho)physiological processes and to identification of potential therapeutic targets. In GVHD, development of novel treatment strategies is urgently needed because it remains the leading diagnostic and therapeutic challenge following allogeneic hematopoietic stem cell transplantation with the consequence of severe morbidity and nonrelapse mortality, especially when nonresponsive to high-dose steroid therapy.<sup>2</sup> In recent years, evidence has emerged that the vascular endothelium could be a target structure for GVHD therapy because an association of the formation of new blood vessels and GVHD has been observed.<sup>3</sup> Per se,

the important role of neovascularization has long been established in GVHD, but the chronological sequence of the establishment of new vasculature and immune cell infiltration has remained unclear.<sup>3</sup> Although it has been previously believed that angiogenesis would be stimulated by secreted mediators from tissue-infiltrating immune cells,<sup>4</sup> Riesner and coworkers observed that the contrary is the case, showing that angiogenesis precedes the infiltration of immune cells into damaged tissues. Importantly, after incipient tissue damage, angiogenesis occurred as early as 2 days after allogeneic hematopoietic stem cell transplantation, indicating that the pathogenesis of GVHD is already commenced early with novel vasculature to tackle a route for immune cells. Their findings suggest a new chronological sequence of GVHD initiation (tissue damage, neovascularization, immune cell infiltration). These observations might help to improve our understanding of the

underlying mechanisms of GVHD. Hence, this new evidence might help explain why GVHD primarily affects tissues that are more susceptible to infectious agents, why GVHD mainly manifests in distinct organs, and how GVHD-mediating T cells are primed and activated, shedding new light on immunobiology in general and GVHD T-cell biology in particular (see figure).

GVHD is believed to be mainly mediated by allogeneic T cells that recognize differences in tissue antigens, which are expressed on recipient cells. These are most importantly major histocompatibility complexes (MHCs, in humans termed HLAs) as a complex with their respective bound peptides.<sup>5</sup> This direct recognition of peptide MHC is avoided by MHC-matched (or HLA-identical) transplantation. However, about 40% of patients who receive HLA-identical grafts will still develop GVHD, which can then be mediated by recognition of minor histocompatibility antigens.<sup>6</sup> These constitute MHC-presented peptides representing protein fragments which are produced during normal cellular metabolism and exhibit amino acid sequence variants based on single-nucleotide polymorphisms between donor and recipient. Further recognition of peptides can also be based on specific sequence variants based on intrinsic alterations in leukemic cells, then mediating graft-versus-leukemia (GVL) effects.<sup>7</sup> Of note, sole recognition of such antigen differences cannot be the only mechanism mediating GVHD or GVL because it has been observed that new autoimmunity (diseases sharing features with “naturally occurring” autoimmune diseases)<sup>8</sup> and graft versus self (GVS; recognition of MHC-restricted autoepitopes)<sup>9</sup> can develop, constituting that further processes must be involved in the genesis of graft-host interaction.

Based on the findings by Riesner et al, neovascularization could hereby be an important prerequisite for priming of naive T cells or for proliferation of tissue-specific memory T cells against alloantigens, autoantigens, and leukemia (or tumor) antigens. This is further underlined by the observation that endothelial cells can also upregulate MHC molecules and function as antigen-presenting cells to naive T cells in inflamed tissue.<sup>10</sup> The data of Riesner and colleagues might in this context also support a further comprehension of in situ T-cell priming. Importantly, they used a



Pathomechanism of GVHD including neovascularization. APC, antigen-presenting cell; CTL, cytotoxic T lymphocyte; IFN, interferon; IL, interleukin; LPS, lipopolysaccharide; Mφ, macrophage; Th1, T helper 1 cell; TNF, tumor necrosis factor. Modified and reprinted with permission from Elsevier (*The Lancet*, 2009, 373, 1550-1561).<sup>6</sup>

chemotherapy-based MHC-matched system in chronological dependency, in contrast to previously used mouse models with MHC mismatch that were designed to induce maximal GVHD. This model could also be used for further studies of GVHD initiation with regard to danger signals, immune-regulatory molecules, and pathogen-associated molecular patterns more similar to transplantation in a human setting. This model could also guide the timing of future therapeutic approaches, as for prevention of T-cell priming by antiangiogenic therapy, the *καιρός* for therapy initiation also has to be early. Naturally, a mismatch model could not mimic development of new autoimmunity or GVS.

In summary, the data of Riesner et al provide promising novel insights into the chronological development of inflammation in general and GVHD in particular. They show that neovascularization paves the way for infiltration of immune cells into damaged tissue and occurs very quickly after allogeneic hematopoietic stem cell transplantation, indicating that priming and antigen exposition of GVHD-mediating T cells also must be an early event in the cascade of GVHD development. These data will therefore contribute to better comprehension of inflammation initiation and T-cell priming in GVHD. As discussed extensively in their

article, the translation of these observations into the clinic in order to improve therapeutic options for GVHD patients, without hampering GVL effects, requires further preclinical evaluation in animal models in order to identify the proper substances and the right *καιρός*.

*Conflict-of-interest disclosure:* The author declares no competing financial interests. ■

## ● ● ● TRANSPLANTATION

Comment on Hiwarkar et al, page 2033

# Guarded promise for adenovirus therapy

John R. Wingard UNIVERSITY OF FLORIDA COLLEGE OF MEDICINE

Adenovirus can cause deadly infection after hematopoietic cell transplantation (HCT). In this issue of *Blood*, Hiwarkar et al present data in pediatric HCT patients with adenoviremia indicating that brincidofovir, a prodrug of cidofovir, was well tolerated, demonstrated substantially more rapid virologic clearance rates compared with cidofovir, was effective even after cidofovir failure, and was efficacious even in the absence of improved lymphopenia.<sup>1</sup> Heretofore, we have lacked effective tools to prevent this life-threatening threat.

**C**idofovir is a deoxycytidine analog that has antiviral activity against the families of double-stranded DNA viruses that are human pathogens, including adenovirus. It has clinical

## REFERENCES

- Riesner K, Shi Y, Jacobi A, et al. Initiation of acute graft-versus-host disease by angiogenesis. *Blood*. 2017;129(14):2021-2032.
- Martin PJ, Inamoto Y, Flowers ME, Carpenter PA. Secondary treatment of acute graft-versus-host disease: a critical review. *Biol Blood Marrow Transplant*. 2012;18(7):982-988.
- Penack O, Socić G, van den Brink MR. The importance of neovascularization and its inhibition for allogeneic hematopoietic stem cell transplantation. *Blood*. 2011;117(16):4181-4189.
- Szekanecz Z, Koch AE. Mechanisms of disease: angiogenesis in inflammatory diseases. *Nat Clin Pract Rheumatol*. 2007;3(11):635-643.
- Felix NJ, Donermeyer DL, Horvath S, et al. Alloreactive T cells respond specifically to multiple distinct peptide-MHC complexes. *Nat Immunol*. 2007;8(4):388-397.
- Ferrara JL, Levine JE, Reddy P, Holler E. Graft-versus-host disease. *Lancet*. 2009;373(9674):1550-1561.
- Wölfel C, Lennerz V, Lindemann E, et al. Dissection and molecular analysis of alloreactive CD8+ T cell responses in allogeneic haematopoietic stem cell transplantation. *Cancer Immunol Immunother*. 2008;57(6):849-857.
- Daikeler T, Labopin M, Di Gioia M, et al; EBMT Autoimmune Disease Working Party. Secondary autoimmune diseases occurring after HSCT for an autoimmune disease: a retrospective study of the EBMT Autoimmune Disease Working Party. *Blood*. 2011;118(6):1693-1698.
- Mirza N, Zierhut M, Korn A, et al. Graft versus self (GvS) against T-cell autoantigens is a mechanism of graft-host interaction. *Proc Natl Acad Sci USA*. 2016;113(48):13827-13832.
- Turesson C. Endothelial expression of MHC class II molecules in autoimmune disease. *Curr Pharm Des*. 2004;10(2):129-143.

DOI 10.1182/blood-2017-02-764696

© 2017 by The American Society of Hematology