

study of the platelet inhibitor prasugrel in children with sickle cell disease was designed to identify drug doses that inhibit platelet function between 30% and 50%, a level thought to balance safety and efficacy,²⁴ similar to the level of inhibition observed here, although the dual effects of RN-1 on platelet function and platelet counts could pose an additional risk for bleeding that will require further monitoring.

Our results show that administration of RN-1 to normal baboons over a prolonged period increases HbF, F cells, and F retics and is generally well tolerated, supporting further development of LSD1 inhibitors as therapeutic agents for SCD. Because LSD1 also has an important functional role in neural stem cell maintenance and proliferation, effects of LSD1 inhibitors on the brain and nervous system should be carefully evaluated.²⁵

*V.I. and K.V. contributed equally to this work.

Acknowledgment: This work was supported by National Institutes of Health, National Heart, Lung, and Blood Institute grant U01 HL117658.

Contribution: V.I. and K.V. performed research and collected, analyzed, and interpreted the data; A.R. and R.M. analyzed data and reviewed the manuscript; S.C. and J.D.E. reviewed the manuscript; J.D. analyzed data and reviewed the manuscript; D.L. designed and performed the research, analyzed the data, and wrote the manuscript.

Conflict-of-interest disclosure: The authors declare no competing financial interests.

ORCID profiles: D.L., 0000-0002-0275-7168.

Correspondence: Donald Lavelle, Jesse Brown Veterans Affairs Medical Center, Room 6215, 820 S Damen Ave, Chicago, IL 60612; e-mail: dlavelle@uic.edu.

References

- Platt OS, Thorington BD, Brambilla DJ, et al. Pain in sickle cell disease. Rates and risk factors. *N Engl J Med*. 1991;325(1):11-16.
- Platt OS, Brambilla DJ, Rosse WF, et al. Mortality in sickle cell disease. Life expectancy and risk factors for early death. *N Engl J Med*. 1994;330(23):1639-1644.
- Schechter AN. Hemoglobin research and the origins of molecular medicine. *Blood*. 2008;112(10):3927-3938.
- DeSimone J, Mueller AL. Fetal hemoglobin synthesis in baboons (*Papio cynocephalus*). *J Lab Clin Med*. 1978;91(6):862-871.
- Tagle DA, Koop BF, Goodman M, Slightom JL, Hess DL, Jones RT. Embryonic epsilon and gamma globin genes of a prosimian primate (*Galago crassicaudatus*). Nucleotide and amino acid sequences, developmental regulation and phylogenetic footprints. *J Mol Biol*. 1988;203(2):439-455.
- DeSimone J, Heller P, Hall L, Zwiars D. 5-Azacytidine stimulates fetal hemoglobin synthesis in anemic baboons. *Proc Natl Acad Sci USA*. 1982;79(14):4428-4431.
- Ley TJ, DeSimone J, Noguchi CT, et al. 5-Azacytidine increases gamma-globin synthesis and reduces the proportion of dense cells in patients with sickle cell anemia. *Blood*. 1983;62(2):370-380.
- Ley TJ, DeSimone J, Anagnou NP, et al. 5-azacytidine selectively increases gamma-globin synthesis in a patient with beta+ thalassemia. *N Engl J Med*. 1982;307(24):1469-1475.
- Koshy M, Dorn L, Bressler L, et al. 2-deoxy 5-azacytidine and fetal hemoglobin induction in sickle cell anemia. *Blood*. 2001;96(7):2379-2384.
- DeSimone J, Koshy M, Dorn L, et al. Maintenance of elevated fetal hemoglobin levels by decitabine during dose interval treatment of sickle cell anemia. *Blood*. 2002;99(11):3905-3908.
- Saunthararajah Y, Hillery CA, Lavelle D, et al. Effects of 5-aza-2'-deoxycytidine on fetal hemoglobin levels, red cell adhesion, and hematopoietic differentiation in patients with sickle cell disease. *Blood*. 2003;102(12):3865-3870.
- Lavelle D, Vaitkus K, Ling Y, et al. Effects of tetrahydropyridine on pharmacokinetics and pharmacodynamics of oral decitabine. *Blood*. 2012;119(5):1240-1247.
- Lavelle D, Gowhari M, Pacini M, et al. Combination with THU to address pharmacologic limitations of decitabine, interim PK/PD from a phase 1/2 clinical trial of oral THU-decitabine in sickle cell disease [abstract]. *Blood*. 2014; 124(21). Abstract 90.
- Cui S, Kolodziej KE, Obara N, et al. Nuclear receptors TR2 and TR4 recruit multiple epigenetic transcriptional corepressors that associate specifically with the embryonic beta-type globin promoters in differentiated adult erythroid cells. *Mol Cell Biol*. 2011;31(16):3298-3311.
- Xu J, Bauer DE, Kerényi MA, et al. Corepressor-dependent silencing of fetal hemoglobin expression by BCL11A. *Proc Natl Acad Sci USA*. 2013;110(16): 6518-6523.
- Shi L, Cui S, Engel JD, Tanabe O. Lysine-specific demethylase 1 is a therapeutic target for fetal hemoglobin induction. *Nat Med*. 2013;19(3):291-294.
- Rivers A, Vaitkus K, Ruiz MA, et al. RN-1, a potent and selective lysine-specific demethylase 1 inhibitor, increases gamma-globin expression, F reticulocytes, and F cells in a sickle cell disease mouse model. *Exp Hematol*. 2015;43(7):546-553.
- Cui S, Lim KC, Shi L, et al. The LSD1 inhibitor RN-1 induces fetal hemoglobin synthesis and reduces disease pathology in sickle cell mice. *Blood*. 2015; 126(3):386-396.
- Rivers A, Vaitkus K, Ibanez V, et al. The LSD1 inhibitor RN-1 recapitulates the fetal pattern of hemoglobin synthesis in baboons (*P. anubis*). *Haematologica*. 2016;101(6):688-697.
- Hack CA, Gleiser CA. Hematologic and serum chemical reference values for adult and juvenile baboons (*Papio* sp). *Lab Anim Sci*. 1982;32(5):502-505.
- Goodall AH, Appleby J. Flow-cytometric analysis of platelet-membrane glycoprotein expression and platelet activation. *Methods Mol Biol*. 2004;272(1): 225-253.
- Saleque S, Kim J, Rooke HM, Orkin SH. Epigenetic regulation of hematopoietic differentiation by Gfi-1 and Gfi-1b is mediated by the cofactors CoREST and LSD1. *Mol Cell*. 2007;27(4):562-572.
- Sprüssel A, Schulte JH, Weber S, et al. Lysine-specific demethylase 1 restricts hematopoietic progenitor proliferation and is essential for terminal differentiation. *Leukemia*. 2012;26(9):2039-2051.
- Styles L, Heiselman D, Heath LE, et al. Prasugrel in children with sickle cell disease: pharmacokinetic and pharmacodynamic data from an open-label, adaptive-design, dose-ranging study. *J Pediatr Hematol Oncol*. 2015;37(1):1-9.
- Maes T, Mascaró C, Ortega A, et al. KDM1 histone lysine demethylases as targets for treatments of oncological and neurodegenerative disease. *Epigenomics*. 2015;7(4):609-626.

DOI 10.1182/blood-2016-10-746727

To the editor:

Zika virus in asymptomatic blood donors in Martinique

Pierre Gallian,^{1,2} André Cabié,^{3,4} Pascale Richard,⁵ Laure Paturel,⁵ Rémi N. Charrel,¹ Boris Pastorino,¹ Isabelle Leparç-Goffart,^{1,6} Pierre Tiberghien,^{2,7} and Xavier de Lamballerie¹

¹Aix-Marseille Université, French Institute of Research for Development, INSERM, Institut Hospitalo-Universitaire Méditerranée Infection, Unité Mixte de Recherche_D190 "Emergence des Pathologies Virales," Marseille, France; ²Etablissement Français du Sang, La Plaine Saint-Denis, France; ³Service de maladies infectieuses et tropicales, Centre Hospitalo-Universitaire de Fort-de-France, Fort-de-France, Martinique; ⁴Université des Antilles, Centre d'Investigation Clinique 1424, INSERM, Equipe d'Accueil 4537, Fort-de-France, Martinique; ⁵Etablissement Français du Sang, Fort-de-France, Martinique; ⁶National Reference Laboratory for Arboviruses, Institut de Recherche Biomédicale des Armées, Marseille, France; and ⁷Université de Franche-Comté, INSERM, Etablissement Français du Sang UMR 1098, Besançon, France

Zika fever is an *Aedes*-borne disease caused by a flavivirus (Zika virus [ZIKV], genus *Flavivirus*). During the past years, ZIKV has spread in Polynesia, South America, and the Caribbean.¹ Most ZIKV infections are asymptomatic or result in mild febrile disease with rash and conjunctivitis. Attention was recently drawn to nonvectored ZIKV transmission,² including sexual,^{3,4} probable blood-borne,⁵ and mother-to-fetus transmission, and to severe forms such as Guillain-Barré syndrome,^{6,7} acute neurological infections,^{8,9} and fetal abnormalities¹⁰ (including microcephaly).

Many aspects of Zika fever natural history remain unknown (eg, the proportion of asymptomatic cases and the duration of viremia). Estimating the prevalence of Zika infections is difficult because a large proportion of infected individuals do not seek medical attention, and seroprevalence studies are hampered by antigenic cross-reactivity with other flaviviruses (eg, dengue virus).¹¹ Here, we present a study of ZIKV infection in blood donors from Martinique island (French West Indies, Caribbean region), with novel epidemiological, biological, and clinical information that refines the picture of Zika fever in adults.

After the ZIKV outbreak in 2013 to 2014 in French Polynesia and the evidence that a significant proportion (3%) of donors were viremic,¹² ZIKV has been recognized in Martinique since the end of 2015, leading to implementation by the French Blood Bank of a systematic individual nucleic acid testing (NAT) of blood donations. Virus detection was performed on plasma samples, using a semi-automated platform consisting of a Microlab-STARlet (Hamilton, Bonaduz, Switzerland) and the NucleoSpin 96 Virus Extraction Kit (Macherey-Nagel, Duren, Germany) for nucleic acids extraction, and a CFX96 thermocycler (Bio-Rad Laboratories, California) and the RealStar Zika Virus RT-PCR Kit_1.1 (Altona Diagnostics, Hamburg, Germany) for reverse transcriptase polymerase chain reaction testing.

Between January 19 and June 10, 2016, 4129 consecutive blood donations were tested (mean age, 41.9 years; sex ratio [M/F], 0.88). Positive NAT detection occurred in 76 blood donations (1.84%), with the most intense detection rate (3%) during weeks 17-20 (mean age, 41.8 years; sex ratio, 1.2). Postdonation inquiry consisted of a telephone call at day 7 postdonation to identify symptoms compatible with ZIKV infection. When the donor declared no sign, a new call was organized 14 days after donation. This information was obtained from 75 viremic donors: 34 (45.3%) remained asymptomatic, and 41 (54.7%) reported symptoms (1-6 days postdonation; Figure 1A) such as fever, conjunctivitis, myalgia, arthralgia, and rash (Figure 1B). There was a trend for higher values of molecular viral load in symptomatic vs asymptomatic donors (mean values, 5.36 vs 4.93 log₁₀ RNA genomic equivalents per milliliter; independent Wilcoxon test, *P* value = .0013; Figure 1C). This difference does not imply that symptomatic donors reached higher viremia than asymptomatic donors: sampling could occur in the former only during the early steps of viremia, whereas in the latter, it could also occur during the decreasing phase of viremia. The range of viral loads (2.09-6.50 log₁₀ RNA copies/mL) was comparable to that previously described in French Polynesian asymptomatic blood donors.¹³ Men declared symptoms less frequently than women (45.2% vs 66.7%; χ^2 test *P* value = .06), but viral load and time to declaration of symptoms were not significantly different in men and women. No significant association of viremia with ABO, Rhesus, and Kell blood groups was detected.

We performed seroprevalence analyses at 2 time points: in 418 donors sampled in early March (March 9-23), and in 176 donors sampled in early June (June 6-13). Samples were tested using an anti-Zika virus NS1 immunoglobulin G enzyme-linked immunosorbent assay (ELISA) kit (anti-Zika Virus ELISA immunoglobulin G; Euroimmun, Germany), and positives were further processed by microneutralization in a 96-well format, using Vero cells, 50 pfu of




Figure 1. Clinical symptoms and Zika viral loads in blood donors. The distribution of symptomatic donors according to time of inaugural symptoms (A) and the frequency of reported symptoms (B) are presented. Zika viral loads (log₁₀ copies/mL) are provided in both asymptomatic and symptomatic blood donors (C). Zika viral load was determined by real-time reverse transcription polymerase chain reaction in 41 donors reporting clinical signs, and 34 donors who did not. The diagram reports for each series the quantitative distribution of viral load, together with the median value \pm SD.

the MRS_OPY_Martinique_PaRi_2015 strain,¹⁴ and a threshold titer of 40, as recommended by the French National Reference Centre for Arboviruses. The seroprevalence after seroneutralization was 13.5% in early March and 42.2% in early June. An ELISA ratio greater than 4 was associated with a positive seroneutralization in more than 95% of cases. Sera with an ELISA ratio greater than 5 were associated with seroneutralization in 100% of cases.




Figure 2. Epidemiological dynamics of Zika outbreak in Martinique. Figure 2A reports the cumulative incidence of Zika virus NAT detection in blood donors (normalized for 100 weekly blood donations). Figure 2B shows the simulated prevalence of Zika virus infections in blood donors at different points, based on incident cases and on different hypotheses for the duration of asymptomatic viremia (4-6 days). It also presents the actual seroprevalence data obtained by seroneutralization at weeks 8 and 22.

During the same period, the cumulative curve of incident cases detected by NAT in blood donors was almost linear (Figure 2A). This allowed performance of a simulation analysis and determination, on the basis of the incident cases detected among blood donors, of the number of donors infected by ZIKV at different points. Starting from the hypothesis of a random distribution of the first day of detectable viremia in infected donors during all days of the period studied, and the day of donation during the days open to collection during the same period, we estimated the number of cases for which donation would occur during the phase of asymptomatic viremia detectable by NAT. We deduced the expected prevalence of ZIKV infection in donors for theoretical durations of asymptomatic viremia in plasma of 4 to 6 days (Figure 2B), by reference to dengue¹⁵ and chikungunya fever.¹⁶ A faithful correlation with actual seroprevalence results was observed for a duration of asymptomatic viremia of 6 days (prevalence estimated at 12.5% vs 13.5% by seroneutralization in early March, and at 43.5% vs 42.2% in early June). This is consistent with the observation that the longest period reported between blood donation and symptoms (presymptomatic viremia) was 6 days (Figure 1A). Because detection methods using whole blood samples have allowed detecting virus RNA up to 68 days after the onset of symptoms,¹⁷ it will be important to determine whether such samples contain infectious virus, and to reassess ZIKV transfusion risk accordingly.

The estimated number of clinically suspect cases in Martinique (French Public Health Institute) was ~7600, or ~2% of the population, in early March (week 8), and ~28 900, or ~7.6% of the population, in early June (week 22).¹⁸ With reference to seroprevalence results, this suggests that the proportion of cases that did not seek medical attention was 80% to 85% and did not change significantly along the study period. Our analyses suggest, therefore, that ZIKV has spread in the general population silently, with a minor effect of the acute disease on medical structures (eg, 10-35 weekly consultations for Zika syndromes in emergency departments, and 10-20 in obstetric departments). The most significant medical effect consisted of 19 Zika-related GBS confirmed cases and Zika infection in 342 pregnant women at the end of our study (June 2016).¹⁹ Therefore, the situation is significantly different from that previously observed in Martinique during the 2014 Chikungunya outbreak, characterized by a high proportion of cases seeking medical attention.²⁰

We conclude that ZIKV individual NAT screening in Martinique during 5 months of circulation of the virus allowed the detection of approximately 2% of contaminated blood donations. The proportion of truly asymptomatic cases of Zika disease among Martiniquan blood donors infected by ZIKV was approximately 45%, and the proportion of cases that did not require medical attention was even higher (80%-85%). The duration of plasma asymptomatic and presymptomatic viremia was estimated to be close to 6 days. We suggest that, once the duration of asymptomatic viremia has been estimated (eg., from early analysis of seroprevalence and PCR detection datasets), NAT incidence studies in blood donors could improve prevalence estimates in the general population in the case of a disease with frequent mild or asymptomatic clinical presentations such as Zika fever. The French Blood Bank experience of individual ZIKV NAT screening and epidemiological studies performed from French blood donor population may be of interest at the time the US Food and Drug Administration is recommending that blood banks screen all blood donations for ZIKV or use an approved pathogen reduction technology.

There is an Inside *Blood* Commentary on this article in this issue.

Acknowledgments: This work was supported by the Etablissement Français du Sang, INSERM, and Aix-Marseille University. P.G., A.C., R.N.C., B.P., I.L.-G., and X.d.L. are members of the ZIKAlliance project, funded by the European Commission.

Contribution: P.R. and L.P. collected samples and data from blood donors. A.C. collected and interpreted data from general population. P.G. and P.T. established the RT-PCR platform and performed NAT screening. B.P. and I.L.-G. designed and performed serological detection. P.G., P.T., R.N.C., and X.d.L. designed the study, interpreted the data, and wrote and edited the manuscript.

Conflict-of-interest disclosure: The authors declare no competing financial interests.

ORCID profiles: P.G., 0000-0002-3310-5808; R.N.C., 0000-0002-7675-8251; X.de.L., 0000-0001-7895-2720.

Correspondence: Pierre Gallian, Etablissement Français du Sang Alpes-Méditerranée, 149 Bd Baille, 13005 Marseille, France; e-mail: pierre.gallian@efs.sante.fr.

References

- Kindhauser MK, Allen T, Frank V, Santhana RS, Dye C. Zika: the origin and spread of a mosquito-borne virus. *Bull World Health Organ.* 2016;94(9):675-686C.
- Grischott F, Puhan M, Hatz C, Schlagenhauf P. Non-vector-borne transmission of Zika virus: A systematic review. *Travel Med Infect Dis.* 2016;14(4):313-330.
- D'Ortenzio E, Matheron S, Yazdanpanah Y, et al. Evidence of sexual transmission of Zika virus. *N Engl J Med.* 2016;374(22):2195-2198.

4. Davidson A, Slavinski S, Komoto K, Rakeman J, Weiss D. Suspected female-to-male sexual transmission of Zika virus - New York City, 2016. *MMWR Morb Mortal Wkly Rep.* 2016;65(28):716-717.
5. Barjas-Castro ML, Angerami RN, Cunha MS, et al. Probable transfusion-transmitted Zika virus in Brazil. *Transfusion.* 2016;56(7):1684-1688.
6. Oehler E, Watrin L, Larre P, et al. Zika virus infection complicated by Guillain-Barre syndrome—case report, French Polynesia, December 2013. *Euro Surveill.* 2014;19(9):20720.
7. Schuler-Faccini L, Ribeiro EM, Feitosa IM, et al; Brazilian Medical Genetics Society–Zika Embryopathy Task Force. Possible association between Zika virus infection and microcephaly - Brazil, 2015. *MMWR Morb Mortal Wkly Rep.* 2016; 65(3):59-62.
8. Mécharles S, Herrmann C, Poullain P, et al. Acute myelitis due to Zika virus infection. *Lancet.* 2016;387(10026):1481.
9. Carreaux G, Maquart M, Bedet A, et al. Zika virus associated with meningoencephalitis. *N Engl J Med.* 2016;374(16):1595-1596.
10. Mlakar J, Korva M, Tul N, et al. Zika virus associated with microcephaly. *N Engl J Med.* 2016;374(10):951-958.
11. Charrel RN, Leparac-Goffart I, Pas S, de Lamballerie X, Koopmans M, Reusken C. Background review for diagnostic test development for Zika virus infection. *Bull World Health Organ.* 2016;94(8):574-584D.
12. Musso D, Nhan T, Robin E, et al. Potential for Zika virus transmission through blood transfusion demonstrated during an outbreak in French Polynesia, November 2013 to February 2014. *Euro Surveill.* 2014;19(14):20761.
13. Aubry M, Richard V, Green J, Brout J, Musso D. Inactivation of Zika virus in plasma with amotosalen and ultraviolet A illumination. *Transfusion.* 2016;56(1): 33-40.
14. Piorowski G, Richard P, Baronti C, et al. Complete coding sequence of Zika virus from Martinique outbreak in 2015. *New Microbes New Infect.* 2016;11: 52-53.
15. Gubler DJ, Suharyono W, Tan R, Abidin M, Sie A. Viraemia in patients with naturally acquired dengue infection. *Bull World Health Organ.* 1981;59(4): 623-630.
16. Gallian P, de Lamballerie X, Salez N, et al. Prospective detection of chikungunya virus in blood donors, Caribbean 2014. *Blood.* 2014;123(23):3679-3681.
17. Lustig Y, Mendelson E, Paran N, Melamed S, Schwartz E. Detection of Zika virus RNA in whole blood of imported Zika virus disease cases up to 2 months after symptom onset, Israel, December 2015 to April 2016. *Euro Surveill.* 2016; 21(26).
18. Santé publique France. Epidemiological bulletin of CIRE Antilles Guyane N°22/ 2016. Available at: http://www.ars.martinique.sante.fr/fileadmin/MARTINIQUE/Actualites/Autres_actu/2016/ZIKA/PE/PE_Zika_2016-22_court.pdf. Accessed 31 August 2016.
19. Santé publique France. Epidemiological bulletin of CIRE Antilles Guyane N°23/ 2016. Available at: <http://invs.santepubliquefrance.fr/fr/Publications-et-outils/Points-epidemiologiques/Tous-les-numeros/Antilles-Guyane/2016/Situation-epidemiologique-du-virus-Zika-aux-Antilles-Guyane.-Point-au-16-juin-2016>. Accessed 31 August 2016.
20. Santé publique France. Epidemiological bulletin of CIRE Antilles Guyane N°1/2015. Available at: http://www.ars.martinique.sante.fr/fileadmin/MARTINIQUE/Actualites/Autres_actu/2015/fin_epidemie_CHIK/PE_DFA_2015-01_Chik_antilles_lg.pdf. Accessed 19 September 2016.

DOI 10.1182/blood-2016-09-737981

© 2017 by The American Society of Hematology