CLINICAL TRIALS AND OBSERVATIONS

Safety and tolerability of pembrolizumab in patients with relapsed/refractory primary mediastinal large B-cell lymphoma

Pier Luigi Zinzani,¹ Vincent Ribrag,² Craig H. Moskowitz,³ Jean-Marie Michot,² John Kuruvilla,⁴ Arun Balakumaran,⁵ Yayan Zhang,⁵ Sabine Chlosta,⁵ Margaret A. Shipp,⁶ and Philippe Armand⁶

¹Institute of Hematology "L. e A. Seràgnoli," University of Bologna, Bologna, Italy; ²Gustave Roussy, Université Paris-Saclay, Département d'hématologie, INSERM U1170, Villejuif, France; ³Memorial Sloan Kettering Cancer Center, New York, NY; ⁴Princess Margaret Cancer Centre and University of Toronto, Toronto, ON, Canada; ⁵Merck & Co., Inc., Kenilworth, NJ; and ⁶Dana-Farber Cancer Institute, Boston, MA

Key Points

- Treatment options for relapsed/refractory PMBCL are limited, and prognosis is generally poor.
- Pembrolizumab had a manageable safety profile and promising antitumor activity in heavily pretreated rrPMBCL patients.

Treatment options for relapsed/refractory primary mediastinal large B-cell lymphoma (rrPMBCL) are limited, and prognosis is generally poor (overall response rate [ORR] 0% to 25%; 2-year overall survival 15%). PMBCL frequently involves PD-1 ligand overexpression, potentially making PMBCL particularly susceptible to PD-1 blockade. We evaluated safety and antitumor activity of pembrolizumab, an anti–PD-1 antibody, in rrPMBCL as part of the KEYNOTE-013 multicohort phase 1b trial. At time of data cutoff, 18 patients (median age 30 years; median 3 prior lines of therapy) had been enrolled and treated, of whom 17 were included in the efficacy analyses. Eleven patients (61%) experienced drug-related adverse events (mostly grade 1-2); none discontinued treatment due to adverse events. ORR was 41% (7/17); 6 additional patients (35%) had stable disease. Of patients evaluable by imaging, 13 out of 16 (81%) had decreases in target lesions. With a median follow-up of 11.3 months, median duration of response was not reached. Two patients reached the maximum 2-year treatment duration and

remain in remission. Median overall survival was not reached for treated patients overall; all responders were still alive at data cutoff. These results in heavily pretreated rrPMBCL patients demonstrate that PD-1 blockade with pembrolizumab has a manageable safety profile and promising antitumor activity. This trial was registered at www.clinicaltrials.gov as #NCT01953692. (*Blood.* 2017;130(3):267-270)

Introduction

Primary mediastinal B-cell lymphoma (PMBCL) is a diffuse large Bcell lymphoma (DLBCL) subtype occurring mostly in young women.¹ While most patients with PMBCL are cured with standard frontline therapy, $^{2} \sim 200$ patients per year in the United States are diagnosed with relapsed/refractory PMBCL (rrPMBCL),³ which has a poor prognosis with limited treatment options.^{2,4} The rarity of rrPMBCL limits the ability to conduct clinical trials, and no standard of care has been identified; rrPMBCL is generally treated like other relapsed/refractory DLBCL (rrDLBCL) subtypes.^{2,4} National Comprehensive Cancer Network guidelines recommend that patients who relapse after autologous stem cell transplantation (SCT) or are ineligible for high-dose therapy be enrolled in clinical trials or receive second-line chemotherapy regimens, palliative radiation, or supportive care. These treatment options generally yield poor outcomes. In a study of 106 PMBCL patients, outcomes in both primary refractory and relapsed PMBCL after doxorubicin-containing therapy were poor: 0% of 35 primary refractory and 4 out of 18 relapsed patients (22%) responded to salvage treatment.⁵ In a retrospective study among 180 patients relapsed/refractory to 1 line of anthracycline-based chemotherapy, salvage chemotherapy resulted in significantly worse overall response rate (ORR) in rrPMBCL (25%) than in rrDLBCL overall (48%); 2-year overall survival after diagnosis of relapsed/refractory disease was 15% and 34%, respectively (also significantly different).⁶ There is an urgent medical need for more effective treatment options in rrPMBCL.

Like classical Hodgkin lymphoma, PMBCL frequently exhibits 9p24.1/*PD-L1*/*PD-L2* copy-number alterations and rearrangements and associated PD-L1 and/or PD-L2 overexpression, potentially facilitating immune evasion.⁷⁻¹¹ The genetics of PMBCL could thus make it particularly susceptible to PD-1 blockade.¹¹ Pembrolizumab is a humanized anti–PD-1 monoclonal antibody blocking interaction of PD-1 with its ligands, PD-L1 and PD-L2. Pembrolizumab has demonstrated efficacy against solid tumors^{12,13} and promising antitumor activity in Hodgkin lymphoma.¹⁴ Here, we report results from the PMBCL cohort of an ongoing phase 1 trial of pembrolizumab in patients with hematologic malignancies.

There is an Inside *Blood* Commentary on this article in this issue.

The publication costs of this article were defrayed in part by page charge payment. Therefore, and solely to indicate this fact, this article is hereby marked "advertisement" in accordance with 18 USC section 1734.

The online version of this article contains a data supplement.

Society of Hematology, San Diego, CA, 3-6 December 2016.

© 2017 by The American Society of Hematology

Submitted 27 December 2016; accepted 24 April 2017. Prepublished online as *Blood* First Edition paper, 10 May 2017; DOI 10.1182/blood-2016-12-758383. Presented in abstract form and orally at the 58th annual meeting of the American

Table 1. Clinical outcomes in patients evaluated for efficacy: response to treatment

	n (%) (N = 17)	90% confidence interval
ORR	7 (41.2)	21.2, 63.6
CR	2 (11.8)	12.4, 52.2
Partial response	5 (29.4)	2.1, 32.6
Stable disease	6 (35.3)	16.6, 58.0
Progressive disease	3 (17.6)	5.0, 39.6
No assessment	1 (5.9)*	0.3, 25.0

*Patient discontinued pembrolizumab therapy prior to the first radiologic tumor assessment because of clinical disease progression.

Study design

KEYNOTE-013 is a multicenter, international, multicohort, open-label phase 1b trial evaluating safety, tolerability, and antitumor activity of pembrolizumab in patients with selected hematologic malignancies. Adults with rrPMBCL who failed, were ineligible for, or refused autologous SCT formed an independent trial cohort. Key eligibility criteria were Eastern Cooperative Oncology Group performance status 0-1, no active autoimmune disease, no allogeneic SCT within the past 5 years, no symptomatic central nervous system disease, no active infection requiring intravenous therapy, and no prior therapy with agents targeting T-cell costimulation or checkpoint pathways. The study was conducted in accordance with principles of Good Clinical Practice and was approved by the appropriate institutional review boards and regulatory agencies. (The complete Study Protocol is available in the supplemental Material, available on the *Blood* Web site.)

The first 11 patients were to receive intravenous pembrolizumab 10 mg/kg every 2 weeks for the trial's duration (1 patient withdrew shortly after enrollment and was not dosed); after a study amendment, the subsequent 8 patients received a fixed dose of IV pembrolizumab 200 mg every 3 weeks, based on pharmacokinetic/pharmacodynamic evaluations that demonstrated the regimens provided similar exposure.¹⁵ Treatment continued up to 2 years, until unacceptable toxicity, or until confirmed disease progression. Treatment response was evaluated using International Working Group 2007 criteria for malignant lymphoma.¹⁶ Adverse events (AEs) were collected throughout the study and for 30 days thereafter (90 days for serious AEs) and graded per NCI CTCAE v4.0.17 Primary end points were safety and ORR by investigator assessment. Key secondary end points included complete response (CR) rate, duration of response (DOR), and overall survival. Patients were censored for DOR at the time of starting another anticancer therapy. Safety was evaluated in all patients who received ≥ 1 dose of study drug; those patients who also had \geq 1 postbaseline tumor assessment, or disease progression before first tumor assessment, were included into the efficacy analyses.



Figure 1. Treatment response in the 16 pembrolizumab-treated patients with rrPMBCL who were evaluable (by imaging) for efficacy at the time of data cutoff. (A) Best percentage change from baseline in target lesions. (Number at the end of each bar indicates the patient identification number.) (B) Transplant eligibility, treatment duration, investigator-assessed response to treatment, and duration of response. (Unless indicated otherwise, patients had stable disease.) PD, progressive disease; PR, partial response; SD, stable disease.

Table 2. Clinical outcomes in patients evaluated for efficacy: duration of response to prior therapy and to pembrolizumab

	Prior lines of therapy	Duration of response to prior therapy, mo (type of response)	Duration of response to pembrolizumab, mo
Complete responders			
Patient 2 (SCT ineligible)	ACVBP	2 (PR)	20.5* (response ongoing)
	R-DHAX	3 (PR)	—
	R-CHOP	— (SD)	_
	BV	— (PD)	_
Patient 7 (prior autologous SCT)	R-CHOP	— (SD)	2.4 (response ongoing)
	R-ACVB	— (PR)	—
	MTX	— (PR)	—
	R-IE	— (PR)	—
	AraC	— (PD)	_
	R-DHAP	— (PR)	—
Partial responders			
Patient 1 (prior autologous SCT)	R-VACOP-B	— (SD)	22.4 (response ongoing)
	R-DHAP	1 (PR)	_
	CCNU-EAM	— (SD)	—
	DA-EPOCH	— (PD)	_
Patient 4 (SCT ineligible)	BV	— (PD)	_
	R-CHOP	3 (PR)	6.9 (disease progression)
	R-DAC	— (SD)	—
Patient 3 (SCT ineligible)	IP-NVB	— (SD)	_
	R-EPOCH	— (SD)	6.5 (response ongoing)
Patient 8 (prior autologous SCT)	R-DHAP	— (PD)	_
	CHO-MTX	1 (PR)	4.6 (response ongoing)
	IE	5 (CR)	—
Patient 5 (SCT ineligible)	BEAM	— (PD)	_
	R-EPOCH	1 (PR)	4.0 (response ongoing)
	R-DHAP	— (PD)	—

Patient numbers refer to Figure 1.

ACVBP, bleomycin, cyclophosphamide, doxorubicin, prednisone, and vindesine; AraC, cytarabine; BEAM, melphalan, carmustine, cytarabine, and etoposide; BV, brentuximab vedotin; CCNU-EAM, cytarabine, etoposide, lomustine, and melphalan; CHO-MTX, doxorubicin, cyclophosphamide, vincristine sulfate, methotrexate, etoposide, and ifosfamide; DA-EPOCH, cyclophosphamide, doxorubicin, etoposide, prednisone, and vincristine sulfate; IE, etoposide and ifosfamide; IP-NVB, ifosfamide, methylprednisolone, and vinorelbine tartrate; MTX, methotrexate; PD, progressive disease; PR, partial response; R-IE, rituximab, etoposide, and ifosfamide; R-AVCB, rituximab, doxorubicin, vindesine, bleomycin, and cyclophosphamide; R-CHOP, cyclophosphamide, doxorubicin, prednisone, rituximab, and vincristine sulfate; R-DAC, carboplatin, cytarabine, dexamethasone, and rituximab; R-DHAP, rituximab, cytarabine, cisplatin, and dexamethasone; R-DHAX, cytarabine, dexamethasone, oxaliplatin, and rituximab; R-EPOCH, cyclophosphamide, doxorubicin, etoposide, prednisone, rituximab, and vincristine sulfate; R-VACOP-B, etoposide, leucovorin, doxorubicin, cyclophosphamide, vincristine, prednisone, and bleomycin.

*Patient initially presented with partial response, and the time shown represents the total duration of response since the date partial response was first observed.

Results and discussion

By the analysis cutoff date (27 May 2016), 19 patients had been enrolled, and 18 received ≥ 1 dose of study drug. One patient had not yet reached the first response assessment at time of data cutoff; the remaining 17 comprised the efficacy analysis population. Median age was 30 years, and 72% were female (supplemental Table 1). Patients were heavily pretreated and had relapsed on multiple treatments; pembrolizumab was the median fourth line of therapy (range, 3rd to 7th). Most patients (94%) previously received rituximab, 33% had prior autologous SCT, and 61% had prior radiation. Twelve patients were transplant ineligible because of chemotherapy-resistant disease.

The safety and tolerability profile of pembrolizumab in rrPMBCL appeared manageable. Eleven patients (61%) experienced drugrelated adverse events (DRAEs), most of which were grade 1 or 2 (supplemental Table 2). Most commonly reported DRAEs were hypothyroidism, diarrhea, nausea, fatigue, pyrexia, and decreased appetite (n = 2 patients each). The only grade 3 DRAE was neutropenia in 1 patient. The only grade 4 DRAE was venoocclusive liver disease subsequent to allogeneic SCT during the follow-up period after pembrolizumab was discontinued; the patient recovered from veno-occlusive liver disease. Potential immune-related AEs (n = 1 each) were grade 2 aggravated diarrhea and grade 2 radiation pneumonitis. No patient discontinued treatment due to AEs. There were no treatment-related deaths.

At best response, the ORR was 41% (7/17), with 2 patients achieving a CR and 5 a partial response; an additional 35% (6/17) had stable disease as best response (Table 1). This ORR was higher than reported in prior retrospective studies, where it ranged from 0% to 25% in patients primary refractory to or relapsed after anthracycline-based chemotherapy.^{5,6} In our study, 81% (13/16) of patients evaluable by imaging had an overall decrease in target lesions; in 2 of these patients, reductions were <5% (Figure 1A). With a median follow-up duration of 11.3 months (range, 3.4 to 27.4 months), median DOR was not reached, and 6 of the 7 responses in patients evaluable by imaging were ongoing at the time of data cutoff (Figure 1B). DOR ranged from 2.3+ to 22.5+ months; DOR in the 2 patients with CR was 2.3+ and 20.5+ months (patients 7 and 2 in Figure 1, respectively). Remissions achieved with pembrolizumab appeared to be long-lasting, while prior treatment responses to high-dose chemotherapy, including to first-line therapy, lasted a few months at most (Table 2). None of the responders received autologous SCT. Two patients, both with stable disease, received allogeneic SCT after pembrolizumab discontinuation (patients 9 and 10). Ten patients discontinued treatment: 5 due to progressive disease based on imaging (patients 4, 10, 11, 13, and 14), 4 for clinical progression (patients 12, 15, and 16 and 1 patient not depicted in Figure 1), and 1 due to physician decision (patient 9).

Two patients with objective response (CR or partial response) reached the maximum 2 years of treatment shortly before data cutoff and remain in remission (follow-up: 25.4 months [patient 1], 23.8 months [patient 2]). All patients who had an objective response to pembrolizumab were alive at time of data cutoff. While these outcomes are very encouraging overall, there exist few published studies specifically conducted in this rare patient population that can serve as accurate historical controls for efficacy comparisons with other treatment options for rrPMBCL.^{5,6} A recent phase 2 trial with brentuximab vedotin in this specific setting was terminated early due to low ORR (2/15 [13%]).¹⁸

These preliminary efficacy and safety findings indicate that pembrolizumab has potential to provide substantial clinical benefit in heavily pretreated patients with rrPMBCL, particularly given the lack of effective alternatives. A global, multicenter, phase 2 trial (KEYNOTE-170) is currently evaluating single-agent pembrolizumab in this setting (www.clinicaltrials.gov #NCT02576990).

Acknowledgments

The authors thank the patients and their families and caregivers for participating in this study, along with all investigators and site personnel, Patricia Marinello (Merck & Co., Inc.) for her contributions to study design, Karen Giallella and Jennifer Anderson (both of Merck & Co., Inc.) for providing study support, Roger Dansey (Merck & Co., Inc.) for critical review of an earlier draft of the manuscript, and Seth Thompson (Merck & Co., Inc.) for providing statistical guidance and support. Medical writing assistance was provided by Dominik Wolf, an employee of Merck & Co., Inc. Editorial support was provided by Sheila Erespe, an employee of Merck & Co., Inc. All principal investigators participating in this trial are also co-authors on this publication.

Funding for this research was provided by Merck & Co., Inc. (Kenilworth, NJ).

References

- Bhatt VR, Mourya R, Shrestha R, Armitage JO. Primary mediastinal large B-cell lymphoma. *Cancer Treat Rev.* 2015;41(6):476-485.
- Martelli M, Di Rocco A, Russo E, Perrone S, Foà R. Primary mediastinal lymphoma: diagnosis and treatment options. *Expert Rev Hematol.* 2015; 8(2):173-186.
- National Cancer Institute. Surveillance, Epidemiology, and End Results Program website. Available from URL: http://seer.cancer.gov/. Accessed 20 February 2017.
- Petković I. Current trends in the treatment of primary mediastinal large B-cell lymphoma - an overview. *Contemp Oncol (Pozn).* 2015;19(6):428-435.
- Lazzarino M, Orlandi E, Paulli M, et al. Treatment outcome and prognostic factors for primary mediastinal (thymic) B-cell lymphoma: a multicenter study of 106 patients. *J Clin Oncol.* 1997;15(4):1646-1653.
- Kuruvilla J, Pintilie M, Tsang R, Nagy T, Keating A, Crump M. Salvage chemotherapy and autologous stem cell transplantation are inferior for relapsed or refractory primary mediastinal large B-cell lymphoma compared with diffuse large B-cell lymphoma. *Leuk Lymphoma*. 2008; 49(7):1329-1336.
- Green MR, Monti S, Rodig SJ, et al. Integrative analysis reveals selective 9p24.1 amplification, increased PD-1 ligand expression, and further

induction via JAK2 in nodular sclerosing Hodgkin lymphoma and primary mediastinal large B-cell lymphoma. *Blood*. 2010;116(17):3268-3277.

- Rosenwald A, Wright G, Leroy K, et al. Molecular diagnosis of primary mediastinal B cell lymphoma identifies a clinically favorable subgroup of diffuse large B cell lymphoma related to Hodgkin lymphoma. J Exp Med. 2003;198(6):851-862.
- Shi M, Roemer MG, Chapuy B, et al. Expression of programmed cell death 1 ligand 2 (PD-L2) is a distinguishing feature of primary mediastinal (thymic) large B-cell lymphoma and associated with PDCD1LG2 copy gain. *Am J Surg Pathol.* 2014;38(12):1715-1723.
- Twa DD, Chan FC, Ben-Neriah S, et al. Genomic rearrangements involving programmed death ligands are recurrent in primary mediastinal large B-cell lymphoma. *Blood*. 2014;123(13): 2062-2065.
- Van Roosbroeck K, Ferreiro JF, Tousseyn T, et al. Genomic alterations of the JAK2 and PDL loci occur in a broad spectrum of lymphoid malignancies. *Genes Chromosomes Cancer*. 2016;55(5):428-441.
- Reck M, Rodríguez-Abreu D, Robinson AG, et al; KEYNOTE-024 Investigators. Pembrolizumab versus Chemotherapy for PD-L1-Positive Non-Small-Cell Lung Cancer. N Engl J Med. 2016; 375(19):1823-1833.

Authorship

Contribution: P.A. and M.A.S. contributed to study inception and design; P.L.Z., P.A., V.R., C.H.M., J.-M.M., and J.K. enrolled patients into this study; Y.Z. conducted the statistical analysis of the data; P.A., J.K., M.A.S., P.L.Z., A.B., and S.C. contributed to data interpretation; P.A., P.L.Z., S.C., and A.B. contributed to writing the first draft of the manuscript; and all authors critically revised the manuscript for important intellectual content, reviewed the submitted version of the manuscript, and agree with its content and submission.

Conflict-of-interest disclosure: P.L.Z. has been a scientific advisor to Merck Sharp & Dohme, Bristol-Myers Squibb, Celgene, Gilead, Infinity, Janssen, Pfizer, Roche, and Takeda. V.R. has been a scientific advisor to Bristol-Myers Squibb, Eisai, Gilead, Infinity, Pharmamar, and NanoString; has received research funding from ArgenX; and has received consulting fees from Servier. C.H.M. has been a scientific advisor to and received research funding from Merck & Co., Inc., BMS, and Seattle genetics. J.-M.M. has received consulting fees from Bristol-Myers Squibb. J.K. has received research funding from Roche; consultancy fees from Bristol-Myers Squibb, Gilead, Janssen, Roche, and Seattle Genetics; and honoraria from Merck & Co., Inc., Amgen, Bristol-Myers Squibb, Celgene, Gilead, Roche, Janssen, Lundbeck, and Seattle Genetics. A.B., Y.Z., and S.C. are employees of Merck Sharp & Dohme Corp., a subsidiary of Merck & Co., Inc., and may own stock and/or hold stock options in the company. M.A.S. has been a scientific advisor to Merck & Co., Inc., and received research funding and consulting fees from Bristol-Myers Squibb. P.A. received institutional research funding from Merck & Co., Inc., Affimed, Bristol-Myers Squibb, Roche, and Pfizer and consultancy fees from Merck & Co., Inc., Bristol-Myers Squibb, Infinity Pharmaceuticals, and Pfizer.

Correspondence: Pier Luigi Zinzani, Institute of Hematology "L. e A. Seràgnoli," University of Bologna, Via Massarenti 9, 40138 Bologna, Italy; e-mail: pierluigi.zinzani@unibo.it.

- Robert C, Schachter J, Long GV, et al; KEYNOTE-006 investigators. Pembrolizumab versus Ipilimumab in Advanced Melanoma. N Engl J Med. 2015;372(26):2521-2532.
- Armand P, Shipp MA, Ribrag V, et al. Programmed Death-1 Blockade With Pembrolizumab in Patients With Classical Hodgkin Lymphoma After Brentuximab Vedotin Failure. J Clin Oncol. 2016;34(31):3733-3739.
- Freshwater T, de Greef R, Kondic A, et al. Assessment of pembrolizumab (MK-3475) dosing strategy based on population pharmacokinetics and exposure-response models. J Pharmacokinet Pharmacodynam. 2015;42:S15.
- Cheson BD, Pfistner B, Juweid ME, et al; International Harmonization Project on Lymphoma. Revised response criteria for malignant lymphoma. *J Clin Oncol.* 2007;25(5):579-586.
- National Cancer Institute. Common terminology criteria for adverse events (CTCAE) v4.0. Available at: https://ctep.cancer.gov/ protocolDevelopment/electronic_applications/ctc. htm. Accessed 20 February 2017.
- Zinzani PL, Pellegrini C, Chiappella A, et al. Brentuximab vedotin in relapsed primary mediastinal large B-cell lymphoma: results from a phase 2 clinical trial. *Blood*. 2017;129(16): 2328-2330.