IMMUNOBIOLOGY AND IMMUNOTHERAPY

Patients with *CD3G* mutations reveal a role for human CD3 γ in T_{reg} diversity and suppressive function

Jared H. Rowe,^{1,*} Ottavia M. Delmonte,^{2,*} Sevgi Keles,³ Brian D. Stadinski,⁴ Adam K. Dobbs,² Lauren A. Henderson,⁵ Yasuhiro Yamazaki,² Luis M. Allende,⁶ Francisco A. Bonilla,⁵ Luis I. Gonzalez-Granado,⁷ Seyma Celikbilek Celik,³ Sukru N. Guner,³ Hasan Kapakli,³ Christina Yee,⁵ Sung-Yun Pai,^{1,8} Eric S. Huseby,⁴ Ismail Reisli,^{3,†} Jose R. Regueiro,^{9,†} and Luigi D. Notarangelo^{2,†}

¹Division of Hematology-Oncology, Boston Children's Hospital, Boston, MA; ²Laboratory of Clinical Immunology and Microbiology, Division of Intramural Research, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Bethesda, MD; ³Division of Pediatric Immunology and Allergy, Meram Medical Faculty, Necmettin Erbakan University, Konya, Turkey; ⁴Department of Pathology, University of Massachusetts Medical School, Worcester, MA; ⁵Division of Immunology, Boston Children's Hospital, Boston, MA; ⁶Department of Immunology and ⁷Primary Immunodeficiencies Unit, Department of Pediatrics, Hospital Universitario 12 de Octubre, Madrid, Spain; ⁸Department of Pediatric Oncology, Dana-Farber Cancer Institute, Boston, MA; and ⁹Department of Immunology, Complutense University School of Medicine, imas12 Research Institute, Madrid, Spain

KEY POINTS

- CD3γ-deficient patients manifest T-cell phenotypic and functional defects that are especially prominent in T_{reg} cells.
- The peripheral T-cell repertoire of CD3γdeficient patients is restricted, with molecular signatures of self-reactivity.

Integrity of the T-cell receptor/CD3 complex is crucial for positive and negative selection of T cells in the thymus and for effector and regulatory functions of peripheral T lymphocytes. In humans, *CD3D*, *CD3E*, and *CD3Z* gene defects are a cause of severe immune deficiency and present early in life with increased susceptibility to infections. By contrast, *CD3G* mutations lead to milder phenotypes, mainly characterized by autoimmunity. However, the role of CD3 γ in establishing and maintaining immune tolerance has not been elucidated. In this manuscript, we aimed to investigate abnormalities of T-cell repertoire and function in patients with genetic defects in *CD3G* associated with autoimmunity. High throughput sequencing was used to study composition and diversity of the T-cell receptor β (TRB) repertoire in regulatory T cells (T_{reg}s), conventional CD4⁺ (T_{conv}), and CD8⁺ T cells from 6 patients with *CD3G* mutations and healthy controls. T_{reg} function was assessed by studying its ability to suppress proliferation of T_{conv} cells. T_{reg} cells of patients with *CD3G* defects had reduced diversity, increased clonality, and reduced suppressive function. The TRB

repertoire of T_{conv} cells from patients with *CD3G* deficiency was enriched for hydrophobic amino acids at positions 6 and 7 of the CDR3, a biomarker of self-reactivity. These data demonstrate that the T-cell repertoire of patients with *CD3G* mutations is characterized by a molecular signature that may contribute to the increased rate of autoimmunity associated with this condition. (*Blood*. 2018;131(21):2335-2344)

Introduction

Integrity of the T-cell receptor/CD3 (TCR/CD3) complex is crucial for T-cell maturation. In particular, the strength of TCR signaling plays a critical role in governing positive and negative selection in the thymus as well as responses of effector and regulatory T (T_{reg}) cells in the periphery. Before reaching the membrane, TCRα/β heterodimers associate with 3 invariant dimers (CD3δ/ε, CD3γ/ε, and CD3ζ/ζ) that compose the CD3 complex.¹ Following localization on the cell surface, the CD3 proteins convert ligand recognition by α/β TCR chains into intracellular signals.²

Both in humans and in mice, genetic defects that cause complete absence of CD3 δ or CD3 ϵ chain expression lead to a block in the development of TCR $\alpha\beta^+$ T cells and are a cause of severe combined immune deficiency.³⁻⁵ Human CD3 ζ deficiency is characterized by a reduced number of circulating T cells that are nonfunctional and display a restricted T-cell repertoire, thereby causing severe immunodeficiency.⁶ Although a severe block in T-cell development is also observed in CD3 γ -deficient mice,⁷ it has been shown that the loss of CD3 γ protein in humans allows the development of polyclonal T cells with impaired, but not abolished, TCR/CD3 signaling, and is associated with a milder clinical phenotype characterized by a variable degree of susceptibility to infections and the frequent occurrence of auto-immune manifestations.⁸⁻¹⁰

A similar phenotype has been also reported in patients and mice with hypomorphic mutations in genes that encode for signaling molecules downstream of the TCR/CD3 complex.^{11,12} Altogether, these observations are consistent with the notion that TCR signaling strength plays a critical role both in T-cell development and function and in establishing and maintaining central and peripheral tolerance.

To gain novel insights into how *CD3G* mutations in humans affect T-cell development and homeostasis, we have analyzed TCR

diversity and composition, T-cell proliferation, and $T_{\rm reg}$ number and function in 6 patients with CD3 γ deficiency and in healthy controls.

Methods

Human subjects

Deidentified patients' clinical and immunologic data were provided by an international network of physicians in the United States, Europe, and Asia. All human subject samples were consented under protocols approved by the institutional review boards at the participating institutions. The study was approved by the institutional review board at Boston Children's Hospital (protocol 0409113R) and at the National Institute of Allergy and Infectious Diseases, National Institutes of Health, Bethesda (protocol 16-I-N139). The study met the institutional review board standards for ethical conduct of research with human subjects.

Flow cytometry/fluorescent activated cell sorting

Peripheral blood mononuclear cells (PBMCs) were processed using Ficoll (GE Healthcare, Malborough, MA) to form a single cell suspension and then stained with the following monoclonal antibodies directed against cell surface antigens: CD4-AlexaFluor700, CD8a-PE/ Dazzle594, CD19-PerCPCy5.5, CD127-PECy7, and CD25-PE (all from Biolegend, San Diego, CA). Intranuclear Foxp3-eFluor450 (eBioscience, San Diego, CA) or Ki67 (Becton Dickinson San Jose, CA) staining was performed using the Foxp3 staining buffers (eBioscience). For further characterization of T_{reg} cells, PBMC were also stained with monoclonal antibodies against CTLA-4 (clone BNI3, eBioscience), ICOS/CD278 (clone DX29, eBioscience), and HELIOS (clone 22F6, eBioscience), along with CD4 and FOXP3. For intracellular staining, fixation/permeabilization buffer (eBioscience) was used according to the manufacturer's instructions. Upon washing, cells were analyzed by flow cytometry using LSRFortessa, and results were analyzed using FlowJo software, version 8.8.6 (Tree Star Ashland, OR). In parallel, PBMC were stained with CD3-eFluor450 (OKT3, eBioscience), CD4-FITC, CD8a-APC, TCRαβ-APC (clone IP26), mouse IgG1-eFluor450, and mouse IgG1-APC (all from Biolegend), followed by cell sorting using FACSAria (Becton Dickinson).

Cell proliferation assay

PBMC were isolated to form a single cell suspension and then stained with carboxyfluorescein succinimidyl ester (CFSE) (Thermo Fisher, Carlsbad, CA) at a concentration of 5 μ M/mL for 10 minutes at 37°C in phosphate-buffered saline. Cells where then washed, resuspended in RPMI containing 10% fetal bovine serum, plated in a 48-well plate at a concentration of 5 \times 10⁵ cells per well, and then cultured with either medium alone (no stimulation), 5 μ g/mL phytohemagglutinin (PHA; Sigma Aldrich, St. Louis, MO), or 100 ng/mL anti-CD3 (eBioscience) either alone or in combination with 100 ng/mL anti-CD28 (eBioscience) for 4 days. For patient 6, these standard doses were titrated by log increments to create the dose-response curves. The cells were then stained for flow cytometry as indicated previously, and proliferation was assessed by CFSE dilution as described.¹³

T_{reg} suppression assay

For evaluating T_{reg} suppressive potency, CD4⁺ cells were enriched by magnetic bead separation (Miltenyi Biotec, Germany) followed by fluorescent activated cell sorting into the following subsets: CD4⁺ CD25^{hi} CD127^{low} cells (T_{reg} cells) and CD4⁺ CD25^{low} CD127^{hi} T effector (T_{eff}) cells. In each experiment, CD4⁺ T_{eff} cells were stained using Cell Trace Violet (Thermo Fisher) according to the manufacturer's instructions. T_{reg} and T_{eff} cells were then mixed at the indicated ratios based on a conventional CD4+ (T_{conv}) count of 5000 cells per well, then stimulated with anti-CD2/CD3/CD28 beads (Treg Suppression Inspector, Miltenyi Biotec) at a 1:1 T_{conv} : bead ratio in complete RPMI media for 4 days. In this assay, T-cell proliferation is indicated by dilution of Cell Trace Violet. The relative T_{reg} suppressive activity was determined by plotting the percentage of dividing cells at each T_{reg} : T_{eff} ratio.

High throughput sequencing (HTS) of TCR $\!\beta$

PBMCs were collected and sorted into CD3⁺ CD4⁺ CD8⁻ $CD25^{hi}\,CD127^{low}\,T_{reg},\,CD3^{+}\,CD4^{+}\,CD8^{-}CD25^{low}\,CD127^{hi}\,T_{conv},$ and CD3⁺ CD8⁺ CD4⁻ T-cell populations using FACSAria cell sorter (Becton Dickinson). The TCRβ (TRB) rearranged genomic products were amplified by multiplex polymerase chain reaction (PCR), using DNA as the template (Adaptive Biotechnologies Seattle, WA). To amplify all possible VDJ combinations, 52 forward primers for the TRBV gene segments and 13 reverse primers to the TRBJ gene segments were used. Adaptive Biotechnologies uses assay-based and computational techniques to minimize PCR amplification bias. The assay is quantitative, and the frequency of a given TRB sequence is representative of the frequency of that clonotype in the original sample. The PCR products were sequenced using the Illumina HiSeq platform. Custom algorithms were used to filter the raw sequences for errors and align the sequences to reference genome sequences. Subsequently, the data were analyzed using the ImmunoSeq set of online tools. The frequency of productive and nonproductive TRB rearrangements was analyzed within both unique and total TRB sequences obtained from sorted T-cell subsets. Distribution of the frequency of individual clonotypes (including TRBV to TRBJ pairing) and use of individual TRBV genes were analyzed within unique sequences. Heat map representation of the freguencies of individual TRBV to TRBJ gene pairs and sequence overlap was produced using Morpheus software (https:// software.broadinstitute.org/morpheus/). TCR self-reactivity was analyzed by analyzing amino acid composition and hydrophobicity at positions 6 and 7 of the TRB CDR3, as previously described.¹⁴ TRB sequences from patients and controls are available at: https://doi.org/10.21417/B7XS7J.

Statistical analysis

The number and percentage of cells and of *TRB* HTS reads were first analyzed and found to follow a Gaussian distribution. The differences between groups were analyzed using either Student t test (for individual comparisons with gaussian distributions), Mann-Whitney test (for individual comparisons with nongaussian distributions), or 2-way analysis of variance (for multiple comparisons) with statistical significance indicating a 95% confidence interval >0, *P < .05, **P < .01, ***P < .001, and ****P < .0001. All statistical analysis was performed using Prism Software.

Results

Patients' molecular, clinical, and immunological features

Six patients carrying biallelic *CD3G* mutations were studied; their molecular, clinical, and laboratory features are reported in Table 1. Five patients were homozygous, and 1 patient was compound heterozygous for *CD3G* mutations. Patient 2 (P2) and P3, and

Table 1. Clinical and immunological phenotype of CD3G-mutated patients

Patient	P1	P2	P3	P4	P5	P6
Age at testing	27 mo	19 y	6 у	15 y	23 y	34 y
CD3G mutation (allele 1)	c.1A>G	c.80-1G>C	c.80-1G>C	c.80-1G>C	c.80-1G>C	c.1A>G
CD3G mutation (allele 2)	c.1A>G	c.80-1G>C	c.80-1G>C	c.80-1G>C	c.80-1G>C	c.80-1G>C
CD3, cells/µL	2018 (850-4300)	630 (850-3200)	2800 (2400-8300)	380 (850-3200)	1200 (570-2400)	1051 (850-3200)
CD4, cells/μL Of which are naïve, %	627 (500-2700) 39.4	520 (400-2100) 19.3	1700 (1300-7100) 41.3	210 (400-2100) 5.4	800 (430-1800) 13.0	842 (400-2100) 3.1
CM, % EM, % EMRA, %	40.3 19.8 0.5	25.5 53.5 1.7	20.7 34.4 3.6	61.7 31.5 1.5	67.2 19.7 0.1	87 9.5 0.1
CD8, cells/μL Of which are naïve, %	1139 (200-1800) 7.4	40 (300-1300) 7.7	822 (400-4100) 26.0	116 (300-1300) 5.3	270 (210-1200) 11.3	200 (300-1300) 5.0
CM, % EM, % EMRA, %	1.6 41.4 49.6	0.7 57.5 34.1	1.2 44.9 27.9	3.0 55.4 36.4	9.0 53.5 26.1	45.5 44.7 4.7
CD19, cells/µL	1988 (180-1300)	250 (120-740)	170 (110-7700)	135 (120-740)	800 (91-610)	195 (120-740)
CD16/56, cells/µL	945 (61-510)	920 (92-1200)	1500 (71-3500)	370 (92-1200)	500 (92-1200)	137 (74-620)
lgG, mg/dL	338 (407-1009)	1240 (768-1632)	881 (331-1164)	491 (768-1632)	1660 (768-1632)	632 (768-1632)
lgA, mg/dL	20 (14-122)	180 (68-408)	18 (14-105)	6.6 (68-408)	165 (68-408)	364 (68-408)
lgM, mg/dL	41 (46-160)	110 (35-263)	213 (41-164)	50.4 (35-263)	219 (35-263)	59 (35-263)
lgE, kU/L	<1 (≤199)	239 (≤696)	6 (≤97)	6 (≤696)	6 (≤214)	ND
Infections	Klebsiella pneumoniae, MSSA, severe EBV	Acute bronchitis (×3)	none	RRTI, bronchiectasis	none	Soft tissue abscesses, viral meningitis
Autoimmunity	Autoimmune enteropathy, GLILD, AIHA	Hypothyroidism	Hypothyroidism	Hypothyroidism, AIHA, ITP, nephropathy, ALPS	Thyroiditis	Vitiligo, hypothyroidism
Skin rash	No	Atopic dermatitis	No	No	Atopic dermatitis	Atopic dermatitis

Reference values for age are shown in parentheses.

AIHA, autoimmune hemolytic anemia; ALPS, autoimmune lymphoproliferative syndrome; CM, central memory (CD45RA⁻ CCR7⁺); EBV, Epstein-Barr virus; EM, effector memory (CD45RA⁻ CCR7⁻); EMRA, effector memory CD45RA⁺ (CD45RA⁺ CCR7⁻); GLILD, granulomatous lymphocytic interstitial lung disease; ITP, idiopathic thrombocytopenic purpura; MSSA, methicillinsensitive *Staphylococcus aureus*; ND, no data; RRTI, recurrent respiratory tract infections.

P4 and P5 were 2 pairs of siblings. All patients had evidence of autoimmunity, which was more severe in P1 and P4 (supplemental Table 1, available on the *Blood* Web site). A history of infections was present in 4 patients, and was more severe in P1, P4, and P6. Significant variability of the immunological profile was observed. In particular, a low CD4⁺ cell count was observed in P4, and decreased CD8⁺ cell counts were detected in P2, P4, and P6. All patients had a reduced proportion of naïve T cells and increased percentage of memory CD4⁺ and CD8⁺ cells, including accumulation of a large proportion of CD45RA⁺ CCR7⁻ T_{EMRA} "exhausted" CD8⁺ T cells in 5 (P1-P5) (Table 1). Low IgG serum levels were present in P1, P4, and P6.

CD3 γ is essential for optimal expression of surface CD3/TCR $\alpha\beta$ complex

To determine the impact of *CD3G* mutations on the expression of the TCR/CD3 complex, we analyzed CD8⁺ T cells, CD4⁺ T_{conv}, and CD4⁺ T_{reg} cells from patients P1 to P5 and controls, and found that expression of both CD3 (as indicated by staining with a monoclonal antibody specific for the CD3 ϵ chain) and TCR $\alpha\beta$ was markedly reduced in each of these subsets in all patients with *CD3G* mutations (Figure 1). These findings confirm and extend previous observations^{10,15-17} and demonstrate that CD3 γ is required to maintain optimal levels of surface TCR $\alpha\beta$ and CD3 complex expression in all T-cell subsets.



Downloaded from http://ashpublications.net/blood/article-pdf/131/21/2335/1468756/blood835561.pdf by guest on 07 June 2024

Figure 1. CD3 and TCR $\alpha\beta$ expression on the surface of T-cell subsets. (A) Expression of CD3 (red line) or isotype (solid gray) on the surface of CD4⁺ FOXP3⁻ T_{conv}, CD4⁺ FOXP3⁺ T_{reg}, and CD8⁺ T cells. (B) Cumulative data of geometric mean fluorescence intensity of CD3 surface expression among T_{conv}, T_{reg}, or CD8⁺ cells. (C) Expression of TCR $\alpha\beta$ (red line) or isotype (solid gray) on the surface of T_{conv}, T_{reg}, and CD8⁺ cells. (D) Expression of TCR $\alpha\beta$ (red line) or isotype (solid gray) on the surface of T_{conv}, T_{reg}, and CD8⁺ cells. (D) Expression among T_{conv}, T_{reg}, and CD8⁺ cells. (B) Cumulative data of geometric mean for mean fluorescence intensity of TCR $\alpha\beta$ surface expression among T_{conv}, T_{reg}, and CD8⁺ cells. (B) Five patients (P1-P5) were studied.

Figure 2. T-cell activation and cell division following mitogen stimulation. (A) Representative plots and (B) cumulative percentage of (top) CD25 expression, (middle) Ki67 expression, and (bottom) CFSE staining in $\rm CD4^{+}$ T cells from controls (solid gray) or CD3G patients (red) in unstimulated samples or following activation with PHA, anti-CD3, or anti-CD3/ CD28. (C) Representative plots and (D) cumulative data for (top) CD25 expression, (middle) Ki67 expression, and (bottom) CFSE staining in CD8⁺ T cells from controls (solid gray) or patients (red) among the indicated groups. (A-D) Response to anti-CD3+anti-CD28 was studied in 5 patients (P1-P5) because not enough cells were available from P6. Dose titration curve of CFSE dilution among (E) CD4+ and (F) CD8+ T cells following stimulation with PHA (top) or anti-CD3 (bottom) for patient P6 (red line) and control sample (blue line). Unstim, unstimulated.





Figure 3. Flow cytometric analysis of peripheral blood lymphocyte populations in patients with CD3G mutations. (A) Representative flow cytometry plots for circulating CD4⁺ and CD8⁺ T lymphocytes in controls and CD3G-mutated patients. (B) Percentage of CD4⁺ (left) and CD8⁺ (right) T cells among PBMCs for control and CD3G-mutated patients. (C). Representative flow cytometry for T_{reg} identification by FOXP3 and CD25 expression. (D) Percentage (left) of FOXP3⁺ T_{reg} cells and FOXP3 mean fluorescence intensity (right) for controls and patients.

CD3 γ deficiency affects T-cell proliferation in response to mitogen stimulation

Next, we analyzed whether the impaired surface expression of CD3/TCR $\alpha\beta$ complex in T cells from *CD3G*-mutated patients would also affect cellular activation and proliferation. When stimulated with phytohemagglutinin (PHA), both CD4⁺ and CD8⁺ T cells in *CD3* γ -deficient patients demonstrated reduced cellular division (as shown by CFSE dilution); a similar trend was observed when analyzing cellular activation (by staining for CD25) and proliferation (by Ki-67 staining) (Figure 2A-D).

Similar, albeit less pronounced, defects in division were observed in CD8⁺, but not in CD4⁺ T cells, from *CD3G*-mutated patients upon in vitro stimulation with anti-CD3. Stimulation with both anti-CD3 and anti-CD28 resulted in overall equivalent activation, proliferation, and cell division in both CD4⁺ and CD8⁺ T cells from patients and controls (Figure 2A-D).

Next, we interrogated whether the impaired cell activation and proliferation observed in response to optimal concentrations of PHA (5 µg/mL) could be overcome by using higher PHA concentrations. Peripheral blood CD8⁺ and CD4⁺ T cells from P6 and a healthy control were stimulated in vitro with increasing concentrations of PHA. As shown in Figure 2E-F, a lower percentage of dividing CD8⁺ and CD4⁺ T cells was detected in P6 than in the healthy control at any PHA concentration. A subtle defect was also observed on stimulation with various concentrations of anti-CD3. Finally, the peak of cell division was reached at the same concentration of PHA or anti-CD3 both in the healthy control and in P6. Taken together, these results demonstrate the impaired surface expression of the TCR/CD3 complex in CD3y-deficient T cells is associated with reduced cellular proliferation after mitogen stimulation, and that this defect cannot be circumvented by increasing the strength of the stimulating condition.

${\sf T}_{\sf reg}$ cells from patients with CD3 γ deficiency have decreased suppressive activity

Given the importance of TCR signaling in T_{reg} development and homeostasis,¹⁸⁻²⁰ we hypothesized that this subpopulation of cells would be disproportionately affected by reduced TCR/CD3 signaling in patients with *CD3G* mutations. Among CD4⁺ T cells, the proportion of CD25^{hi} FOXP3⁺ T_{reg} cells was significantly lower in *CD3G*-mutated patients than in controls. However, levels of FOXP3 protein expression were similar in T_{reg} cells from patients and controls (Figure 3). Other phenotypic markers of T_{reg} cells (CTLA-4, HELIOS, ICOS) were equally expressed both in patient-and in control-derived T_{reg} cells (supplemental Figure 2).

Next, we sought to determine whether the impairment of TCR/CD3 signaling observed in CD3y-deficient T cells would also affect the capacity of T_{reg} cells to suppress the activation of T_{conv} cells. CD4⁺ T cells from 4 patients (P2-P5) and controls were sort-purified into CD25^{hi} CD127^{low} T_{reg} and CD25^low CD127^{hi} T_{eff} cells (Figure 4A). The T_{eff} cells from controls were assayed for cell division following coculture with varying ratios of T_{reg} cells isolated from either controls or patients. Within each experiment, the percentage of dividing cells was normalized to the percentage of dividing $T_{\rm eff}$ cells upon stimulation with anti-CD2/CD3/CD28 beads in the absence of T_{req} cells (Figure 4B; solid gray area). T_{eff} cells cocultured with control T_{reg} cells demonstrated a linear reduction in cell proliferation as the number of T_{reg} cells was uptitrated to a final T_{reg} : T_{eff} cell ratio of 1:1 (Figure 4B-C). By contrast, T_{reg} cells from CD3G-mutated patients completely failed to suppress proliferation of control T_{eff} cells up to a 1:2 T_{reg} : T_{eff} cell ratio; reduced suppressive activity was also detected at a 1:1 T_{reg}:T_{conv} ratio (Figure 4B-C). Together, these findings demonstrate that T_{reg} cells isolated from patients with CD3G mutations are unable to suppress the proliferation of T_{eff} cells upon stimulation, suggesting that loss of T_{reg} suppressive function may be an

Figure 4. T_{reg} suppression assay from fluorescent activated cell sorting purified $T_{\rm reg}$ cells in CD3Gmutated patients. (A) CD25^{hi}CD127^{lo} T_{reg} sort purification strategy from control (left) and patient (right) samples. Intranuclear FOXP3 staining was performed on CD25^{Io}CD127^{hi} T_{eff} cells (solid blue) and CD25^{hi}CD127^{Io} T_{reg} cells (red line). (B) Representative plots of Cell Trace Violet dilution in experiments in which CD4+ CD25low CD127^{hi} T_{eff} cells from controls were either left unstimulated (solid blue) or stimulated with anti-CD3 + anti-CD28 + anti-CD2, in the absence (solid gray) or presence (red line) of T_{reg} cells from controls (left) or from CD3G-mutated patients (right). (C) Cumulative percentage of dividing cells in T_{reg} suppression assay, normalized to the no T_{reg} group within each individual experiment. T_{reg} function was tested in 4 patients (P2-P5) and 4 controls. Results are expressed as mean \pm standard deviation. *P < .05; **P < .01; ***P < .001.



important component of altered peripheral T-cell tolerance in patients with CD3 γ deficiency.

Skewed use of TRBV and TRBJ genes and abnormalities of TRB CD3 composition in patients with CD3G mutations

Reduced TCR/CD3 expression and signaling may also affect mechanisms of positive and negative selection in the thymus,^{21,22} with CD3y playing an important role in this process.23 To investigate in greater detail how human CD3y deficiency may alter peripheral T-cell repertoire diversity and composition, we performed HTS to analyze TRB diversity, CDR3 characteristics, and TRB V-J pair gene use in patient T cells. Peripheral blood T-cell subsets were sort-purified into CD8⁺ T cells, CD4⁺ CD25^{hi} CD127^{lo} T_{reg} cells, and CD4+ CD25 $^{\rm low}$ CD127 $^{\rm hi}$ T_{conv} cells. Upon HTS of TRB rearrangements, there was a \sim log10 reduction in the number of unique productive reads identified in CD4+ T_{reg} cells from CD3 $\gamma\text{-}$ deficient patients compared with controls, whereas no differences were observed within CD4 $^+$ T_{conv} and CD8 $^+$ T cells (Figure 5A). Consistent with previous studies, 24,25 Shannon entropy index revealed equivalent diversity of $T_{\rm reg}$ and $T_{\rm conv}$ cells in controls samples (Figure 5B). However, T_{req} cells from CD3_Y-deficient patients had reduced diversity compared with autologous T_{conv} cells and with T_{reg} cells from healthy controls (Figure 5B). There was a trend toward reduced Shannon entropy index for patient-derived CD8 $^+$ T cells compared with control CD8 $^+$ cells, although the difference was not statistically significant.

Consistent with the reduced diversity, T_{reg} cells from CD3 γ deficient patients demonstrated a marked restriction in the number of unique *TRB VJ* pairs identified (Figure 5C; supplemental Figure 1). A similar, albeit less pronounced, restriction was present also in CD4⁺ T_{conv} and in CD8⁺ T cells from patients compared with controls (Figure 5C; supplemental Figure 1). This restriction in *TRB* diversity was driven by skewed usage of both the *TRB V* and *J* genes (supplemental Table 1). Interestingly, similar differences were observed in T_{reg} and T_{conv} cells for individual *V* and *J* genes that were under- or overrepresented in patients vs controls (supplemental Table 1). Together, these results indicate that mutations in *CD3G* result in skewing of the *TRB* repertoire through alterations in the frequency of V and *J* gene use, which results in restriction of the overall TCR repertoire.

Because patients with *CD3G* mutations develop multiple autoimmune manifestations, we sought to further characterize the



Figure 5. HTS analysis of TRB repertoire in CD4⁺ T_{reg}, CD4⁺ T_{conv}, and CD8⁺ T-cell subsets. (A) Number of unique reads identified in each patient for T_{reg}, T_{conv}, and CD8⁺ T cells and (B) Shannon entropy index calculated among unique reads for T_{reg}, T_{conv}, and CD8⁺ T cells. (A-B) *P < .05; **P < .01. (C) Representative heat maps depicting V β and J β gene pairing in unique sequences from (left) CD4⁺ T_{conv} (middle) CD4⁺ T_{reg}, and (right) CD8⁺ T cells from controls and patient 1. (D) Hydrophobicity analysis of amino acid residues at positions 6 and 7 of the TRB-CDR3 in (left) CD4⁺ T_{conv} (middle) CD4⁺ T_{reg}, and (right) CD8⁺ T cells from CD3G-mutated patients (n = 6) compared with controls (n = 8). * $P < 10^{-4}$.

TRB repertoire for properties of the CDR3 region that may associate with self-reactivity. In particular, the presence of hydrophobic amino acids at positions 6 and 7 of the TRB-CDR3 has been previously shown to promote self-reactivity in both mouse and human TCRs.¹⁴ Among T_{reg} cells, there were no significant changes in the relative hydrophobicity of amino acid positions 6 and 7 when comparing *CD3G*-mutated patients with controls (Figure 5D). However, the CD4⁺ T_{conv} cell population from *CD3G*-mutated patients contained significantly more reads enriched in hydrophobic amino acid residues at positions 6 and 7 of the *TRB* CDR3, compared with controls (Figure 5D), suggesting enrichment in self-reactive specificities among T_{conv} cells.

Discussion

Autoimmunity is a frequent complication of primary immune deficiencies. In particular, patients with genetic defects reducing the strength of TCR/CD3 signaling are exquisitely susceptible to the development of autoimmunity.¹¹ The number, diversity, and function of T_{reg} cells have each been demonstrated to be critical factors for sustaining immune tolerance,²⁶⁻³⁰ and changes in the relative suppressive potency of T_{reg} cells have been demonstrated to lead to autoimmunity in patients and mouse models with autoimmunity.³¹⁻³⁶ However, the relative contribution of abnormalities of T_{reg} cell number, diversity, and function to the development of clinical autoimmune disease in patients with

genetic defects of TCR signaling remains largely unknown. Herein, we performed a comprehensive analysis of peripheral T-cell number, phenotype, function, and repertoire in patients with disease-causing *CD3G* mutations. We have demonstrated that genetic defects in *CD3G* lead to impaired expression of CD3 and TCR molecules on the cell surface and overall defective T-cell response to mitogenic signals, which was especially profound upon stimulation with PHA. Furthermore, impaired cell division was also observed for CD8⁺ (but not CD4⁺) T cells from CD3 γ -deficient patients upon in vitro activation with anti-CD3. Similarly, a more profound impairment of T-cell proliferation in a mixed leukocyte reaction had been previously reported for CD8⁺ than for CD4⁺ T cells from CD3 γ -deficient patients.¹⁶

 T_{reg} cells are dependent on TCR signaling for their development and suppressive function. In this study, we have shown that defects of TCR/CD3 expression and signaling in patients with *CD3G* mutations are associated with a reduced proportion of T_{reg} cells, restriction of T_{reg} repertoire diversity, and impaired T_{reg} function. Previous studies have shown that diversity of TCR repertoire is maximal in naïve T cells, lower in T memory, and lowest in T_{EMRA} cells,^{37,38} and that among memory T cells, richness of T-cell repertoire is higher in CD4⁺ than in CD8⁺ T cells.³⁹ We observed a trend for reduced *TRB* repertoire diversity among patient-derived CD8⁺ T cells. This may reflect the accumulation of effector memory and T_{EMRA} CD8⁺ cells in

the patients. Although the reduced number of naïve T cells and the slightly reduced diversity of CD8⁺ T cells may play contribute to increased susceptibility to infections in CD3 γ deficiency, a significant restriction of repertoire diversity was observed only in the T_{reg} compartment, and was more prominent in patients P1 and P4. Interestingly, these were also the patients with the most severe autoimmune manifestations. Overall, these data strongly indicate that abnormalities of T_{reg} diversity play an important role in determining the severity of immune dysregulation in CD3 γ deficient patients, and suggest that similar abnormalities may be at play also in other forms of PID with defective TCR signaling.

Analysis of amino acid composition at positions 6 and 7 of the TRB-CDR3 disclosed a molecular signature indicative of increased self-reactivity in CD4 $^+$ T_{conv} cells. TCR affinity for peptides and TCR signaling strength dictate the fate of developing CD4⁺ T cells in the thymus medulla. In particular, single positive CD4⁺ cells that recognize with high affinity the selfpeptides that are presented by medullary thymic epithelial cells or by dendritic cells, undergo negative selection. Alternatively, self-reactive T cells may be diverted to natural T_{reg} cells.⁴⁰ This determines important differences in the antigen specificity of T_{reg} vs CD4⁺ T_{conv} cells, with the former being enriched for selfreactive specificities.¹⁴ Abnormalities of TCR signaling strength, and in particular reduced signaling as in patients with CD3y deficiency, may affect this process, leading to an increase in selfreactive clones within the CD4+ $T_{\rm conv}$ population. Our results showing enrichment for amino acids that promote self-reactivity, and reduced usage of those limiting self-reactivity at positions 6 and 7 of the TRB-CDR3 in CD4 $^+$ T_{conv} cells from CD3 γ -deficient patients are consistent with this hypothesis. We have recently documented similar abnormalities of TRB composition in CD4⁺ T_{conv} cells from patients with hypomorphic RAG mutations,⁴¹ another condition that is characterized by increased occurrence of autoimmune manifestations. In that case, however, it is likely that the altered fate of self-reactive CD4⁺ thymocytes are secondary to reduced expression of AIRE and of AIRE-dependent tissue-restricted peptides in the thymic medulla, but the ultimate effect would be similar, with an excess of self-reactive specificities in CD4 $^+$ T_{conv} cells that are exported to the periphery. Interestingly, no molecular signature of enhanced self-reactivity was found among CD8⁺ T cells. In this regard, the molecular mechanisms controlling T-cell tolerance in thymus medulla (including generation of T_{reg} cells from CD25⁺ FOXP3⁻ or CD25⁻ FOXP3⁺ progenitor cells) are better defined for CD4⁺ cells, but remain largely unknown for CD8⁺ cells.⁴²

Finally, although analysis of the T_{reg} TRB-CDR3 did not reveal significant changes in overall self-reactivity in CD3 γ -deficient patients compared with controls, reduced signaling via the TCR/CD3 complex may have important consequences for the survival and function of T_{reg} cells in the periphery. Indeed, the relative amount of TCR signaling is essential to maintaining survival and function of T_{reg} cells.^{19,20} We have demonstrated that T_{reg} cells from CD3 γ -deficient patients have defective

suppressive activity. Whether their survival is also affected, and whether this may contribute to the marked restriction of *TRB* repertoire in this subset, requires further investigation.

In summary, our data suggest a model whereby human CD3 γ deficiency impairs surface expression of CD3 and TCR molecules, reducing TCR signaling strength and thereby affecting T-cell development, selection, and fate. Ultimately, this results in enrichment for self-reactive specificities within CD4⁺ T_{conv} cells, which may undergo peripheral activation and expansion because of the reduced number and impaired function of T_{reg} cells. Altogether, these abnormalities create an environment that promotes the development of autoimmunity in this condition.

Acknowledgment

This work was supported by the Division of Intramural Research, National Institute of Allergy and Infectious Diseases, National Institutes of Health.

Authorship

Contribution: J.H.R. and O.M.D. performed the experiments, analyzed the data, and wrote the manuscript; B.D.S. and E.S.H. performed analysis of self-reactivity index; A.K.D., S.C.C., and Y.Y. performed experiments; L.A.H. contributed to the analysis of TRB repertoire; S.K., L.M.A., F.A.B., L.I.G.-G., S.N.G., H.K., C.Y., S.-Y.P., I.R., and J.R.R. provided patient samples and clinical information; and L.D.N. conceived the study, supervised all experiments, and wrote the manuscript.

Conflict-of-interest disclosure: The authors declare no competing financial interests.

ORCID profiles: S.K., 0000-0002-4351-8200; B.D.S., 0000-0002-3831-6751; A.K.D., 0000-0002-3432-3137; L.A.H., 0000-0001-6603-763X; Y.Y., 0000-0002-1695-8614; L.M.A., 0000-0001-9586-8539; F.A.B., 0000-0002-5553-2077; L.I.G.-G., 0000-0001-6917-8980; S.N.G., 0000-0002-8860-6132; C.Y., 0000-0002-0025-6335; S.-Y.P., 0000-0002-0158-8147; J.R.R., 0000-0001-8442-7762; L.D.N., 0000-0002-8335-0262.

Correspondence: Luigi D. Notarangelo, Laboratory of Clinical Immunology and Microbiology, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Building 10, Room 5-3950, 10 Centre Dr, Bethesda, MD 20892; e-mail: luigi.notarangelo2@nih.gov.

Footnotes

Submitted 23 February 2018; accepted 9 April 2018. Prepublished online as *Blood* First Edition paper, 13 April 2018; DOI 10.1182/blood-2018-02-835561.

*J.H.R. and O.M.D. contributed equally to this study.

†I.R., J.R.R., and L.D.N. contributed equally to this study.

The online version of this article contains a data supplement.

The publication costs of this article were defrayed in part by page charge payment. Therefore, and solely to indicate this fact, this article is hereby marked "advertisement" in accordance with 18 USC section 1734.

REFERENCES

- Call ME, Wucherpfennig KW. The T cell receptor: critical role of the membrane environment in receptor assembly and function. Annu Rev Immunol. 2005;23(1): 101-125.
- Alarcón B, Gil D, Delgado P, Schamel WW. Initiation of TCR signaling: regulation within CD3 dimers. *Immunol Rev.* 2003;191(1):38-46.
- Picard C, Bobby Gaspar H, Al-Herz W, et al. International Union of Immunological Societies: 2017 Primary Immunodeficiency

Diseases Committee Report on Inborn Errors of Immunity. *J Clin Immunol*. 2018;38(1): 96-128.

 Dadi HK, Simon AJ, Roifman CM. Effect of CD3delta deficiency on maturation of alpha/ beta and gamma/delta T-cell lineages in severe combined immunodeficiency. *N Engl J Med*. 2003;349(19):1821-1828.

- de Saint Basile G, Geissmann F, Flori E, et al. Severe combined immunodeficiency caused by deficiency in either the delta or the epsilon subunit of CD3. *J Clin Invest.* 2004;114(10): 1512-1517.
- Rieux-Laucat F, Hivroz C, Lim A, et al. Inherited and somatic CD3zeta mutations in a patient with T-cell deficiency. N Engl J Med. 2006; 354(18):1913-1921.
- Haks MC, Cordaro TA, van den Brakel JH, et al. A redundant role of the CD3 gammaimmunoreceptor tyrosine-based activation motif in mature T cell function. *J Immunol.* 2001;166(4):2576-2588.
- Arnaiz-Villena A, Timon M, Corell A, Perez-Aciego P, Martin-Villa JM, Regueiro JR. Brief report: primary immunodeficiency caused by mutations in the gene encoding the CD3gamma subunit of the T-lymphocyte receptor. N Engl J Med. 1992;327(8):529-533.
- Tokgoz H, Caliskan U, Keles S, Reisli I, Guiu IS, Morgan NV. Variable presentation of primary immune deficiency: two cases with CD3 gamma deficiency presenting with only autoimmunity. *Pediatr Allergy Immunol.* 2013; 24(3):257-262.
- Gokturk B, Keles S, Kirac M, et al. CD3G gene defects in familial autoimmune thyroiditis. Scand J Immunol. 2014;80(5):354-361.
- Notarangelo LD. Immunodeficiency and immune dysregulation associated with proximal defects of T cell receptor signaling. *Curr Opin Immunol.* 2014;31:97-101.
- Liston A, Enders A, Siggs OM. Unravelling the association of partial T-cell immunodeficiency and immune dysregulation. Nat Rev Immunol. 2008;8(7):545-558.
- Jabara HH, Ohsumi T, Chou J, et al. A homozygous mucosa-associated lymphoid tissue 1 (MALT1) mutation in a family with combined immunodeficiency. J Allergy Clin Immunol. 2013;132(1):151-158.
- Stadinski BD, Shekhar K, Gómez-Touriño I, et al. Hydrophobic CDR3 residues promote the development of self-reactive T cells. Nat Immunol. 2016;17(8):946-955.
- Torres PS, Alcover A, Zapata DA, et al. TCR dynamics in human mature T lymphocytes lacking CD3 gamma. J Immunol. 2003; 170(12):5947-5955.
- Timón M, Arnaiz-Villena A, Rodríguez-Gallego C, Pérez-Aciego P, Pacheco A, Regueiro JR. Selective disbalances of peripheral blood T lymphocyte subsets in human CD3 gamma deficiency. *Eur J Immunol.* 1993;23(7): 1440-1444.
- 17. Ozgür TT, Asal GT, Cetinkaya D, et al. Hematopoietic stem cell transplantation in a CD3 gamma-deficient infant with

inflammatory bowel disease. *Pediatr Transplant*. 2008;12(8):910-913.

- Feng Y, van der Veeken J, Shugay M, et al. A mechanism for expansion of regulatory T-cell repertoire and its role in self-tolerance. *Nature*. 2015;528(7580):132-136.
- Levine AG, Arvey A, Jin W, Rudensky AY. Continuous requirement for the TCR in regulatory T cell function. *Nat Immunol.* 2014; 15(11):1070-1078.
- Vahl JC, Drees C, Heger K, et al. Continuous T cell receptor signals maintain a functional regulatory T cell pool. *Immunity*. 2014;41(5): 722-736.
- Melichar HJ, Ross JO, Herzmark P, Hogquist KA, Robey EA. Distinct temporal patterns of T cell receptor signaling during positive versus negative selection in situ. *Sci Signal.* 2013; 6(297):ra92.
- Fu G, Rybakin V, Brzostek J, Paster W, Acuto O, Gascoigne NR. Fine-tuning T cell receptor signaling to control T cell development. *Trends Immunol.* 2014;35(7):311-318.
- Haks MC, Pépin E, van den Brakel JH, et al. Contributions of the T cell receptor-associated CD3gamma-ITAM to thymocyte selection. *J Exp Med.* 2002;196(1):1-13.
- Kasow KA, Chen X, Knowles J, Wichlan D, Handgretinger R, Riberdy JM. Human CD4+ CD25+ regulatory T cells share equally complex and comparable repertoires with CD4+CD25- counterparts. *J Immunol.* 2004; 172(10):6123-6128.
- Stewart JJ, Lee CY, Ibrahim S, et al. A Shannon entropy analysis of immunoglobulin and T cell receptor. *Mol Immunol.* 1997;34(15): 1067-1082.
- Bacchetta R, Passerini L, Gambineri E, et al. Defective regulatory and effector T cell functions in patients with FOXP3 mutations. *J Clin Invest*. 2006;116(6):1713-1722.
- Föhse L, Suffner J, Suhre K, et al. High TCR diversity ensures optimal function and homeostasis of Foxp3+ regulatory T cells. Eur J Immunol. 2011;41(11):3101-3113.
- Ferreira C, Singh Y, Furmanski AL, Wong FS, Garden OA, Dyson J. Non-obese diabetic mice select a low-diversity repertoire of natural regulatory T cells. Proc Natl Acad Sci USA. 2009;106(20):8320-8325.
- Bourgeois C, Stockinger B. CD25+CD4+ regulatory T cells and memory T cells prevent lymphopenia-induced proliferation of naive T cells in transient states of lymphopenia. J Immunol. 2006;177(7):4558-4566.
- Nishio J, Baba M, Atarashi K, et al. Requirement of full TCR repertoire for regulatory T cells to maintain intestinal homeostasis. *Proc Natl Acad Sci USA*. 2015;112(41): 12770-12775.

- 31. Longhi MS, Ma Y, Grant CR, et al. T-regs in autoimmune hepatitis-systemic lupus erythematosus/mixed connective tissue disease overlap syndrome are functionally defective and display a Th1 cytokine profile. J Autoimmun. 2013;41:146-151.
- 32. Liberal R, Grant CR, Holder BS, et al. In autoimmune hepatitis type 1 or the autoimmune hepatitis-sclerosing cholangitis variant defective regulatory T-cell responsiveness to IL-2 results in low IL-10 production and impaired suppression. *Hepatology*. 2015;62(3): 863-875.
- Valencia X, Yarboro C, Illei G, Lipsky PE. Deficient CD4+CD25high T regulatory cell function in patients with active systemic lupus erythematosus. J Immunol. 2007;178(4): 2579-2588.
- Humblet-Baron S, Sather B, Anover S, et al. Wiskott-Aldrich syndrome protein is required for regulatory T cell homeostasis. J Clin Invest. 2007;117(2):407-418.
- Fontenot JD, Rasmussen JP, Gavin MA, Rudensky AY. A function for interleukin 2 in Foxp3-expressing regulatory T cells [published correction appears in Nat Immunol. 2006;7:427]. Nat Immunol. 2005;6(11): 1142-1151.
- Allenspach E, Torgerson TR. Autoimmunity and primary immunodeficiency disorders. J Clin Immunol. 2016;36(suppl 1):57-67.
- 37. Oakes T, Heather JM, Best K, et al. Quantitative characterization of the T cell receptor repertoire of naïve and memory subsets using an integrated experimental and computational pipeline which is robust, economical, and versatile. *Front Immunol.* 2017;8: 1267.
- Qi Q, Liu Y, Cheng Y, et al. Diversity and clonal selection in the human T-cell repertoire. *Proc Natl Acad Sci USA*. 2014;111(36): 13139-13144.
- Wang C, Sanders CM, Yang Q, et al. High throughput sequencing reveals a complex pattern of dynamic interrelationships among human T cell subsets. *Proc Natl Acad Sci USA*. 2010;107(4):1518-1523.
- Perry JS, Hsieh CS. Development of T-cell tolerance utilizes both cell-autonomous and cooperative presentation of self-antigen. *Immunol Rev.* 2016;271(1):141-155.
- Rowe JH, Stadinski BD, Henderson LA, et al. Abnormalities of T-cell receptor repertoire in CD4⁺ regulatory and conventional T cells in patients with RAG mutations: Implications for autoimmunity. J Allergy Clin Immunol. 2017; 140(6):1739-1743.e7.
- 42. Abramson J, Anderson G. Thymic epithelial cells. Annu Rev Immunol. 2017;35(1):85-118.