

## THROMBOSIS AND HEMOSTASIS

## A novel role for the macrophage galactose-type lectin receptor in mediating von Willebrand factor clearance

Soracha E. Ward,<sup>1,\*</sup> Jamie M. O'Sullivan,<sup>1,\*</sup> Clive Drakeford,<sup>1</sup> Sonia Aguila,<sup>1</sup> Christopher N. Jondle,<sup>2</sup> Jyotika Sharma,<sup>2</sup> Padraic G. Fallon,<sup>3</sup> Teresa M. Brophy,<sup>1</sup> Roger J. S. Preston,<sup>1</sup> Paul Smyth,<sup>4</sup> Orla Sheils,<sup>4</sup> Alain Chion,<sup>1</sup> and James S. O'Donnell<sup>1,5</sup>

<sup>1</sup>Irish Centre for Vascular Biology, Molecular and Cellular Therapeutics, Royal College of Surgeons in Ireland, Dublin, Ireland; <sup>2</sup>Department of Basic Biomedical Sciences, University of North Dakota School of Medicine and Health Sciences, Grand Forks, ND; <sup>3</sup>Inflammation and Immunity Research Group, Trinity Translational Medicine Institute, and <sup>4</sup>Department of Histopathology, Trinity Translational Medicine Institute, Trinity College Dublin, St. James's Hospital, Dublin, Ireland; and <sup>5</sup>National Centre for Coagulation Disorders, St. James's Hospital, Dublin, Ireland

## KEY POINTS

- VWF sialylation modulates *in vivo* clearance through Ashwell-Morrell independent pathways.
- VWF binding to MGL plays a novel role in facilitating VWF clearance.


Previous studies have shown that loss of terminal sialic acid causes enhanced von Willebrand factor (VWF) clearance through the Ashwell-Morrell receptor (AMR). In this study, we investigated (1) the specific importance of *N*- vs *O*-linked sialic acid in protecting against VWF clearance and (2) whether additional receptors contribute to the reduced half-life of hyposialylated VWF.  $\alpha$ 2-3-linked sialic acid accounts for <20% of total sialic acid and is predominantly expressed on VWF *O*-glycans. Nevertheless, specific digestion with  $\alpha$ 2-3 neuraminidase ( $\alpha$ 2-3Neu-VWF) was sufficient to cause markedly enhanced VWF clearance. Interestingly, *in vivo* clearance experiments in dual  $VWF^{-/-}/Asgr1^{-/-}$  mice demonstrated enhanced clearance of  $\alpha$ 2-3Neu-VWF even in the absence of the AMR. The macrophage galactose-type lectin (MGL) is a C-type lectin that binds to glycoproteins expressing terminal *N*-acetylgalactosamine or galactose residues. Importantly, the markedly enhanced clearance of hyposialylated VWF in  $VWF^{-/-}/Asgr1^{-/-}$  mice was significantly attenuated in the presence of an anti-MGL inhibitory antibody. Furthermore, dose-dependent binding of human VWF to purified recombinant human MGL was confirmed using surface plasmon resonance. Additionally, plasma VWF:Ag levels were significantly elevated in  $MGL1^{-/-}$  mice compared with controls. Collectively, these findings identify MGL as a novel macrophage receptor for VWF that significantly contributes to the clearance of both wild-type and hyposialylated VWF. (*Blood*. 2018;131(8):911-916)

Similarly, genetic inactivation of ST3GalIV sialyltransferase in a transgenic mouse model causes enhanced VWF clearance.<sup>9</sup> Several studies have reported significantly reduced VWF sialylation levels in patients with type 1 VWD.<sup>7,9</sup> Furthermore, van Schooten et al reported an inverse correlation between aberrant sialylation of T antigen and plasma VWF:Ag levels, suggesting that *O*-linked sialylation on VWF may be of particular importance.<sup>7</sup>

## Introduction

Although substantial progress has been achieved in understanding von Willebrand factor (VWF) structure and function, the biological mechanisms underpinning VWF clearance from the plasma remain poorly understood.<sup>1</sup> Nevertheless, studies have demonstrated that enhanced VWF clearance plays an important role in the etiology of both type 1 and type 2 von Willebrand disease (VWD).<sup>1,2</sup> During biosynthesis, VWF undergoes complex posttranslational modification, including significant *N*- and *O*-linked glycosylation. Mass spectrometry studies have shown that sialylated biantennary complex-type chains constitute the commonest *N*-linked glycans expressed on VWF, whereas a disialylated core 1 tetrasaccharide structure (known as the T antigen) accounts for 70% of the total *O*-glycan population.<sup>3,4</sup> Thus, the majority of *N*- and *O*-linked glycans of human VWF are capped by negatively charged sialic acid residues.<sup>5</sup> In keeping with other plasma glycoproteins, terminal sialic acid expression plays an important role in protecting VWF against clearance.<sup>6,7</sup> Consequently, enzymatic removal of terminal sialylation from VWF has been associated with a markedly reduced plasma half-life *in vivo*.<sup>8</sup>

Current evidence suggests that the enhanced clearance of hyposialylated VWF occurs via the Ashwell-Morrell receptor (AMR).<sup>10</sup> This C-type lectin is expressed on hepatocytes and is composed of 2 transmembrane protein subunits (Asgr-1 and Asgr-2). Grewal et al previously demonstrated that plasma VWF clearance is significantly attenuated in *Asgr-1* knockout mice.<sup>10</sup> Nevertheless, important questions regarding the biological mechanisms through which VWF sialylation regulates its clearance *in vivo* remain unclear. In particular, the relative importance of *N*-linked vs *O*-linked sialylation in regulating physiological and/or pathological clearance of VWF has not been defined. In addition to the AMR, a number of other lectin receptors have been shown to bind with enhanced affinity to hyposialylated



**Figure 1. Clearance of hyposialylated VWF proceeds independently of AMR.** (A) To study the effects of N- and O-linked sialylation on VWF clearance, purified human pd-VWF was treated with either  $\alpha$ 2-3,6,8,9 or  $\alpha$ 2-3 neuraminidase. In vivo clearance for each glycoform was then assessed in *VWF<sup>-/-</sup>* mice and compared with that of wild-type pd-VWF. At each time point, residual circulating VWF concentration was determined by VWF:Ag enzyme-linked immunosorbent assay. All results are plotted as percentage residual VWF:Ag levels relative to the amount injected. Three to five mice were used per time point. Data are represented as mean  $\pm$  SEM. In some cases, the SEM cannot be seen because of its small size. (B) In the presence of ASOR, the enhanced in vivo clearance of both  $\alpha$ 2-3 Neu-VWF and  $\alpha$ 2-3,6,8,9 Neu-VWF was significantly attenuated ( $\alpha$ 2-3 Neu-VWF  $t_{1/2}$  =  $8.2 \pm 1.4$  minutes vs  $12.4 \pm 2.4$  minutes,  $P < .05$ ; and  $\alpha$ 2-3,6,8,9 Neu-VWF  $t_{1/2}$  =  $3.7 \pm 0.7$  minutes vs  $14.4 \pm 2.7$  minutes,  $P < .005$ , respectively). (C) To determine whether AMR-independent pathways contribute to the enhanced clearance of hyposialylated VWF, in vivo clearance studies were repeated in *VWF<sup>-/-</sup>/Asgr1<sup>-/-</sup>* mice. Importantly, the markedly enhanced clearance of both  $\alpha$ 2-3 Neu-VWF and  $\alpha$ 2-3,6,8,9 Neu-VWF was still evident in the absence of the AMR ( $t_{1/2}$  =  $8.2 \pm 0.6$  and  $3.2 \pm 0.4$  compared with  $50.6 \pm 2$  minutes for pd-VWF;  $P < .05$ ). Furthermore, the reduced half-life observed for  $\alpha$ 2-3 Neu-VWF (D) and  $\alpha$ 2-3,6,8,9 Neu-VWF (E) were not significantly different in the presence or absence of the AMR ( $\alpha$ 2-3 Neu-VWF  $t_{1/2}$  =  $8.2 \pm 1.4$  minutes vs  $8.2 \pm 0.6$  minutes,  $P = .96$ ; and  $\alpha$ 2-3,6,8,9 Neu-VWF  $t_{1/2}$  =  $3.7 \pm 0.7$  minutes vs  $3.2 \pm 0.4$  minutes,  $P = .42$ , respectively).




Figure 2.

glycoproteins.<sup>11</sup> In this study, we demonstrate a critical role for VWF O-linked sialylation in modulating in vivo clearance and further define a novel role for the macrophage galactose-type lectin (MGL) in regulating VWF clearance in a sialic acid-dependent manner.

## Study design

### Isolation and digestion of human plasma-derived VWF

As described in the supplemental Methods (available on the *Blood* Web site), plasma-derived VWF (pd-VWF) was purified from commercial concentrate Fandhi (Grifols, Barcelona, Spain) and subsequently treated with  $\alpha$ 2-3 neuraminidase (*Streptococcus pneumoniae*; Sigma Aldrich, Ireland) or  $\alpha$ 2-3,6,8,9 neuraminidase (*Arthrobacter ureafaciens*; New England Biolabs, United Kingdom) as previously described.<sup>5</sup> VWF glycan expression was analyzed using lectin enzyme-linked immunosorbent assays (see supplemental Methods and supplemental Figure 1).<sup>8</sup>

### VWF clearance studies

VWF<sup>-/-</sup> and *Asgr1*<sup>-/-</sup> mice on a C57BL/6J background were obtained from the Jackson Laboratory (Sacramento, CA) and crossbred to generate novel VWF<sup>-/-</sup>/*Asgr1*<sup>-/-</sup> double-knockout mice. *MGL1*<sup>-/-</sup> mice were also obtained from the Jackson Laboratory. Where indicated, clearance studies were repeated in the presence of either clodronate or asialo-orosomucoid (ASOR) as previously described.<sup>8,12</sup> Specific clearance studies were performed after inhibition of MGL using a polyclonal goat anti-mouse MGL1/2 antibody. All in vivo clearance experiments were performed as detailed in the supplemental Methods in accordance with the Health Product Regulatory Authority, Ireland.

### In vitro VWF binding studies

As described in the supplemental Methods, surface plasmon resonance (SPR) was used to evaluate MGL binding to VWF.<sup>13</sup> Briefly, purified pd-VWF was immobilized on a CM5 chip, and binding to recombinant MGL (R&D Systems, United Kingdom) was determined. Furthermore, proximity ligation assay (Duolink-PLA; Sigma Aldrich, Ireland) was performed to evaluate colocalization of VWF and MGL on THP1 macrophages.

### Data presentation and statistical analysis

Experimental data were analyzed with GraphPad Prism version 5.0 (GraphPad Software, San Diego, CA). Data were expressed as mean values  $\pm$  standard error of the mean (SEM). Data were

analyzed with Student unpaired 2-tailed t test, and *P* values of  $<.05$  were deemed significant.

## Results and discussion

### VWF sialylation modulates clearance through Ashwell-Morrell independent pathways

In keeping with previous studies, we observed that combined removal of N- and O-linked sialic acid by digestion with  $\alpha$ 2-3,6,8,9 neuraminidase resulted in markedly enhanced clearance of pd-VWF in VWF<sup>-/-</sup> mice (Figure 1A). Specific removal of  $\alpha$ 2-3-linked sialic acid was sufficient to markedly enhance VWF clearance ( $t_{1/2} = 9.0 \pm 1$  minutes;  $P < .05$ ) (Figure 1A). In fact, clearance of  $\alpha$ 2-3 Neu-VWF was almost as rapid as that of  $\alpha$ 2-3,6,8,9 Neu-VWF ( $t_{1/2} = 4.0 \pm 0.3$  minutes). This finding is interesting because  $\alpha$ 2-3-linked sialic acid is predominantly located on the O-linked glycans of human VWF and accounts for  $<20\%$  of total sialic acid expression.<sup>5</sup> Currently, the AMR is the only receptor described to regulate clearance of hyposialylated VWF.<sup>10</sup> We found that the enhanced clearance of  $\alpha$ 2-3 Neu-VWF and  $\alpha$ 2-3,6,8,9 Neu-VWF were both significantly attenuated in the presence of a hyposialylated inhibitor glycoprotein (ASOR) (Figure 1B). However, ASOR has a short plasma half-life and is not a specific AMR inhibitor.<sup>14</sup> Previous studies have shown that AMR demonstrates significantly greater affinity for exposed galactose residues on tri- and tetra-antennary galactoses (as present on VWF N-glycan) compared with terminal galactose moieties on mono- or biantennary galactoses (as present on VWF O-glycan).<sup>15,16</sup> We therefore hypothesized that other lectin receptors may contribute to the enhanced clearance of hyposialylated VWF and be of particular importance in modulating the effects of O-linked sialylation on VWF clearance. To address this, in vivo clearance studies were repeated in dual VWF<sup>-/-</sup>/*Asgr1*<sup>-/-</sup> knockout mice. Critically, we observed that markedly enhanced clearance of both  $\alpha$ 2-3 Neu-VWF and  $\alpha$ 2-3,6,8,9 Neu-VWF persisted in VWF<sup>-/-</sup>/*Asgr1*<sup>-/-</sup> mice ( $t_{1/2} = 8.2 \pm 0.6$  and  $3.2 \pm 0.4$  vs  $50.6 \pm 2$  minutes for pd-VWF;  $P < .05$ ) (Figure 1C). Furthermore, the enhanced clearance rates observed for  $\alpha$ 2-3 Neu-VWF and  $\alpha$ 2-3,6,8,9 Neu-VWF were not significantly different in the presence or absence of the AMR (Figure 1D-E). Collectively, these data confirm that reductions in N- and/or O-linked sialylation have major effects on VWF half-life and demonstrate that  $\alpha$ 2-3-linked sialic acid expressed on O-linked glycans may be of particular importance in regulating pd-VWF clearance. Furthermore, our findings suggest that previously unrecognized AMR-independent pathways contribute to the enhanced clearance of hyposialylated VWF in vivo.

**Figure 2. MGL facilitates VWF clearance in vivo.** To investigate other receptors and/or cell types that modulate the enhanced clearance of hyposialylated VWF,  $\alpha$ 2-3 Neu-VWF (A) and  $\alpha$ 2-3,6,8,9 Neu-VWF (B) clearance studies in VWF<sup>-/-</sup>/*Asgr1*<sup>-/-</sup> mice were repeated in the presence of ASOR, or following clodronate-induced macrophage depletion. The enhanced clearance of both  $\alpha$ 2-3 Neu-VWF and  $\alpha$ 2-3,6,8,9 Neu-VWF was still inhibited by ASOR (blue lines) even in the absence of the AMR ( $\alpha$ 2-3 Neu-VWF  $t_{1/2} = 8.2 \pm 1.4$  minutes vs  $16.8 \pm 1.6$  minutes,  $P < .005$ ; and  $\alpha$ 2-3,6,8,9 Neu-VWF  $t_{1/2} = 3.7 \pm 0.7$  minutes vs  $7.7 \pm 1.3$  minutes,  $P < .05$ , respectively). In addition, clodronate-induced macrophage depletion (green lines) also significantly attenuated the enhanced clearance of hyposialylated VWF ( $\alpha$ 2-3 Neu-VWF  $t_{1/2} = 8.2 \pm 1.4$  minutes vs  $24.0 \pm 1.1$  minutes,  $P < .05$ ; and  $\alpha$ 2-3,6,8,9 Neu-VWF  $t_{1/2} = 3.7 \pm 0.7$  minutes vs  $9.6 \pm 4.1$  minutes,  $P < .05$ , respectively). Three to 5 mice were used per time point, and data are represented as mean  $\pm$  SEM. (C) SPR was used to evaluate the binding of immobilized purified pd-VWF to recombinant human MGL. Dose-dependent binding were observed, with  $K_D$  (app) of  $18.4 \pm 3$   $\mu$ g/mL. (D) In mice, there are 2 homologs of human MGL, mMGL1 and mMGL2. Murine MGL1 shares significant sequence homology with human MGL and binds oligosaccharides with multiple terminal Gal residues including the T antigen. Interestingly, the markedly enhanced clearance of both  $\alpha$ 2-3 Neu-VWF and  $\alpha$ 2-3,6,8,9 Neu-VWF in VWF<sup>-/-</sup>/*Asgr1*<sup>-/-</sup> mice was significantly attenuated in the presence of an mMGL blocking antibody vs isotype immunoglobulin G (IgG) control antibody, respectively ( $\alpha$ 2-3 Neu-VWF  $t_{1/2} = 21.9 \pm 11.8$  minutes vs  $9.1 \pm 1.5$  minutes,  $P < .05$ ; and  $\alpha$ 2-3,6,8,9 Neu-VWF  $t_{1/2} = 24.4 \pm 8.1$  minutes vs  $5.7 \pm 2.1$  minutes,  $P < .05$ , respectively). (E) THP1 macrophages incubated with VWF demonstrated VWF-MGL colocalization detected by Duolink-PLA, visualized as red spots via immunofluorescence microscopy. No signal was observed from cells incubated with phosphate-buffered saline (PBS) alone. (F) Plasma VWF:Ag levels were significantly elevated in *MGL1*<sup>-/-</sup> mice compared with wild-type (WT) littermate controls ( $P < .05$ ). (G) The clearance of endogenous murine VWF in *MGL1*<sup>-/-</sup> mice was significantly attenuated compared with wild-type controls at all time points measured ( $P < .05$ ). (H) In vivo clearance of wild-type pd-VWF in VWF<sup>-/-</sup>/*Asgr1*<sup>-/-</sup> mice was significantly attenuated in the presence of an mMGL blocking antibody compared with isotype control IgG ( $t_{1/2} = 64.6 \pm 18.4$  minutes vs  $42.8 \pm 10.7$  minutes;  $P < .005$ ). A minimum of 3 mice were used per time point; data are plotted as mean  $\pm$  SEM.

## The macrophage galactose receptor regulates in vivo clearance of VWF

To investigate other receptors and/or cell types that modulate the enhanced clearance of hyposialylated VWF,  $\alpha$ 2-3 Neu-VWF and  $\alpha$ 2-3,6,8,9 Neu-VWF clearance studies in  $VWF^{-/-}/Asgr1^{-/-}$  mice were repeated in the presence of ASOR, or following clodronate-induced macrophage depletion (Figure 2A-B). The enhanced clearance of both  $\alpha$ 2-3 Neu-VWF and  $\alpha$ 2-3,6,8,9 Neu-VWF was still inhibited by ASOR even in the absence of the AMR. Interestingly, in vivo macrophage depletion also significantly attenuated the enhanced clearance of hyposialylated VWF. Finally, in vitro binding studies demonstrated enhanced binding of asialo-VWF to differentiated THP1 macrophages (supplemental Figure 2). Collectively, these data demonstrate that additional asialo-receptors, at least in part expressed on macrophages, regulate the enhanced clearance of hyposialylated VWF in vivo.

MGL is a C-type lectin receptor expressed as a homo-oligomer on antigen-presenting cells such as macrophages and dendritic cells (supplemental Figure 3).<sup>17</sup> The carbohydrate recognition domain of MGL binds with high affinity to glycoproteins expressing terminal *N*-acetylglucosamine or galactose (Gal) residues, and thus MGL can regulate glycoprotein endocytosis.<sup>18-20</sup> MGL binding to oligosaccharide chains is attenuated by terminal sialylation.<sup>21</sup> Importantly, given the putative role of VWF O-linked glycans in modulating clearance, MGL also recognizes the T antigen.<sup>22</sup> In mice, there are 2 homologs of human MGL, mMGL1 and mMGL2.<sup>23</sup> Murine MGL1 shares significant sequence homology with human MGL and has been shown to bind oligosaccharides with terminal Gal residues including the so called T antigen.<sup>23</sup> Of note, previous studies have demonstrated that ~70% of the O-glycans of VWF are composed of this sialylated tumor-associated T antigen structure.<sup>4,7</sup> Interestingly, the enhanced clearance of both  $\alpha$ 2-3 Neu-VWF and  $\alpha$ 2-3,6,8,9 Neu-VWF in  $VWF^{-/-}/Asgr1^{-/-}$  mice was significantly attenuated in the presence of anti-mMGL1/2 inhibitory antibody, suggesting a novel role for MGL in regulating macrophage-mediated clearance of hyposialylated VWF (Figure 2D). Importantly, we observed dose-dependent binding of human pd-VWF to purified recombinant human MGL using SPR (Figure 2C). Moreover, Duolink-PLA analysis demonstrated that VWF colocalizes with MGL on the surface of THP1 macrophages, as indicated by the distinct red fluorescent dots (Figure 2E). Plasma VWF:Ag levels were significantly elevated in  $MGL1^{-/-}$  mice compared with wild-type controls ( $152.6 \pm 15.7\%$  vs  $100 \pm 16.9\%$ ;  $P < .05$ ) (Figure 2F). Furthermore, in vivo clearance of endogenous murine VWF was attenuated in  $MGL1^{-/-}$  mice (Figure 2G), suggesting that MGL-mediated VWF clearance is important even in the presence of AMR. Finally, clearance of pd-VWF in  $VWF^{-/-}/Asgr1^{-/-}$  mice was attenuated in the presence of mMGL1/2 inhibitory antibody (Figure 2H). Collectively, these findings reveal MGL as a novel macrophage lectin receptor for VWF that contributes to the clearance of both wild-type

and hyposialylated VWF. Further studies will be required to determine the importance of MGL compared with other recently described receptors involved in regulating VWF clearance.<sup>24</sup> Nevertheless, the role of MGL in modulating VWF clearance has direct translational relevance in that quantitative variations in *N*- and *O*-linked sialylation have been described in patients with type 1 VWD.<sup>7,9</sup> In addition, desialylation of VWF has been described with glycoprotein ageing in plasma<sup>25</sup> and can also occur during infections with specific pathogens that are associated with significantly enhanced neuraminidase activity (eg, *Streptococcus pneumoniae*).<sup>10</sup>

## Acknowledgments

The authors thank Nico van Rooijen of the Foundation Clodronate Liposomes (Haarlem, The Netherlands) for generously providing the liposome-clodronate.

This work was supported by a Science Foundation Ireland Principal Investigator Award (11/PI/1066, J.S.O.).

## Authorship

Contribution: S.E.W., J.M.O., S.A., C.D., C.N.J., J.S., P.G.F., T.M.B., P.S., O.S., and A.C. performed experiments; S.E.W., J.M.O., S.A., C.D., C.N.J., J.S., P.G.F., T.M.B., R.J.S.P., O.S., A.C., and J.S.O. designed the research and analyzed the data; and all authors were involved in writing and reviewing the manuscript.

Conflict-of-interest disclosure: J.S.O. has served on the speaker's bureau for Baxter, Bayer, Novo Nordisk, Boehringer Ingelheim, Leo Pharma, and Octapharma; has served on the advisory boards of Baxter, Bayer, Octapharma CSL Behring, Daiichi Sankyo, Boehringer Ingelheim, and Pfizer; and has received research grant funding awards from Baxter, Bayer, Pfizer, and Novo Nordisk. The remaining authors declare no competing financial interests.

Correspondence: Jamie M. O'Sullivan, Department of Molecular and Cellular Therapeutics, Irish Centre for Vascular Biology, Royal College of Surgeons in Ireland, 123 St Stephen's Green, Dublin 2, Ireland; e-mail: jamieosullivan@rcsi.ie.

## Footnotes

Submitted 6 June 2017; accepted 12 December 2017. Prepublished online as *Blood* First Edition paper, 27 December 2017; DOI 10.1182/blood-2017-06-787853.

\*S.E.W. and J.M.O. contributed equally to this study.

The online version of this article contains a data supplement.

There is a *Blood* Commentary on this article in this issue.

The publication costs of this article were defrayed in part by page charge payment. Therefore, and solely to indicate this fact, this article is hereby marked "advertisement" in accordance with 18 USC section 1734.

## REFERENCES

1. Casari C, Lenting PJ, Wohner N, Christophe OD, Denis CV. Clearance of von Willebrand factor. *J Thromb Haemost*. 2013;11(suppl 1):202-211.
2. Rawley O, O'Sullivan JM, Chion A, et al. von Willebrand factor arginine 1205 substitution results in accelerated macrophage-

dependent clearance in vivo. *J Thromb Haemost*. 2015;13(5):821-826.

3. Canis K, McKinnon TAJ, Nowak A, et al. Mapping the N-glycome of human von Willebrand factor. *Biochem J*. 2012;447(2):217-228.
4. Canis K, McKinnon TAJ, Nowak A, et al. The plasma von Willebrand factor O-glycome

comprises a surprising variety of structures including ABH antigens and disialosyl motifs. *J Thromb Haemost*. 2010;8(1):137-145.

5. McGrath RT, McKinnon TAJ, Byrne B, et al. Expression of terminal alpha2-6-linked sialic acid on von Willebrand factor specifically enhances proteolysis by ADAMTS13. *Blood*. 2010;115(13):2666-2673.

6. Sodetz JM, Pizzo SV, McKee PA. Relationship of sialic acid to function and in vivo survival of human factor VIII/von Willebrand factor protein. *J Biol Chem*. 1977;252(15): 5538-5546.
7. van Schooten CJM, Denis CV, Lisman T, et al. Variations in glycosylation of von Willebrand factor with O-linked sialylated T antigen are associated with its plasma levels. *Blood*. 2007; 109(6):2430-2437.
8. O'Sullivan JM, Aguila S, McRae E, et al. N-linked glycan truncation causes enhanced clearance of plasma-derived von Willebrand factor. *J Thromb Haemost*. 2016;14(12): 2446-2457.
9. Ellies LG, Ditto D, Levy GG, et al. Sialyltransferase ST3Gal-IV operates as a dominant modifier of hemostasis by concealing asialoglycoprotein receptor ligands. *Proc Natl Acad Sci USA*. 2002;99(15): 10042-10047.
10. Grewal PK, Uchiyama S, Ditto D, et al. The Ashwell receptor mitigates the lethal coagulopathy of sepsis. *Nat Med*. 2008;14(6): 648-655.
11. Preston RJS, Rawley O, Gleeson EM, O'Donnell JS. Elucidating the role of carbohydrate determinants in regulating hemostasis: insights and opportunities. *Blood*. 2013; 121(19):3801-3810.
12. Chion A, O'Sullivan JM, Drakeford C, et al. N-linked glycans within the A2 domain of von Willebrand factor modulate macrophage-mediated clearance. *Blood*. 2016;128(15): 1959-1968.
13. O'Sullivan JM, Jenkins PV, Rawley O, et al. Galectin-1 and galectin-3 constitute novel-binding partners for factor VIII. *Arterioscler Thromb Vasc Biol*. 2016;36(5):855-863.
14. Wall DA, Wilson G, Hubbard AL. The galactose-specific recognition system of mammalian liver: the route of ligand internalization in rat hepatocytes. *Cell*. 1980; 21(1):79-93.
15. Ashwell G, Harford J. Carbohydrate-specific receptors of the liver. *Annu Rev Biochem*. 1982;51(1):531-554.
16. Lodish HF. Recognition of complex oligosaccharides by the multi-subunit asialoglycoprotein receptor. *Trends Biochem Sci*. 1991; 16(10):374-377.
17. Higashi N, Morikawa A, Fujioka K, et al. Human macrophage lectin specific for galactose/N-acetylgalactosamine is a marker for cells at an intermediate stage in their differentiation from monocytes into macrophages. *Int Immunol*. 2002;14(6):545-554.
18. van Vliet SJ, van Liempt E, Saeland E, et al. Carbohydrate profiling reveals a distinctive role for the C-type lectin MGL in the recognition of helminth parasites and tumor antigens by dendritic cells. *Int Immunol*. 2005; 17(5):661-669.
19. van Vliet SJ, Aarnoudse CA, Broks-van den Berg VC, Boks M, Geijtenbeek TB, van Kooyk Y. MGL-mediated internalization and antigen presentation by dendritic cells: a role for tyrosine-5. *Eur J Immunol*. 2007;37(8): 2075-2081.
20. Oo-puthinan S, Maenuma K, Sakakura M, et al. The amino acids involved in the distinct carbohydrate specificities between macrophage galactose-type C-type lectins 1 and 2 (CD301a and b) of mice. *Biochim Biophys Acta*. 2008; 1780(2):89-100.
21. Yamamoto K, Ishida C, Shinohara Y, et al. Interaction of immobilized recombinant mouse C-type macrophage lectin with glycopeptides and oligosaccharides. *Biochemistry*. 1994;33(26):8159-8166.
22. Mortezaei N, Behnken HN, Kurze AK, et al. Tumor-associated Neu5Ac-Tn and Neu5Gc-Tn antigens bind to C-type lectin CLEC10A (CD301, MGL). *Glycobiology*. 2013;23(7): 844-852.
23. Tsuiji M, Fujimori M, Ohashi Y, et al. Molecular cloning and characterization of a novel mouse macrophage C-type lectin, mMGL2, which has a distinct carbohydrate specificity from mMGL1. *J Biol Chem*. 2002;277(32): 28892-28901.
24. Pipe SW, Montgomery RR, Pratt KP, Lenting PJ, Lillicrap D. Life in the shadow of a dominant partner: the FVIII-VWF association and its clinical implications for hemophilia A. *Blood*. 2016;128(16):2007-2016.
25. Yang WH, Aziz PV, Heithoff DM, Mahan MJ, Smith JW, Marth JD. An intrinsic mechanism of secreted protein aging and turnover. *Proc Natl Acad Sci USA*. 2015;112(44): 13657-13662.