#### THROMBOSIS AND HEMOSTASIS

# Dysfunctional endogenous FIX impairs prophylaxis in a mouse hemophilia B model

Brian Cooley,<sup>1</sup> George J. Broze Jr,<sup>2</sup> David M. Mann,<sup>3</sup> Feng-Chang Lin,<sup>4</sup> Lee G. Pedersen,<sup>5</sup> and Darrel W. Stafford<sup>6</sup>

<sup>1</sup>Department of Pathology and Laboratory Medicine, University of North Carolina, Chapel Hill, NC; <sup>2</sup>Department of Medicine, Division of Medicine, Washington University in St. Louis, St. Louis, MO; <sup>3</sup>Mann BioConsulting, Gaithersburg, MD; <sup>4</sup>Department of Biostatistics, <sup>5</sup>Department of Chemistry, and <sup>6</sup>Department of Biology and Pathology, University of North Carolina at Chapel Hill, NC

#### KEY POINTS

- An endogenous, dysfunctional (CRM<sup>+</sup>)
  FIX molecule affects prophylactic FIX efficacy.
- Recovery studies indicate the amount of extravascular, Col4bound FIX is several fold greater than the FIX in plasma.

Factor IX (FIX) binds to collagen IV (Col4) in the subendothelial basement membrane. In hemophilia B, this FIX-Col4 interaction reduces the plasma recovery of infused FIX and plays a role in hemostasis. Studies examining the recovery of infused BeneFix (FIX<sub>WT</sub>) in null (cross-reactive material negative, CRM<sup>-</sup>) hemophilia B mice suggest the concentration of Col4 readily available for binding FIX is ~405 nM with a 95% confidence interval of 374 to 436 nM. Thus, the vascular cache of FIX bound to Col4 is several-fold the FIX level measured in plasma. In a mouse model of prophylactic therapy (testing hemostasis by saphenous vein bleeding 7 days after infusion of 150 IU/kg FIX), FIX<sub>WT</sub> and the increased half-life FIXs Alprolix (FIX<sub>FC</sub>) and Idelvion (FIX<sub>Alb</sub>) produce comparable hemostatic results in CRM<sup>-</sup> mice. In bleeding CRM<sup>-</sup> hemophilia B mice, the times to first clot at a saphenous vein injury site after the infusions of the FIX agents are significantly different, at FIX<sub>WT</sub> < FIX<sub>FC</sub> < FIX<sub>Alb</sub>. Dysfunctional forms of FIX, however, circulate in the majority of patients with hemophilia B (CRM<sup>+</sup>). In the mouse

prophylactic therapy model, none of the FIX products improves hemostasis in CRM<sup>+</sup> mice expressing a dysfunctional FIX, FIX<sub>R333Q</sub>, that nevertheless competes with infused FIX for Col4 binding and potentially other processes involving FIX. The results in this mouse model of CRM<sup>+</sup> hemophilia B demonstrate that the endogenous expression of a dysfunctional FIX can deleteriously affect the hemostatic response to prophylactic therapy. (*Blood.* 2019;133(22):2445-2451)

# Introduction

Individuals with hemophilia B are deficient in coagulation factor IX (FIX). In the body, FIX distributes between plasma and another readily accessible extravascular location. This extravascular binding initially attributed to an unknown molecule on the surface of the endothelium,<sup>1,2</sup> was subsequently shown to be type IV collagen (Col4).<sup>3</sup> Col4 is the only collagen restricted exclusively to, and a prominent component of, the basement membrane of all cells.<sup>4</sup> The equilibration between FIX in plasma and FIX bound to vascular Col4 is fast and reversible.<sup>5,6</sup> Therefore, the recovery in plasma of FIX infused into a patient with hemophilia B is only 40% to 50% of that anticipated in the absence of the vascular Col4 binding sites.<sup>7</sup>

We demonstrated that the  $\gamma$ -carboxyglutamic acid (Gla) domain of FIX is solely responsible for its interaction with Col4,<sup>8</sup> and that lysine at position 5 and valine at position 10 within the Gla domain are crucial for this interaction. For example, mutations at lysine 5 (FIX<sub>K5A</sub>) or valine 10 (FIX<sub>V10K</sub>) substantially reduce the affinity of FIX for Col4, whereas FIX<sub>K5R</sub> binds Col4 with 3- to 5-fold increased affinity.<sup>9</sup> Remarkably, all these tested mutant FIX molecules have normal or increased activity in in vitro phospholipiddependent coagulation assays. That the extravascular binding of FIX to Col4 plays a functional role in hemostasis is demonstrated by 3 observations: postinfusion recovery in mice of FIX<sub>K5A</sub> > FIX<sub>WT</sub> > FIX<sub>K5R</sub>,<sup>10</sup> bleeding in FIX null mice after a saphenous vein injury is significantly less 7 days after infusion of FIX<sub>K5R</sub> than after infusion of FIX<sub>K5A</sub> (when plasma FIX levels are undetectable),<sup>11</sup> and knock-in mice expressing FIX<sub>K5A</sub> have a clear, modest bleeding diathesis.<sup>12,13</sup>

There is considerable interpatient variability in the recovery of infused FIX in patients with hemophilia B, leading to the recommendation that FIX recovery be empirically determined in individual patients. Missense mutations in the FIX gene occur in 75% of patients with hemophilia B and in 60% of those with severe hemophilia B.<sup>14,15</sup> These missense mutations frequently associate with the production of dysfunctional FIX proteins that are detectable by immunologic antigen assays as cross-reactive material (CRM<sup>+</sup>). Therefore, most patients with hemophilia B are CRM<sup>+</sup>, rather than null for FIX expression (CRM<sup>-</sup>). There is evidence for enhanced plasma recovery of FIX in some CRM<sup>+</sup> patients compared with CRM<sup>-</sup> patients.<sup>16-18</sup> Increased recovery is expected if the endogenous, dysfunctional FIX occupies Col4 binding sites, thereby competing with the infused FIX for binding to Col4.

Here, we compare the hemostatic effects of a variety of FIX products in mouse models of FIX CRM^- (null), and CRM^+

hemophilia B. Empiric results for the recovery of human FIX in hemophilia mice were used to estimate the quantity of readily accessible vascular Col4 capable of binding FIX.

# Methods

## Mice

Mouse experiments were approved by the Institutional Animal Care and Use Committee at the University of North Carolina at Chapel Hill and followed Public Health Service guidelines for animal care and use. FIX null (CRM<sup>-</sup>)<sup>19</sup> and hFIX<sub>R333Q</sub> knock-in<sup>6</sup> mice (CRM<sup>+</sup>) with hemophilia B and normal control mice from Jackson Labs were all in the C57BL/6 genetic background.

## Proteins

The Pfizer Corporation provided BeneFix (FIX<sub>WT</sub>). Alprolix (FIX<sub>FC</sub>) and Idelvion (FIX<sub>Alb</sub>) were from Diversified Biologicals (Miami, FL). The proteins were reconstituted per the manufacturer's instructions immediately before infusion, and a new vial was used for each experiment.<sup>20</sup> FXIa was from Haematologic Technologies, Essex Junction, VT.

#### Human FIX ELISA

FIX was quantified by sandwich enzyme-linked immunosorbent assay (ELISA) performed at Green Mountain Antibodies (Burlington, VT). Ninety-six-well plates were coated overnight at 4°C with 100  $\mu$ L per well of a mouse monoclonal antibody to the heavy chain of human FIX (GMA-102) at 2 µg/mL in 0.2 M in carbonate-bicarbonate buffer at pH 9.4 (Pierce #28382). Coated plates were blocked for 1 hour at 4°C with blocking buffer (0.1% bovine serum albumin, 20 mM sodium phosphate, 0.15 M sodium chloride, 0.05% Tween20), 300  $\mu$ L per well. Plates were washed 3 times in PBST (20 mM sodium phosphate at pH 7.4, 0.15 M sodium chloride, 0.05% Tween20), and serial dilutions of mouse plasma in blocking buffer (100 µL per well) were applied to blocked plates for 1 hour at room temperature. After washing as described earlier, plates were incubated for 1 hour at room temperature with 100 µL per well biotinylated mouse monoclonal antibody to the light chain of FIX (GMA-184-BIO) at 1 µg/mL in blocking buffer. Plates were washed as described earlier and incubated for 30 minutes at room temperature with 100 µL per well streptavidin HRP (Jackson ImmunoResearch, catalog #016-030-084) diluted 1:5000 in blocking buffer. After a final washing step, plates were developed with 100 µL per well o-phenylenediamine dihydrochloride in the dark for 5 minutes, stopped with 50  $\mu L$  per well 2 M  $H_2SO_4,$  and read at 490 nm on a microplate reader. FIX concentrations in mouse plasma samples were determined by comparison with a standard curve produced with FIX (Enzyme Research Laboratories, HFIX 1009) in human FIX-deficient plasma (George King Biomedical 0900). All samples were tested at least twice, and the mean values were reported.

# Estimation of total Col4 available for binding human $\ensuremath{\mathsf{FIX}_{\mathsf{WT}}}$ in FIX null mice

Blood samples were obtained from the saphenous vein 10 minutes after intravenous infusion of variable quantities of  $FIX_{WT}$ . Our calculations assume that the FIX remaining in the plasma is that not bound to Col4, and that the FIX lost during the initial 10 minutes has formed a complex with Col4. In our model, concentrations of infused FIX, plasma FIX, and Col4-bound FIX are calculated as moles



Figure 1. Estimation of vascular Col4 available for binding FIX. Blood samples were taken 10 minutes after infusion of different doses (8, or more, mice per dose) of FIX<sub>WT</sub> into CRM<sup>-</sup> mice. The data were fit to FIXt-C (the quadratic equation for equilibrium) and were plotted as the total infused FIX (x-axis) vs measured plasma FIX (y-axis). The empirically determined FIX concentrations were determined by ELISA (*Methods*). The plotted line is the fit of the experimental data (FIXt-C), where C is the calculated complex. The error bars represent the standard deviation of the mean for 8 or more mice for each level of infused FIX. The 2 light red points in the figure represent data collected in our laboratory about 5 years ago, and is only included as in internal check. The table above the figure represents the statistics of the quadratic fit.

per plasma volume.<sup>21</sup> The data were fit to FIXt-C by the quadratic equation for chemical equilibrium:

$$\begin{split} C &= \frac{1}{2} \bigg( Kd + FIXt + Col4t \\ &- \sqrt{-4 \, x \, FIXt \, x \, Col4t + \left(Kd + FIXt + Col4t\right)^2} \, \bigg) \end{split}$$

where C represents the calculated complex between FIX and Col4, FIXt is total FIX, Col4t is total Col4 available to bind FIX,  $K_d$  is the equilibrium dissociation constant for the FIX-Col4 interaction, and the plasma concentration is FIXt-C (FIXt is calculated as though all infused FIX remained in the plasma volume). The least squares fit was obtained using Mathematica version 11.3, by several different methods, Quasi-Newton, PrincipalAxis, Levenburg-Marquardt, and GradientAxis; all yielded the same results. The error bars in Figure 1 represent the standard deviations of the mean calculated individually from all the data points for each infused FIX concentration (n = at least 8). The confidence interval was calculated as described.<sup>22</sup>

#### Saphenous vein bleeding assay

The saphenous vein bleeding assay (SVBA) was used to assess hemostasis as previously described.<sup>23</sup> FIX was infused into the left saphenous vein of hemophilia B mice and, after 7 days (in most cases), the right saphenous vein was used for the assay. In the SVBA, after initial vein injury, each time a clot forms it is mechanically disrupted, and the number of clots that occur over the course of 30 minutes recorded. An increased number of clots equates with better hemostasis.

## Times to first clot

The right saphenous vein of CRM<sup>-</sup> mice was injured as described for the SVBA, and 15 minutes was allowed for any baseline clotting to occur. In CRM<sup>-</sup> mice, clot formation ceases by 8 minutes postinjury; any clots formed during the first 8 minutes were physically removed. Fifteen minutes after the initial injury, the different FIX agents were infused into the left saphenous vein and the time until the first clot formed at the right saphenous vein injury site, determined during a 20-minute window. There is insufficient blood loss during this time to affect clotting.

### **Activation of FIX molecules**

FIX was activated with FXIa. To remove the FXIa activator, the generated FIXas were isolated by binding to a HiTrap Q column, washing with 250 mM salt, and eluting with 10 mM CaCl<sub>2</sub> in 250 mM salt (method courtesy of Dougald Monroe, University of North Carolina).

## Statistics

The Mann-Whitney test was used to compare SVBA results between groups. The statistics for samples comparing CRM<sup>+</sup> and CRM<sup>-</sup> mice used a median test with *P* value derived by a Bootstrap method<sup>24</sup> that adjusts for the different backgrounds. Except for the Bootstrap method, all statistical calculations were done with Mathematica, version 11.3.

# Results

# Estimation of vascular Col4 available for FIX binding

Previous work demonstrated that the rapid initial disappearance of FIX<sub>WT</sub> from plasma after infusion into a CRM<sup>-</sup> hemophilia B mouse<sup>6</sup> is predominantly a result of its binding to extravascular Col4.<sup>10,11</sup> There is a significant amount of Col4 in the body, but the amount capable of binding to FIX is unknown. We therefore estimated the amount of Col4 readily available for binding plasma FIX by infusing increasing doses of FIX (8-20 mice per dose) and taking plasma samples after 10 minutes for ELISA analysis. The following assumptions were made: a fraction of total body Col4 is readily accessible to the plasma;<sup>5,6</sup> the binding of rFIX to Col4 has 1:1 stoichiometry, is reversible, and reaches near equilibrium by 10 minutes; the plasma volume in the mouse is 40 mL/kg<sup>21</sup>; and clearance of FIX from the plasma through other mechanisms other than by binding Col4 is negligible within the first 10 minutes after infusion.

Mathematica (version 11.3) NonLinearModelFit was used to fit the data to the quadratic equation for chemical equilibrium,<sup>22,25</sup> and the total Col4 (Col4t) available for binding FIX and the in vivo value for  $K_d$  were estimated. The best least squares fits for Col4t and  $K_d$  were predicted to be 405 and 40 nM, respectively, with 95% confidence intervals of 342 to 467 and -7.1 to 87.0 nM (Figure 1). If, after simultaneous fitting of both parameters to the data, the result of Col4t is input into the quadratic equation and again fit to the data, the same parameters are achieved. However, *P* values and confidence intervals are much improved. For example, if 404 nM is input as Col4t, the predicted confidence interval for  $K_d$  is 16.5 to 63.5



Figure 2. Comparison of prophylactic FIX<sub>WT</sub> and FIX<sub>Alb</sub> treatment in CRM<sup>-</sup> mice. SVBA 7 days after infusion of FIX<sub>WT</sub> (green) or FIX<sub>Alb</sub> (blue) into CRM<sup>-</sup> mice at 50, 100, 150, and 200 IU/kg. Results for untreated CRM<sup>-</sup> (purple) and WT (red) mice are also shown. There is no significant difference between the median SVBA values for the FIX agents. Box plot rendition of the results showing the minimum, first quartile, median (waist), third quartile, and maximum values. KO, knockout.

## Hemostatic efficacy of FIX<sub>Alb</sub> in CRM<sup>-</sup> mice

Recombinant FIX products with extended plasma half-lives are marketed to simplify prophylactic therapy in patients with hemophilia B. These molecules include FIX<sub>FC</sub> (Alprolix, a fusion of FIX and the Fc domain of immunoglobulin G)<sup>26</sup> and FIX<sub>Alb</sub> (Idelvion, a fusion of FIX and albumin).<sup>27</sup> We previously demonstrated that the hemostatic effects of FIX<sub>WT</sub> and FIX<sub>FC</sub> are equivalent in CRM<sup>-</sup> hemophilia B mice.<sup>11,12</sup> Here, similar studies were performed using FIX<sub>Alb</sub> to test whether other longer circulating FIX's behaved similarly. Similar to FIX<sub>FC</sub>, FIX<sub>Alb</sub> is not statistically superior to FIX<sub>WT</sub> in CRM<sup>-</sup> mice at any of the doses tested (Figure 2).

# Effect of endogenous, dysfunctional (CRM<sup>+</sup>) FIX on hemostasis after FIX<sub>WT</sub> infusion in mice

The majority of patients with hemophilia B are CRM<sup>+</sup>: that is. they express a dysfunctional FIX protein. For example, 69 of the individuals with hemophilia B in the FIX variant database<sup>15</sup> have the missense mutation R333Q. These patients generally have functional and antigenic levels of FIX in plasma of less than 3% and 100%, respectively, a moderate to severe bleeding diathesis, and a form of dysfunctional FIX that is expected to interact with Col4 normally. We hypothesized that although a dysfunctional FIX that binds Col4 would increase the plasma recovery of infused FIX, it would decrease the amount of infused FIX bound to Col4, and thereby potentially reduce hemostatic effectiveness.<sup>13</sup> To test this hypothesis, hemostasis was compared 7 days after the infusion of FIX<sub>WT</sub> in CRM<sup>-</sup> and hFIX(R333Q) knock-in CRM<sup>+</sup> hemophilia B mice. These CRM<sup>+</sup> mice are reported to have a circulating hFIX(R333Q) plasma level of 1.2  $\mu$ g/mL (21.4 nM), and similar to their human counterpart, a severe hemophilia phenotype.<sup>6</sup> As predicted, the results (Figure 3) confirm a deleterious effect of the endogenous, dysfunctional FIX on hemostasis in  $CRM^+$  mice when compared with the same dose of  $FIX_{WT}$  in CRM<sup>-</sup> mice (median SVBA values of 4 and 12, respectively; *P* < .001).



Figure 3. Reduced efficacy of prophylactic FIX<sub>WT</sub> treatment in CRM<sup>+</sup> compared with CRM<sup>-</sup> mice. SVBA evaluation of hemostasis 7 days after the infusion of 150 IU/kg of FIX<sub>WT</sub> in CRM<sup>-</sup> and CRM<sup>+</sup> mice. Baseline SVBA in untreated CRM<sup>-</sup> (black) and CRM<sup>+</sup> (orange) mice, CRM<sup>+</sup> mice infused with FIX<sub>WT</sub> (green), CRM<sup>-</sup> mice infused with FIX<sub>WT</sub> (light green), and wild-type mice (red). Box plot rendition of the results showing the minimum, first quartile, median (waist), third quartile, and maximum values.

# Comparison of FIX<sub>WT</sub>, FIX<sub>FC</sub>, and FIX<sub>Alb</sub> in CRM<sup>+</sup> mice

Figure 4A shows that 7 days after infusion of 150 IU/kg (340 nM) into the CRM<sup>+</sup> mice, FIX<sub>WT</sub>, FIX<sub>FC</sub>, FIX<sub>Alb</sub> did not produce SVBA results significantly different from those of untreated CRM<sup>+</sup> mice. In addition, FIX<sub>FC</sub> and FIX<sub>Alb</sub> were not statistically different from  $FIX_{WT}$ , nor were they different from each other. The observation that 6 of the 20 CRM $^+$  mice infused with FIX<sub>FC</sub> had SVBA results within the range for WT mice is notable, suggesting that FIX<sub>FC</sub> might be superior to the other FIX molecules in CRM<sup>+</sup> mice. This led us to investigate whether the potential benefits of  $FIX_{FC}$  or  $FIX_{Alb}$ , compared with  $FIX_{WT}$ , would become apparent at higher doses in CRM<sup>+</sup> mice. Figure 4B shows that hemostasis in CRM<sup>+</sup> mice 7 days after infusion of very high doses of  $FIX_{FC}$ (500 IU/kg) was significantly better than that in untreated CRM<sup>+</sup> mice (median SVBA, 10.5 and 3.0, respectively; P = .014). Infusions of FIX<sub>WT</sub> (800 IU/kg) or FIX<sub>Alb</sub> (500 IU/kg), however, did not produce results significantly better than the CRM<sup>+</sup> mouse background; moreover, the results with FIX<sub>WT</sub> and FIX<sub>Alb</sub> were not statistically different from one another (P = .38). Thus, of the 2 longer circulating FIXs tested, only  $FIX_{FC}$  appears to be marginally superior to FIX<sub>WT</sub> in CRM<sup>+</sup> mice.

# Time course of the hemostatic effects of $FIX_{\rm WT}$ and $FIX_{\rm FC}$ in CRM $^-$ and CRM $^+$ mice.

After the infusion of 150 IU/kg of FIX<sub>WT</sub> or FIX<sub>FC</sub>, the SVBA was used to assess hemostasis at 1 and 7 days in CRM<sup>-</sup> mice (Figure 5A), and at 1, 3, 5, and 7 days in CRM<sup>+</sup> mice (Figure 5B). In Figure 5C, the median SVBA results in Figure 5A-B, after adjustment for the backgrounds in untreated mice, are shown.

The hemostatic effects of  $FIX_{WT}$  and  $FIX_{FC}$  were not statistically different at any point in either the CRM<sup>-</sup> or CRM<sup>+</sup> mice (Figure 5A and 5B). As assessed by the SVBA median results, however, the decline in the protective hemostatic effect of the infusions was much faster in CRM<sup>+</sup> than in CRM<sup>-</sup> mice (Figure 5C). At 7 days after infusion, the difference between the results in CRM<sup>-</sup> and CRM<sup>+</sup> mice was statistically significant for both FIX<sub>WT</sub> (P = .001) and FIX<sub>FC</sub> (P = .02) treated mice.

#### Effect of FIX reagents on an extant hemorrhage

The various FIXs were infused into CRM<sup>-</sup> mice 15 minutes after saphenous vein injury to determine the time necessary for the first clot to form in an ongoing hemorrhage (Figure 6). The mean times to first clot (TTFC) were 5.9, 7.3, and 11.8 minutes for FIX<sub>WT</sub>, FIX<sub>FC</sub>, and FIX<sub>Alb</sub>, respectively, with the TTFCs for both FIX<sub>FC</sub> and FIX<sub>Alb</sub> significantly prolonged compared with that for FIX<sub>WT</sub> (P = .007 and P < .001, respectively). FIX<sub>Alb</sub> displayed extreme variation in its TTFC; indeed, 3 of 15 mice infused with FIX<sub>Alb</sub> failed to clot during the 20-minute observation period.

Part of the delay in TTFC may be explained by the time required to activate each zymogen molecule. Therefore, the TTFC assay was repeated with the activated proteins  $FIX_{WTAct}$  and  $FIX_{FCAct}$ . (The creation of  $FIX_{AlbAct}$  is not possible because the C-terminal albumin is proteolytically removed during activation.) TTFCs were significantly faster for both activated  $FIX_{WT}$  and  $FIX_{FC}$  (1.9 and 2.9 minutes, respectively)



Figure 4. Comparison of prophylactic treatment with the different FIX agents in CRM<sup>+</sup> mice. (A) SVBA7 days after the infusion of 150 IU/kg FIX<sub>WT</sub>, FIX<sub>FC</sub>, or FIX<sub>Alb</sub> in CRM<sup>+</sup> mice. Results for untreated CRM<sup>+</sup> and wild-type mice are also shown. (B) SVBA7 days after the infusion of extremely high doses of FIX<sub>WT</sub>, FIX<sub>FC</sub>, and FIX<sub>Alb</sub> (Blue), WT mice. Untreated CRM<sup>+</sup> mice (orange), FIX<sub>WT</sub> (green), FIX<sub>FC</sub> (purple), FIX<sub>Alb</sub> (blue), WT mice (red). Box plot rendition of the results showing the minimum, first quartile, median (waist), third quartile, and maximum values. Note no difference between agents at the 150 IU/kg dose, but significantly increased efficacy with FIX<sub>FC</sub> at very high doses.



Figure 5. Time course of hemostasis after infusions of FIX<sub>WT</sub> and FIX<sub>FC</sub> in CRM<sup>-</sup> and CRM<sup>+</sup> mice. (A) SVBA at 1 and 7 days after the infusion of 150 IU/kg FIX<sub>WT</sub> (green) or FIX<sub>FC</sub> (purple) in CRM<sup>-</sup> mice. (B) SVBA at 1, 3, 5, and 7 days after the infusion of 150 IU/kg FIX<sub>WT</sub> (green) or FIX<sub>FC</sub> (purple) in CRM<sup>+</sup> mice. (C) Plot of median SVBA values for FIX<sub>WT</sub> (red) and FIX<sub>FC</sub> (blue) in CRM<sup>-</sup> mice and for FIX<sub>WT</sub> (green) and FIX<sub>FC</sub> (orange) in CRM<sup>+</sup> mice from panels A and B after adjustment for the median SVBA baseline values of untreated CRM<sup>-</sup> (1) and CRM<sup>+</sup> (3) mice. Note no significant difference in the hemostatic effects of the agents in CRM<sup>+</sup> or CRM<sup>-</sup> mice at 24 hours, but the rate of decay in hemostatic effectiveness is much faster in CRM<sup>+</sup> compared with CRM<sup>-</sup> mice. At 7 days, the differences between SVBA results in CRM<sup>-</sup> and CRM<sup>+</sup> mice are statistically significant for both FIX<sub>WT</sub> (P=.001) and FIX<sub>FC</sub> (P = .02) treatments.

than their zymogen forms, but the TTFC for activated  $FIX_{FC}$  was still significantly slower than that for activated  $FIX_{WT}$  (P = .037).

# Discussion

Large amounts of Col4 are present in the body, but not all Col4 binds FIX. For example, despite its access to Col4 in the liver sinusoids, FIX is not bound.<sup>12</sup> Presumably, lack of FIX binding to the Col4 in the liver sinusoids is the result of an unknown posttranslational modification of the Col4, rendering it unable to bind FIX. One example could be prolyl hydroxylation.<sup>28,29</sup> Most of the subendothelium, however, contains Col4 that binds FIX avidly, thus producing a cache of FIX in basement membranes that is severalfold greater than the FIX circulating in plasma. That this FIX-Col4 complex plays an essential role in hemostasis is best illustrated by the bleeding diathesis evident in knock-in mice expressing  $FIX_{K5A}$ , which has normal in vitro coagulant activity, but binds Col4 poorly.<sup>13,30</sup>

Our studies suggest the concentration of extravascular Col4 capable of binding plasma FIX in the mouse is 405 nM with a 95% confidence interval of 342 to 467 nM. This value is consistent with previously reported studies examining the competitive release of endogenous extravascular FIX in baboons.<sup>5</sup> Binding to Col4 also appears to explain the 40% to 50% expected recovery of infused FIX observed in patients with hemophilia B.<sup>7</sup> The majority of patients with hemophilia B are CRM<sup>+</sup> and have variable circulating levels of a dysfunctional FIX. Depending on the level of the dysfunctional FIX and its ability to bind Col4, this CRM<sup>+</sup> FIX could increase the plasma recovery of infused FIX<sub>WT</sub> bound to vascular Col4 by competing for Col4 binding.

The Gla domain of FIX is responsible for its binding to Col4. For example, targeted mutation of 6 specific residues within the Gla domain of FVII is sufficient to produce a molecule with the same  $K_d$  for Col4 as FIX<sub>WT</sub>. Further, replacing the Gla domain of protein C with that of FIX produces molecules that binds Col4 with the same  $K_d$  as FIX.<sup>9,31</sup> The majority of the described missense mutations in CRM<sup>+</sup> patients with hemophilia B should not affect the sequence<sup>15</sup> or function<sup>32</sup> of the Gla domain and will, therefore, produce FIX proteins that presumably compete with infused FIX for Col4 binding. To examine coagulation in



Figure 6. Times to first clot for the different FIX agents in CRM<sup>-</sup> mice. The saphenous vein of CRM<sup>-</sup> mice was injured and the mice were allowed to bleed for 15 minutes. Each FIX agent (500 IU/kg) was then infused and the TTFC (see Methods) determined. FIX<sub>WT</sub> (green), FIX<sub>FC</sub> (purple) and FIX<sub>Alb</sub> (blue). FIX<sub>WTAct</sub> and FIX<sub>FCAct</sub> are the activated forms of the respective zymogens. Box plot rendition of the results showing the minimum, first quartile, median (waist), third quartile, and maximum values.

a situation mimicking prophylactic therapy, we tested hemophilia B mice with the SVBA assay 7 days after the infusion of the FIX. Hemostasis after the injection of FIX<sub>WT</sub> was substantially better in CRM<sup>-</sup> than in CRM<sup>+</sup> (FIXR<sub>333Q</sub>) mice (Figure 3). Under the same experimental conditions, infusion of FIX<sub>Alb</sub> into either CRM<sup>-</sup> or CRM<sup>+</sup> mice produced results indistinguishable from those after the infusion of FIX<sub>WT</sub>, including the reduction in hemostasis found in CRM<sup>+</sup> compared with CRM<sup>-</sup> recipients (Figure 4A-B).

We had previously demonstrated that infusions of FIX<sub>WT</sub> and FIX<sub>FC</sub> also produced equivalent hemostatic results in CRM<sup>-</sup> mice, using this same model.<sup>12</sup> At extremely high levels of FIX infusion ( $\geq$ 500 IU/kg), however, enhanced hemostasis is apparent when FIX<sub>FC</sub> is compared with FIX<sub>WT</sub> and FIX<sub>Alb</sub> in CRM<sup>+</sup> mice (Figure 4B). The longer plasma half-life of FIX<sub>FC</sub> may contribute to its improved effect over FIX<sub>WT</sub> at these high levels of the infused FIX. The reason for the disparity in the results between FIX<sub>FC</sub> and FIX<sub>Alb</sub>, another long half-life agent, however, is not apparent (see the following paragraph).

The hemostatic effects measured with the SVBA are remarkably preserved 7 days after the infusion of  $\ensuremath{\mathsf{FIX}_{\mathsf{WT}}}$  into  $\ensuremath{\mathsf{CRM}^-}$  mice, when plasma levels are not measurable<sup>11</sup> (Figure 5A), consistent with a functional role for the FIX-Col4 vascular complex in hemostasis. In contrast, we observe a progressive decrease in hemostatic effectiveness during the same period in CRM<sup>+</sup> mice (Figure 5B). The differences in hemostatic efficacy when the mouse has endogenous FIX may reflect the continuing expression of dysfunctional FIX, which competes for Col4 binding, in the CRM<sup>+</sup> animals. Interestingly, we observe a linear kinetics loss of hemostatic protection over time (Figure 5C). This linear loss of efficacy contrasts with the welldocumented exponential kinetics loss of infused FIX from the plasma of many animals.<sup>6,32,33</sup> This suggests that hemostatic protection is attributable to something other than simply the plasma level of FIX and is consistent with our argument that extravascular FIX plays a functional role in hemostasis. Thus, in a CRM<sup>+</sup> background because of a dysfunctional FIX that binds Col4, treatment with a FIX molecule with enhanced Col4 binding such as FIX<sub>K5R</sub><sup>11</sup> might be advantageous.

To evaluate further the different FIX molecules, their hemostatic effectiveness was tested in bleeding CRM<sup>-</sup> mice as the TTFC. Mouse blood completely circulates about 6 times per minute, and an inert molecule is mixed entirely by 25 seconds.34 Compared with FIX<sub>WT</sub>, the TTFC of both longer half-life FIX molecules is significantly delayed (Figure 6). The cause or causes of the apparent slower in vivo onset of coagulation of the zymogen  $FIX_{FC}$  and  $FIX_{AIb}$  forms in the mouse are likely multifactorial and could include differences in their rates of activation, the activities of their activated FIXa forms once generated, and perhaps their binding to Col4. In the latter regard, it is of interest that dosing recommendations in the package inserts of BeneFix (FIX<sub>WT</sub>), Alprolix (FIX<sub>FC</sub>), and Idelvion (FIX<sub>Alb</sub>) suggest plasma recoveries of the infused products to be 38%, 50%, and 65%, respectively. Given the dominant effect of vascular Col4 binding on the plasma recovery of the infused FIX, the dosing recommendations suggest that the relative degree of Col4 binding of the agents is  $FIX_{WT} > FIX_{FC} > FIX_{Alb}$ . Of the longer half-life molecules, the efficacy of  $FIX_{FC}$  appeared to be superior to that of FIX<sub>Alb</sub> in our mouse hemophilia B models.

On the basis of plasma FIX levels, the longer half-life FIX products are administered at a reduced frequency (1/wk) for

prophylactic treatment and were rapidly accepted by the hemophilia B community. Studies supporting the efficacy of the less frequent prophylactic doses of these products, however, do not consider the potential procoagulant effect of the extravascular FIX bound to Col4. FIX<sub>WT</sub>, FIX<sub>FC</sub>, and FIX<sub>Alb</sub> all produce equivalent results in our mouse model of once-per-week prophylactic therapy, but clinical studies directly comparing the various FIX agents have not been performed. Of note in this regard, once-per-week dosing of 100 IU/kg BeneFix (FIX<sub>WT</sub>) in patients with hemophilia B appears to produce outcomes similar to those reported for Alprolix (FIX<sub>FC</sub>).<sup>35</sup>

We believe the deleterious hemostatic effect of the dysfunctional FIX tested in our mouse studies (FIX<sub>R333O</sub>) is primarily a result of its competition with infused FIX for Col4 binding. FIX bound to Col4 in the extracellular matrix of the subendothelium is in an ideal location to contribute to the hemostatic response to vascular injury. Tissue factor, which is responsible for the initiation of coagulation, is expressed by pericytes, which share the extracellular matrix of microvascular endothelial cells.<sup>36</sup> Plasma von Willebrand factor carries FVIII and also binds Col4 specifically.<sup>36,37</sup> Further, platelets bind collagens of various types, including Col4, so the possibility that all these procoagulant elements combine locally on a subendothelial scaffold to augment clotting is appealing.

It must be pointed out, however, that the endogenous expression of a dysfunctional FIX could affect hemostasis through other mechanisms as well, including competition with FIX for activation by FVIIa/tissue factor or FXIa and competition with FIXa for FVIIIa binding. The latter mechanism is unlikely in our tested CRM<sup>+</sup> mouse, as the hemostatic defect with FIX<sub>R333Q</sub> is a result of its much-reduced affinity for FVIIIa.<sup>38</sup>

The presence of CRM<sup>+</sup> material may explain some of the variability in disease severity in patients with hemophilia B with the same level of plasma FIX functional activity. For example, a patient whose FIX is 100% active with a plasma antigen level of only 2% may have very different bleeding expectations than a CRM<sup>+</sup> patient expressing a dysfunctional FIX with 2% activity and an antigen level of 100%.

In sum, these studies in murine hemophilia B models suggest a potential, unappreciated role of the endogenous, dysfunctional FIX in CRM<sup>+</sup> individuals in attenuating their hemostatic response to prophylactic therapy. The extent of the deleterious effect would depend on the level of expression of the dysfunctional FIX and its specific molecular abnormality.

# Acknowledgments

The authors thank Katherine Stafford and Sheue-Mei Wu for editing and discussions.

This work was supported by a trust fund of D.W.S.

# Authorship

Contribution: D.W.S. originated the work and wrote the first draft; D.W.S. and D.M.M. planned most of the experiments; B.C. performed the experiments and helped plan the time to first clot experiments; G.J.B. analyzed the data and helped write (and rewrite) the manuscript; F.-C.L. evaluated most of the data and performed some of the statistical calculations; L.G.P. was also involved in planning and especially spent hours understanding and helping with the mathematical evaluations and also

spent considerable time with revisions; and all authors carefully read the final version and made helpful comments.

Conflict-of-interest disclosure: D.W.S. has a patent on the VKOR enzyme and, in Europe, on  $FIX_{KSR/R338L}$  (the latter is under consideration in the United States). The remaining authors declare no competing financial interests.

Correspondence: Darrel W. Stafford, Department of Biology and Pathology, University of North Carolina at Chapel Hill, 442 Wilson Hall, South Road, Chapel Hill, NC; e-mail: dws@e-mail.unc.edu.

#### REFERENCES

- Stern DM, Nawroth PP, Kisiel W, Vehar G, Esmon CT. The binding of factor IXa to cultured bovine aortic endothelial cells. Induction of a specific site in the presence of factors VIII and X. J Biol Chem. 1985;260(11):6717-6722.
- Stern DM, Drillings M, Nossel HL, Hurlet-Jensen A, LaGamma KS, Owen J. Binding of factors IX and IXa to cultured vascular endothelial cells. *Proc Natl Acad Sci USA*. 1983;80(13):4119-4123.
- Cheung WF, van den Born J, Kühn K, Kjellén L, Hudson BG, Stafford DW. Identification of the endothelial cell binding site for factor IX. Proc Natl Acad Sci USA. 1996;93(20):11068-11073.
- Khoshnoodi J, Pedchenko V, Hudson BG. Mammalian collagen IV. *Microsc Res Tech*. 2008;71(5):357-370.
- Stern DM, Knitter G, Kisiel W, Nawroth PP. In vivo evidence of intravascular binding sites for coagulation factor IX. *Br J Haematol.* 1987; 66(2):227-232.
- Jin DY, Zhang TP, Gui T, Stafford DW, Monahan PE. Creation of a mouse expressing defective human factor IX. *Blood*. 2004;104(6): 1733-1739.
- Poon MC, Lillicrap D, Hensman C, Card R, Scully MF. Recombinant factor IX recovery and inhibitor safety: a Canadian post-licensure surveillance study. *Thromb Haemost.* 2002; 87(3):431-435.
- Toomey JR, Smith KJ, Roberts HR, Stafford DW. The endothelial cell binding determinant of human factor IX resides in the gammacarboxyglutamic acid domain. *Biochemistry*. 1992;31(6):1806-1808.
- Cheung WF, Hamaguchi N, Smith KJ, Stafford DW. The binding of human factor IX to endothelial cells is mediated by residues 3-11. *J Biol Chem.* 1992;267(29):20529-20531.
- Gui T, Lin HF, Jin DY, et al. Circulating and binding characteristics of wild-type factor IX and certain Gla domain mutants in vivo. *Blood.* 2002;100(1):153-158.
- Feng D, Stafford KA, Broze GJ, Stafford DW. Evidence of clinically significant extravascular stores of factor IX. J Thromb Haemost. 2013; 11(12):2176-2178.
- Cooley B, Funkhouser W, Monroe D, et al. Prophylactic efficacy of BeneFIX vs Alprolix in hemophilia B mice. *Blood*. 2016;128(2): 286-292.
- Gui T, Reheman A, Ni H, et al. Abnormal hemostasis in a knock-in mouse carrying a variant of factor IX with impaired binding to collagen type IV. J Thromb Haemost. 2009; 7(11):1843-1851.

- DiMichele D. Inhibitor development in haemophilia B: an orphan disease in need of attention. Br J Haematol. 2007;138(3):305-315.
- Rallapalli PM, Kemball-Cook G, Tuddenham EG, Gomez K, Perkins SJ. An interactive mutation database for human coagulation factor IX provides novel insights into the phenotypes and genetics of hemophilia B. J Thromb Haemost. 2013;11(7):1329-1340.
- Briet E. Three problems of haemophilia B [PhD Thesis]. The University of Amsterdam Medical School; 1977.
- Roth DA, Kessler CM, Pasi KJ, Rup B, Courter SG, Tubridy KL; Recombinant Factor IX Study Group. Human recombinant factor IX: safety and efficacy studies in hemophilia B patients previously treated with plasma-derived factor IX concentrates. *Blood*. 2001;98(13): 3600-3606.
- Ewenstein BM, Joist JH, Shapiro AD, et al; Mononine Comparison Study Group. Pharmacokinetic analysis of plasma-derived and recombinant F IX concentrates in previously treated patients with moderate or severe hemophilia B. *Transfusion*. 2002;42(2):190-197.
- Lin HF, Maeda N, Smithies O, Straight DL, Stafford DW. A coagulation factor IX-deficient mouse model for human hemophilia B. *Blood*. 1997;90(10):3962-3966.
- Hamaguchi N, Charifson PS, Pedersen LG, Brayer GD, Smith KJ, Stafford DW. Expression and characterization of human factor IX. Factor IXthr-397 and factor IXval-397. J Biol Chem. 1991;266(23):15213-15220.
- Eisner C, Ow H, Yang T, et al. Measurement of plasma volume using fluorescent silica-based nanoparticles. J Appl Physiol (1985). 2012; 112(4):681-687.
- Johnson ML. Why, when, and how biochemists should use least squares. Anal Biochem. 1992;206(2):215-225.
- Pastoft AE, Lykkesfeldt J, Ezban M, Tranholm M, Whinna HC, Lauritzen B. A sensitive venous bleeding model in haemophilia A mice: effects of two recombinant FVIII products (N8 and Advate(®)). *Haemophilia*. 2012;18(5): 782-788.
- Carpenter J, Bithell J. Bootstrap confidence intervals: when, which, what? A practical guide for medical statisticians. *Stat Med.* 2000;19(9): 1141-1164.
- Pollard TD. A guide to simple and informative binding assays. *Mol Biol Cell*. 2010;21(23): 4061-4067.
- Powell JS, Pasi KJ, Ragni MV, et al; B-LONG Investigators. Phase 3 study of recombinant factor IX Fc fusion protein in hemophilia B. N Engl J Med. 2013;369(24):2313-2323.

# Footnotes

Submitted 2 November 2018; accepted 9 April 2019. Prepublished online as *Blood* First Edition paper, 16 April 2019; DOI 10.1182/ blood.2018884015.

The publication costs of this article were defrayed in part by page charge payment. Therefore, and solely to indicate this fact, this article is hereby marked "advertisement" in accordance with 18 USC section 1734.

- Martinowitz U, Lissitchkov T, Lubetsky A, et al. Results of a phase I/II open-label, safety and efficacy trial of coagulation factor IX (recombinant), albumin fusion protein in haemophilia B patients. *Haemophilia*. 2015; 21(6):784-790.
- Pokidysheva E, Boudko S, Vranka J, et al. Biological role of prolyl 3-hydroxylation in type IV collagen. Proc Natl Acad Sci USA. 2014;111(1):161-166.
- Hudson DM, Joeng KS, Werther R, et al. Posttranslationally abnormal collagens of prolyl 3-hydroxylase-2 null mice offer a pathobiological mechanism for the high myopia linked to human LEPREL1 mutations. J Biol Chem. 2015;290(13):8613-8622.
- Cooley BC. In vivo fluorescence imaging of large-vessel thrombosis in mice. Arterioscler Thromb Vasc Biol. 2011;31(6):1351-1356.
- Geng JP, Christiansen WT, Plow EF, Castellino FJ. Transfer of specific endothelial cellbinding properties from the procoagulant protein human factor IX into the anticoagulant protein human protein C. *Biochemistry*. 1995; 34(26):8449-8457.
- Peters RT, Low SC, Kamphaus GD, et al. Prolonged activity of factor IX as a monomeric Fc fusion protein. *Blood.* 2010;115(10):2057-2064.
- Björkman S, Carlsson M, Berntorp E. Pharmacokinetics of factor IX in patients with haemophilia B. Methodological aspects and physiological interpretation. Eur J Clin Pharmacol. 1994;46(4):325-332.
- Welsher K, Sherlock SP, Dai H. Deep-tissue anatomical imaging of mice using carbon nanotube fluorophores in the second nearinfrared window. Proc Natl Acad Sci USA. 2011;108(22):8943-8948.
- Kavakli K, Smith L, Kuliczkowski K, et al. Onceweekly prophylactic treatment vs. on-demand treatment with nonacog alfa in patients with moderately severe to severe haemophilia B. Haemophilia. 2016;22(3):381-388.
- Flood VH, Schlauderaff AC, Haberichter SL, et al; Zimmerman Program Investigators. Crucial role for the VWF A1 domain in binding to type IV collagen. *Blood.* 2015;125(14):2297-2304.
- 37. Slobodianuk TL, Kochelek C, Foeckler J, Kalloway S, Weiler H, Flood VH. Defective collagen binding and increased bleeding in a murine model of von Willebrand disease affecting collagen IV binding. J Thromb Haemost. 2019;17(1):63-71.
- Bajaj SP, Schmidt AE, Mathur A, et al. Factor IXa:factor VIIIa interaction. helix 330-338 of factor ixa interacts with residues 558-565 and spatially adjacent regions of the a2 subunit of factor VIIIa. J Biol Chem. 2001;276(19):16302-16309.