

LYMPHOID NEOPLASIA

Cirmtuzumab blocks Wnt5a/ROR1 stimulation of NF- κ B to repress autocrine STAT3 activation in chronic lymphocytic leukemia

Yun Chen,^{1,2,*} Liguang Chen,^{1,*} Jian Yu,¹ Emanuela M. Ghia,¹ Michael Y. Choi,¹ Ling Zhang,¹ Suping Zhang,^{1,3} Elsa Sanchez-Lopez,^{4,5} George F. Widhopf II,¹ Karen Messer,¹ Laura Z. Rassenti,¹ Catriona Jamieson,^{1,6} and Thomas J. Kipps^{1,3}

¹Moore's Cancer Center, University of California, San Diego, La Jolla, CA; ²Edmond H. Fischer Translational Medical Research Laboratory, Scientific Research Center, The Seventh Affiliated Hospital, Sun Yat-sen University, Shenzhen, Guangdong, China; ³Guangdong Key Laboratory for Genome Stability & Disease Prevention, Department of Pharmacology, International Cancer Center, Shenzhen University Health Science Center, Shenzhen, Guangdong, China; and ⁴Department of Pharmacology and ⁵Department of Pathology, School of Medicine, and ⁶Division of Regenerative Medicine, University of California, San Diego, La Jolla, CA

KEY POINTS

- Nurse-like cells express Wnt5a to induce ROR1-dependent stimulation of NF- κ B, which leads to autocrine IL-6-induced STAT3 activation in CLL cells.
- Cirmtuzumab inhibits Wnt5a-induced, ROR1-dependent stimulation of NF- κ B, and thereby represses autocrine IL-6-dependent STAT3 activation in CLL.

Coculture of nurse-like cells (NLCs) with chronic lymphocytic leukemia (CLL) cells induced leukemia cell phosphorylation of STAT3 (pSTAT3), which could be blocked by anti-Wnt5a antibodies or the anti-ROR1 monoclonal antibody, cirmtuzumab. Time-course studies revealed Wnt5a could induce activation of NF- κ B within 30 minutes, but required more than 3 hours to induce pSTAT3. Culture of isolated CLL cells for 24 hours revealed Wnt5a-induced expression of interleukin 6 (IL-6), IL-8, CCL2, CCL3, CCL4, and CXCL1, which in turn could induce pSTAT3 in unstimulated CLL cells within 30 minutes. We found that Wnt5a could induce CLL cell expression of NF- κ B target genes, including IL-6, and that this effect could be blocked by cirmtuzumab or drugs that inhibit NF- κ B. Examination of CLL cells and plasma collected from patients treated with cirmtuzumab revealed reduced levels of phosphorylated p65 and diminished expression of NF- κ B and STAT3 target genes in CLL cells, as well as lower plasma levels of IL-6, in the samples after therapy. Collectively, these studies indicate that Wnt5a/ROR1-dependent signaling contributes to CLL cell activation of NF- κ B, which in turn causes autocrine IL-6-induced activation of pSTAT3. As such, this study demonstrates that cirmtuzumab can inhibit leukemia cell activation of both NF- κ B and STAT3 in patients with CLL. (*Blood*. 2019;134(13):1084-1094)

Introduction

Receptor tyrosine kinase-like orphan receptor 1 (ROR1) is a developmentally restricted oncoembryonic surface protein that is expressed on neoplastic cells of many types of cancer,¹ including chronic lymphocytic leukemia (CLL), but not on most normal postpartum tissues.² ROR1 can serve as a receptor for Wnt5a, which can promote leukemia cell growth and survival,²⁻⁴ potentially accounting for the observation that high-level CLL cell expression of ROR1 is associated with early disease progression and shorter overall survival.⁵ These properties of ROR1 make it an attractive target for therapy of patients with CLL, prompting development of a humanized monoclonal antibody, called cirmtuzumab, which targets ROR1 and inhibits ROR1-signaling in vitro.^{4,6-10} A phase I study of cirmtuzumab in patients with CLL demonstrated that this antibody also could inhibit ROR1-signaling in vivo, suppressing leukemia cell activation of ρ -GTPases and phosphorylation of HS1.¹¹

In addition to activation of ρ -GTPases,⁴ ROR1 signaling also can induce recruitment and activation of 14-3-3 ζ ,⁸ HS1,⁷ DOCK2,⁶

and cortactin,¹² each of which also contributes to Wnt5a-induced, ROR1-dependent enhancement of CLL cell proliferation and/or migration. Wnt5a also has been reported to induce activation of STAT3 in melanoma cells¹³ and activation of NF- κ B in the human embryonic kidney cell line HEK293 in a ROR1-dependent-manner.² Activation of NF- κ B and STAT3 also can enhance proliferation and/or survival of CLL cells.¹⁴⁻¹⁷ Furthermore, activation of STAT3 may enhance expression of ROR1,¹⁸ potentially providing a positive feedback loop in which Wnt5a could upregulate the expression of its receptor. However, it is not known whether Wnt5a could induce activation of STAT3 or NF- κ B in ROR1-expressing CLL cells.

Also not clear is the principle cellular source or sources for Wnt5a. Although hydrophobic amino acids and posttranslational palmitoylation provides mature Wnt proteins the capacity to act primarily as surface proteins tethered to the plasma membrane,¹⁹ Wnt5a can be found at high levels in the plasma of patients with CLL relative to that of age-matched healthy adults.^{4,11} The cell

source responsible for the high levels of Wnt5a in plasma of patients with CLL is not clear. Although CLL cells themselves have been noted to express Wnt5a, only 38% of patients have detectable transcripts of *Wnt5a* in their leukemia cells.²⁰ Other candidates include nurse-like cells (NLCs), the nonmalignant accessory cells residing in the proliferation centers of lymphoid tissue that are derived from monocytes,^{21,22} which may express Wnt5a.²³ Expression of Wnt5a by NLCs implicates that there most likely are higher concentrations of Wnt5a in lymphoid tissues than in the circulation, potentially leading to amplified Wnt5a/ROR1 signaling and upregulated expression of ROR1 through the positive feedback loop within the microenvironment.

Among circulating CLL cells, the relative expression CD5 and CXCR4 can be used to distinguish between leukemia cells that recently have exited from the lymphoid tissue microenvironment (CD5^{high} CXCR4^{dim}) and leukemia cells that may be poised to reenter the lymphoid compartments (CXCR4^{high} CD5^{dim}).^{24,25} Prior studies demonstrated that circulating CXCR4^{dim} CD5^{high} CLL cells express higher levels of genes upregulated in leukemia cells present in lymphoid tissues than blood CXCR4^{high} CD5^{dim} CLL cells of the same patient,²⁴ providing a means with which to assess how the lymphoid microenvironment influences CLL cell gene-expression. We examined whether CXCR4^{dim} CD5^{high} CLL cells have higher levels of activated STAT3 and ROR1 compared with CXCR4^{high} CD5^{dim} CLL cells.

Methods

CLL specimens

Blood samples were collected from Moores Cancer Center patients who provided written informed consent using a protocol approved by the Institutional Review Board of the University of California, San Diego (approval number 090401), in accordance with the Declaration of Helsinki. Peripheral blood mononuclear cells were isolated by density-gradient centrifugation with Ficoll-Paque PLUS (GE Healthcare Life Sciences), followed by purification with human B Cell Isolation Kit (130-091-151, Miltenyi Biotec Inc.). CLL cells were segregated into "ROR1^{high}" and "ROR1^{low}" subgroups, as described.⁵ Plasma was collected from blood samples that had undergone centrifugation for 10 minutes at 187g, and stored at -80°C. Plasma samples and CLL cells were collected from patients treated with cirmtuzumab in the phase 1 clinical study of cirmtuzumab (registered at www.clinicaltrials.gov as #NCT02222688).¹⁰

Generation of nurse-like cells

Peripheral blood mononuclear cells isolated by Ficoll-Paque PLUS from patients with CLL were suspended in RPMI with 20% fetal bovine serum to a final concentration of 2×10^7 /mL, as described.²¹ After 14 days, the nonadherent CLL cells were removed by vigorously pipetting the contents of the well, leaving the adherent cells untouched. We observed these cells to have the morphology typical of NLC via microscopy.

Materials

We obtained BAY 11-7082 (S2913) from Selleckchem. Human recombinant Wnt5a (645-WN) was from R&D Systems. Wnt5a was suspended in sterile PBS containing 0.1% bovine serum albumin at 100 µg/mL.

Immunoblot analysis

Immunoblot analysis was performed as described.² Primary monoclonal antibody (mAb) for ROR1 (#4102), Phospho-STAT3 (Tyr705; #9145), STAT3 (#4904), phospho-p65 (#3033), phospho-IκBα (#9246), p65 (#8242), and IκBα (#4814) were obtained from Cell Signaling Technology. Secondary antibody conjugated with horseradish peroxidase was from Cell Signaling Technology.

Flow cytometry

Flow cytometry analysis was performed as described.^{2,11} Anti-ROR1 mAb (4A5) conjugated with Alexa Fluor 647 (4A5-Alexa Fluor 647) was generated in our laboratory. 4A5 is a non-cross-blocking mAb specific for a ROR1 epitope distinct from that recognized by cirmtuzumab.² Anti-CD19-PerCP, anti-CD5-FITC, anti-pSTAT3 Y705-PE, and anti-CXCR4-APC antibodies were from BD Biosciences. The molecules of ROR1 per cell were determined with the Quantum MESF (Molecules of Equivalent Soluble Fluorochrome) microspheres. Alexa Fluor 647 conjugated beads were run on the same day as stained cell samples to establish a calibration curve relating channel values to MESF units, or ΔMFI.

ELISA

IL-6 in plasma or medium was detected by Human IL-6 ELISA (enzyme-linked immunosorbent assay) kit (430504; BioLegend). Wnt5a was detected by Human WNT5A ELISA kit (OKEH00723; Aviva Systems Biology).

Human cytokine assay

CLL cells were cultured with or without Wnt5a (200 ng/mL) or cirmtuzumab (10 µg/mL) or anti-Wnt5a antibody (MAB645; R&D Systems) (5 µg/mL) in RPMI culture-media, which were analyzed by Proteome Profiler Human Cytokine Array Kit (ARY005B; R&D Systems), as per the manufacturer's instructions.

NF-κB target gene expression

We used the RT2 Profiler PCR (polymerase chain reaction) Array for human NF-κB signaling targets (PAHS-225ZA; Qiagen) to profile the expression of 83 NF-κB target genes. Real-time PCR was performed per the manufacturer's instructions. The threshold (C_T) value for each gene in the array was normalized to the average C_T value of 4 housekeeping genes. The Volcano Plots and ClusterGram Heatmap were generated using the Qiagen data analysis center (<https://www.qiagen.com/us/shop/genes-and-pathways/data-analysis-center-overview-page/>), as per the vendor's instructions.

Quantitative reverse-transcription PCR

Total RNA was extracted using Trizol (Life Technologies). A 10-µg volume of total RNA was incubated with 10 U RNase-free DNase I (Life Technologies) for 30 minutes at 37°C. RNA was further purified with an RNeasy Mini Kit (QIAGEN, Hilden, Germany). The purified total RNA (2 µg) was reverse transcribed with Superscript III Reverse Transcriptase (Life Technologies). Taq 2× Master Mix (NEB) was used for the PCR, per the manufacturer's protocol. SYBR Green PCR Master Mix (Thermo Fisher Scientific) and 7900HT Fast Real-Time PCR System (Applied Biosystems) were used for real-time PCR, per the manufacturer's protocol. The expression level of *WNT5A* transcripts was calculated using *ACTB*.

Gene set enrichment analysis

We performed gene set enrichment analysis (GSEA) of STAT3 target genes in matched lymph nodes and peripheral blood CLL cells collected from 17 patients previously included in a study by Herishanu et al.¹⁶ Peripheral blood mononuclear cells of 3 CLL samples were collected before therapy (Pre-Rx) and at day 28 (D28) of cirtumzumab treatment. Each D28 sample was collected after patients had received 2 doses of 16 mg/kg cirtumzumab. Negative isolation of CLL cells to more than 95% purity was performed before RNA isolation. Total RNA was extracted using Trizol reagent (Life Technologies). RNAseq data were analyzed as described.¹¹ Using GSEA software,²⁶ we conducted GSEA on the primary RNAseq data from CLL cells isolated from patients before therapy (Pre-Rx) and at D28 of cirtumzumab treatment. We focused our attention on changes in the NF- κ B²⁷ and STAT3²⁸ target genes. Each gene set was considered significant when the false discovery rate (FDR) was less than 25%.²⁶ The FDR q value was adjusted for gene set size and multiple hypothesis testing.

Statistical analyses

Data are presented as the mean \pm standard deviation. Differences between 2 groups were determined by 2-tailed Student *t* test. Significance was analyzed by GraphPad Prism 6.0 (GraphPad Software Inc.), and *P* < .05 was considered significant.

Results

The expression of ROR1 and phosphorylation of STAT3 are higher in CD5^{high} CXCR4^{dim} CLL cells than their CD5^{dim} CXCR4^{high} counterparts

We examined 24 CLL cell samples that expressed high (ROR1^{high}, *N* = 12) or low (ROR1^{low}, *N* = 12) levels of surface ROR1 (supplemental Table 1, available on the *Blood* Web site), as per a previously defined threshold,⁴ which subsequently was noted to correspond to $\approx 4 \times 10^3$ molecules of ROR1 per cell.¹¹ CLL cells were stained with antibodies specific for CD5 or CXCR4. CD5^{high} CXCR4^{dim} and CD5^{dim} CXCR4^{high} subpopulations were electronically gated for analysis of the relative expression of other proteins by these subpopulations. The levels of ROR1 and pSTAT3 were determined in each gated subpopulation. We noted that CD5^{high} CXCR4^{dim} CLL cells, which presumably had recently exited the lymph node, had higher levels of ROR1 and pSTAT3 than the CD5^{dim} CXCR4^{high} counterpart CLL cells (Figure 1A-C; supplemental Table 1), although the difference in the levels of ROR1 between such subpopulations did not reach statistical significance for ROR1^{low} samples (Figure 1B). Consistent with the notion that CLL cells exiting the lymph nodes may have higher levels of activated STAT3 than those circulating in the blood is our gene set enrichment analyses of the published transcriptome data on CLL cells isolated from the lymph node and blood of 17 patients.¹⁶ This analysis revealed that the expression levels of STAT3-target genes (*N* = 97) were significantly higher in CLL cells isolated from the lymph node than in CLL cells isolated from the blood of the same patients (NES = 1.30; FDR *q* = 0.08; Figure 1D).

NLCs induce CLL cell activation of STAT3 and upregulation of ROR1 via Wnt5a

We cultured NLCs with CLL cells from each of 6 patients with ROR1^{high} CLL and 6 patients with ROR1^{low} CLL, using an established coculture system.²⁹ The levels of ROR1 and pSTAT3


were determined by flow cytometry. CLL cells cocultured with NLC had increased pSTAT3 in both ROR1^{high} and ROR1^{low} samples (Figure 2A-C). Although ROR1^{high} CLL cells cocultured with NLC had increased levels of ROR1, the upregulation of ROR1 in ROR1^{low} CLL cells cocultured with NLC was not statistically significant (Figure 2B). Activation of STAT3 and upregulation of ROR1 in ROR1^{high} CLL cells were blocked respectively by adding the anti-ROR1 mAb, cirtumzumab, or neutralizing antibodies to Wnt5a at the initiation of culture (Figure 2A-C). Consistent with the notion that NLCs were the principle source of Wnt5a, we noted that isolated NLC expressed significantly higher amounts of WNT5A than the CLL cells of the same patient (Figure 2D). Moreover, we detected higher amounts Wnt5a in the culture supernatants of isolated NLC than in the supernatants of cultures containing similar numbers of CLL cells (Figure 2E).

Wnt5a/ROR1 induces production of proinflammatory factors that induce phosphorylation of STAT3 and upregulation of ROR1 in CLL cells

We stimulated CLL cells with exogenous Wnt5a for 0 to 6 and 24 hours in the presence of control IgG or cirtumzumab. Cell lysates were analyzed by immunoblot for pSTAT3 (Y705) and total STAT3. This revealed that Wnt5a induced phosphorylation of STAT3 in ROR1-positive CLL, and that this effect could be blocked by cirtumzumab (Figure 3A). The capacity of Wnt5a to induce pSTAT3 was less apparent in CLL cells that lacked expression of ROR1 (supplemental Figure 1). Wnt5a also enhanced expression of ROR1 on ROR1-positive CLL cells, but did not induce expression of ROR1 on CLL cells that lacked ROR1 (supplemental Figure 2). Time course studies revealed latent induction of pSTAT3, which first was detected 3 hours after stimulation with Wnt5a. Because of this latent induction, we hypothesized that the capacity of Wnt5a to induce activation of STAT3 in CLL cells was indirect, being caused by a factor or factors made by isolated CLL cells on stimulation by Wnt5a.

We examined harvested supernatants of CLL cells cultured with or without Wnt5a, using a human cytokine array. Multiple proinflammatory factors including IL-6, IL-8, CCL2, CCL3, CCL4, and CXCL1 were detected in the medium of CLL cells cultured for 24 hours with Wnt5a that either were not detected or were found at significantly lower levels in the medium of CLL cells cultured without Wnt5a (Figure 3B). Cirtumzumab inhibited the capacity of Wnt5a to induce CLL cells to generate most of these factors (Figure 3C). The medium harvested from CLL cells cultured with Wnt5a, but not the medium of CLL cells cultured without Wnt5a, could induce pSTAT3 within 30 minutes in serum-starved CLL cells; this effect could be inhibited by the anti-IL-6-receptor mAb, tocilizumab (Figure 3D). The conditioned medium also induced upregulation of ROR1 in ROR1^{high} CLL cells; this effect also could be inhibited by tocilizumab, but not by neutralizing antibodies specific for Wnt5a (supplemental Figure 3). We used exogenous Wnt5a, IL-6, IL-8, CCL2, CCL3, CCL4, or CXCL1 to stimulate serum-starved CLL cells for 30 minutes, and found IL-6 was the most potent early inducer of pSTAT3 (supplemental Figure 4). IL-6 could induce pSTAT3 in CLL cells independent of the expression of ROR1 (supplemental Figure 5), but it could only upregulate the level of ROR1 on CLL

Figure 1. CD5^{high} CXCR4^{dim} CLL cells express higher levels of ROR1 and phosphorylated STAT3 than CD5^{dim} CXCR4^{high} counterparts. (A) Flow cytometry analysis of pSTAT3 (Y705) and ROR1 in CD5^{high} CXCR4^{dim} and CD5^{dim} CXCR4^{high} CLL cells. (B) Dot figures representing ROR1 intensity in CD5^{high} CXCR4^{dim} and CD5^{dim} CXCR4^{high} CLL cells from ROR1^{high} (n = 12) and ROR1^{low} (n = 12) patients. Error bars denote standard deviation. (C) Bar figures representing pSTAT3 Δ MFI in CD5^{high} CXCR4^{dim} and CD5^{dim} CXCR4^{high} CLL cells from ROR1^{high} (n = 12) and ROR1^{low} (n = 12) patients. Error bars denote standard deviation. (D) GSEA plot of STAT3 target genes in lymph nodes vs peripheral blood CLL cells collected from 17 patients. The gene set size was 97, the normalized enrichment score (NES) was 1.30, and the false discovery rate (FDR) q value was 0.08.



cells that already expressed ROR1 (supplemental Figure 6). Because IL-6 is a target gene of STAT3, we considered whether the increased expression of IL-6 might be a consequence of STAT3 activation, rather than a cause. Using reverse transcription PCR on RNA isolated from cultured CLL cells, we detected increased levels of *IL6* within 1 hour of treatment with Wnt5a before the noted phosphorylation of STAT3. As such, the induced expression of *IL6* preceded the 3 or more hours required for Wnt5a to induce CLL cell activation of STAT3 (Figure 3E). Similarly, we detected IL-6 by ELISA in the media of CLL cells within a few hours after adding exogenous Wnt5a (Figure 3F). These data indicate that the activation of STAT3 largely was caused by autocrine IL-6, which was expressed after Wnt5a-induced activation of ROR1 signaling.

Wnt5a induces production of proinflammatory factors via activation of NF- κ B

Prior studies noted that NLC could stimulate CLL cells to activate NF- κ B,²⁰⁻³² which in turn could influence expression of the genes encoding many of the cytokines and chemokines we found were induced by Wnt5a.³³ We hypothesized that Wnt5a expressed by NLC could contribute to CLL cell activation of NF- κ B. First, we found that Wnt5a indeed could induce phosphorylation of p65 and I κ B α in CLL cells that expressed ROR1 (Figure 4A). Moreover, the capacity of Wnt5a to induce activation of NF- κ B in CLL cells could be blocked by cirmtuzumab or neutralizing antibodies specific for Wnt5a (Figure 4A). As anticipated, we found that NLC also could induce phosphorylation of p65 in both




Figure 2. NLC Express Wnt5a, which can induce ROR1-dependent CLL cell activation of STAT3 and upregulation of ROR1. (A) Flow cytometry analysis of pSTAT3 (Y705) and ROR1 in CLL cells cocultured with or without NLC for 16 hours with control IgG or anti-Wnt5a or cirmtuzumab. (B) Bar figures represent the intensity of ROR1 staining of CLL cells from ROR1^{high} (n = 6) and ROR1^{low} (n = 6) patients cocultured with or without NLC for 16 hours in the presence of control IgG, anti-Wnt5a, or cirmtuzumab. Error bars denote standard deviation. (C) Bar figures represent the levels of pSTAT3 ΔMFI in CLL cells from ROR1^{high} (n = 6) and ROR1^{low} (n = 6) patients cocultured with or without NLC for 16 hours in the presence of control IgG, anti-Wnt5a, or cirmtuzumab. Error bars denote standard deviation. (D) Quantitative real-time PCR analysis and comparison of WNT5A transcript levels in NLC vs CLL cells of the same patient (n = 11). Error bars denote standard deviation. (E) ELISA of Wnt5a in the medium of cell cultures containing 1 × 10⁴/mL NLC and 1 × 10⁴/mL CLL cells for 0, 1, 2, and 5 hours. Error bars denote standard deviation (n = 3).

ROR1^{high} and ROR1^{low} CLL cells (Figure 4B; supplemental Figure 7). Unexpectedly, however, we found that cirmtuzumab or anti-Wnt5a antibodies caused near-complete suppression of the p65 phosphorylation in ROR1^{high} CLL cells induced by co-culture with NLC, which elaborate factors, such as APRIL or BAFF, that also could activate NF-κB (Figure 4B; supplemental Figure 7).³⁰⁻³² As such, these studies reveal that Wnt5a may be a primary factor involved in NLC-induced activation of NF-κB in ROR1-expressing CLL cells.

We reasoned that the capacity of anti-Wnt5a or cirmtuzumab to inhibit the activation of STAT3 in CLL cells was a result of the production of cytokines induced by activation of NF-κB. Consistent with this notion, we observed that levels of IL-6 released into media were elevated in ROR1^{high} CLL cells cocultured with NLC, which could be inhibited by anti-Wnt5a or cirmtuzumab, but not in ROR1^{low} CLL cells (supplemental Figure 8). Also, a small molecule inhibitor of NF-κB, BAY 11-7082, could inhibit Wnt5a-induced CLL cell production of IL-6 and activation of

STAT3 (Figure 4C-D). Furthermore, we found that Wnt5a could induce CLL cells to enhance expression of NF-κB target genes, including *IL6*, and that this effect could be blocked by cirmtuzumab (Figure 4E-F). We also examined this effect in MEC1, which is a human cell line derived from CLL that does not express ROR1. Previously, we demonstrated that MEC1 cells transfected to express ROR1 (MEC1-ROR1 cells) had higher rates of cell proliferation and more effective chemokine-directed migration than parental MEC1 cells.⁴ In this study, we found that MEC1-ROR1 had higher levels of phosphorylated p65 and STAT3, and secreted higher amounts of IL-6 into the media, than MEC1 cells (supplemental Figure 9A-B). Furthermore, knockout of *IL6R* reduced the levels of activated STAT3 in MEC1-ROR1 cells (supplemental Figure 9C-D).

Treatment with cirmtuzumab inhibits CLL cell activation of NF-κB and STAT3

To examine the in vivo significance of our findings, we collected plasma and CLL cells from patients treated with cirmtuzumab in




Figure 3. Wnt5a/ROR1 induces production of proinflammatory factors that lead to phosphorylation of STAT3 in CLL cells. (A) Immunoblot analysis of anti-pSTAT3(Y705) or anti-STAT3 in lysates of CLL cells treated with Wnt5a (200 ng/ml) for 0-6 hours or 24 hours in the presence of control IgG or cirtumzumab. Bar figures represent ratios of density (pSTAT3/Total STAT3). (B) Wnt5a/ROR1 induced production of cytokines in CLL cells. The medium of CLL cells cultured with or without Wnt5a for 16 hours was analyzed by the human cytokine array. (C) Cirtumzumab or neutralizing anti-Wnt5a antibodies blocked Wnt5a-induced cytokine production in CLL cells. The medium of CLL cells cultured with or without Wnt5a, in the presence of control IgG, cirtumzumab, or anti-Wnt5a was analyzed by the human cytokine array. (D) Immunoblot analysis of anti-pSTAT3 (Y705) or anti-STAT3 in lysates of CLL cells treated with control medium, conditioned medium, or IL-6. Tocilizumab or control IgG was added to the CLL cultures 2 hours before treatment. The numbers on the top indicate the relative density of pSTAT3/Total STAT3. The density of the sample treated by conditioned medium only (the 3rd lane) was considered as 1.0. (E) Reverse transcript PCR analysis of IL-6 and β -actin in RNA isolated from purified CLL cells treated with Wnt5a (200 ng/mL) in 0 to 5 hours. (F) ELISA of IL-6 in medium of isolated CLL cells cultured with Wnt5a for 0 to 8 hours in the presence of cirtumzumab or control IgG (n = 5).

a phase 1 clinical trial.¹¹ Immunoblot and RNAseq, respectively, were used to examine for activation of NF- κ B and the transcriptomes of negatively selected CLL cells. ELISA was used to evaluate the concentrations of IL-6 in plasma collected before and after treatment with cirtumzumab. Consistent with our in vitro findings, cirtumzumab downregulated CLL cell expression of NF- κ B and STAT3 target genes in vivo (Figure 5A) and reduced the levels phosphorylated p65 and STAT3 detected in the CLL cells of treated patients (Figure 5B). The levels of IL-6 in plasma also were decreased significantly in patients after treatment with this antibody (Figure 5C; supplemental Table 2).

Discussion

In this study, we found that NLC could induce Wnt5a/ROR1-dependent activation of NF- κ B in CLL, which in turn elicits

autocrine IL-6-induced activation of STAT3. First, we found that levels of ROR1 and pSTAT3 were higher in circulating CD5^{high} CXCR4^{dim} CLL cells than in CD5^{dim} CXCR4^{high} CLL cells in any 1 patient, suggesting that ROR1 and pSTAT3 may be higher in CLL cells within the leukemia microenvironment present in lymphoid tissues. We found NLC-induced Wnt5a/ROR1-dependent activation of STAT3 and upregulation of ROR1 in ROR1⁺ CLL cells by inducing the leukemia cells to produce proinflammatory cytokines, predominantly IL-6, through the activation of NF- κ B.

Prior studies revealed that NF- κ B may be constitutively activated in patients with CLL, thereby contributing to disease progression and CLL-cell resistance to apoptosis.¹⁵ CLL cells present in the microenvironment are more likely to have activated NF- κ B than the CLL cells in the circulation.¹⁶ Activation of NF- κ B in CLL was considered a consequence of CLL-cell exposure to factors such




Figure 4. Wnt5a induces ROR1-dependent production of proinflammatory factors through activation of NF-κB in CLL cells. (A) Immunoblot analysis of lysates from CLL cells treated by Wnt5a (200 ng/mL) for 30 minutes, with or without cirmtuzumab, and probed using anti-phospho-p65, anti-p65, anti-phospho-IkBα, anti-IkBα or anti-β actin. (B) Immunoblot analysis of lysates from CLL cells cocultured with NLC overnight, with or without cirmtuzumab or anti-Wnt5a, and probed with anti-phospho-p65 or anti-p65. (C) ELISA of IL-6 in medium of CLL cells cultured with Wnt5a and BAY 11-7082 (5-20 μM) for 8 hours, as indicated in the text. (D) Immunoblot analysis of lysates from CLL cells treated by Wnt5a and BAY 11-7082 (5-20 μM) for 8 hours that was probed with anti-pSTAT3 (Y705) or anti-STAT3, as indicated in the text. (E) Volcano plot of transcripts of NF-κB target genes expressed by CLL cells stimulated with Wnt5a in the presence of control IgG or cirmtuzumab ($P < .05$ was considered statistically significant). (F) Heat map of transcripts of NF-κB target genes expressed by CLL cells stimulated with Wnt5a in the presence of control IgG or cirmtuzumab.

as BAFF and/or APRIL, as well as B-cell receptor signaling, which are more prominent in the microenvironments of lymphoid tissues than in the blood.^{22,30,31,34} In this study, we found that cirmtuzumab, a mAb that could block ROR1 signaling, inhibited CLL cell phosphorylation of p65 and expression of NF-κB target genes and proteins, such as IL-6, in patients treated with this mAb. These findings indicate that ROR1 signaling is at least

a major contributor to the activation of NF-κB in ROR1-positive CLL cells in vivo.

The in vitro effects of IL-6 on the survival of CLL cells appear context-based.³⁵ Prior studies showed that IL-6 could inhibit the proliferation of CLL cells when stimulated with tumor necrosis factor α or Toll-like receptor 7.^{36,37} However, other studies found




Figure 5. Cirtuzumab inhibits expression of NF-κB and STAT3 target genes in the CLL cells of patients treated with cirtuzumab in phase I clinical trial. (A) GSEA plots of NF-κB and STAT3 target genes on pretreatment vs paired postcirtuzumab CLL cells at D28 of therapy ($n = 3$). Gene set size (Size), enrichment score, normalized ES (NES), and FDR q values are indicated. (B) Immunoblot and flow cytometry analysis of CLL cells isolated from patients before and after treatment with cirtuzumab for phospho-p65 and phospho-STAT3, respectively. (C) The paired concentrations of IL-6 in the plasma of patients before and after treatment with cirtuzumab ($n = 5$).

that IL-6 could enhance the CLL-cell proliferation and resistance to spontaneous apoptosis.^{38,39} A more recent study found that high-level CLL-cell production of IL-6 apparently was associated with high levels of CLL-cell STAT activation and more aggressive disease.⁴⁰ Conceivably, targeting signaling pathways that induce higher levels of IL-6 may have beneficial clinical effects, as have drugs that inhibit B-cell receptor signaling, which also leads to higher levels of JAK/STAT activation in CLL cells.⁴¹ Because drugs that inhibit B-cell receptor signaling, such as ibrutinib, do not inhibit Wnt5a-induced ROR1 signaling in neoplastic B cells,^{9,42} there may be an added advantage to targeting both signaling pathways in patients with CLL.

In addition to IL-6, we observed that Wnt5a/ROR1 signaling also induced CLL cells to make other proinflammatory factors, such as IL-8, CCL3, and CCL4, which also may play a role in promoting disease progression in patients with CLL.⁴³⁻⁴⁵ Moreover, the current study demonstrates that cirtuzumab also could inhibit Wnt5a-induced CLL-cell expression of IL-6 and these other cytokines/chemokines, which help recruit nonneoplastic accessory cells to the leukemia microenvironment.⁴⁶

Wnt5a not only is expressed by accessory cells in the microenvironment but also is present at high levels in the blood of patients with CLL relative to that of age-matched healthy adults.⁴ Although amphipathic Wnt factors typically are thought of being




Figure 6. A model for NLC-Wnt5a-induced proinflammatory factors through ROR1-dependent NF- κ B activation, which induces production of IL-6 and autocrine activation of STAT3.

involved primarily in cognate cell-cell interactions, Wnt5a may be released from NLC and carried on plasma proteins to effect activation of CLL cells within the circulation. This may account in part for the high levels of activated NF- κ B found in the circulating CLL cells of some patients, whose leukemia cells are relatively resistant to spontaneous or drug-induced apoptosis.¹⁴ We showed that the CLL cells of patients treated by cirmtuzumab had reduced activation of NF- κ B and reduced expression of NF- κ B target genes. Our data support a model proposing that Wnt5a expressed by NLC induces leukemia cell activation of NF- κ B, which may enhance CLL-cell survival and induce expression of proinflammatory factors by CLL cells in vivo that in turn could activate CLL cells in an autocrine fashion (Figure 6). Consistent with the notion that NLC expression of Wnt5a contributes to a ROR1-dependent survival stimulus for CLL cells, we note that cirmtuzumab or neutralizing antibodies to Wnt5a can mitigate the survival advantage of CLL cells cocultured with NLC cells in vitro (supplemental Figure 10).

Collectively, these data indicate that Wnt5a/ROR1-dependent signaling induced by NLC contributes to the noted activation of NF- κ B, leading to cytokine production and activation of STAT3 in CLL. As such, agents such as cirmtuzumab that can block ROR1 signaling can suppress leukemia-cell activation of NF- κ B and production of factors that contribute to

disease progression, providing another rationale for evaluating the potential benefit of cirmtuzumab in patients with this disease.

Acknowledgments

The authors thank the California Institute for Regenerative Medicine (grant DR3-06924) for supporting us in the generation of anti-ROR1 mAb, cirmtuzumab. This work was supported by the UC San Diego Foundation Blood Cancer Research Fund, a SCOR grant (7005-14) from the Leukemia and Lymphoma Society, and a grant (R01-CA236361 [T.J.K.]) from the National Institutes of Health, National Cancer Institute. The RNA sequencing was conducted at the IGM Genomics Center, University of California, San Diego, La Jolla, CA (National Institutes of Health, National Cancer Institute grant P30CA023100).

Authorship

Contribution: Y.C. and T.J.K. conceived the project; Y.C., L.C., J.Y., M.Y.C., and L.Z. performed the experiments; Y.C., L.C., J.Y., E.M.G., M.Y.C., S.Z., E.S.-L., G.F.W., K.M., L.Z.R., C.J., and T.J.K. analyzed data; and Y.C. and T.J.K. wrote the manuscript.

Conflict-of-interest disclosure: Cirmtuzumab was developed by T.J.K. and licensed from the University of California, San Diego, to Oncternal Therapeutics, Inc., which provided stock/options and funding to the University of California, San Diego, and to T.J.K. The remaining authors declare no competing financial interests.

Footnotes

Submitted 31 January 2019; accepted 21 June 2019. Prepublished online as *Blood* First Edition paper, 13 August 2019; DOI 10.1182/blood.2019001366.

*Y.C. and L.C. contributed equally to this work.

This work was presented in part at the 60th annual meeting of the American Society of Hematology, San Diego, CA, 1-4 December 2018.

The online version of this article contains a data supplement.

The publication costs of this article were defrayed in part by page charge payment. Therefore, and solely to indicate this fact, this article is hereby marked "advertisement" in accordance with 18 USC section 1734.

REFERENCES

- Zhang S, Chen L, Wang-Rodriguez J, et al. The onco-embryonic antigen ROR1 is expressed by a variety of human cancers. *Am J Pathol*. 2012;181(6):1903-1910.
- Fukuda T, Chen L, Endo T, et al. Antisera induced by infusions of autologous Ad-CD154-leukemia B cells identify ROR1 as an oncofetal antigen and receptor for Wnt5a. *Proc Natl Acad Sci USA*. 2008;105(8):3047-3052.
- Widhopf GF II, Cui B, Ghia EM, et al. ROR1 can interact with TCL1 and enhance leukemogenesis in Eμ-TCL1 transgenic mice. *Proc Natl Acad Sci USA*. 2014;111(2):793-798.
- Yu J, Chen L, Cui B, et al. Wnt5a induces ROR1/ROR2 heterooligomerization to enhance leukemia chemotaxis and proliferation. *J Clin Invest*. 2016;126(2):585-598.
- Cui B, Ghia EM, Chen L, et al. High-level ROR1 associates with accelerated disease progression in chronic lymphocytic leukemia. *Blood*. 2016;128(25):2931-2940.
- Hasan MK, Yu J, Widhopf GF II, et al. Wnt5a induces ROR1 to recruit DOCK2 to activate Rac1/2 in chronic lymphocytic leukemia. *Blood*. 2018;132(2):170-178.
- Hasan MK, Yu J, Chen L, et al. Wnt5a induces ROR1 to complex with HS1 to enhance migration of chronic lymphocytic leukemia cells. *Leukemia*. 2017;31(12):2615-2622.
- Yu J, Chen L, Chen Y, et al. Wnt5a induces ROR1 to associate with 14-3-3ζ for enhanced chemotaxis and proliferation of chronic lymphocytic leukemia cells. *Leukemia*. 2017;31(12):2608-2614.
- Yu J, Chen L, Cui B, et al. Cirmetuzumab inhibits Wnt5a-induced Rac1 activation in chronic lymphocytic leukemia treated with ibrutinib. *Leukemia*. 2017;31(6):1333-1339.
- Choi MY, Widhopf GF II, Wu CC, et al. Pre-clinical specificity and safety of UC-961, a first-in-class monoclonal antibody targeting ROR1. *Clin Lymphoma Myeloma Leuk*. 2015;15(suppl):S167-S169.
- Choi MY, Widhopf GF 2nd, Ghia EM, et al. Phase I trial: cirmetuzumab inhibits ROR1 signaling and stemness signatures in patients with chronic lymphocytic leukemia. *Cell Stem Cell*. 2018;22(6):951-959.
- Hasan MK, Rassenti L, Widhopf GF II, Yu J, Kipps TJ. Wnt5a causes ROR1 to complex and activate cortactin to enhance migration of chronic lymphocytic leukemia cells. *Leukemia*. 2019;33(3):653-661.
- Dissanayake SK, Olkhanud PB, O'Connell MP, et al. Wnt5A regulates expression of tumor-associated antigens in melanoma via changes in signal transducers and activators of transcription 3 phosphorylation. *Cancer Res*. 2008;68(24):10205-10214.
- Liu FT, Jia L, Wang P, Wang H, Farren TW, Agrawal SG. STAT3 and NF-κB cooperatively control in vitro spontaneous apoptosis and poor chemo-responsiveness in patients with chronic lymphocytic leukemia. *Oncotarget*. 2016;7(22):32031-32045.
- Mansouri L, Papakonstantinou N, Ntousa S, Stamatopoulos K, Rosenquist R. NF-κB activation in chronic lymphocytic leukemia: A point of convergence of external triggers and intrinsic lesions. *Semin Cancer Biol*. 2016;39:40-48.
- Herishanu Y, Pérez-Galán P, Liu D, et al. The lymph node microenvironment promotes B-cell receptor signaling, NF-kappaB activation, and tumor proliferation in chronic lymphocytic leukemia. *Blood*. 2011;117(2):563-574.
- Hazan-Halevy I, Harris D, Liu Z, et al. STAT3 is constitutively phosphorylated on serine 727 residues, binds DNA, and activates transcription in CLL cells. *Blood*. 2010;115(14):2852-2863.
- Li P, Harris D, Liu Z, Liu J, Keating M, Estrov Z. Stat3 activates the receptor tyrosine kinase like orphan receptor-1 gene in chronic lymphocytic leukemia cells. *PLoS One*. 2010;5(7):e11859.
- Janda CY, Waghay D, Levin AM, Thomas C, Garcia KC. Structural basis of Wnt recognition by Frizzled. *Science*. 2012;337(6090):59-64.
- Janovska P, Poppova L, Plevova K, et al. Autocrine signaling by Wnt-5a deregulates chemotaxis of leukemic cells and predicts clinical outcome in chronic lymphocytic leukemia. *Clin Cancer Res*. 2016;22(2):459-469.
- Tsukada N, Burger JA, Zvaifler NJ, Kipps TJ. Distinctive features of "nurselike" cells that differentiate in the context of chronic lymphocytic leukemia. *Blood*. 2002;99(3):1030-1037.
- Kipps TJ, Stevenson FK, Wu CJ, et al. Chronic lymphocytic leukaemia. *Nat Rev Dis Primers*. 2017;3(1):16096.
- Shao Y, Zheng Q, Wang W, Xin N, Song X, Zhao C. Biological functions of macrophage-derived Wnt5a, and its roles in human diseases. *Oncotarget*. 2016;7(41):67674-67684.
- Calissano C, Damle RN, Marsilio S, et al. Intracloal complexity in chronic lymphocytic leukemia: fractions enriched in recently born/divided and older/quiescent cells. *Mol Med*. 2011;17(11-12):1374-1382.
- Cui B, Chen L, Zhang S, et al. MicroRNA-155 influences B-cell receptor signaling and associates with aggressive disease in chronic lymphocytic leukemia. *Blood*. 2014;124(4):546-554.
- Subramanian A, Tamayo P, Mootha VK, et al. Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles. *Proc Natl Acad Sci USA*. 2005;102(43):15545-15550.
- Feuerhake F, Kutok JL, Monti S, et al. NF-kappaB activity, function, and target-gene signatures in primary mediastinal large B-cell lymphoma and diffuse large B-cell lymphoma subtypes. *Blood*. 2005;106(4):1392-1399.
- Dauer DJ, Ferraro B, Song L, et al. Stat3 regulates genes common to both wound healing and cancer. *Oncogene*. 2005;24(21):3397-3408.
- Burger JA, Tsukada N, Burger M, Zvaifler NJ, Dell'Aquila M, Kipps TJ. Blood-derived nurse-like cells protect chronic lymphocytic leukemia B cells from spontaneous apoptosis through stromal cell-derived factor-1. *Blood*. 2000;96(8):2655-2663.
- Nishio M, Endo T, Tsukada N, et al. Nurselike cells express BAFF and APRIL, which can promote survival of chronic lymphocytic leukemia cells via a paracrine pathway distinct from that of SDF-1α. *Blood*. 2005;106(3):1012-1020.
- Endo T, Nishio M, Enzler T, et al. BAFF and APRIL support chronic lymphocytic leukemia B-cell survival through activation of the canonical NF-kappaB pathway. *Blood*. 2007;109(2):703-710.
- Zhang W, Kater AP, Widhopf GF II, et al. B-cell activating factor and v-Myc myelocytomatosis viral oncogene homolog (c-Myc) influence progression of chronic lymphocytic leukemia. *Proc Natl Acad Sci USA*. 2010;107(44):18956-18960.
- Pahl HL. Activators and target genes of Rel/NF-kappaB transcription factors. *Oncogene*. 1999;18(49):6853-6866.
- Rozovski U, Harris DM, Li P, et al. Activation of the B-cell receptor successively activates NF-κB and STAT3 in chronic lymphocytic leukemia cells. *Int J Cancer*. 2017;141(10):2076-2081.
- Lapalombella R. Interleukin-6 in CLL: accelerator or brake? *Blood*. 2015;126(6):697-698.

36. Li Y, Shi Y, McCaw L, et al. Microenvironmental interleukin-6 suppresses toll-like receptor signaling in human leukemia cells through miR-17/19A. *Blood*. 2015;126(6):766-778.
37. Aderka D, Maor Y, Novick D, et al. Interleukin-6 inhibits the proliferation of B-chronic lymphocytic leukemia cells that is induced by tumor necrosis factor-alpha or -beta. *Blood*. 1993;81(8):2076-2084.
38. Moreno A, Villar ML, Cámara C, et al. Interleukin-6 dimers produced by endothelial cells inhibit apoptosis of B-chronic lymphocytic leukemia cells. *Blood*. 2001;97(1):242-249.
39. Hahn T, Shulman L, Karov Y, Vorst E, Berrebi A. Involvement of interleukin-6 in the autocrine stimulation of chronic lymphocytic leukemia B cells by tumor necrosis factor. *Leuk Lymphoma*. 1991;5(suppl 1):65-69.
40. Wang HQ, Jia L, Li YT, Farren T, Agrawal SG, Liu FT. Increased autocrine interleukin-6 production is significantly associated with worse clinical outcome in patients with chronic lymphocytic leukemia. *J Cell Physiol*. 2019; 234(8):13994-14006.
41. Rozovski U, Wu JY, Harris DM, et al. Stimulation of the B-cell receptor activates the JAK2/STAT3 signaling pathway in chronic lymphocytic leukemia cells. *Blood*. 2014; 123(24):3797-3802.
42. Yu J, Chen Y, Chen L, et al. Cirtumzumab inhibits ibrutinib-resistant, Wnt5a-induced Rac1 activation and proliferation in mantle cell lymphoma. *Oncotarget*. 2018;9(37): 24731-24736.
43. Yoon JY, Lafarge S, Dawe D, et al. Association of interleukin-6 and interleukin-8 with poor prognosis in elderly patients with chronic lymphocytic leukemia. *Leuk Lymphoma*. 2012;53(9):1735-1742.
44. Yan XJ, Dozmorov I, Li W, et al. Identification of outcome-correlated cytokine clusters in chronic lymphocytic leukemia. *Blood*. 2011; 118(19):5201-5210.
45. Sivina M, Hartmann E, Kipps TJ, et al. CCL3 (MIP-1α) plasma levels and the risk for disease progression in chronic lymphocytic leukemia. *Blood*. 2011;117(5):1662-1669.
46. Burger JA, Quiroga MP, Hartmann E, et al. High-level expression of the T-cell chemokines CCL3 and CCL4 by chronic lymphocytic leukemia B cells in nurselike cell cocultures and after BCR stimulation. *Blood*. 2009; 113(13):3050-3058.