

THROMBOSIS AND HEMOSTASIS

Prognostic factors for VTE and bleeding in hospitalized medical patients: a systematic review and meta-analysis

Andrea J. Darzi,^{1,2} Samer G. Karam,^{1,2} Rana Charide,³ Itziar Etxeandia-Ikobaltzeta,^{1,2} Mary Cushman,^{4,5} Michael K. Gould,⁶ Lawrence Mbuagbaw,^{1,2,7,8} Frederick A. Spencer,⁹ Alex C. Spyropoulos,¹⁰ Michael B. Streiff,¹¹ Scott Woller,¹² Neil A. Zakai,^{4,5} Federico Germini,^{1,2,9,13} Marta Rigoni,¹⁴ Arnav Agarwal,^{1,2,15} Rami Z. Morsi,³ Alfonso Iorio,^{1,2,9} Elie A. Akl,^{1-3,16} and Holger J. Schünemann^{1,2,9}

¹Michael G. DeGroot Cochrane Canada and ²McMaster GRADE (Grading of Recommendations Assessment, Development, and Evaluation) Centres, Department of Health Research Methods, Evidence, and Impact, McMaster University, Hamilton, ON, Canada; ³American University of Beirut (AUB) GRADE Center, Clinical Research Institute, American University of Beirut, Beirut, Lebanon; ⁴Department of Medicine and ⁵Department of Pathology and Laboratory Medicine, Larner College of Medicine, University of Vermont, Burlington, VT; ⁶Department of Research and Evaluation, Kaiser Permanente Southern California, Pasadena, CA; ⁷Biostatistics Unit, Father Sean O'Sullivan Research Centre, St Joseph's Healthcare, Hamilton, ON, Canada; ⁸Centre for Development of Best Practices in Health (CDBPH), Yaoundé Central Hospital, Yaoundé, Cameroon; ⁹Department of Medicine, McMaster University, Hamilton, ON, Canada; ¹⁰The Donald and Barbara Zucker School of Medicine at Hofstra/Northwell, Department of Medicine, Northwell Health at Lenox Hill Hospital, Manhasset, NY; ¹¹Division of Hematology, Department of Medicine, The Johns Hopkins University School of Medicine, Baltimore, MD; ¹²Department of Medicine, Intermountain Medical Center, University of Utah School of Medicine, Salt Lake City, UT; ¹³Department of Health Sciences, Università degli Studi di Milano, Milan, Italy; ¹⁴Healthcare Research and Innovation Program, Health Technology Assessment Unit, Fondazione Bruno Kessler, Trento, Italy; ¹⁵Faculty of Medicine, University of Toronto, Toronto, ON, Canada; and ¹⁶Department of Internal Medicine, AUB, Beirut, Lebanon

KEY POINTS

- Using a systematic approach, we identified 23 prognostic factors for venous thromboembolism and 15 for bleeding.
- We identified several prognostic factors for VTE and bleeding that are not considered in most of the widely used risk assessment models.

There may be many predictors of venous thromboembolism (VTE) and bleeding in hospitalized medical patients, but until now, systematic reviews and assessments of the certainty of the evidence have not been published. We conducted a systematic review to identify prognostic factors for VTE and bleeding in hospitalized medical patients and searched Medline and EMBASE from inception through May 2018. We considered studies that identified potential prognostic factors for VTE and bleeding in hospitalized adult medical patients. Reviewers extracted data in duplicate and independently and assessed the certainty of the evidence using the Grading of Recommendations Assessment, Development, and Evaluation approach. Of 69 410 citations, we included 17 studies in our analysis: 14 that reported on VTE, and 3 that reported on bleeding. For VTE, moderate-certainty evidence showed a probable association with older age; elevated C-reactive protein (CRP), D-dimer, and fibrinogen levels; tachycardia; thrombocytosis; leukocytosis; fever; leg edema; lower Barthel Index (BI) score; immobility; paresis; previous history of VTE; thrombophilia; malignancy; critical illness; and infections. For bleeding, moderate-certainty evidence showed a probable association with older age, sex, anemia, obesity, low

hemoglobin, gastroduodenal ulcers, rehospitalization, critical illness, thrombocytopenia, blood dyscrasias, hepatic disease, renal failure, antithrombotic medication, and presence of a central venous catheter. Elevated CRP, a lower BI, a history of malignancy, and elevated heart rate are not included in most VTE risk assessment models. This study informs risk prediction in the management of hospitalized medical patients for VTE and bleeding; it also informs guidelines for VTE prevention and future research. (*Blood*. 2020;135(20):1788-1810)

Introduction

Venous thromboembolism (VTE), comprising deep vein thrombosis (DVT) and pulmonary embolism (PE), has an annual incidence of ~1 per 1000 in adult populations.^{1,2} VTE is a major burden in hospitalized medical patients. Medical patients can be classified as having acute, critical, or chronic medical illness, and their risk for both VTE and bleeding may depend on the severity of their medical illness. The incidence of VTE in hospitalized acutely ill medical patients detected by screening is up to 14.9%.³ From 50% to 70% of symptomatic

VTEs and 70% to 80% of fatal PEs occur in acute medically ill patients.⁴⁻⁷

The risk of hospital-acquired VTE is reduced by using pharmacological and nonpharmacological interventions, but these interventions are not without potential patient harms. Risk assessment models (RAMs) have been used in hospitalized medically ill patients to stratify the different subsets of patients by their risk of having a VTE or clinically significant bleeding event.⁸ This stratification may then support optimized management for the

prevention of either outcome.⁹ A RAM is a formal combination of multiple predictors from which risks of a specific end point can be calculated for individuals. The value of using RAMs include generation of specific baseline risks to inform recommendations for a strata of patients and calculation of a predicted risk of an outcome for an individual patient (eg, VTE or bleeding) based on the patient's characteristics (ie, the prognostic factors). Implementation of RAMs in the field of VTE prevention can be accomplished by embedding them in clinical encounters or decision aids to individualize the use of guideline recommendations. However, the application of RAMs is variable in current practice.³

Most RAMs are developed using data registries that are not based on a systematic review of all potential prognostic factors.¹⁰ However, guiding principles for developing RAMs describe the importance of identifying prognostic factors through systematic reviews.¹⁰ We identified only one systematic review, conducted 11 years ago, that evaluated VTE as an outcome in medical patients, but the effect sizes of the prognostic factors were not subjected to meta-analysis, and bleeding risk, critical for balancing benefits and harms in these patients, was not included as an outcome.¹¹

Therefore, our aim was to conduct a systematic review of prognostic factors for VTE and bleeding in hospitalized medical patients that may inform management, future guideline recommendations, and the development of RAMs in hospitalized medical patients.

Methods

We conducted a systematic review using Cochrane methodology to identify studies that reported on prognostic factors for VTE and bleeding in hospitalized medical patients.¹² We developed a protocol that was reviewed and revised by the co-authors, but we did not register it because of confidentiality clauses in the research contract.

Data sources and searches

We searched Medline and EMBASE from inception through May 2018 with the assistance of an information scientist. Supplemental Table 1 (available on the *Blood* Web site) provides detailed descriptions of the search strategy. The search included both medical subject heading (MeSH) terms and text-word terms. It combined VTE-related terms with primary prevention terms and 2 search blocks defining prognosis and prediction guide filters. We used no language restrictions or time limits.

Study selection

Four teams of 2 reviewers participated in training and calibration exercises before starting the screening processes. The teams screened independently and in duplicate the titles and abstracts of all the retrieved citations. They then retrieved the full texts of all citations judged as potentially eligible by at least 1 of the reviewers on each team. The reviewers screened the full texts independently and in duplicate and compared their results. A third senior reviewer resolved disagreements when necessary. Reviewers used a standardized screening form and conducted calibration exercises before the screening process.

The eligibility criteria for study selection comprised the following characteristics.

Population We included studies that evaluated adult medical patients who were acutely, critically, or chronically ill. We also included studies in which the population included nonmedical patients or medical patients with a recent history of surgery or trauma if the final regression model adjusted for these factors. We included studies if less than 10% of the population was receiving thromboprophylaxis or if the statistical analysis adjusted for the use of thromboprophylaxis. Thromboprophylaxis included the use of anticoagulation therapy (ie, warfarin, low-molecular-weight heparin, and unfractionated heparin), antiplatelet therapy (ie, aspirin), or mechanical prophylaxis (ie, elastic stockings or intermittent pneumatic compression).

We excluded studies if the population did not reflect the general population of interest, such as studies of only selected types of cancer patients.^{13,14} We defined acutely ill medical patients as those hospitalized for a medical illness including heart failure, respiratory insufficiency, stroke, and infectious or inflammatory diseases requiring urgent care.² Critically ill patients were those who had an immediately life-threatening condition and were admitted to an intensive or critical care unit.² Chronically ill medical patients included those with acute exacerbations of chronic medical conditions who required hospitalization.²

Exposure We investigated all prognostic factors reported in individual studies.

Comparisons We investigated the absence or different levels of the prognostic factor.

Outcomes Studies had to report on the outcomes VTE or bleeding. VTE was defined as any symptomatic or asymptomatic DVT or PE within 90 days after discharge. Bleeding included major or nonmajor but clinically significant bleeding within 90 days after discharge.⁸

Setting Studies that included patients who were admitted to a nonsurgical inpatient ward.

Type of study We included prognostic factor and RAM studies that are based on typologies of prognosis proposed by Iorio and colleagues,¹⁵ founded on the PROGNosis REsearch Strategy (PROGRESS) Group framework.¹⁶

Data extraction

Two reviewers abstracted data independently and in duplicate from all eligible studies using standardized forms. Reviewers compared and discussed results and consulted a third reviewer in case of any disagreement. We conducted calibration exercises and piloting of all forms before the start of the data abstraction process. All eligible studies were published in English.

For all identified studies, RAMs, and prognostic factor studies, the reviewers abstracted data on the following characteristics:

- Study context (eg, country and year of publication)
- Type of prediction model study (development, validation, and impact)

- Study design (eg, cohort or case-control; duration of follow-up)
- Population and their demographics (eg, sample size, age, number of centers, and administration of prophylaxis and what type)
- Outcomes (VTE and bleeding)
- Prognostic factors, definitions, and measurement methods (including thresholds used for continuous predictors)
- Measures of association (eg, odds ratio [OR], risk ratio, and hazard ratio)

Quality assessment

Risk-of-bias assessment

We assessed the risk of bias in the included studies by using the Prediction Study Risk of Bias Assessment Tool (PROBAST) for RAM studies¹⁷ and the Quality in Prognosis Studies tool (QUIPS) for prognostic factor studies.¹⁸⁻²⁰

Synthesis of findings and certainty of evidence assessment

We presented the results of the studies, including the individual prognostic factors, in both tabular and narrative formats. We also described the identified prognostic factor studies and the measure of association with the outcomes of interest. We performed an assessment of the certainty of evidence for each of the prognostic factors per outcome, based on the GRADE approach.¹⁵ The approach considers the following domains: risk of bias, indirectness, inconsistency, imprecision, and publication bias. We developed evidence profiles and rated the overall certainty of evidence as high, moderate, and low or very low, depending on the grading of the individual domains.¹⁵ We narratively described the strength of the association using the terms “there is,” “there probably is,” or “there may be,” depending on whether the quality of the evidence was “high,” “moderate,” or “low/very low,” respectively.

Data synthesis and analysis

We standardized the units of measurement for each prognostic factor, unifying the direction of the predictors, adjusting the weights of the studies, and calculating crude-effect estimates when not provided.²¹ When possible, we performed meta-analysis of all prognostic factors associated with the outcomes VTE and bleeding that were reported by more than 1 study. We then presented the effect estimate as OR and the corresponding 95% confidence interval (CI). In studies that reported the measure of association as a hazard ratio or risk ratio, we converted them to ORs using the baseline risk (incidence of those not on prophylaxis having VTE or bleeding out of the total sample) reported in the studies.^{22,23} We conducted a meta-analysis of associations using the generic inverse variance-based method to produce an overall measure of association. We used the crude effect estimates when the adjusted estimates were not provided. We explored consistency of the associations between the results of our meta-analysis and studies reporting the same predictors that could not be pooled. All analyses used random-effect models applying the prognosis module in Review Manager, version 5.3.²⁴

Results

Figure 1 is a Preferred Reporting Items for Systematic Reviews and Meta-Analyses flowchart. Our search identified 69 410 citations, of which we included 807 studies for full text assessment. Seventeen studies fulfilled the inclusion criteria for evaluating VTE or bleeding outcomes or both.^{7,8,25-39}

Description of included studies

Table 1 describes the characteristics of the included studies reporting on the outcomes VTE and/or bleeding. Eight studies were prognostic factor studies: 4 were prognostic model development studies, and 5 were external validation studies. Five studies were retrospective case-control studies,²⁵⁻²⁹ 2 of which were multicenter^{25,26}; 5 were retrospective cohorts,³⁰⁻³⁴ 3 of which were multicenter.^{31,32,34} Seven studies were prospective cohorts,^{7,8,35-39} 4 of which were multicenter.^{7,8,36,39} The included studies were conducted in the United States (n = 9), China (n = 3), Canada (n = 2), United Kingdom (n = 1), Poland (n = 1), and Japan (n = 1). Of the 14 included studies for VTE (defined as proximal DVT or PE), 9 reported on symptomatic VTE only.^{7,25-31,34} The other 5 studies reported on both symptomatic and asymptomatic VTE.³⁵⁻³⁹ The follow-up time was up to 3 months in 12 of 14 studies reporting on VTE. The other 2 studies, Zhou et al²⁹ and Yi et al,³⁹ had a follow-up time of 6 months and 1 year, respectively, but also reported the occurrence of VTE during hospitalization. In accordance with our protocol, we used the incidence of VTE during hospitalization from those studies. The 3 studies that reported on bleeding (major or clinically relevant) had a follow-up time up to 1 month.^{8,32,33} Of the 14 studies reporting on VTE, 12 studies included patients who received thromboprophylaxis in 0.4% to 67% of the patients. Of those, 2 studies^{35,36} included less than 10% of patients on thromboprophylaxis, and 10 studies adjusted for prophylaxis in the statistical analysis.^{7,26-31,34,38,39} As for bleeding, all 3 studies reported prophylaxis use in 9% to 70%, which was accounted for in their analysis.^{8,32,33}


Risk-of-bias assessment

Risk of bias was serious across all identified studies, each presenting risk of bias in at least 1 domain or item (Tables 2 and 3). Among the 17 included studies, 10 were retrospective, which may have introduced classification bias.²⁵⁻³⁴ Seven of the 8 prognostic factor studies included only the variables significant in bivariable analysis in their final regression model and did not present any data for nonsignificant predictors in their adjusted analysis.^{30,32,33,36-39} Two of the 8 prognostic factor studies^{32,33} and 4 of the 9 prognostic model development or validation studies did not have a clear description of appropriate outcome measurement.²⁶⁻²⁹ We detected no evidence of publication bias through visual assessment of asymmetry of the funnel plot for each pooled predictor in those that included at least 10 studies (Tables 2 and 3). Supplemental Table 2 provides the detailed judgements for each of the risk-of-bias domain criteria.

Prognostic factors for VTE in hospitalized medical patients

Investigated were 29 candidate prognostic factors for VTE from 14 studies including 15 1714 patients. Table 2 provides the evidence profile for VTE-related prognostic factors. Supplemental Figure 1 provides the forest plots of the meta-analysis of each of the prognostic factors.

Figure 1. Preferred Reporting Items for Systematic Reviews and Meta-Analysis flowchart.



Demographic factors We found moderate-certainty evidence that there is probably an association between risk of any VTE and age ≥ 60 y (OR, 1.34; 95% CI, 1.17-1.55)^{7,25-27,29,31,34-37,39} and that there is probably little to no association between risk of any VTE and sex (males vs females: OR, 1.03; 95% CI, 0.80-1.33).^{27,34-37}

Functional factors There was moderate-certainty evidence for a probable association between risk of any VTE and lower Barthel Index scores (BI ≤ 9 ; OR, 8.30; 95% CI, 2.70-25.52)³⁷; immobility, defined as confinement to bed for >72 hours or >7 days or bedridden or nonambulatory (OR, 3.17; 95% CI, 2.18-4.62)^{7,25,26,28,29,31,37,39}; and paresis (OR, 2.97; 95% CI, 1.20-7.36).^{7,25,26,37}

Medical illness and patient history factors We identified moderate-certainty evidence for an association between risk of any VTE and history of VTE (OR, 6.08; 95% CI, 3.71-9.97)^{7,25-27,29-31,36}; thrombophilia, defined as familial or acquired disorder of the hemostatic system (OR, 5.88; 95% CI, 2.80-12.35)^{7,25,26,29,34}; history of malignancy (OR, 3.20; 95% CI, 2.14-4.79)²⁷; active malignancy defined as the presence of cancer on admission or within the past year (OR, 2.65; 95% CI, 1.79-3.91)^{7,25,27,28,30,31,34,36,37}; critical illness, defined as intensive care unit (ICU) or coronary care unit (CCU) stay or need for resuscitation (OR, 1.65; 95% CI, 1.39-1.95)^{7,25-27,29,34,36}; and infections including cellulitis, pneumonia, and sepsis (OR, 1.48; 95% CI, 1.16-1.89).^{27-29,31,36}

We found low-certainty evidence that there may be an association between risk of any VTE and history of heart failure (OR, of 2.68; 95% CI, 1.11-6.44)^{27,29,35}; autoimmune diseases, including rheumatological diseases and inflammatory diseases (OR, 2.33; 95% CI, 1.13-4.83)^{31,34,36,39}; central venous catheter (CVC) use (OR, 2.05; 95% CI, 0.74-5.65)^{31,34}; and severe stroke, defined as acute ischemic stroke (OR, 1.79; 95% CI, 0.77-4.18).^{29,31,32,35,36} The findings for severe stroke when assessed using a diagnostic tool, the National Institutes of Health Stroke Scale (NIHSS), were consistent with our results (Table 2).³⁵ We also identified low-certainty evidence that there may be an association between risk of any VTE and current tobacco use (OR, 1.59; 95% CI, 0.28-9.03); however, there may be little to no association between risk of any VTE and previous tobacco use (OR, 0.97; 95% CI, 0.24-3.92).

Furthermore, we identified low-certainty evidence that there may be little to no association between the risk of any VTE and respiratory failure (OR, 1.04; 95% CI, 0.69-1.58), coronary artery disease (CAD; OR, 1.01; 95% CI, 0.33-3.09), acute heart failure (OR, 0.82; 95% CI, 0.42-1.60)^{31,36} and hormone use (OR, 0.8; 95% CI, 0.36-1.78).^{29,31,36,37}

We found very-low-certainty evidence that there may be little to no association between risk of any VTE and chronic renal failure (OR, 0.76; 95% CI, 0.18-3.18).³⁶

Laboratory and physical examination factors There is moderate-certainty evidence of an association between risk of any VTE and C-reactive protein (CRP) > 10 mg/L (OR, 10.10; 95% CI, 1.93-52.85)³⁵

Table 1. Study characteristics

First author, year (country)	Population (sample size)	Time frame (y)	Mean age, y (SD)*	Study type (number of centers and study design)	Prophylaxis (%)	Outcome (number of events)	Diagnostic methods	Follow-up time	Variables in multivariate logistic regression, when applicable
Venous thromboembolism									
Spyropoulos et al ¹⁷ (US)	Acutely ill medical patients (N = 15 156)	2002-2006	68 (52-79)*	Prognostic model development (multicenter prospective cohort)	VTE prophylaxis (adjusted-dose warfarin, elastic stockings, LMWH, unfractionated heparin, intermittent pneumatic compression, and aspirin): 44% The model was adjusted for VTE prophylaxis.	Any symptomatic VTE (lower extremity DVT, and PE) (n = 184)	Clinically observed VTE DVT verified by positive venogram or compression ultrasonography. PE verified by positive lung scan, pulmonary angiogram, or spiral CT scan. Fatal PE was defined as PE diagnosed at autopsy or, in the absence of autopsy, when PE was considered the most likely cause of death.	92 d	IMPROVE RAM-related factors: Age >60 y; prior cancer, prior VTE, ICU/CCU stay, lower limb paralysis, immobility Known thrombophilia
Mahan et al ²⁵ (Canada)	Acute medical patients (N = 417). VTE cases: n = 139; non-VTE controls: n = 278)	2005-2011	Cases: 68; controls: 65	External validation (multicenter retrospective case-control)	VTE prophylaxis, 0%	Any symptomatic VTE (lower extremity DVT, and PE; n = 139)	PE, verified by a positive pulmonary angiogram, spiral computed tomography, or high-probability ventilation/perfusion scan or at autopsy. Lower extremity DVT verified by positive compression ultrasonography, computed tomography, or magnetic resonance imaging, or at autopsy.	92 d	IMPROVE RAM-related factors: Previous VTE; known thrombophilia; lower limb paralysis; current cancer; immobilization ≥7; ICU/CCU stay; age >60 y

BI, Barthel Index; BMI, body mass index; COPD, chronic obstructive pulmonary disease; CTA, computed tomographic angiography; GI, gastrointestinal; ICD-9-CM, International Classification of Diseases, Ninth Revision, Clinical Modification; IVc, inferior vena cava; LMWH, low-molecular-weight heparin; mRS, modified Rankin Score; NYHA, New York Heart Association; RBC, red blood cell.

*Caprini factors: stroke; acute spinal cord injury or paralysis (<1 mo); hip, pelvis, or leg fracture (<1 mo); multiple trauma (<1 mo); age ≥75; history of VTE; family history of VTE; history of thrombophilia; heparin-induced thrombocytopenia; age (41-60); age (61-74); positive history of cancer; immobilizing plaster cast; congestive heart failure; COPD or pulmonary function; inflammatory bowel disease; severe long disease (including pneumonia); acute myocardial infarction; sepsis (<1 mo); surgery (<1 mo); postpartum (<1 mo); history of unexpected stillborn infant, recurrent spontaneous abortion (≥ 3) or premature birth; varicose veins; BMI >25 kg/m²; swollen legs (current); CVC present on admission; immobile or not ambulating; and hormone replacement therapy or oral contraceptives.

†Padua factors: active cancer; previous VTE; reduced mobility; known thrombophilia; recent trauma and/or surgery; elderly age; heart and/or respiratory failure; acute myocardial infarction or ischemic stroke; acute infection and/or rheumatologic disorder; obesity; ongoing hormone treatment, and VTE prophylaxis.

Table 1. (continued)

First author, year (country)	Population (sample size)	Time frame (y)	Mean age, y (SD)*	Study type (number of centers and study design)	Prophylaxis (%)	Outcome (number of events)	Diagnostic methods	Follow-up time	Variables in multivariate logistic regression, when applicable
Rosenberg et al ¹⁶ (US)	Medical patients (N = 539; VTE cases: n = 135 and non-VTE controls: n = 404)	2009-2013	67	External validation (multicenter retrospective case-control)	Any prophylaxis in VTE cases: 49% (of those 44% pharmacological VTE and 5% mechanical) Any prophylaxis in non-VTE controls: 45% (of those 40% pharmacological VTE and 5% mechanical) The results were essentially unchanged when the cases and controls were stratified into groups that received VTE prophylaxis, including pharmacological prophylaxis during hospitalization and those that did not.	Any symptomatic VTE (n = 135)	VTE events identified using ICD-9 codes	Within 90 d following the index admission	IMPROVE RAM-related factors: same as those in Spyropoulos et al ⁷
Zakai et al ¹⁷ (US)	Patients admitted to medical services (N = 900; VTE cases: 299; and non-VTE controls: 601)	2002-2009	cases: 63 ¹⁷ ; controls: 66 ¹⁵	Prognostic model development (single-center retrospective case-control)	Pharmacological prophylaxis in VTE cases: 64.6% and in non-VTE controls: 62.2%; mechanical prophylaxis in VTE cases: 31.8% and in non-VTE controls: 27.6%; the model was adjusted for both mechanical and pharmacological prophylaxis.	Any symptomatic VTE (upper and lower extremity DVT and PE; n = 299)	VTE events identified using ICD-9 VTE discharge codes. Codes confirmed by clinician review. Records were reviewed by a research nurse and all hospital-acquired VTE cases and 20% of noncases were reviewed by a physician.	Discharge or transfer from medical service	Venous thrombosis prophylaxis (mechanical; pharmacologic); demographics (age, sex, BMI); medical history (myocardial infarction, COPD, diabetes, chronic kidney disease); conditions active on admission (fever, COPD, pneumonia, any infection).

BI, Barthel Index; BMI, body mass index; COPD, chronic obstructive pulmonary disease; CTA, computed tomographic angiography; GI, gastrointestinal; ICD-9-CM, International Classification of Diseases, Ninth Revision, Clinical Modification; IVC, inferior vena cava; LWMH, low-molecular-weight heparin; mRS, modified Rankin Score; NYHA, New York Heart Association; RBC, red blood cell.

*Caprini factors: stroke; acute spinal cord injury or paralysis (<1 mo); hip, pelvis, or leg fracture (<1 mo); multiple trauma (<1 mo); history of thrombophilia; heparin-induced thrombocytopenia; age (41-60); age (61-74); positive history of cancer; immobilizing plaster cast; congestive heart failure; COPD or pulmonary function; inflammatory bowel disease; severe long disease (including pneumonia); acute myocardial infarction; sepsis (<1 mo); postpartum (<1 mo); history of unexpected stillborn infant; recurrent spontaneous abortion (≥ 3) or premature birth; varicose veins; BMI >25 kg/m²; swollen legs (current); CVC present on admission; immobile or not ambulating; and hormone replacement therapy or oral contraceptives.

†Padua factors: active cancer; previous VTE; reduced mobility; known thrombophilia; recent trauma and/or surgery; elderly age; heart and/or respiratory failure; acute myocardial infarction or ischemic stroke; acute infection and/or rheumatologic disorder; obesity; ongoing hormone treatment, and VTE prophylaxis.

Table 1. (continued)

First author, year (country)	Population (sample size)	Time frame (y)	Mean age, y (SD)*	Study type (number of centers and study design)	Prophylaxis (%)	Outcome (number of events)	Diagnostic methods	Follow-up time	Variables in multivariate logistic regression, when applicable
Zakai et al ²⁸ (US)	Medical patients (N = 188; VTE cases: 65 and non-VTE controls: 123)	2000-2002	68	External validation (single-center retrospective case-control)	VTE prophylaxis (including warfarin, unfractionated heparin, low molecular weight heparin or intermittent pneumatic compression devices): in VTE cases: 59%; and non-VTE controls 47%. The model was adjusted for prophylaxis.	Any symptomatic VTE (upper and lower extremity DVT and PE; n = 65)	VTE events identified using ICD-9 VTE discharge codes.	LOS case: 16 (10-28); controls: 6 (4-10)	Trauma last 3 mo; active cancer past year; admission fever; leg edema on admission; immobility >72 h; bacterial infection (cellulitis, pneumonia, sepsis, other); platelet count >350 × 10 ⁹ /L; use of VTE prophylaxis
Zhou et al ²⁹ (China)	Medically ill patients (N = 1804; VTE cases: 902 and non-VTE controls: 902)	2013-2016	Cases: 60 [†] ; controls: 57 ^{††}	External validation (single-center retrospective Case-control)	Any prophylaxis: in VTE cases: 4.1% and in non-VTE controls: 6.1% VTE prophylaxis included any mechanical use (intermittent pneumatic compression devices or sole vein pump) or pharmacological use (unfractionated heparin, LMWH, warfarin, fondaparinux sodium, etc.) The model was adjusted for VTE prophylaxis	Any symptomatic VTE (defined as DVT or PE; n = 902)	DVT verified by positive compression ultrasonography and/or contrast venography. PE verified by positive pulmonary angiogram, spiral computed tomography, or high probability ventilation/perfusion scanning or at autopsy.	6 mo after discharge	Caprini RAM factors* Padua RAM factors†
Barclay et al ³⁰ (US)	Chronic Liver disease (N = 1581)	2008-2011	51(11)	Prognostic factor (single-center retrospective cohort)	Pharmacological VTE prophylaxis: 24.8% (Unfractionated heparin: 9.7%; low molecular weight heparin: 88.0%; or both: 2.3%) The model was adjusted for pharmacological prophylaxis	Any symptomatic VTE (including DVT, PE or portal vein thrombosis-PVT) (n = 23)	VTE event identified in the medical record. VTE confirmed with radiologic testing.	4-7 d	VTE prophylaxis Active malignancy Trauma or surgery during hospitalization History of VTE

BI, Barthel Index; BMI, body mass index; COPD, chronic obstructive pulmonary disease; CTA, computed tomographic angiography; GI, gastrointestinal; ICD-9-CM, International Classification of Diseases, Ninth Revision, Clinical Modification; IVC, inferior vena cava; LMWH, low-molecular-weight heparin; mRS, modified Rankin Score; NYHA, New York Heart Association; RBC, red blood cell.

*Caprini factors: stroke; acute spinal cord injury or paralysis (<1 mo); hip, pelvis, or leg fracture (<1 mo); multiple trauma (<1 mo); age ≥75; history of VTE; family history of VTE; history of thrombophilia; heparin-induced thrombocytopenia; age (41-60); age (61-74); positive history of cancer; immobilizing plaster cast; congestive heart failure; COPD or pulmonary function; inflammatory bowel disease; severe long disease (including pneumonia); acute myocardial infarction; sepsis (<1 mo); surgery (<1 mo); postpartum (<1 mo); history of unexpected stillborn infant; recurrent spontaneous abortion (≥ 3) or premature birth; varicose veins; BMI >25 kg/m²; swollen legs (current); CVC present on admission; immobile or not ambulating; and hormone replacement therapy or oral contraceptives.

†Padua factors: active cancer; previous VTE; reduced mobility; known thrombophilia; recent trauma and/or surgery; elderly age; heart and/or respiratory failure; acute myocardial infarction or ischemic stroke; acute infection and/or rheumatologic disorder; obesity; ongoing hormone treatment, and VTE prophylaxis.

Table 1. (continued)

First author, year (country)	Population (sample size)	Time frame (y)	Mean age, y (SD)*	Study type (number of centers and study design)	Prophylaxis (%)	Outcome (number of events)	Diagnostic methods	Follow-up time	Variables in multivariate logistic regression, when applicable
Grant et al ³¹ (US)	Hospitalized medical patients (N = 63 548)	2011-2014	66	External validation (multicenter retrospective cohort)	Pharmacological VTE prophylaxis: 60.9% The model was adjusted for the pharmacological prophylaxis	Any symptomatic VTE (defined as proximal upper- or proximal lower-extremity DVT and PE) (n = 670)	VTE was clinically suspected. VTE was Image confirmed. Majority of events were identified by medical record review, 44 (6.6%) of the events were confirmed via telephone follow-up.	90 d	Caprini RAM factors*
Rothberg et al ³⁴ (US)	Medical Patients (N = 46 503)	2004-2005	NR	Prognostic model development (multicenter retrospective cohort)	Pharmacological VTE prophylaxis: 30% There was no difference in the model estimates for the factors when the model was adjusted for prophylaxis.	Any symptomatic VTE (n = 1 052)	VTE verified by lower extremity ultrasonography, venography, CT angiogram, ventilation/perfusion scan, or pulmonary angiogram) on hospital day 3 or later Secondary diagnosis of VTE provided using ICD-9 diagnoses.	30 30 d	Any prophylaxis; female; length of stay ≥ 6 d; age (18-49; 50-64, and >65 y); primary diagnosis (pneumonia, COPD, stroke, congestive heart failure, urinary tract infection, respiratory failure, septicemia) Comorbidities (inflammatory bowel disease, obesity, inherited thrombophilia); cancer (18-49, 50-64, and >65 y). Treatments (CVC, mechanical ventilation, urinary catheter, chemotherapy, steroids)

BI, Barthel Index; BMI, body mass index; COPD, chronic obstructive pulmonary disease; CTA, computed tomographic angiography; GI, gastrointestinal; ICD-9-CM, International Classification of Diseases, Ninth Revision, Clinical Modification; IV, inferior vena cava; LWMH, low-molecular-weight heparin; mRS, modified Rankin Score; NYHA, New York Heart Association; RBC, red blood cell.

*Caprini factors: stroke; acute spinal cord injury or paralysis (< 1 mo); hip, pelvis, or leg fracture (< 1 mo); multiple trauma (< 1 mo); age ≥ 75 ; history of VTE; family history of VTE; history of thrombophilia; heparin-induced thrombocytopenia; age (41-60); age (61-74); positive history of cancer; immobilizing plaster cast; congestive heart failure; COPD or pulmonary function; inflammatory bowel disease; severe long disease (including pneumonia); acute myocardial infarction; sepsis (< 1 mo); surgery (< 1 mo); postpartum (< 1 mo); history of unexpected stillborn infant, recurrent spontaneous abortion (≥ 3) or premature birth; varicose veins; BMI > 25 kg/m²; swollen legs (current); CVC present on admission; immobile or not ambulating; and hormone replacement therapy or oral contraceptives.

†Padua factors: active cancer; previous VTE; reduced mobility; known thrombophilia; recent trauma and/or surgery; elderly age; heart and/or respiratory failure; acute myocardial infarction or ischemic stroke; acute infection and/or rheumatologic disorder; obesity; ongoing hormone treatment, and VTE prophylaxis.

Table 1. (continued)

First author, year (country)	Population (sample size)	Time frame (y)	Mean age, y (SD)*	Study type (number of centers and study design)	Prophylaxis (%)	Outcome (number of events)	Diagnostic methods	Follow-up time	Variables in multivariate logistic regression, when applicable
Bembenek et al ³⁵ (Poland)	Early stroke patients (N = 299)	2007-2009	75 (64-82)*	Prognostic factor (single-center, prospective cohort)	Oral anticoagulation: 7.1%. The model was not adjusted for oral anticoagulation, but less than 10% of the included patients received prophylaxis.	Any symptomatic or asymptomatic DVT (n = 9; 7 of which were distal)	The first ultrasonography was performed within the first 7 d and then 8-10 d after stroke onset by a trained physician blinded to patients' baseline health status, to identify patients in whom DVT occurred early in the course of stroke.	Days 3 and 9 after stroke	Age for each additional 10 y; female; hypertension; congestive heart failure; atrial fibrillation; diabetes; smoking status (current and previous); prestroke disability (mRS 0-1 patient and mRS 0-2 patients); stroke severity (each additional 4 patients; NIHSS, NIHSS > 7 patients, NIHSS > 14 patients); decreased consciousness (≥ 1 patient in NIHSS, ≥ 2 patients in NIHSS); inflammatory markers (CRP > 10 mg/L, fibrinogen > 4 mg/dL)
Fan et al ³⁶ (China)	Acutely ill medical patients (N = 458)	2006-2007	77 (7)	Prognostic factor (multicenter prospective cohort)	Pharmacological VTE prophylaxis: 0%. Mechanical VTE prophylaxis (graduated compression stockings): 0.4%. The model was not adjusted for the mechanical thromboprophylaxis.	Any symptomatic or asymptomatic VTE (DVT or PE; n = 45: 30 symptomatic and 15 asymptomatic)	VTE verified by compression ultrasonography at enrollment and 3-wk follow-up. Symptomatic cases were all screened by lower limb color duplex ultrasonography.	90 d follow-up for symptomatic and 3 wk for asymptomatic	Univariate model with results provided; a multivariate analysis was conducted but results of each factor were not reported.

BI, Barthel Index; BMI, body mass index; COPD, chronic obstructive pulmonary disease; CTA, computed tomographic angiography; GI, gastrointestinal; ICD-9-CM, International Classification of Diseases, Ninth Revision, Clinical Modification; IVC, inferior vena cava; LWMH, low-molecular-weight heparin; mRS, modified Rankin Score; NYHA, New York Heart Association; RBC, red blood cell.

*Caprini factors: stroke; acute spinal cord injury or paralysis (<1 mo); hip, pelvis, or leg fracture (<1 mo); multiple trauma (<1 mo); age ≥ 75 ; history of VTE; family history of VTE; history of thrombophilia; heparin-induced thrombocytopenia; age (41-60); age (61-74); positive history of cancer; immobilizing plaster cast; congestive heart failure; COPD or pulmonary function; inflammatory bowel disease; severe long disease (including pneumonia); acute myocardial infarction; sepsis (<1 mo); surgery (<1 mo); postpartum (<1 mo); history of unexpected stillborn infant, recurrent spontaneous abortion (≥ 3) or premature birth; varicose veins; BMI > 25 kg/m²; swollen legs (current); CVC present on admission; immobile or not ambulating; and hormone replacement therapy or oral contraceptives.

†Padua factors: active cancer; previous VTE; reduced mobility; known thrombophilia; recent trauma and/or surgery; elderly age; heart and/or respiratory failure; acute myocardial infarction or ischemic stroke; acute infection and/or rheumatologic disorder; obesity; ongoing hormone treatment, and VTE prophylaxis.

Table 1. (continued)

First author, year (country)	Population (sample size)	Time frame (y)	Mean age, y (SD)*	Study type (number of centers and study design)	Prophylaxis (%)	Outcome (number of events)	Diagnostic methods	Follow-up time	Variables in multivariate logistic regression, when applicable
Kelly et al. ³⁸ (United Kingdom)	Acute ischemic stroke (N = 102)	Not reported	70 (12)	Prognostic factor (single-center prospective cohort)	VTE prophylaxis: 0%	Any asymptomatic or asymptomatic VTE (defined as proximal DVT or PE; n = 41).	<p>Patients were assessed weekly for clinical evidence of VTE. New increases in calf circumference from initial assessment of ≥ 3 cm (based on the Wells scoring system), local pain or tenderness for DVT, and oxygen saturations $\leq 92\%$ and/or respiratory rate > 20/min in a patient otherwise asymptomatic for PE. VTE was classified as "unrecognized clinical" if associated with the aforementioned signs or symptoms that went unrecognized by the attending team.</p> <p>Magnetic resonance direct thrombus imaging was performed. If DVT was identified, thoracic imaging was performed to detect PE. All scans were reviewed independently by 2 reviewers who reached a consensus.</p> <p>Clinical events diagnosed conventionally and data from postmortem examinations were included.</p>	21 d	Age > 70 ; BI ≤ 9 ; total anterior circulation infarcts, malignancy, and atrial fibrillation

BI, Barthel index; BMI, body mass index; COPD, chronic obstructive pulmonary disease; CTA, computed tomographic angiography; GI, gastrointestinal; ICD-9-CM, International Classification of Diseases, Ninth Revision, Clinical Modification; IVc, inferior vena cava; LWMH, low-molecular-weight heparin; mRS, modified Rankin Score; NYHA, New York Heart Association; RBC, red blood cell.

*Caprini factors: stroke; acute spinal cord injury or paralysis (< 1 mo); hip, pelvis, or leg fracture (< 1 mo); multiple trauma (< 1 mo); age ≥ 75 ; history of VTE; family history of VTE; history of thrombophilia; heparin-induced thrombocytopenia; age (41–60); age (61–74); positive history of cancer; immobilizing plaster cast; congestive heart failure; COPD or pulmonary function; inflammatory bowel disease; severe long disease (including pneumonia); acute myocardial infarction; sepsis (< 1 mo); surgery (< 1 mo); postpartum (< 1 mo); history of unexpected stillborn infant; recurrent spontaneous abortion (≥ 3) or premature birth; varicose veins; BMI > 25 kg/m²; swollen legs (current); CVC present on admission; immobile or not ambulating; and hormone replacement therapy or oral contraceptives.

†Padua factors: active cancer; previous VTE; reduced mobility; known thrombophilia; recent trauma and/or surgery; elderly age; heart and/or respiratory failure; acute myocardial infarction or ischemic stroke; acute infection and/or rheumatologic disorder; obesity; ongoing hormone treatment, and VTE prophylaxis.

Table 1. (continued)

First author, year (country)	Population (sample size)	Time frame (y)	Mean age, y (SD)*	Study type (number of centers and study design)	Prophylaxis (%)	Outcome (number of events)	Diagnostic methods	Follow-up time	Variables in multivariate logistic regression, when applicable
Ota et al ³⁸ (Japan)	Congestive heart failure (N = 161)	2003-2008	69.3 (10.8)	Prognostic factor (single-center, prospective cohort)	Anticoagulant therapy in DVT cases: 38.9%; non-DVT cases: 44.1% Antiplatelet therapy in DVT cases: 66.7% and in the non-DVT: 62.9% The model was adjusted for anticoagulant therapy.	Any symptomatic or asymptomatic DVT (no PE was detected; n = 18).	DVT verified by standardized sonography criterion of venous noncompressibility. PE verified by pulmonary angiography.	11.8 ± 11.5 d	NYHA functional class; poor IVC collapsibility; no anticoagulation therapy
Yi et al ³⁹ (China)	Acute stroke patients (N = 1380)	2009-2010	69.8 (11.6)	Prognostic factor (multicenter, prospective cohort)	Pharmacological VTE prophylaxis with warfarin or LMWH: 15% The model was adjusted for prophylaxis.	Any symptomatic or asymptomatic VTE (any PE and any DVT) (n = 62; 32 symptomatic DVT and 30 asymptomatic DVT)	DVT verified by VDU, venous angiography or venous CTA examination. PE verified by chest CTA or pulmonary angiography.	12 mo	For PE as an outcome: Age ≥ 70 y, bedridden, incidence of DVT For DVT as an outcome: Age ≥ 70 y; bedridden, Wells score ≥ 2, NIHSS score of lower limbs ≥ 3, BI score, rehabilitative therapy, anticoagulant therapy, concentration of D-dimer evaluated at admission.

BI, Barthel Index; BMI, body mass index; COPD, chronic obstructive pulmonary disease; CTA, computed tomographic angiography; GI, gastrointestinal; ICD-9-CM, International Classification of Diseases, Ninth Revision, Clinical Modification; IVC, inferior vena cava; LMWH, low-molecular-weight heparin; mRS, modified Rankin Score; NYHA, New York Heart Association; RBC, red blood cell.

*Caprini factors: stroke; acute spinal cord injury or paralysis (<1 mo); hip, pelvis, or leg fracture (<1 mo); multiple trauma (<1 mo); age ≥ 75; history of VTE; family history of VTE; history of thrombophilia; heparin-induced thrombocytopenia; age (41-60); age (61-74); positive history of cancer; immobilizing plaster cast; congestive heart failure; COPD or pulmonary function; inflammatory bowel disease; severe long disease (including pneumonia); acute myocardial infarction; sepsis (<1 mo); surgery (<1 mo); postpartum (<1 mo); history of unexpected stillborn infant, recurrent spontaneous abortion (≥ 3) or premature birth; varicose veins; BMI > 25 kg/m²; swollen legs (current); CVC present on admission; immobile or not ambulating; and hormone replacement therapy or oral contraceptives.

†Padua factors: active cancer; previous VTE; reduced mobility; known thrombophilia; recent trauma and/or surgery; elderly age; heart and/or respiratory failure; acute myocardial infarction or ischemic stroke; acute infection and/or rheumatologic disorder; obesity; ongoing hormone treatment, and VTE prophylaxis.

Table 1. (continued)

First author, year (country)	Population (sample size)	Time frame (y)	Mean age, y (SD)*	Study type (number of centers and study design)	Prophylaxis (%)	Outcome (number of events)	Diagnostic methods	Follow-up time	Variables in multivariate logistic regression, when applicable
Bleeding									
Decousus et al ^b (Canada)	Acutely ill medical patients (N = 15 156)	2002-2006	68.2 (51.8-78.9)*	Prognostic model development (multicenter, prospective cohort)	Pharmacological VTE prophylaxis: 48% (LWMH, 38.4%; unfractionated heparin, 11.1%, aspirin, 0.7%) Mechanical VTE prophylaxis: 9% (elastic stockings, 5.4%; intermittent pneumatic compression, 3.8%) There were no differences in the estimates of associations when the model was adjusted for pharmacologic prophylaxis use.	Major or clinically relevant bleeding (n = 230; 83 major and 147 nonmajor, but clinically relevant, bleeding)	Major bleeding was defined as a bleeding event contributing to death, clinically overt bleeding associated with a decrease in hemoglobin level of ≥ 2 g/dL or leading to transfusion of at least 2 units of packed RBCs, or bleeding within a critical organ (including intracranial, retroperitoneal, intraocular, adrenal gland, spinal, or pericardial bleeding). Nonmajor but clinically relevant bleeding was defined as overt gastrointestinal bleeding (except for insignificant hemorroidal bleeding), gross hematuria (macroscopic and lasting longer than 24 h), substantial epistaxis that required intervention and was recurrent and/or lasted at least 5 min, extensive hematoma or bruising (>5 cm in diameter), intra-articular bleeding (documented by aspiration), menorrhagia or metrorrhagia (increased quantity or duration), or other bleeding important enough to be recorded on the hospital chart.	14 d	Active gastroduodenal ulcer; bleeding in 3 mo before admission; platelet count $<50 \times 10^9/L$; age ≥ 85 vs <40 y; hepatic failure; severe renal failure GFR <30 vs ≥ 60 mL/min/m ² ; ICU/CCU; CVC; rheumatic disease; current cancer; age 40-84 vs <40 y; male sex; moderate renal failure (GFR 30-59 vs ≥ 60 mL/min/m ²).

Bl, Barthel Index; BMI, body mass index; COPD, chronic obstructive pulmonary disease; CTA, computed tomographic angiography; GI, gastrointestinal; ICD-9-CM, International Classification of Diseases, Ninth Revision, Clinical Modification; IVC, inferior vena cava; LWMH, low-molecular-weight heparin; mRS, modified Rankin Score; NYHA, New York Heart Association; RBC, red blood cell.
 *Caprini factors: stroke; acute spinal cord injury or paralysis (<1 mo); hip, pelvis, or leg fracture (<1 mo); multiple trauma (<1 mo); age ≥ 75 ; history of thrombophilia; heparin-induced thrombocytopenia; age (41-60); age (61-74); positive history of cancer; immobilizing plaster cast; congestive heart failure; COPD or pulmonary function; inflammatory bowel disease; severe long disease (including pneumonia); acute myocardial infarction; sepsis (<1 mo); surgery (<1 mo); postpartum (<1 mo); history of unexpected stillborn infant; recurrent spontaneous abortion (≥ 3) or premature birth; varicose veins; BMI >25 kg/m²; swollen legs (current); CVC present on admission; immobile or not ambulating; and hormone replacement therapy or oral contraceptives.
 †Padua factors: active cancer; previous VTE; reduced mobility; known thrombophilia; recent trauma and/or surgery; elderly age; heart and/or respiratory failure; acute myocardial infarction or ischemic stroke; acute infection and/or rheumatologic disorder; obesity; ongoing hormone treatment; and VTE prophylaxis.

Table 1. (continued)

First author, year (country)	Population (sample size)	Time frame (y)	Mean age, y (SD)*	Study type (number of centers and study design)	Prophylaxis (%)	Outcome (number of events)	Diagnostic methods	Follow-up time	Variables in multivariate logistic regression, when applicable
Mahan et al ³² (US)	Medical patients (N = 327 578)	2005-2009	69	Prognostic factor (multicenter retrospective cohort)	All antithrombotic agent use: 9.4% Anticoagulants: 3.9% (warfarin, 3.6%; enoxaparin, 0.4%; heparin, 0.1% and other, <0.0%) Antiplatelets: 5.7% (clopidogrel: 4.6%; aspirin-dipyridamole, 0.9%; other, 0.3%) Anticoagulants and antiplatelets: 0.2% The model was adjusted for antithrombotic use.	Major or clinically relevant bleeding (n = 29 264; 5 951 major and 23 313 minor bleeding)	Bleeding events were identified through the ICD-9-CM diagnosis codes.	Within 30 d after hospitalization	Age (40-54, 55-64, 65-74, ≥75); male; preindex risk factors (insufficient renal function, cancer, rheumatoid arthritis, gastrooduodenal ulcer, blood dyscrasias, thrombocytopenia, liver disease, CVC, thromboembolic stroke, estrogen use); postindex risk factors (postdischarge antithrombotic meds use, rehospitalization, length of stay (2, 3-5, and ≥6 d))

BI, Barthel Index; BMI, body mass index; COPD, chronic obstructive pulmonary disease; CTA, computed tomographic angiography; GI, gastrointestinal; ICD-9-CM, International Classification of Diseases, Ninth Revision, Clinical Modification; IVC, inferior vena cava; LWMH, low-molecular-weight heparin; mRS, modified Rankin Score; NYHA, New York Heart Association; RBC, red blood cell.

*Caprini factors: stroke; acute spinal cord injury or paralysis (<1 mo); hip, pelvis, or leg fracture (<1 mo); multiple trauma (<1 mo); history of VTE; family history of VTE; history of thrombophilia; heparin-induced thrombocytopenia; age (41-60); age (61-74); positive history of cancer; immobilizing plaster cast; congestive heart failure; COPD or pulmonary function; inflammatory bowel disease; severe long disease (including pneumonia); acute myocardial infarction; sepsis (<1 mo); surgery (<1 mo); postpartum (<1 mo); history of unexpected stillborn infant, recurrent spontaneous abortion (≥ 3) or premature birth; varicose veins; BMI >25 kg/m²; swollen legs (current); CVC present on admission; immobile or not ambulating; and hormone replacement therapy or oral contraceptives.

†Padua factors: active cancer; previous VTE; reduced mobility; known thrombophilia; recent trauma and/or surgery; elderly age; heart and/or respiratory failure; acute myocardial infarction or ischemic stroke; acute infection and/or hematologic disorder; obesity; ongoing hormone treatment, and VTE prophylaxis.

Table 1. (continued)

First author, year (country)	Population (sample size)	Time frame (y)	Mean age, y (SD)*	Study type (number of centers and study design)	Prophylaxis (%)	Outcome (number of events)	Diagnostic methods	Follow-up time	Variables in multivariate logistic regression, when applicable
Patell et al ³³ (US)	Cancer patients (N = 3358)	2012-2014	62(19.98)*	Prognostic factor (single- center retrospective cohort)	Antiplatelets: 14% Anticoagulants: 67% Antiplatelet agents on day of admission were not found to be statistically significant in univariate analysis, so were not added to the multivariate regression analysis mode. Anticoagulation exposure on admission was noted to be associated with a decreased risk of bleeding (OR, 0.5; 95% CI, 0.3-0.8; P = 0.004) although this was not significant in multivariable analysis (but the model adjusted for it).	Major or clinically relevant bleeding (n = 69; 51 major and 18 nonmajor but clinically relevant bleeding)	Bleeding was assessed using the International Society on Thrombosis Hemostasis definitions of major bleeding and clinically relevant nonmajor bleeding Bleeding events were identified from discharge summaries of admissions being studied. To obtain details of event, documentation including diagnostic tests (imaging and procedures) and clinical notes was used. All bleeding events were confirmed manually by 2 investigators (third-year internal medicine residents at the time of the study). When unclear, individual cases were cross-reviewed, discussed and included if both agreed. No separate training was performed, and no coding was used to extract bleeding information.	Median length of stay was 5 d (range, 0-152 days).	Reason for admission (anemia), BMI \geq 40; cancer site: GI; low hemoglobin ($<$ 13 g/dL for men and $<$ 11.5 g/dL for women); low platelets ($<$ 150 000/ μ L)

BI, Barthel Index; BMI, body mass index; COPD, chronic obstructive pulmonary disease; CTA, computed tomographic angiography; GI, gastrointestinal; ICD-9-CM, International Classification of Diseases, Ninth Revision, Clinical Modification; IVC, inferior vena cava; LWMH, low-molecular-weight heparin; mRS, modified Rankin Score; NYHA, New York Heart Association; RBC, red blood cell.

*Caprini factors: stroke; acute spinal cord injury or paralysis ($<$ 1 mo); hip, pelvis, or leg fracture ($<$ 1 mo); multiple trauma ($<$ 1 mo); history of VTE; family history of VTE; history of thrombophilia; heparin-induced thrombocytopenia; age (41-60); age (61-74); positive history of cancer; immobilizing plaster cast; congestive heart failure; COPD or pulmonary function; inflammatory bowel disease; severe long disease (including pneumonia); acute myocardial infarction; sepsis ($<$ 1 mo); surgery ($<$ 1 mo); postpartum ($<$ 1 mo); history of unexpected stillborn infant, recurrent spontaneous abortion (\geq 3) or premature birth; varicose veins; BMI $>$ 25 kg/m²; swollen legs (current); CVC present on admission; immobile or not ambulating; and hormone replacement therapy or oral contraceptives.

†Padua factors: active cancer; previous VTE; reduced mobility; known thrombophilia; recent trauma and/or surgery; elderly age; heart and/or respiratory failure; acute myocardial infarction or ischemic stroke; acute infection and/or rheumatologic disorder; obesity; ongoing hormone treatment, and VTE prophylaxis.

Table 2. Evidence profile for VTE-related prognostic factors

No. of studies	Certainty assessment domains						Overall certainty in the evidence about this prognostic factor	Relative effect (95% CI)
	Study design	Risk of bias	Indirect	Inconsistent	Imprecise	Publication bias		
Age (>60 y vs <60 y)^{7,25-27,29,31,34,37,39}								
9*	Observational	Serious†	Not serious	Not serious	Not serious	Undetected	⊕⊕⊕○ MODERATE	OR, 1.34 (95% CI, 1.17-1.55)
Sex (male vs female)^{27,34-37}								
5	Observational	Serious	Not serious	Not serious	Not serious	Undetected	⊕⊕⊕○ MODERATE	OR, 1.03 (95% CI, 0.80-1.33)
CRP (>10 mg/L vs <10 mg/L)³⁵								
1	Observational	Serious	Not serious	Not serious	Not serious	Undetected	⊕⊕⊕○ MODERATE	OR, 10.10 (95% CI, 1.93-52.85)
D-dimer (>500 ng/mL vs <500 ng/mL at baseline; and increase vs no increase)^{36,39}								
2	Observational	Serious†	Not serious	Not serious	Not serious	Undetected	⊕⊕⊕○ MODERATE	Categorical: OR, 2.46 (95% CI, 1.19-5.10) Continuous: OR, 3.45 (95% CI, 2.01-5.92)
Heart rate (elevated >100 beats per minute vs nonelevated <100 beats per minute)²⁷								
1	Observational	Serious	Not serious	Not serious	Not serious	Undetected	⊕⊕⊕○ MODERATE	OR, 2.48 (95% CI, 1.66-3.71)
Thrombocytosis (platelet count >350 × 10⁹/L vs <350 × 10⁹/L)^{27,28}								
2	Observational	Serious	Not serious	Not serious	Not serious	Undetected	⊕⊕⊕○ MODERATE	OR, 2.16 (95% CI, 1.40-3.35)
Leukocytosis (WBC ≥11 × 10⁹/L vs <11 × 10⁹/L)²⁷								
1	Observational	Serious	Not serious	Not serious	Not serious	Undetected	⊕⊕⊕○ MODERATE	OR, 1.91 (95% CI, 1.24-2.94)
Fever (body temperature >38°C-38.5°C vs <38°C-38.5°C)^{27,28}								
2	Observational	Serious	Not serious	Not serious	Not serious	Undetected	⊕⊕⊕○ MODERATE	OR, 1.88 (95% CI, 1.10-3.21)
Leg edema (presence vs absence)^{28,31}								
2	Observational	Serious	Not serious	Not serious	Not serious	Undetected	⊕⊕⊕○ MODERATE	OR, 1.88 (95% CI, 1.23-2.90)

Question: prognostic factors for medical patients; outcome: VTE; setting: inpatient.

BI, Barthel Index; WBC, white blood cell count.

*Certainty in evidence was downgraded for risk of bias, given a follow-up time of more than 3 months in the included studies that may cause an overestimation of the magnitude of the association (Zhou et al²⁹: 6 mo after discharge and Yi 2012: 12 mo after discharge).

†Certainty in evidence was downgraded for imprecision, given that the confidence interval suggests that there may be no association.

‡Certainty in evidence was downgraded for risk of bias, given that the results of each prognostic factor in the multivariate analysis were not reported and we therefore had to rely on the unadjusted measures of association. Also, the multivariate analysis included only factors statistically significant in the univariate analysis.

§Certainty in evidence was downgraded for inconsistency but not imprecision, given that the inconsistency is the likely cause for the imprecision.

||Certainty in evidence was downgraded for imprecision, given the small number of events (n = 32).

¶Bembenek et al³⁵ assessed the severity of a stroke experienced by an individual by using the NIHSS, a diagnostic tool; results were consistent with the meta-analysis that severe stroke may result in an increase in risk of any DVT (OR, 2.11; 95% CI, 0.50-8.90) for NIHSS >7 compared with a NIHSS score <7. Also, severe stroke may result in an increase in risk of any DVT (OR, 1.34; 95% CI, 0.25-7.18) for NIHSS >14 compared with a NIHSS <14. When NIHSS was assessed continuously, results showed that severe stroke may result in an increase in risk of any DVT (OR, 1.21; 95% CI, 0.86-1.70) for each additional 4 points on the NIHSS.

#Fan et al,³⁶ with 458 patients older than 60 y of which 45 patients had any VTE, presented age as a continuous variable and showed no association between age and any VTE (OR, 1.03; 95% CI, 0.98-1.08). Bembenek et al,³⁵ with 299 patients of which 9 had any DVT, 7 of which were distal, presented age per 10-y increase and showed a decrease in risk per 10-y increase in age with any DVT (OR, 0.64; 95% CI, 0.33-1.24).

Table 2. (continued)

No. of studies	Certainty assessment domains						Overall certainty in the evidence about this prognostic factor	Relative effect (95% CI)
	Study design	Risk of bias	Indirect	Inconsistent	Imprecise	Publication bias		
Varicose veins (presence vs absence)^{29,31}								
2	Observational	Serious†	Not serious	Not serious	Serious‡	Undetected	⊕⊕○○ LOW	OR, 1.53 (95% CI, 0.85-2.76)
Obesity (BMI >30 kg/m² vs <30 kg/m²)^{27,29,34}								
3	Observational	Serious†	Not serious	Not serious	Serious‡	Undetected	⊕⊕○○ LOW	OR, 1.34 (95% CI, 0.94-1.91)
Fibrinogen levels (elevated levels >400 mg/dL) vs no elevated levels³⁵								
1	Observational	Serious	Not serious	Not serious	Not serious	Undetected	⊕⊕⊕○ MODERATE	OR, 0.18 (95% CI, 0.04-0.81)
BI score ≤9 vs >9^{37,39}								
2	Observational	Serious†	Not serious	Not serious	Not serious	Undetected	⊕⊕⊕○ MODERATE	OR, 8.30 (95% CI, 2.70-25.52)
Immobility: defined as confinement to bed for >72 h or >7 d or bedridden or nonambulatory (yes vs no)^{7,26-29,31,37,39}								
8	Observational	Serious†	Not serious	Not serious	Not serious	Undetected	⊕⊕⊕○ MODERATE	OR, 3.17 (95% CI, 2.18-4.62)
Paresis (yes vs no)^{7,25,26,37}								
4	Observational	Serious	Not serious	Not serious	Not serious	Undetected	⊕⊕⊕○ MODERATE	OR, 2.97 (95% CI, 1.20-7.36)
Previous VTE (yes vs no)^{7,25-27,29-31,36}								
8	Observational	Serious†	Not serious	Not serious	Not serious	Undetected	⊕⊕⊕○ MODERATE	OR, 6.08 (95% CI, 3.71-9.97).
Thrombophilia (familial or acquired disorder of the hemostatic system; yes vs no)^{7,25,26,29,34}								
5	Observational	Serious†	Not serious	Not serious	Not serious	Undetected	⊕⊕⊕○ MODERATE	OR, 5.88 (95% CI, 2.80-12.35)
Malignancy (active malignancy, defined as the presence of cancer on admission or within the past year, vs no active malignancy; and history vs no history of malignancy)^{7,25-28,30,31,34,36,37}								
10	Observational	Serious	Not serious	Not serious	Not serious	Undetected	⊕⊕⊕○ MODERATE	Active cancer: OR, 2.65 (95% CI, 1.79-3.91) History of cancer: OR, 3.20 (95% CI, 2.14-4.79)

Question: prognostic factors for medical patients; outcome: VTE; setting: inpatient.

BI, Barthel Index; WBC, white blood cell count.

*Certainty in evidence was downgraded for risk of bias, given a follow-up time of more than 3 months in the included studies that may cause an overestimation of the magnitude of the association (Zhou et al²⁹: 6 mo after discharge and Yi 2012: 12 mo after discharge).

†Certainty in evidence was downgraded for imprecision, given that the confidence interval suggests that there may be no association.

‡Certainty in evidence was downgraded for risk of bias, given that the results of each prognostic factor in the multivariate analysis were not reported and we therefore had to rely on the unadjusted measures of association. Also, the multivariate analysis included only factors statistically significant in the univariate analysis.

§Certainty in evidence was downgraded for inconsistency but not imprecision, given that the inconsistency is the likely cause for the imprecision.

||Certainty in evidence was downgraded for imprecision, given the small number of events (n = 32).

¶Bembenek et al³⁵ assessed the severity of a stroke experienced by an individual by using the NIHSS, a diagnostic tool; results were consistent with the meta-analysis that severe stroke may result in an increase in risk of any DVT (OR, 2.11; 95% CI, 0.50-8.90) for NIHSS >7 compared with a NIHSS score <7. Also, severe stroke may result in an increase in risk of any DVT (OR, 1.34; 95% CI, 0.25-7.18) for NIHSS >14 compared with a NIHSS <14. When NIHSS was assessed continuously, results showed that severe stroke may result in an increase in risk of any DVT (OR, 1.21; 95% CI, 0.86-1.70) for each additional 4 points on the NIHSS.

#Fan et al,³⁴ with 458 patients older than 60 y of which 45 patients had any VTE, presented age as a continuous variable and showed no association between age and any VTE (OR, 1.03; 95% CI, 0.98-1.08). Bembenek et al,³⁵ with 299 patients of which 9 had any DVT, 7 of which were distal, presented age per 10-y increase and showed a decrease in risk per 10-y increase in age with any DVT (OR, 0.64; 95% CI, 0.33-1.24).

Table 2. (continued)

No. of studies	Certainty assessment domains						Overall certainty in the evidence about this prognostic factor	Relative effect (95% CI)
	Study design	Risk of bias	Indirect	Inconsistent	Imprecise	Publication bias		
Critical illness: defined as ICU or CCU stay, or need for resuscitation (yes vs no)^{7,25-27,29,34,36}								
7	Observational	Serious†	Not serious	Not serious	Not serious	Undetected	⊕⊕⊕○ MODERATE	OR, 1.65 (95% CI, 1.39-1.95)
Infections: including cellulitis, pneumonia, and sepsis (yes vs no)^{28,31}								
5	Observational	Serious†	Not serious	Not serious	Not serious	Undetected	⊕⊕⊕○ MODERATE	Any infection: OR, 1.48 (95% CI, 1.16-1.89) Acute infection: OR, 1.59 (95% CI, 1.23-2.06) Sepsis: OR, 1.07 (95% CI, 0.70-1.62)
Heart failure (HF) (acute HF vs no acute HF; history of HF vs no history of HF)^{28,29,31,35,36}								
5	Observational	Serious	Not serious	Not serious	Serious‡	Undetected	⊕⊕○○ LOW	Acute heart failure: OR, 0.82 (95% CI, 0.42-1.60) History of heart failure: OR, 2.68 (95% CI, 1.11-6.44)
Autoimmune disease: including rheumatological diseases and inflammatory diseases (yes vs no)^{27,29,31,35}								
4	Observational	Serious†	Not serious	Serious	Not serious	Undetected	⊕⊕○○ LOW	OR, 2.33 (95% CI, 1.13-4.83)
CVC (presence vs absence)^{31,34}								
2	Observational	Serious	Not serious	Serious§	Not serious	Undetected	⊕⊕○○ LOW	OR, 2.05 (95% CI, 0.74-5.65)
Severe stroke: defined as acute ischemic stroke (yes vs no)^{29,31,32,36}								
4	Observational	Serious†	Not serious	Not serious	Serious‡	Undetected	⊕⊕○○ LOW	Acute ischemic stroke: OR, 1.79 (95% CI, 0.77-4.18) When stroke was assessed in terms of the NIHSS, we found consistent results.¶

Question: prognostic factors for medical patients; outcome: VTE; setting: inpatient.

BI, Barthel Index; WBC, white blood cell count.

*Certainty in evidence was downgraded for risk of bias, given a follow-up time of more than 3 months in the included studies that may cause an overestimation of the magnitude of the association (Zhou et al²⁹: 6 mo after discharge and Yi 2012: 12 mo after discharge).

†Certainty in evidence was downgraded for imprecision, given that the confidence interval suggests that there may be no association.

‡Certainty in evidence was downgraded for risk of bias, given that the results of each prognostic factor in the multivariate analysis were not reported and we therefore had to rely on the unadjusted measures of association. Also, the multivariate analysis included only factors statistically significant in the univariate analysis.

§Certainty in evidence was downgraded for inconsistency but not imprecision, given that the inconsistency is the likely cause for the imprecision.

¶Certainty in evidence was downgraded for imprecision, given the small number of events (n = 32).

¶Bembenek et al³⁵ assessed the severity of a stroke experienced by an individual by using the NIHSS, a diagnostic tool; results were consistent with the meta-analysis that severe stroke may result in an increase in risk of any DVT (OR, 2.11; 95% CI, 0.50-8.90) for NIHSS >7 compared with a NIHSS score <7. Also, severe stroke may result in an increase in risk of any DVT (OR, 1.34; 95% CI, 0.25-7.18) for NIHSS >14 compared with a NIHSS <14. When NIHSS was assessed continuously, results showed that severe stroke may result in an increase in risk of any DVT (OR, 1.21; 95% CI, 0.86-1.70) for each additional 4 points on the NIHSS.

#Fan et al³⁶ with 458 patients older than 60 y of which 45 patients had any VTE, presented age as a continuous variable and showed no association between age and any VTE (OR, 1.03; 95% CI, 0.98-1.08). Bembenek et al³⁵ with 299 patients of which 9 had any DVT, 7 of which were distal, presented age per 10-y increase and showed a decrease in risk per 10-y increase in age with any DVT (OR, 0.64; 95% CI, 0.33-1.24).

Table 2. (continued)

No. of studies	Certainty assessment domains						Overall certainty in the evidence about this prognostic factor	Relative effect (95% CI)
	Study design	Risk of bias	Indirect	Inconsistent	Imprecise	Publication bias		
Tobacco (current use vs no current use; previous use vs no previous use)³⁵								
1	Observational	Serious	Not serious	Not serious	Serious‡	Undetected	⊕⊕○○ LOW	Current tobacco use: OR, 1.59 (95% CI, 0.28-9.03) Previous tobacco use: OR, 0.97 (0.24-3.92)
Hormone use: defined as estrogen intake (yes vs no)³¹								
1	Observational	Serious	Not serious	Not serious	Serious‡	Undetected	⊕⊕○○ LOW	OR, 0.80 (95% CI, 0.36-1.78).
Renal failure (yes vs no)³⁶								
1	Observational	Serious	Not serious	Not serious	Very Serious‡,#	Undetected	⊕○○○ VERY LOW	OR, 0.76 (95% CI, 0.18-3.18)
Respiratory failure (yes vs no)^{27,29,31,36}								
4	Observational	Serious‡	Not serious	Not serious	Serious‡	Undetected	⊕⊕○○ LOW	Any respiratory failure: OR, 1.04 (95% 0.69-1.58). Acute respiratory failure: OR, 1.18 (95% CI, 0.76-1.84) Chronic respiratory failure: OR, 0.58 (95% 0.30-1.10)
Coronary artery disease (yes vs no)^{29,31,36,37}								
4	Observational	Serious‡	Not serious	Serious§	Not serious	Undetected	⊕⊕○○ LOW	OR, 1.01 (95% CI, 0.33-3.09)

Question: prognostic factors for medical patients; outcome: VTE; setting: inpatient.

BI, Barthel Index; WBC, white blood cell count.

*Certainty in evidence was downgraded for risk of bias, given a follow-up time of more than 3 months in the included studies that may cause an overestimation of the magnitude of the association (Zhou et al²⁹: 6 mo after discharge and Yi 2012: 12 mo after discharge).

†Certainty in evidence was downgraded for imprecision, given that the confidence interval suggests that there may be no association.

‡Certainty in evidence was downgraded for risk of bias, given that the results of each prognostic factor in the multivariate analysis were not reported and we therefore had to rely on the unadjusted measures of association. Also, the multivariate analysis included only factors statistically significant in the univariate analysis.

§Certainty in evidence was downgraded for inconsistency but not imprecision, given that the inconsistency is the likely cause for the imprecision.

||Certainty in evidence was downgraded for imprecision, given the small number of events (n = 32).

||Bembenek et al³⁵ assessed the severity of a stroke experienced by an individual by using the NIHSS, a diagnostic tool; results were consistent with the meta-analysis that severe stroke may result in an increase in risk of any DVT (OR, 2.11; 95% CI, 0.50-8.90) for NIHSS >7 compared with a NIHSS score <7. Also, severe stroke may result in an increase in risk of any DVT (OR, 1.34; 95% CI, 0.25-7.18) for NIHSS >14 compared with a NIHSS <14. When NIHSS was assessed continuously, results showed that severe stroke may result in an increase in risk of any DVT (OR, 1.21; 95% CI, 0.86-1.70) for each additional 4 points on the NIHSS.

#Fan et al,³⁶ with 458 patients older than 60 y of which 45 patients had any VTE, presented age as a continuous variable and showed no association between age and any VTE (OR, 1.03; 95% CI, 0.98-1.08). Bembenek et al,³⁵ with 299 patients of which 9 had any DVT, 7 of which were distal, presented age per 10-y increase and showed a decrease in risk per 10-y increase in age with any DVT (OR, 0.64; 95% CI, 0.33-1.24).

and D-dimer >500 ng/mL at baseline (OR, 2.46; 95% CI, 1.19-5.10).³⁶ The findings for D-dimer concentration, when assessed as a continuous variable, were consistent with our results (Table 2).³⁹ Also, there is probably an association between risk of any VTE and elevated heart rate (>100 beats per minute; OR, 2.48; 95% CI, 1.66-3.71),²⁷ thrombocytosis (platelet count, >350 × 10⁹/L; OR, 2.16; 95% CI, 1.40-3.35),^{27,28} leukocytosis (white blood cell count, ≥11 × 10⁹/L; OR, 1.91; 95% CI: 1.24-2.94),²⁷ fever (body temperature >38°C-38.5°C; OR, 1.88; 95% CI, 1.10-3.21),^{27,28} leg edema (OR, 1.88; 95% CI, 1.23-2.90),^{28,31} and elevated fibrinogen level (>400 mg/dL; OR, 0.18; 95% CI, 0.04-0.81).³⁵

We identified low-certainty evidence that there may be an association between risk of any VTE and varicose veins (OR, 1.53; 95% CI, 0.85-2.76)^{29,31} and obesity (BMI >30 kg/m²; OR, 1.34; 95% CI, 0.94-1.91).^{27,29,31,34,36}

Additional analyses

We performed a sensitivity analysis, including studies that reported on immobility, to compare the association between immobility >72 hours and >7 days with risk of VTE. We found similar effect estimates for both categories with

Table 3. Evidence profile for bleeding-related prognostic factors

No. of studies	Certainty assessment domains						Overall certainty in the evidence about this prognostic factor	Relative effect (95% CI)
	Study design	Risk of bias	Indirect	Inconsistent	Imprecise	Publication bias		
Age (≥ 65 vs < 65)^{8,33}								
2	Observational	Serious*,†	Not serious	Not serious	Not serious	Undetected	⊕⊕⊕○ MODERATE	Age ≥ 65 : OR, 1.95 (95% CI, 1.59-2.38)
Sex (male vs female)^{8,33}								
2	Observational	Serious*,†	Not serious	Not serious	Not serious	Undetected	⊕⊕⊕○ MODERATE	OR, 1.27 (95% CI, 1.09-1.47).
Anemia as a reason for admission (presence vs absence)³⁴								
2	Observational	Serious*,‡	Not serious	Not serious	Not serious	Undetected	⊕⊕⊕○ MODERATE	OR, 5.15 (95% CI, 2.45-10.81)
Morbid obesity (BMI ≥ 40 kg/m² vs BMI < 40 kg/m²)³⁴								
1	Observational	Serious‡	Not serious	Not serious	Not serious	Undetected	⊕⊕⊕○ MODERATE	OR, 3.08 (95% CI, 1.35-7.02)
Low hemoglobin: defined as < 13 g/dL in men and < 11.5 g/dL in women (yes vs no)³⁴								
1	Observational	Serious‡	Not serious	Not serious	Not serious	Undetected	⊕⊕⊕○ MODERATE	OR, 2.33 (95% CI, 1.04-5.22)
Gastroduodenal ulcers (yes vs no)^{8,33}								
2	Observational	Serious*,†	Not serious	Not serious	Not serious	Undetected	⊕⊕⊕○ MODERATE	OR, 2.74 (95% CI, 1.42-5.26)
Rehospitalization (yes vs no)³³								
1	Observational	Serious†	Not serious	Not serious	Not serious	Undetected	⊕⊕⊕○ MODERATE	OR, 2.39 (95% CI, 2.25-2.54)
Critical illness (yes vs no)⁸								
1	Observational	Serious*	Not serious	Not serious	Not serious	Undetected	⊕⊕⊕○ MODERATE	OR, 2.10 (95% CI, 1.42-3.11).
Thrombocytopenia (yes vs no)^{8,33,34}								
3	Observational	Serious*,†,‡	Not serious	Not serious	Not serious	Undetected	⊕⊕⊕○ MODERATE	All: OR, 1.79 (95% CI, 0.97-3.29) <50 $\times 10^9/L$: OR, 3.37 (95% CI, 1.84-6.18) <150 $\times 10^9/L$: OR, 1.30 (95% CI, 0.92-1.82)
Blood dyscrasias defined as the presence of any bleeding disorders on admission (presence vs absence)³³								
1	Observational	Serious†	Not serious	Not serious	Not serious	Undetected	⊕⊕⊕○ MODERATE	OR, 1.70 (95% CI, 1.60-1.81)

Question: prognostic factors for medical patients; outcome: bleeding; setting: inpatient.

*Certainty in evidence was downgraded for risk of bias, given that patients were enrolled both prospectively and retrospectively in Decousus et al.⁸ The retrospective enrollment of patients may have introduced classification bias.

†Certainty in evidence was downgraded for risk of bias, given that the authors evaluated bleeding risk in medical patients after hospitalization, that may overestimate the magnitude of the association. This is possibly due to patients being discharged on thromboprophylaxis without proper risk stratification for bleeding placing unmonitored patients at a higher risk of having a bleeding event.

‡Certainty in evidence was downgraded for risk of bias, given that the population is specific to hospitalized cancer patients who are at a higher risk of VTE and may be given thromboprophylaxis, placing them at a higher risk of having a bleeding event. This in turn may overestimate the magnitude of the association.

§Certainty in evidence was downgraded for imprecision, given that the CI suggests that there may be no association.

Table 3. (continued)

No. of studies	Certainty assessment domains						Overall certainty in the evidence about this prognostic factor	Relative effect (95% CI)
	Study design	Risk of bias	Indirect	Inconsistent	Imprecise	Publication bias		
Hepatic disease (yes vs no)^{8,33}								
2	Observational	Serious*,†	Not serious	Not serious	Not serious	Undetected	⊕⊕⊕○ MODERATE	OR, 1.53 (95% CI, 1.09-2.15)
Renal failure (yes vs no)^{8,33}								
2	Observational	Serious*,†	Not serious	Not serious	Not serious	Undetected	⊕⊕⊕○ MODERATE	Total: OR, 1.43 (95% CI, 1.06-1.93) Any renal failure (RF): OR, 1.23 (95% CI, 0.92-1.65). Moderate RF (GFR 30-59 mL/min/m ²): OR, 1.37(95% CI, 0.84-2.23) Severe RF (GFR <30 mL/min/m ²): OR, 2.14 (95% CI, 1.22-3.75)
Antithrombotic medication (yes vs no)^{8,33}								
2	Observational	Serious*,†	Not serious	Not serious	Not serious	Undetected	⊕⊕⊕○ MODERATE	OR, 1.28 (95% CI, 1.01-1.64)
CVC (yes vs no)^{8,33}								
2	Observational	Serious*,†	Not serious	Not serious	Not serious	Undetected	⊕⊕⊕○ MODERATE	OR, 1.37 (95% CI, 0.83-2.26)
Autoimmune disease (yes vs no)⁸								
2	Observational	Serious*	Not serious	Not serious	Serious§	Undetected	⊕⊕○○ LOW	OR, 1.30 (95% CI, 0.77-2.19)
Hormone use: defined as estrogen intake (yes vs no)³³								
1	Observational	Serious†	Not serious	Not serious	Not serious	Undetected	⊕⊕⊕○ MODERATE	OR, 0.95 (95% CI, 0.82-1.10)
Malignancy (yes vs no)^{8,33}								
2	Observational	Serious *,†	Not serious	Not serious	Serious§	Undetected	⊕⊕○○ LOW	OR, 1.08 (95% CI, 0.42-2.77)

Question: prognostic factors for medical patients; outcome: bleeding; setting: inpatient.

*Certainty in evidence was downgraded for risk of bias, given that patients were enrolled both prospectively and retrospectively in Decousus et al.⁸ The retrospective enrollment of patients may have introduced classification bias.

†Certainty in evidence was downgraded for risk of bias, given that the authors evaluated bleeding risk in medical patients after hospitalization, that may overestimate the magnitude of the association. This is possibly due to patients being discharged on thromboprophylaxis without proper risk stratification for bleeding placing unmonitored patients at a higher risk of having a bleeding event.

‡Certainty in evidence was downgraded for risk of bias, given that the population is specific to hospitalized cancer patients who are at a higher risk of VTE and may be given thromboprophylaxis, placing them at a higher risk of having a bleeding event. This in turn may overestimate the magnitude of the association.

§Certainty in evidence was downgraded for imprecision, given that the CI suggests that there may be no association.

a slightly stronger association between immobility, defined as bed rest for >7 days, and risk of VTE (OR, 3.67; 95% CI, 0.85-15.93) compared with immobility, defined as bed rest for >72 hours, and risk of VTE (OR, 3.18; 95% CI, 1.10-9.16).

We also conducted a sensitivity analysis, including studies that reported on symptomatic VTE only, to evaluate the influence of the studies that reported on both symptomatic and asymptomatic VTE. The results of the sensitivity analysis showed similar effect estimates across prognostic factors except for CAD.

The association between CAD and risk of symptomatic VTE was somewhat stronger in the sensitivity analysis of the 9 studies (OR, 2.02; 95% CI, 0.32-12.64), compared with little to no association with risk of VTE in the primary analysis (OR, 1.01; 95% CI, 0.33-3.09; supplemental Table 4).

Prognostic factors for bleeding in medical hospitalized patients

Three studies including 160 142 patients investigated 17 candidate prognostic factors for bleeding. Table 3 provides the evidence profile for bleeding-related prognostic factors. Supplemental Figure 2 provides the forest plots of the meta-analyses of each of the prognostic factors.

Demographic factors We found moderate-certainty evidence that there is probably an association between risk of bleeding and age ≥ 65 (OR, 1.95; 95% CI, 1.59-2.38)^{8,32} and sex (males vs females; OR, 1.27; 95% CI, 1.09-1.47).^{8,32}

Medical illness and patient history factors There was moderate-certainty evidence of a probable association between risk of bleeding and gastroduodenal ulcers (OR, 2.74; 95% CI, 1.42-5.26),^{8,32} rehospitalization (OR, 2.39; 95% CI, 2.25- 2.54),³² critical illness including ICU or CCU stay (OR, 2.10; 95% CI, 1.42-3.11),⁸ and thrombocytopenia (OR, 1.79; 95% CI 0.97-3.29).^{8,32,33} When cutoffs for thrombocytopenia were assessed separately, results showed that there probably is a greater magnitude of association between risk of bleeding and platelet count $< 50 \times 10^9/L$ compared with a platelet count $\geq 50 \times 10^9/L$ (OR, 3.37; 95% CI, 1.84-6.18),²⁶ whereas a smaller magnitude of association between risk of bleeding and platelet count $< 150 \times 10^9/L$ compared with platelet count $\geq 150 \times 10^9/L$ (OR, 1.30; 95% CI, 0.92-1.82).^{30,32} We also found moderate-certainty evidence that there is probably an association between risk of bleeding and blood dyscrasia, defined as the presence of any bleeding disorder on admission (OR, 1.70; 95% CI, 1.60-1.81),³² hepatic disease (OR, 1.53; 95% CI, 1.09-2.15),^{8,32} and renal failure (OR, 1.43; 95% CI, 1.06-1.93).^{8,32} One study assessed renal failure by severity and the results showed that there probably is a greater magnitude of association between risk of bleeding and severe renal failure (glomerular filtration rate [GFR] < 30 mL/min/m²) vs no severe renal failure (OR, 2.14; 95% CI, 1.22-3.75)⁸ and a smaller magnitude of association between risk of bleeding and moderate renal failure (GFR, 30-59 mL/min/m²) compared with no moderate renal failure (OR, 1.37; 95% CI, 0.84-2.23).⁸ We also identified moderate-certainty evidence that there is probably an association between risk of bleeding and CVC use (OR, 1.37; 95% CI, 0.83-2.26)^{8,32} and antithrombotic medication use (OR, 1.28; 95% CI, 1.01-1.64).^{8,32}

We found moderate-certainty evidence that there is probably little to no association between risk of bleeding and hormone use, defined as estrogen intake (OR, 0.95; 95% CI, 0.82-1.10).³²

Low-quality evidence showed that there may be an association between risk of bleeding and autoimmune disease (OR, 1.30; 95% CI, 0.77-2.19).^{8,32} However, we identified low-certainty evidence that there may be little to no association between risk of bleeding and malignancy (OR, 1.08; 95% CI, 0.42-2.77).^{8,32}

Laboratory and physical examination factors There is moderate-certainty evidence of a probable association between risk of bleeding and anemia as the reason for admission (OR, 5.15; 95% CI, 2.45-10.81),^{8,33} morbid obesity (BMI ≥ 40 kg/m²; OR, 3.08; 95% CI, 1.35-7.02),³³ and low hemoglobin (< 13 g/dL in men and < 11.5 g/dL in women; OR, 2.33; 95% CI, 1.04-5.22).³³

Discussion

Summary of findings

We evaluated prognostic factors for VTE and bleeding in hospitalized medical patients. We identified 23 prognostic factors for VTE and 15 for bleeding, some supported by moderate certainty of the evidence. Age, critical illness, CVC use, and autoimmune disease were prognostic of both outcomes. Obesity (BMI > 30 kg/m²) and morbid obesity (BMI > 40 kg/m²) were associated with VTE and bleeding, respectively. However, only age, critical illness, and autoimmune disease had the same quality of evidence for the 2 outcomes. This study is unique in many aspects including its comprehensiveness, the novelty of its findings, and its transparent approach.

Strengths

Methodologically, our study benefits from the rigorous methods, the breadth of our search, our duplicate and independent screening, the data abstraction process used, and our assessment of the certainty of evidence on the basis of a structured framework. Also, we conducted sensitivity analyses to compare different duration cutoffs for immobility and to address differences in type of outcome. Other strengths include the involvement of a number of content and methodological experts.

Limitations and challenges

A potential limitation in terms of the search strategy was the focus on prevention, which we applied to restrict the rather large number of citations that we identified in our searches and because we believed that we would not miss relevant studies. To confirm this, we checked a large random sample ($n = 3000$) of citations obtained from a search not including a restriction to the topic of prevention. We did not identify any study that would have fulfilled the inclusion criteria and, therefore, our original search was unlikely to have missed eligible studies. Also, time bias may be a potential limitation, as we identified some of the prognostic factors from older studies. This may overestimate VTE events when considering overall trends in the reduction of VTE events in hospitalized medical patients over time.

Potential limitations of the included studies relate to the inconsistency and variability across eligibility criteria in the original studies and variability in study design, study type, sample size, and definitions of the prognostic factors. Other challenges include inconsistency in methods of measurement used across studies and contamination of the population with nonmedical hospitalized patients.

Implications for practice

Our study identified candidate prognostic factors for VTE and bleeding that have been considered in the analysis of some developed and widely used RAMs in daily practice, such as the Caprini, IMPROVE (International Medical Prevention Registry on Venous Thromboembolism) VTE, IMPROVE bleed, and Padua models.^{7,40-42} However, some factors that we identified as having

a probable association with VTE, based on our meta-analysis results, were not included or considered in the development of most of the RAMs, such as elevated CRP >10 mg/L (OR, 10.10), lower BI scores (BI ≤ 9; OR, 8.30), history of malignancy (OR, 3.20), and elevated heart rate (>100 beats/min; OR, 2.48). In addition, we found that an elevated fibrinogen level was inversely associated with DVT risk in patients with early stroke.³⁵ This observation was opposite to the finding that elevated CRP, another acute-phase prognostic factor, showed an association with DVT risk. The authors speculated that this finding may be a result of fibrinogen depletion due to active clot formation.³⁵ We believe that such reverse causation, given the study design, may be plausible. However, given the small sample size, the finding warrants further investigations in primary studies. In terms of bleeding, the candidate prognostic factor antithrombotic use showed mixed results in the different studies and was included only in 1 final model.³² However, we identified a probable association with the outcome when the individual results were subjected to meta-analysis,^{8,25,33} perhaps because of the limitation of the databases used that may not include all potential prognostic factors. Another reason that their findings may have been limited is the methods used in the development of RAMs for the multivariate analysis, such as automated procedures (eg, backward or forward) used for prognostic factor selection or selection of factors based on statistical significance at the univariate analysis stage. Therefore, the findings in our study ensure the consideration of all identified potential prognostic factors in the literature during the development of a RAM, a better assessment of the databases being used, and the comprehensiveness of the factors included in the databases. Studies in this systematic review included patients who received thromboprophylaxis, which may have altered the risk estimates. However, we controlled for the use of thromboprophylaxis in several ways. We selected studies only if they included a small fraction of patients receiving thromboprophylaxis (less than 10%) or if it was controlled for in the statistical analysis. Beyond that, given the general assumption that the relative risk related to a prognostic factor remains largely unaffected by administration of thromboprophylaxis (although the baseline risk of course might change), we used relative estimates of the risk. This assumption is supported by the observation that, in several studies, the relative estimates of risk were not influenced by adjustment for prophylaxis use in the statistical analyses. We believe that these measures should address the influence of thromboprophylaxis on the prognostic factor we addressed herein.

Implications for future research

Research may be needed to reevaluate existing RAMs, as the developers of the models may not have been able to use the variables we identified, given the limitations in the existing databases. However, a full development or improvement of a RAM that supports clinical practice requires further investigation of all the prognostic factors we identified in our study.

Conclusion

In this systematic review, we identified all reported relevant prognostic factors for VTE and bleeding in hospitalized medical patients. Some of these factors are not part of current risk prediction for VTE and bleeding in hospitalized medical patients. Our findings will help inform experts in developing population-

based guidelines and accurate, user-friendly RAMs to better guide individual patient prophylactic management.

Acknowledgments

This work was supported by a subcontract (200-2016-92458) from the US Centers for Disease Control and Prevention (CDC), through Karna LLC.

The CDC, through Karna LLC, had no role in the detailed design and conduct of the study; the collection, management, analysis, and interpretation of the data; or the preparation of the manuscript. The findings and conclusions of this report are those of the authors and do not necessarily represent the official position of the CDC.

Authorship

Contribution: A.J.D. and H.J.S. were responsible for the conception and design of the study; A.J.D., S.G.K., R.C., I.E.-I., F.G., M.R., A.A., R.Z.M., and H.J.S. performed data acquisition; A.J.D., M.R., L.M., and H.J.S. performed data analyses; A.J.D., S.G.K., I.E.-I., M.C., M.K.G., L.M., F.A.S., A.C.S., M.B.S., S.W., N.A.Z., F.G., M.R., A.A., R.Z.M., A.I., E.A.A., and H.J.S. interpreted the results; A.J.D. and H.J.S. drafted the manuscript; and all authors critically revised the manuscript and approved the final version.

Conflict-of-interest disclosure: A.J.D., I.E.-I., A.I., E.A.A., and H.J.S. are members of the GRADE working group. A.C.S. has received remuneration for consulting from Bayer, Janssen, and Portola and research support grants from Boehringer Ingelheim, Janssen, and the Center for Medicare and Medicaid Services. A.C.S. also reported an intellectual conflict as the lead in the group that derived and validated the IMPROVE tool for VTE risk assessment in hospitalized medical patients. M.B.S. has received remuneration for consulting from Bayer, Janssen, Pfizer, and Portola and research support grants from Boehringer-Ingelheim, Janssen, Portola, and Roche. M.C. is a former board member (2013-2017) of the American Heart Association and chaired the committee that produced the American Society of Hematology (ASH) 2018 Guidelines for Management of Venous Thromboembolism: Prophylaxis for Hospitalized and Nonhospitalized Medical Patients (ASH 2018 Guidelines for Management of VTE). F.A.S. and N.A.Z. reported participating as panel members for the ASH 2018 Guidelines for Management of VTE. N.A.Z. also reported receiving honoraria in 2017 from ASH for the Highlights of ASH 2017 presentation (Dallas, New York, and Latin America). H.J.S. was cochair of the ASH 2018 Guidelines for Management of VTE committee and procured grant funding from the CDC for this study. The remaining authors declare no competing financial interests.

ORCID profiles: A.J.D., 0000-0002-2498-1697; S.G.K., 0000-0002-5167-4246; M.C., 0000-0002-7871-6143; L.M., 0000-0001-5855-5461; M.B.S., 0000-0002-6429-3493; S.W., 0000-0002-2522-2705; F.G., 0000-0002-0802-3616; M.R., 0000-0002-0530-9491; R.Z.M., 0000-0003-2131-3711; A.I., 0000-0002-3331-8766; H.J.S., 0000-0003-3211-8479.

Correspondence: Holger J. Schünemann, Department of Health Research Methods, Evidence, and Impact, McMaster University, 1280 Main St West, Hamilton, ON L8N 3Z5, Canada; e-mail: schuneh@mcmaster.ca.

Footnotes

Submitted 10 October 2019; accepted 12 February 2020; prepublished online on *Blood* First Edition 24 February 2020. DOI 10.1182/blood.2019003603.

The online version of this article contains a data supplement.

There is a *Blood* Commentary on this article in this issue.

The publication costs of this article were defrayed in part by page charge payment. Therefore, and solely to indicate this fact, this article is hereby marked "advertisement" in accordance with 18 USC section 1734.

REFERENCES

- Cushman M. Epidemiology and risk factors for venous thrombosis. *Semin Hematol*. 2007;44(2):62-69.
- Schünemann HJ, Cushman M, Burnett AE, et al. American Society of Hematology 2018 guidelines for management of venous thromboembolism: prophylaxis for hospitalized and nonhospitalized medical patients. *Blood Adv*. 2018;2(22):3198-3225.
- Cruden P, Cushman M, Repp AB. Hospitalist assessment of venous thromboembolism and bleeding risk: A survey study. *Thromb Res*. 2019;178:155-158.
- Anderson FA Jr, Zayaruzny M, Heit JA, Fidan D, Cohen AT. Estimated annual numbers of US acute-care hospital patients at risk for venous thromboembolism. *Am J Hematol*. 2007;82(9):777-782.
- Goldhaber SZ, Dunn K, MacDougall RC. New onset of venous thromboembolism among hospitalized patients at Brigham and Women's Hospital is caused more often by prophylaxis failure than by withholding treatment. *Chest*. 2000;118(6):1680-1684.
- Kahn SR, Lim W, Dunn AS, et al. Prevention of VTE in nonsurgical patients: antithrombotic therapy and prevention of thrombosis: American College of Chest Physicians evidence-based clinical practice guidelines. *Chest*. 2012;141(2 suppl):e195S-e226S.
- Spyropoulos AC, Anderson FA Jr, FitzGerald G, et al; IMPROVE Investigators. Predictive and associative models to identify hospitalized medical patients at risk for VTE. *Chest*. 2011;140(3):706-714.
- Decousus H, Tapson VF, Bergmann J-F, et al; IMPROVE Investigators. Factors at admission associated with bleeding risk in medical patients: findings from the IMPROVE investigators. *Chest*. 2011;139(1):69-79.
- Spyropoulos AC. Risk assessment of venous thromboembolism in hospitalized medical patients. *Curr Opin Pulm Med*. 2010;16(5):419-425.
- Riley RD, Moons KG, Snell KI, Ensor J, Hooft L, Altman DG, et al. A guide to systematic review and meta-analysis of prognostic factor studies. *BMJ*. 2019;364:k4597.
- Rocha AT, Paiva EF, Lichtenstein A, Milani R Jr, Cavalheiro CF, Maffei FH. Risk-assessment algorithm and recommendations for venous thromboembolism prophylaxis in medical patients. *Vasc Health Risk Manag*. 2007;3(4):533-553.
- Green S, Higgins J, eds. *Cochrane handbook for systematic reviews of intervention*. London: Cochrane; 2005
- Kuderer NM, Poniewierski MS, Culakova E, et al. Predictors of venous thromboembolism and early mortality in lung cancer: results from a Global Prospective Study (CANTARISK). *Oncologist*. 2018;23(2):247-255.
- Lee J-C, Ro YS, Cho J, et al. Characteristics of venous thromboembolism in pancreatic adenocarcinoma in east Asian ethnics: a large population-based observational study. *Medicine (Baltimore)*. 2016;95(17):e3472.
- Iorio A, Spencer FA, Falavigna M, et al. Use of GRADE for assessment of evidence about prognosis: rating confidence in estimates of event rates in broad categories of patients. *BMJ*. 2015;350:h870
- Hemingway H, Croft P, Perel P, et al. Prognosis research strategy (PROGRESS) 1: A framework for researching clinical outcomes. *BMJ* 2013; 346:e5595.
- Wolff RWP, Whiting P, Mallett S, et al, eds. PROBAST: a "risk of bias" tool for prediction modelling studies [abstract]. *Cochrane Colloquium Abstracts*. 2015
- Tang SW, Zhang Y, Tao BL, Yang ZR, Sun F, Zhan SY. Risk of bias assessment: (7) Assessing Bias in Studies of Prognostic Factors [in Chinese]. *Zhonghua Liu Xing Bing Xue Za Zhi*. 2018;39(7):1003-1008.
- Hayden JA, van der Windt DA, Cartwright JL, Côté P, Bombardier C. Assessing bias in studies of prognostic factors. *Ann Intern Med*. 2013;158(4):280-286.
- Reeves BC, Deeks JJ, Higgins JPT, Wells GA. Including non-randomized studies. In: *Cochrane Handbook for Systematic Reviews of Interventions*. 2008:391.
- Ahn S, Fessler JA. Standard errors of mean, variance, and standard deviation estimators. Ann Arbor: Electrical Engineering and Computer Science Department, The University of Michigan; 2003:1-2
- Grant RL. Converting an odds ratio to a range of plausible relative risks for better communication of research findings [published correction appears in *BMJ*. 2014;348:g2124]. *BMJ*. 2014;348(1):f7450.
- Wang Z. Converting odds ratio to relative risk in cohort studies with partial data information. *J Stat Softw*. 2013;55(5):1-11.
- Review manager (RevMan)[computer program]. Version 5.2; 2011.
- Mahan CE, Liu Y, Turpie AG, et al. External validation of a risk assessment model for venous thromboembolism in the hospitalised acutely-ill medical patient (VTE-VALOURR). *Thromb Haemost*. 2014;112(4):692-699.
- Rosenberg D, Eichorn A, Alarcon M, McCullagh L, McGinn T, Spyropoulos AC. External validation of the risk assessment model of the International Medical Prevention Registry on Venous Thromboembolism (IMPROVE) for medical patients in a tertiary health system. *J Am Heart Assoc*. 2014;3(6):e001152.
- Zakai NA, Callas PW, Repp AB, Cushman M. Venous thrombosis risk assessment in medical inpatients: the medical inpatients and thrombosis (MITH) study. *J Thromb Haemost*. 2013;11(4):634-641.
- Zakai NA, Wright J, Cushman M. Risk factors for venous thrombosis in medical inpatients: validation of a thrombosis risk score. *J Thromb Haemost*. 2004;2(12):2156-2161.
- Zhou H, Hu Y, Li X, et al. Assessment of the Risk of Venous Thromboembolism in Medical Inpatients using the Padua Prediction Score and Caprini Risk Assessment Model. *J Atheroscler Thromb*. 2018;25(11):1091-1104.
- Barclay SM, Jeffres MN, Nguyen K, Nguyen T. Evaluation of pharmacologic prophylaxis for venous thromboembolism in patients with chronic liver disease. *Pharmacotherapy*. 2013; 33(4):375-382.
- Grant PJ, Greene MT, Chopra V, Bernstein SJ, Hofer TP, Flanders SA. Assessing the caprini score for risk assessment of venous thromboembolism in hospitalized medical patients. *Am J Med*. 2016;129(5):528-535.
- Mahan CE, Spyropoulos AC, Fisher MD, et al. Antithrombotic medication use and bleeding risk in medically ill patients after hospitalization. *Clin Appl Thromb Hemost*. 2013;19(5):504-512.
- Patell R, Gutierrez A, Rybicki L, Khorana AA. Identifying predictors for bleeding in hospitalized cancer patients: A cohort study. *Thromb Res*. 2017;158:38-43.
- Rothberg MB, Lindenauer PK, Lahti M, Pekow PS, Selker HP. Risk factor model to predict venous thromboembolism in hospitalized medical patients. *J Hosp Med*. 2011;6(4):202-209.
- Bembek J, Karlinski M, Kobayashi A, Czlonkowska A. Early stroke-related deep venous thrombosis: risk factors and influence on outcome. *J Thromb Thrombolysis*. 2011; 32(1):96-102.
- Fan J, Li X, Cheng Y, Yao C, Zhong N; Investigators Group. Measurement of D-dimer as aid in risk evaluation of VTE in elderly patients hospitalized for acute illness: a prospective, multicenter study in China. *Clin Invest Med*. 2011;34(2):E96-E104.
- Kelly J, Rudd A, Lewis RR, Coshall C, Moody A, Hunt BJ. Venous thromboembolism after acute ischemic stroke: a prospective study using magnetic resonance direct thrombus imaging. *Stroke*. 2004;35(10):2320-2325.
- Ota S, Yamada N, Tsuji A, Ishikura K, Nakamura M, Ito M. Incidence and clinical predictors of deep vein thrombosis in patients hospitalized with heart failure in Japan. *Circ J*. 2009;73(8):1513-1517.
- Yi X, Lin J, Han Z, Zhou X, Wang X, Lin J. The incidence of venous thromboembolism following stroke and its risk factors in eastern China. *J Thromb Thrombolysis*. 2012;34(2):269-275.
- Barbar S, Noventa F, Rossetto V, et al. A risk assessment model for the identification of hospitalized medical patients at risk for venous thromboembolism: the Padua Prediction Score. *J Thromb Haemost*. 2010;8(11):2450-2457.
- Caprini JA, Arcelus JI, Reyna J, eds. Effective risk stratification of surgical and nonsurgical patients for venous thromboembolic disease. *Semin Hematol*. 2001;38(suppl 5):12-19.
- Spyropoulos AC, Ageno W, Albers GW, et al; MARINER Investigators. Rivaroxaban for thromboprophylaxis after hospitalization for medical illness. *N Engl J Med*. 2018;379(12):1118-1127.