



## MYELOID NEOPLASIA

Comment on Zhen et al, page 2373

# RUNX1 and *inv*(16) are frenemies in AML

Sridhar Rao | Blood Research Institute, Versiti; Medical College of Wisconsin

**In this issue of *Blood*, Zhen et al<sup>1</sup> provide novel molecular insights into the requirement for *Runx1* in a mouse model of *inv*(16) acute myeloid leukemia (AML).**


Core binding factor (CBF) leukemias are a common subtype of de novo AML and are represented by 2 distinct chromosomal abnormalities, t(8;21) and *inv*(16), which result in the fusion oncoproteins RUNX1-RUNX1T1 and CBFβ-SMMHC, respectively.<sup>2-5</sup> They derive their name from normal CBF, which is a heterodimeric transcription factor that regulates gene

expression and is composed of a DNA-binding α subunit (RUNX1, RUNX2, or RUNX3) and the non-DNA-binding CBFβ subunit. Although the 2 leukemias are treated similarly with high-dose chemotherapy and generally are considered favorable risk, up to 40% of patients relapse and subsequently require additional therapies, including allogeneic bone marrow

transplantation, to achieve long-term survival. Relapses are due to the presence of a small number of leukemia-initiating cells (LICs; also called leukemia stem cells), which are resistant to conventional chemotherapy. Collectively, this illustrates the need for novel targeted therapies to improve survival while reducing toxicities for *inv*(16) AML patients.

*Inv*(16) has generally been considered a classic example of the original Gilliland 2-hit model of AML in which a proliferative signal is provided by a mutation, resulting in an activated signaling molecule, in combination with the loss of a transcription factor, which promotes the expansion of undifferentiated cells.<sup>6</sup> In line with this model, CBFβ-SMMHC has been thought to cause AML by sequestering RUNX1, preventing it from regulating myeloid maturation and resulting in a differentiation block.<sup>3,7</sup> This would suggest that a complete loss of RUNX1 activity is required for *inv*(16)-driven leukemia; however, there are other contradictory lines of evidence suggesting that RUNX1 may play a role in leukemogenesis. One example is that loss-of-function mutations in *RUNX1* are often found in t(8;21), whereas they are rarely seen in *inv*(16), implying that there may be a negative selection bias against additional reduction in RUNX1 activity.<sup>4,5</sup> As such, whether *inv*(16) requires some attenuated amount of RUNX1 to drive leukemia remains an important and unresolved question. This information is critical because it provides an important opportunity for developing targeted therapies.

Zhen et al used a genetically engineered “knock-in” allele that inducibly recapitulates *inv*(16) in mice (*Mx1-Cre; Cbfβ<sup>+/56M</sup>*), in combination with an inducible null allele of *Runx1* (*Runx1<sup>fl/fl</sup>*), to definitively determine whether *Runx1* is required for AML driven by CBFβ-SMMHC. Although the *inv*(16) animals with normal *Runx1* developed AML with a latency of 16 weeks, similar to prior studies,<sup>8</sup> animals deficient in *Runx1* did not develop AML for up to 1 year, demonstrating that loss of



CBFβ-SMMHC requires RUNX1 for leukemia development. (A) Under normal conditions, hematopoietic stem and progenitor cells (HSPC) are converted to LICs by the action of the *inv*(16)-derived fusion oncoprotein, CBFβ-SMMHC, in combination with an activating mutation in a signaling molecule, such as *NRAS*, *KRAS*, or *FLT3*. The existing paradigm has been that CBFβ-SMMHC drives leukemia development by acting as a dominant negative which sequesters RUNX1, preventing it from promoting normal myeloid development. Zhen et al demonstrate that loss of *RUNX1* prevents leukemia development in mice by preventing the development and maintenance of AMPs, a type of LIC specific to *inv*(16). (B) In AMP cells, RUNX1 recruits CBFβ-SMMHC to chromatin, where RUNX1:CBFβ-SMMHC predominantly activate cell cycle gene expression while simultaneously repressing a smaller number of genes that promote differentiation.

*Runx1* blocks CBF $\beta$ -SMMHC–driven leukemia development. Given the block in leukemia development, the investigators hypothesized that LICs were negatively affected by loss of *Runx1*. To test this hypothesis, they quantified a previously described LIC population termed abnormal myeloid progenitors (AMPs; Lin<sup>-</sup>c-Kit<sup>+</sup>Sca1<sup>-</sup>CD34<sup>-</sup>FC $\gamma$ II/III<sup>+</sup>).<sup>8</sup> Importantly, the investigators found that loss of *Runx1* hindered the development and maintenance of the AMP population, implying that *Runx1* is required for CBF $\beta$ -SMMHC–driven LICs (see figure panel A).

Given the prior paradigm that CBF $\beta$ -SMMHC prevents RUNX1 from binding DNA, the investigators performed a series of detailed molecular studies to delineate the requirement for CBF $\beta$ -SMMHC and RUNX1 in the LIC population. Using bulk and single-cell RNA-sequencing, the investigators identified 2 hallmarks of LICs generated by CBF $\beta$ -SMMHC but lacking RUNX1. First, these LICs showed reduced expression of genes known to be critical drivers of the cell cycle, including the G2M checkpoint and targets of c-Myc and E2F. Second, they showed increased expression of genes required for normal myeloid differentiation. Collectively, these data imply that CBF $\beta$ -SMMHC works collaboratively with RUNX1 to mediate the proliferative signal and differentiation block, which are hallmarks of the Gilliland model for AML development.

Next, to definitively establish how CBF $\beta$ -SMMHC and RUNX1 cooperate to control gene expression, the investigators demonstrate that RUNX1 recruits CBF $\beta$ -SMMHC to chromatin, and they work in concert to activate gene expression, including genes that control proliferation (see figure panel B). This is in sharp contrast to prior literature that reported that CBF $\beta$ -SMMHC operates predominantly to repress gene expression. These important mechanistic studies connect how CBF $\beta$ -SMMHC and RUNX1 work collaboratively to directly regulate genes critical to LICs. It is important to note that these data represent 1 of the important ways that CBF $\beta$ -SMMHC operates during leukemia development, but it may be that the existing model of CBF $\beta$ -SMMHC sequestering RUNX1 occurs in parallel and is also critical to leukemogenesis. As such, it may be that, although CBF $\beta$ -SMMHC sequesters most RUNX1 during leukemia development, a small amount of active RUNX1 is required for transformation.

Perhaps most importantly, this work has the potential to inform targeted therapies. Although current treatment regimens rely on high-dose chemotherapy, an ongoing concern remains that conventional therapeutics are not effective at targeting LICs, which can cause relapse even when present in miniscule amounts. Prior studies have identified a small molecular inhibitor of the CBF $\beta$ -SMMHC:RUNX1 protein:protein interaction as a potential therapy for inv(16) AML.<sup>9,10</sup> Zhen et al elevate this potential approach by demonstrating that it could directly target LICs, an important driver of relapse. In doing so, the investigators provide a path forward to develop more effective and less toxic therapies for patients with inv(16) AML.

**Conflict-of-interest disclosure:** The author declares no competing financial interests. ■

## REFERENCES

- Zhen T, Cao Y, Ren G, et al. RUNX1 and CBF $\beta$ -SMMHC transactivate target genes together in abnormal myeloid progenitors for leukemia development. *Blood*. 2020;136(21):2373-2385.
- Schoch C, Kern W, Schnittger S, Buchner T, Hiddemann W, Haferlach T. The influence of age on prognosis of de novo acute myeloid leukemia differs according to cytogenetic subgroups. *Haematologica*. 2004;89(9):1082-1090.
- Goyama S, Mulloy JC. Molecular pathogenesis of core binding factor leukemia: current knowledge and future prospects. *Int J Hematol*. 2011;94(2):126-133.
- Duployez N, Marceau-Renaut A, Boissel N, et al. Comprehensive mutational profiling of core binding factor acute myeloid leukemia. *Blood*. 2016;127(20):2451-2459.
- Faber ZJ, Chen X, Gedman AL, et al. The genomic landscape of core-binding factor acute myeloid leukemias. *Nat Genet*. 2016;48(12):1551-1556.
- Gilliland DG. Hematologic malignancies. *Curr Opin Hematol*. 2001;8(4):189-191.
- Adya N, Stacy T, Speck NA, Liu PP. The leukemic protein core binding factor  $\beta$  (CBF $\beta$ )-smooth-muscle myosin heavy chain sequesters CBF $\alpha$ 2 into cytoskeletal filaments and aggregates. *Mol Cell Biol*. 1998;18(12):7432-7443.
- Kuo YH, Landrette SF, Heilman SA, et al. Cbf beta-SMMHC induces distinct abnormal myeloid progenitors able to develop acute myeloid leukemia. *Cancer Cell*. 2006;9(1):57-68.
- Illendula A, Pulikkan JA, Zong H, et al. A small-molecule inhibitor of the aberrant transcription factor CBF $\beta$ -SMMHC delays leukemia in mice. *Science*. 2015;347(6223):779-784.
- Pulikkan JA, Hegde M, Ahmad HM, et al. CBF $\beta$ -SMMHC inhibition triggers apoptosis by disrupting MYC chromatin dynamics in acute myeloid leukemia [published correction appears in *Cell*. 2018;174(5):1325]. *Cell*. 2018;174(1):172-186.e21.

DOI 10.1182/blood.2020008802

© 2020 by The American Society of Hematology

## CLINICAL TRIALS AND OBSERVATIONS

Comment on Bartlett et al, page 2401

# Hodgkin lymphoma: outsmarting HRS cells

Joseph M. Connors | BC Cancer Centre for Lymphoid Cancer

**In their article in this issue of *Blood* describing the use of a new bispecific antibody, AFM13, plus a checkpoint inhibitor, pembrolizumab, for patients with recurrent or refractory classic Hodgkin lymphoma (cHL), Bartlett and her coauthors provide dramatic evidence of the treatment potency that can be achieved with a 3-pronged therapeutic attack employing 2 biologic agents targeting specific immunologic characteristics of cHL Hodgkin and Reed-Sternberg (HRS) cells.<sup>1</sup> They show how to forcefully recruit the patient's immune system's killer cells while at the same time enhancing their cytotoxicity. Their work nicely demonstrates how recent progress in the treatment of cHL has been carefully built on an increasingly exquisite understanding of the molecular biology underlying the immunology of malignant HRS cells and their dynamic interaction with both specific and nonspecific effector cells in the immune system.**

The universal, persistent expression of the CD30 antigen on the surface of cHL cells provides an attractive target for specially

crafted antibodies, a property that has been effectively exploited by the development of the antibody-drug conjugate