MYELOID NEOPLASIA

Plasmacytoid dendritic cell expansion defines a distinct subset of *RUNX1*-mutated acute myeloid leukemia

Wenbin Xiao,^{1,2} Alexander Chan,¹ Michael R. Waarts,² Tanmay Mishra,² Ying Liu,¹ Sheng F. Cai,^{2,3} Jinjuan Yao,⁴ Qi Gao,¹ Robert L. Bowman,² Richard P. Koche,⁵ Isabelle S. Csete,² Nicole L. DelGaudio,² Andriy Derkach,⁶ Jeeyeon Baik,¹ Sophia Yanis,¹ Christopher A. Famulare,⁷ Minal Patel,⁷ Maria E. Arcila,^{1,4} Maximilian Stahl,³ Raajit K. Rampal,^{2,3} Martin S. Tallman,³ Yanming Zhang,⁸ Ahmet Dogan,¹ Aaron D. Goldberg,³ Mikhail Roshal,¹ and Ross L. Levine^{2,3,5,7}

¹Department of Pathology, Hematopathology Service, ²Human Oncology and Pathogenesis Program, Molecular Cancer Medicine Service, ³Department of Medicine, Leukemia Service, ⁴Department of Pathology, Molecular Diagnostic Laboratory, ⁵Center for Epigenetics Research, ⁶Department of Epidemiology-Biostatistics, ⁷Center for Hematologic Malignancies, and ⁸Department of Pathology, Cytogenetics Laboratory, Memorial Sloan Kettering Cancer Center, New York, NY

KEY POINTS

- pDC-AML is characterized by a high frequency of RUNX1 mutations and increased expression of a pDC transcriptional program.
- CD123 targeting represents a potential treatment approach for pDC-AML.

Plasmacytoid dendritic cells (pDCs) are the principal natural type I interferon–producing dendritic cells. Neoplastic expansion of pDCs and pDC precursors leads to blastic plasmacytoid dendritic cell neoplasm (BPDCN), and clonal expansion of mature pDCs has been described in chronic myelomonocytic leukemia. The role of pDC expansion in acute myeloid leukemia (AML) is poorly studied. Here, we characterize patients with AML with pDC expansion (pDC-AML), which we observe in ~5% of AML cases. pDC-AMLs often possess cross-lineage antigen expression and have adverse risk stratification with poor outcome. *RUNX1* mutations are the most common somatic alterations in pDC-AML (>70%) and are much more common than in AML without pDC expansion and BPDCN. We demonstrate that pDCs are clonally related to, as well as originate from, leukemic blasts in pDC-AML. We further demonstrate that leukemic blasts from *RUNX1*-mutated AML upregulate a pDC transcriptional program, poising the cells toward pDC differentiation and expansion. Finally, tagraxofusp, a targeted therapy directed to CD123, reduces leukemic burden

eliminates pDCs in a patient-derived xenograft model. In conclusion, pDC-AML is characterized by a high frequency of *RUNX1* mutations and increased expression of a pDC transcriptional program. CD123 targeting represents a potential treatment approach for pDC-AML. (*Blood*. 2021;137(10):1377-1391)

Introduction

Plasmacytoid dendritic cells (pDCs) are the principal natural type l interferon, interferon- α (IFN- α)-producing dendritic cells, that play critical roles in the immune response.^{1,2} Human pDCs can be readily identified by flow cytometry based on high expression of CD123, HLA-DR, and CD303/BDCA2 in the absence of other lineage markers.³ pDCs express the cytokine receptor FLT3 (CD135), whose ligand FLT3L is necessary and sufficient for pDC development from myeloid and lymphoid progenitors,^{4,5} suggesting the existence of a robust transcriptional program that drives pDC specification and differentiation.⁶ TCF4, a master transcription factor,^{7,8} acts jointly with MTG16 and additional factors, such as BCL11A, to promote pDC development^{9,10} by activating transcription factors involved in pDC differentiation (SPIB, IRF7, and IRF8).7 IRF family members, particularly IRF7, play a key role in activating IFN-I genes by interacting with other important transcription factors, such as RUNX2, SPIB, and NFATC3.¹¹⁻¹⁴ CXXC5 and TET2 have been shown to promote

IRF7 expression by maintaining hypomethylation of the *IRF7* promoter in pDCs.¹⁵

The role of pDCs in malignancy has just begun to unfold. The nature of IFN- α -producing pDCs has led to the hypothesis that pDCs may possess antitumor activities. pDCs, via their secreted IFN-α, promote natural killer cell-mediated killing of tumor cells in vitro.¹⁶⁻¹⁸ Topical treatment with imiquimod, one of the best characterized imidazoquinoline compounds and a Toll-like receptor 7 agonist, leads to the recruitment of pDCs and subsequent secretion of IFN- α and granzyme B, which correlate with regression of basal cell carcinoma and melanoma.^{19,20} In contrast, recent studies have shown that pDC accumulation is associated with progression in different human cancer contexts.²¹⁻²³ Depletion of pDCs inhibits breast cancer cell growth and metastasis in a mouse model.²² It has been shown that tumor-associated pDCs possess impaired IFN-a production, resulting in a microenvironment that favors regulatory T-cell expansion. In sum, pDCs may have antitumor or protumor

effects, depending upon the malignancy and inflammatory context. $^{\rm 24}$

Normally, pDCs account for <1% of total nucleated cells in bone marrow (BM) and peripheral blood.²⁵⁻²⁷ Neoplastic expansion of pDCs or pDC precursors leads to blastic plasmacytoid dendritic cell neoplasm (BPDCN),^{28,29} and clonal expansion of mature pDCs has been reported in chronic myelomonocytic leukemia (CMML).30 In contrast, the role of pDC expansion in acute myeloid leukemia (AML) remains to be elucidated. A few small patient series have reported skin, nodal, and marrow pDC expansion in AML,³¹⁻³⁷ and a possible correlation with FLT3^{ITD} mutations has been proposed.^{38,39} In 2018, we reported our preliminary analysis of 24 patients with AML with pDC expansion, suggesting that this might represent an entity with distinct genetic and biologic features.⁴⁰ Here, we performed detailed genetic, transcriptional, and functional characterization of patients with AML with pDC expansion (pDC-AML) and investigate potential therapeutic interventions for this specific AML subset.

Patients and methods

Patients

Patients with AML were identified from the patient database at the Memorial Sloan Kettering Cancer Center (MSKCC) between January 2014 and December 2019. The key words "plasmacytoid dendritic cells" or "plasmacytoid dendritic cell" were searched for flow cytometric and pathology reports. The clinical, morphologic, immunophenotypic, and cytogenetic/molecular results were independently reviewed again by 3 hematopathologists (W.X., A.C. and M.R.). This study was approved by the Institutional Review Board at MSKCC.

Additional details about additional patient cohorts, cell culture, flow cytometric sorting and analysis, chromosome analysis, fluorescent in situ hybridization studies, sequencing studies, patient-derived xenograft, and statistics are included in the supplemental Methods (available on the *Blood* Web site).

Results

A subset of AML is characterized by increased levels of pDCs

pDCs were identified by their characteristic immunophenotype on flow cytometry^{3,6}: specifically low side scatter and CD45 (dim), CD123 (bright), and HLA-DR surface expression (Figure 1A-B; supplemental Figure 1; supplemental Tables 1 and 2). CD303 was also evaluated in a subset of subjects. The median proportion of pDCs in normal marrow controls was 0.29% (interquartile range [IQR], 0.19-0.37) (Figure 1C). pDCs were markedly depleted in the BM of the majority of patients with AML, with a median pDC proportion of 0.03% (IQR, 0.006-0.14%), a 10-fold reduction compared with normal controls (P < .001). Although none of the normal controls had >1% pDCs, there have been reports of as high as 1.6% pDCs in other normal control cohorts.²⁶ Therefore, 2% was chosen to delineate a stringent cutoff for pDC expansion, which is equivalent to 10 standard deviations (SDs) from the mean pDC level of normal controls. Among 850 AML patients, we identified 26 AML patients with \geq 2% pDCs in their diagnostic marrow samples. In addition, 16 AML patients had \geq 2% pDCs at a later time point after diagnosis (either with persistent disease or relapse). Therefore, 4.9% (42/850) of total patients with AML in our study cohort had pDC expansion (hereafter referred to as pDC-AML). In these patients, the proportion of pDCs ranged from 2.2% to 35.9%, with a median of 7.7% (IQR, 3.4-9.9; P < .0001 vs normal control) (Figure 1C).

pDCs and blasts from pDC-AML are immunophenotypically distinct from those from AML without pDC expansion

We next performed immunophenotypic characterization of pDCs in pDC-AML samples and compared this with a cohort of BPDCN samples. CD123 was universally expressed on BPDCN cells, albeit at lower levels compared with normal pDCs, as previously reported (median of mean fluorescence intensity [MFI]): 10855, P = .0045 vs normal controls, P = .0015 vs pDC-AML) (Figure 1D).⁴¹ By contrast, we found that pDCs in pDC-AML patients had high levels of CD123 expression, similar to pDCs from normal controls (MFI, 27 613 vs 25 611, P = .7). pDC distribution in AML, as demonstrated by CD123 immunohistochemical staining, showed a predominantly interstitial pattern throughout the marrow intermixed with, but distinct from, CD34⁺ leukemic blasts (Figure 1E-F). Although small loose clusters of pDCs were also noted (Figure 1G-H), large aggregates, as seen in CMML, were not observed. Flow cytometry assisted sorting (FACS)-purified leukemic blasts and pDCs displayed distinct blastic and plasmacytoid morphology, respectively (Figure 1I-J). The pDCs from patients with pDC-AML expressed CD4, CD123, and HLA-DR at levels comparable to normal pDCs. However, they also frequently showed aberrant expression of CD56 (4/30; 13%), CD34 (25/41; 61%), CD5 (5/29, 17%), CD13 (17/41, 41%), and cytoplasmic terminal deoxynucleotidyltransferasede (4/10, 40%). Other markers expressed by normal pDCs were positive in only a subset of patients with pDC-AML, including CD303 (7/10, 70%), CD2 (8/29, 28%), and CD33 (11/33; 33%) (supplemental Figure 1; supplemental Table 2). This immunophenotype appears to be intermediate between normal pDCs and BPDCN⁴¹ and shows maturational continuity with the leukemic blasts.^{36,37}

The leukemic blasts from 13 of 41 (32%) pDC-AML patients showed expression of ≥ 2 crosslineage antigens (including CD2, cytoplasmic CD3, CD5, CD7, CD10, CD19, cytoplasmic CD79a) compared with only 3 of 100 (3%) AML patients without pDC expansion (P < .0001; Table 1). Eight of 41 (20%) patients met the immunophenotypic criteria (based on cytoplasmic CD3, CD19, and cytoplasmic CD79a) of mixed phenotype acute leukemia in contrast to only 2 of 100 (2%) patients with AML without pDC expansion (P < .0001; Table 1, supplemental Figure 1; supplemental Table 3). However, these 10 patients with mixed phenotype in both groups carried a diagnosis of secondary AML, which precludes a diagnosis of de novo mixed phenotype acute leukemia.⁴² Eighteen pDC-AML patients showed monocytic differentiation based on flow cytometric evidence of immature monocytes (≥5%) and/or CD64 expression on the leukemic blasts. After induction therapy, the number of pDCs declined significantly in proportion to the reduction in leukemic blasts (supplemental Figure 2A).

pDC-AML is associated with poor prognosis

Patients with pDC-AML had a median age of 68 years (IQR, 58-76) and male predominance (male/female ratio, 2.8:1) (Table 1;



Figure 1. Increased pDCs in a subset of AML. (A-B) Flow cytometric identification of pDCs (aqua population represents pDCs). (C) pDC proportion (percentage of white blood cells) in BM aspirates from patients with AML and normal controls (median ± IQR). AML, AML without pDC expansion; Controls, normal subjects; pDC-AML, AML with pDC expansion. (D) CD123 levels on CD34⁺ blasts and pDCs (median ± IQR). (E) Hematoxylin and eosin stain of BM biopsy from a representative patient with pDC-AML (inset: anti-CD34 immunostain). (F) Anti-CD123 immunostain of the patient from panel E. (G) Hematoxylin and eosin stain of BM biopsy from another representative patient with pDC-AML (inset: anti-CD34 immunostain). (H) Anti-CD123 immunostain of the patient in panel G. (I-J) Wright-Giemsa stain of flow-sorted leukemic blasts (I) and pDCs (J) from pDC-AML. **P < .01, ***P < .01. Original magnification, ×200 (E-H) and ×1000 (I-J). MFI, mean fluorescence intensity.

supplemental Table 4), similar to patients with AML without pDC expansion (referred to as AML) and patients with BPDCN (data not shown). Secondary AML (AML with myelodysplasia-related changes and therapy-related AML) were \sim 50% in both cohorts (22/42 in pDC-AML vs 45/100 in AML). Patients with pDC-AML had fewer circulating blasts (median 2.5% vs 20% in AML; P < .05). Skin involvement was more common in pDC-AML (6/42 [14%] vs 3/100 [3%] in AML; P = .01) but less frequent than in BPDCN (7/9 [78%]; P = .004). Cytogenetic abnormalities were identified in 25 of 40 (62.5%) patients with pDC-AML (results not available in 2 patients), and 9 of 40 (23%) pDC-AML patients had adverse cytogenetics, not different from the AML cohort (59% and 22%, respectively). The most common abnormality was del7 (5/40; 13%). Only 3 of 40 (7.5%) had trisomy 13. Thirty of 37 patients with pDC-AML (81%; unable to evaluate in 5 patients) had adverse risk based on European LeukemiaNet stratification compared with 35 of 99 (35%) in the AML cohort (P < .0001). Twenty-one of 42 (50%) patients with pDC-AML received induction therapy compared with 100% in the AML cohort (P < .0001). Six of 16 (38%) pDC-AML patients relapsed after allogeneic hematopoietic stem cell transplantation compared with 8 of 61 (13%) in the AML cohort (P = .07). The median survival of patients with pDC-AML was 18.8 months, which was shorter than for the AML cohort (36.7 months; P < .08) (supplemental Figure 2B). When using 2% as a cutoff, increased pDCs at diagnosis were associated with inferior outcome in AML by univariate analysis (hazard ratio, 2.1; 95% confidence interval [CI]: 1.2-3.7; P = .01), but its prognostic value was not significant after controlling for age and European LeukemiaNet risk (hazard ratio, 0.70; 95% CI, 0.4-1.4; P = .3). We also compared patients with pDC-AML with pDC expansion at diagnosis with those who only manifested expanded pDCs after diagnosis. The former had a higher median age (72 vs 59 years; P < .05), higher platelet counts (median 78 \times 10⁹/L vs 44 \times 10⁹/L; P < .05), and fewer circulating blasts (0% vs 28%; P < .05) (Table 1). Patients with pDC-AML with pDC expansion at diagnosis had shorter median overall survival (15.9 vs 30.8 months compared with those with expanded pDCs only after diagnosis; P < .05) (supplemental Figure 2C).

pDC-AML is characterized by frequent *RUNX1* mutations

Targeted sequencing analysis revealed that 32 of 41 (78%) patients with pDC-AML had RUNX1 aberrations, including

Table 1. Clinicopathologic features of pDC-AML

	pDC-AML at diagnosis (n = 26)	pDC-AML late (n = 16)	pDC-AML (combined) (n = 42)	AML without pDC expansion (n = 100)	
Age, median (IQR), y	72 (65-77)*	59 (35-70)	68 (58-76)	62 (50-70)	
Sex, male/female, n	19/7	12/4	31/11	56/44	
Prior therapy, n	7 (5 HMA, 1 lenalidomide, 1 HMA + lenalidomide)	2 (2 HMA)	9 (7 HMA, 1 lenalidomide, 1 HMA + lenalidomide)	22 (20 HMAs, 1 HMA + ruxolitinib, 1 anti-CD33)	
Prior allo-HSCT, n	0	0	0	3	
History of MDS/CMML, n	9/1	4/0	13/1	21/7	
CBC at diagnosis WBC, ×10°/L, median (IQR) ANC, ×10°/L, median (IQR) Absolute monocytes, ×10°/ L, median (IQR) Hb, g/dL, median (IQR) PLT, ×10°/L, median (IQR) Blasts, median (IQR), %	3.5 (1.2-9.1) 0.55 (0.20-2.8) 0.20 (0.0-1.4) 8.5 (7.5-9.8) 78 (39-105)* 0 (0-19)*	6.5 (1.3-26.5) 0.60 (0.18-4.5) 0.10 (0.0-2.3) 8.5 (7.1-11.2) 44 (27-64) 28 (0-63)	4.1 (1.2-11.5) 0.55 (0.20-3.0) 0.15 (0.0-1.6) 8.5 (7.5-12.6) 63 (37-97) 2.5 (0-30)†	3.5 (1.8-8.9) 0.6 (0.2-1.9) 0.10 (0-0.75) 8.9 (7.6-10.4) 53 (27-114) 20 (5-47)	
BM blasts, median (IQR), %	37 (27-66)	42 (22-80)	37 (24-69)	42 (30-65)	
BM cellularity, median (IQR), %	70 (30-80)	60 (30-100)	65 (30-80)†	80 (60-90)	
Extramedullary disease, n (%)/sites	6 (23)/(5 skin, 1 lymph node)	1 (6)/(skin and lymph node)	7 (17)/(6 skin, 2 lymph node)	8 (8)/(3 skin, 1 lymph node, 1 lung, 1 breast, 1 ovary, 1 soft tissue)	
≥2 Cross lineage markers, n (%)	6/25 (24)	7/16 (44)	13/41 (32)‡	3 (3)	
Blast CD123 MFI, median (IQR)	2538 (1146-3563)*	1108 (765-1682)	1773 (1006-3077)‡	1008 (673-1936)	
ECOG status documented, n 0 1 2 3	19 12 5 0 2	10 5 4 1 0	29 17 9 1 2	61 35 24 2 0	
AML WHO classification, n De novo AML Therapy-related AML AML-MRC	12 2 12	8 2 6	20 4 18	55 17 28	
CG risk stratification, n Favorable Intermediate Normal karyotype Unfavorable Complex karyotype Unavailable	0 20 7 5 2 1	0 11 8 4 0 1	0 31 15 9 2 2	8 67 41 21 12 4	
ELN risk stratification, n Favorable Intermediate Adverse Unavailable	0 3 23 0	0 4 7 5	0 7 30‡ 5	23 41 35 1	

allo-HSCT, allogeneic hematopoietic stem cell transplantation; AML-MRC, AML with myelodysplasia-related changes; ANC, absolute neutrophil count; CBC, complete blood count; CG, cytogenetic; CR, complete remission; CRi, incomplete remission; ECOG, Eastern Cooperative Oncology Group; ELN, European LeukemiaNet; Hb, hemoglobin; HMA, hypomethylating agent; HSCT, hematopoietic stem cell; MDS, myelodysplastic syndrome; MLFS, morphologic leukemia-free state; PLT, platelets; PR, partial response; WBC, white blood cell; WHO, World Health Organization.

*P < .05, Student t test, pDC-AML at diagnosis vs pDC-AML late.

 $\dagger P < .05$, Student t test, pDC-AML (combined) vs AML without pDC expansion.

 $\ddagger P <$.0001, Fisher's exact test, pDC-AML (combined) vs AML without pDC expansion.

Table 1. (continued)

	pDC-AML at diagnosis (n = 26)	pDC-AML late (n = 16)	pDC-AML (combined) (n = 42)	AML without pDC expansion (n = 100)	
Response, n (%) CR CRi MLFS PR	6 (23) 1 (4) 1 (4) 2 (8)	7 (44) 2 (13) 1 (6) 2 (13)	13 (31) 3 (7) 2 (5) 4 (10)	62 (62) 9 (9) 5 (5) 5 (5)	
Induction therapy, n	9	12	21	100	
Consolidation therapy, n	6	6	12	39	
HSCT, n (%) Relapse post-HSCT	5 (19) 2 (40)	11 (69) 4 (36)	16 (38) 6 (38)	61 (61) 8 (13)	
Median overall survival from diagnosis, mo	15.9*	30.8	18.8	36.7	

allo-HSCT, allogeneic hematopoietic stem cell transplantation; AML-MRC, AML with myelodysplasia-related changes; ANC, absolute neutrophil count; CBC, complete blood count; CG, cytogenetic; CR, complete remission; CRi, incomplete remission; ECOG, Eastern Cooperative Oncology Group; ELN, European LeukemiaNet; Hb, hemoglobin; HMA, hypomethylating agent; HSCT, hematopoietic stem cell; MDS, myelodysplastic syndrome; MLFS, morphologic leukemia-free state; PLT, platelets; PR, partial response; WBC, white blood cell; WHO, World Health Orcanization.

*P < .05, Student t test, pDC-AML at diagnosis vs pDC-AML late.

P < .05, Student t test, pDC-AML (combined) vs AML without pDC expansion.

 $\pm P < .0001$, Fisher's exact test, pDC-AML (combined) vs AML without pDC expansion.

29 patients with RUNX1 mutations (34 total mutations; Figure 2A). These mutations were split between missense (18) and nonsense/frame shift (16), with the majority within the RUNT DNA binding domain (Figure 2B). Two patients had atypical RUNX1 translocations: t(8;21) with PLAG1 and t(15;21) with SPATA5L1 as the fusion partners; both fusions retained the RHD and TAD domains of RUNX1 at the 3' end of the fusion proteins, and 1 patient had a deletion of the region including RUNX1. By comparison, 14 of 100 (14%) of the patients with AML without pDC expansion and 0 of 7 patients with BPDCN had RUNX1 mutations (P < .0001; Table 1 and data not shown). Twelve of 18 (67%) patients with pDC-AML with monocytic differentiation had RUNX1 mutations. Ten of 29 patients with RUNX1 mutations had germline controls (nail DNA). Among the 19 patients with RUNX1 mutations who did not have germline controls, 14 had variant allele frequencies (VAFs) of RUNX1 mutations <30%, and 5 had VAFs >40%.

Other recurrent mutations observed in patients with pDC-AML included SRSF2 (n = 13; 32%), ASXL1 (n = 10; 24%), TET2 (n = 9; 22%), DNMT3A (n = 7; 17%), NRAS (n = 7; 17%), PHF6 (n = 6; 15%), IDH1 (n = 4; 10%), SF3B1 (n = 4; 10%), FLT3 (n = 4; 10%), and TP53 (n = 4; 10%). Sixteen of 29 (55%) patients with RUNX1 mutations had \geq 1 mutation commonly observed in clonal hematopoiesis (TET2, ASXL1, and DNMT3A). Six patients had TET2 and SRSF2 concurrent mutations, including the 1 patient with a history of CMML. Based on the VAFs and the blast percentage, RUNX1 mutations were present in the dominant clones in 26 of 29 (90%) patients (Figure 2C; supplemental Table 5). Two of 4 patients with pDC-AML with TP53 mutations had concurrent RUNX1 mutations and the VAF of TP53 mutations in the RUNX1-mutant cases was <5%, suggesting a minor TP53-mutant subclone and a dominant RUNX1-mutant clone.

pDCs share identical mutations and cytogenetic abnormalities with leukemic blasts

We next assessed whether pDCs in pDC-AML were clonally derived from the leukemic clone. We used FACS to purify leukemic blasts and pDCs from 9 patients for paired genomic analysis. Monocytes and T cells were also sorted from a subset of patients. Targeted sequencing and fluorescent in situ hybridization studies were performed on sorted cells from 7 and 2 patients, respectively. As shown in Table 2, the sorted pDCs shared identical mutations and/or cytogenetic abnormalities with the leukemic blasts in all 9 patients. Moreover, the VAFs of these mutations were comparable between pDCs and blasts. Monocytes sorted from 3 patients also had the same mutations and abnormalities, whereas sorted T cells from 5 patients were negative for all mutations and abnormalities. Collectively, these results indicate that pDCs are neoplastic in origin and are clonally related to the leukemic blasts.

Leukemic blasts from pDC-AML show increased pDC differentiation potential

Our genetic data suggest the possibility that pDCs can differentiate from leukemic clones. To address this, we established an in vitro culture system that induces pDC differentiation from hematopoietic stem/progenitor cells (HSPCs). We cultured cells in FLT3L (100 ng/mL), stem cell factor (10 ng/mL), and thrombopoietin (50 ng/mL)–containing serum-free media (supplemental Figure 3) and assessed pDC differentiation potential with these basal conditions and with the addition of other cytokines/ factors. Under these conditions, we were able to efficiently differentiate pDCs from cord blood. The pDCs differentiated from cord blood cells showed characteristic surface expression of CD123, HLA-DR, and CD303 (supplemental Figure 3E-F) and plasmacytoid cytomorphology (data not shown). StemRegenin 1, a small molecule that activates the aryl hydrocarbon receptor, has previously been shown to increase hematopoietic stem cell



Figure 2. RUNX1 mutations in pDC-AML. (A) Oncoplot of mutations in pDC-AML. (B) Lollipop graph of RUNX1 mutations in pDC-AML. (C) VAF of the major mutations compared with blast percentages. AML, NOS, AML not otherwise specified; AML-MRC, AML with myelodysplasia-related changes; t-AML: therapy-related AML.

self-renewal,⁴³ platelet production,⁴⁴ and pDC differentiation.^{16,45,46} Consistent with this, pDC output from cord blood CD34⁺ cells was increased by nearly 10-fold by adding StemRegenin 1 to basal culture conditions (supplemental Figure 3A,C). pDC output from granulocyte colony-stimulating factor-mobilized peripheral blood CD34⁺ HSPCs was 10-fold lower than that from cord blood HSPCs (supplemental Figure 3B,D).

We then FACS-purified leukemic blasts from pDC-AML (n = 5), as well as AML without pDC expansion, but with or without RUNX1 mutations (n = 5 and n = 10, respectively) (Figure 3A-C). We used CD34⁺ cord blood cells as controls (n = 10). After 2 weeks of in vitro culture, CD34⁺ cord blood cells generated a mean of 10.8% pDCs (standard error of the mean [SEM], 1.6%; 95% CI, 7.2-14.5). The leukemic blasts from patients with pDC-AML had 2.5-fold higher pDC output than did cord blood-derived cells, with a mean of 26.8% pDCs (SEM, 4.6%; 95% CI, 14.1-39.6; P < .001 vs cord blood) (Figure 3A,D). In contrast, leukemic blasts from patients with AML without RUNX1 mutations and without pDC expansion in vivo failed to produce pDCs in vitro (Figure 3C,D). Interestingly, leukemic blasts from patients with AML with RUNX1 mutations who did not show pDC expansion in their clinical isolates were able to differentiate into pDCs, with a mean of 12.7% pDCs (SEM, 4.6%; 95% CI, 0.4-25.1; P < .04 vs pDC-AML, P < .001 vs AML without RUNX1 mutations), equivalent to pDC differentiation/expansion

potential from CD34⁺ cord blood cells (Figure 3B,D). Leukemic blasts from pDC-AML had higher pDC output at 2 weeks (mean, 23.7%; SEM, 7.6%) than did CD34⁺ cord blood cells (mean, 7.7%; SEM, 5.6%; P < .09), and they continued to generate increased levels of pDCs for 4 weeks (mean, 32.3% at 3 weeks and 30.5% at 4 weeks, P < .018 and 0.018, respectively, vs cord blood mean, mean, 3.7% at 3 weeks and 2.1% at 4 weeks) (Figure 3E). pDCs generated in vitro had a typical pDC immunophenotype and cytomorphology (Figure 3F). In total, these experiments provide in vitro evidence that suggests that leukemic blasts from pDC-AML have an intrinsic propensity to differentiate into pDCs.

Leukemic blasts recapitulate the phenotype of pDC-AML in vivo

We next established a patient-derived xenograft (PDX) model to further characterize pDC-AML. Peripheral blood mononuclear cells isolated from 6 patients with pDC-AML were introduced into sublethally irradiated NSG mice by tail vein injection, and human CD45⁺ cells were detected at >0.1% in the peripheral blood of all recipients at 3 and 6 months (supplemental Table 6).⁴⁷ However, only 1 pDC-AML clinical isolate reproducibly generated overt AML in recipient mice. This PDX showed a nearly identical immunophenotype to the original leukemia, with high levels of CD34⁺ leukemic blasts and increased pDCs (supplemental Figure 4). We then harvested BM

	Shared variants and cytogenetic abnormalities	VAF, %			
Patient		Blasts	pDCs	Monocytes	T cells
1	RUNX1 D198G NRAS G12D WT1 R380Tfs*5 FLT3 I836del FLT3D835Y	48.9 27.9 13.9 10.1 2.2	48.5 35.3 8.2 5.6 0.2	NA	NA
2	SRSF2 P95H ASXL1 E635Rfs*15 KITD816V	39.9 19.0 4.0	52.8 22.2 2.0	NA	NA
3	RUNX1 A435Gfs*165	37.6	44.2	NA	NA
4	RUNX1 L98fs*24 DNMT3A G543C DNMT3AI715Gfs*69	45.2 52.5 19.9	44.7 47.5 20.0	NA	0 0 0
5	RUNX1 D62Rfs*11 TET2 Q743* TET2 R1465* SRSF2 P95L CBL L380P NRAS G12V KRAS G12R Trisomy 8	3.2 82.6 14.9 37.8 27.3 7.8 1.5 22.7	2.0 96.4 3.5 49.3 19.3 1.4 0.1 21	3.9 50.4 50.1 27.3 21.3 17.0 2.7 85	NA 0
6	JARID2 E50Afs*8	86.7	88.3	NA	NA
7	RUNX1 V164_G165reSWGCP PHF6 C283F SRSF2 P96L TET2 Q964* ASXL1 P701fs*2 NF1 S2309T NF1 P2310N	34 97 50 47 46 30 30	35 98 53 52 50 29 29	NA	0 0 0 0 0 0
8	CRLF2 fusion	27.5	25	19.5	0
9	Deletion of 7q31	98	98	99	0

Table 2. Shared genetic abnormalities between sorted blasts and pDCs from pDC-AML

NA, sorted population not available for analysis.

cells from primary recipients and transplanted them into secondary NSG recipients in 3 groups: whole BM, FACS-purified leukemic blasts, and FACS-purified pDCs. Unsorted BM cells (1 million cells per mouse) faithfully recapitulated pDC-AML in secondary recipient mice (Figure 4). Sorted leukemic blasts (0.5 million per mouse) also gave rise to AML with increased pDCs (Figure 4A-D,F). In contrast, sorted pDCs (0.5 million per mouse) displayed nearly undetectable engraftment in secondary NSG recipient mice up to 8 weeks posttransplant (Figure 4A-B,E). Collectively, these data provide in vitro and in vivo evidence indicating that pDCs originate from leukemic blasts but, alone, cannot initiate leukemia in vivo.

Leukemic blasts from pDC-AML upregulate a pDC transcriptional program

Transcriptional profiling was performed to elucidate the generegulatory programs underlying the ability of leukemic blasts from pDC-AML to differentiate into pDCs. CD34⁺ blasts and pDCs were sorted from patients with pDC-AML and from normal controls. Based on our identification of RUNX1 mutations as highly enriched in pDC-AML, we hypothesized that this genetic event may confer a pDC transcriptional program to leukemic blasts. To address this possibility, we sorted CD34⁺ blasts from patients with AML (no pDC expansion), with and without mutations in RUNX1. Principal component analysis revealed that pDCs from pDC-AML clustered independently from normal pDCs and from leukemic blasts, consistent with their neoplastic origin and their retention of a pDC transcriptional program (Figure 5A). To determine whether blasts from pDC-AML or AML with RUNX1 mutations upregulate a pDC transcriptional signature, we calculated a normalized single sample GSEA score based on expression of previously established pDC genes (Figure 5B; supplemental Figure 5A).48,49 This analysis showed that blasts from pDC-AML significantly upregulated a pDC



Figure 3. Leukemic blasts from pDC-AML and AML with *RUNX1* mutations have greater differentiation propensity to pDCs in vitro. (A-C) The sorted leukemic blasts were cultured in vitro for 2 weeks and subjected to immunophenotyping by flow cytometry. (D) pDC proportions in the culture after 2 weeks were compared. Data are mean ± SD. (E) Time course experiments showed that a high pDC proportion persisted from the leukemic blasts of pDC-AML. Data are mean ± SEM. (F) Wright-Giernsa staining of sorted pDCs (were CD303⁺ and had bright CD123 and HLA-DR expression and were negative for CD11b, CD14, CD19, CD3, and CD56) from the in vitro culture. AML *RUNX1*, AML with *RUNX1* mutations or pDC expansion. **P* < .05, ****P* < .001. Original magnification, ×1000 (F).

transcriptional program compared with blasts from AML without RUNX1 mutations (no pDCs). As an orthogonal approach, we identified upregulated genes, with respect to normal marrow CD34⁺ cells, shared between different groups of leukemic blasts and normal pDCs, and confirmed our observations (supplemental Figure 5B). Blasts from AML with RUNX1 mutation (no pDCs) had greater enrichment of a pDC transcriptional program than did blasts from AML without RUNX1 mutation (no pDCs), but this difference was not statistically significant (supplemental Figure 5B). To identify the specific transcriptional program of blasts from pDC-AML and AML with RUNX1 mutations, we focused on those upregulated genes shared between these groups and shared with normal pDCs (Figure 5C-D). Gene ontology analysis of this gene subset or of genes upregulated in pDC-AML blasts identified inflammatory pathways and interferon signaling as highly enriched processes, an intriguing finding because these processes contribute to pDC development (Figure 5E; supplemental Figure 5C). Consistent with these findings, we observed that several genes that are involved in the IFN response were highly upregulated in normal pDCs and neoplastic pDCs, moderately upregulated in blasts from

pDC-AML and AML with *RUNX1* mutations, but not upregulated in blasts from AML without *RUNX1* mutations (Figure 5F). In total, this suggests that leukemic blasts from pDC-AML and AML with *RUNX1* mutations may gain an IFN-driven pDC transcriptional program, potentially contributing to the increased pDC output in these settings.

Anti-CD123-targeted therapy reduces pDCs and leukemic blasts in vivo

Tagraxofusp, a fusion protein consisting of interleukin-3 fused to a truncated diphtheria toxin payload with high affinity and avidity for the interleukin-3 receptor- α (CD123), has shown substantive therapeutic efficacy in BPDCN and, therefore, is approved by the US Food and Drug Administration for its treatment.^{50,51} Because pDCs are increased in patients with pDC-AML, and the leukemic blasts also expressed higher levels of CD123 compared with AML with no pDC expansion (median of MFI, 1773 vs 1008; P = .0003) (Figure 1D), we hypothesized that CD123-targeted therapy may have efficacy in pDC-AML. To evaluate tagraxofusp in pDC-AML, unsorted BM cells harvested from primary PDX NSG mice (supplemental Figure 4; supplemental Table 6) were



Figure 4. Leukemic blasts differentiate into pDCs in vivo. (A-B) BM cells were harvested from the primary NSG mice receiving pDC-AML leukemic cells, as described in supplemental Figure 4 and supplemental Table 6. The unsorted BM cells, purified leukemic blasts, and purified pDCs were injected IV into secondary NSG mice. Engraftment of human CD45⁺ cells was evaluated from peripheral blood 4 weeks (A) and 8 weeks (B) after transplant. (C) pDC-AML phenotype in the secondary NSG mice receiving purified leukemic blasts. Data are mean \pm SD. (D-E) Representative flow plots showing hCD45⁺ cells in the secondary NSG mice receiving purified blasts (D) or pDCs (E). (F) Representative flow plot showing pDC-AML phenotype in hCD45⁺ cells from panel D. ****P* < .001.

propagated into secondary NSG recipients. After 8 weeks, mice were randomized based on comparable engraftment of human (h)CD45 cells (mean value of hCD45⁺ cells: 8.4% in phosphatebuffered saline (PBS) cohort, 9.2% in tagraxofusp 0.1 mg/kg per day cohort, and 10.0% in tagraxofusp 0.2 mg/kg per day cohort, supplemental Figure 6A). After 1 cycle of intraperitoneal injection of tagraxofusp (5 days), the cohorts taking 0.1 mg/kg per day or 0.2 mg/kg per day showed >50% reduction in hCD45⁺ cells (mean: PBS, 62.5%; tagraxofusp 0.1 mg/kg per day, 33.1%; tagraxofusp 0.2 mg/kg per day, 26.8%; P < .001 PBS vs 0.1 mg/kg per day, P < .001 PBS vs 0.2 mg/kg per day) (Figure 6A). Although the proportion of CD34+ blasts in the hCD45⁺ population was not significantly affected by the treatment (Figure 6B,D-F), the leukemic burden overall (as evaluated by the proportion of CD34⁺ blasts in total white blood cells) was twofold to threefold lower in tagraxofusp-treated mice (mean: PBS, 30.7%; tagraxofusp 0.1 mg/kg per day, 14.7%; tagraxofusp 0.2 mg/kg per day, 11.7%; P < .0017 PBS vs 0.1 mg/kg per day, P < .0015 PBS vs 0.2 mg/kg per day) (Figure 6C). Consistent with the high efficacy of tagraxofusp in BPDCN, pDCs were almost completely eliminated by tagraxofusp in the cohorts taking 0.1 mg/kg per day or 0.2 mg/kg per day (mean: PBS, 4.8%; 0.1 mg/kg per day, 1.3%; 0.2 mg/kg per day, 1.4%; P < .0001 PBS vs 0.1 mg/kg per day or PBS vs 0.2 mg/kg per day) (Figure 6B, D-G; supplemental Figure 6B). More importantly, there was a significant reduction in leukemic burden in BM with tagraxofusp treatment (total BM cellularity: 17 million cells in PBS vs 2.9 million cells in tagraxofusp 0.1 mg/kg per day; P < .001; Figure 6G). We did not observe a significant difference in overall survival, this is likely attributed to treatment-associated toxicities that limit the duration of therapy in NSG mice (data not shown). Therefore, tagraxofusp is able to effectively eliminate neoplastic pDCs and reduce leukemic burden in an in vivo model of pDC-AML.

Discussion

Our studies show that a subset of patients with AML, most commonly with *RUNX1* mutations, have pDC expansion in vivo, pDC differentiation capacity in vitro, and a distinct transcriptional program, suggesting that this represents an AML subtype with distinct genetic, transcriptional, and biologic features. Moreover, our studies suggest that CD123 targeting may have therapeutic importance in AML with pDC expansion, alone or in combination with other antileukemic therapies.^{52,53}



Figure 5. Leukemic blasts from pDC-AML upregulate a pDC transcriptional program. (A) Principal component analysis of gene expression in normal marrow CD34⁺ cells, normal pDCs, pDCs from pDC-AML, blasts from pDC-AML, blasts from AML without *RUNX1* mutations or pDC expansion, and blasts from AML with *RUNX1* mutations but no pDC expansion. (B) pDC transcriptional program, as evaluated by normalized single sample GSEA scores for each group. Scores are calculated based on expression levels of pDC genes. (C) Upset plot displaying overlap of upregulated genes among blasts from pDC-AML, blasts from AML without *RUNX1* mutations or pDC expansion, and blasts from AML with a blasts from AML with *RUNX1* mutations or pDC expansion, and blasts from AML with *RUNX1* mutations or pDC expansion, and blasts from AML with *RUNX1* mutations or pDC expansion, and blasts from AML with *RUNX1* mutations or pDC expansion, and blasts from AML with *RUNX1* mutations or pDC expansion, and blasts from AML with *RUNX1* mutations or pDC expansion, pDCs from pDC-AML, and normal pDCs. All groups are compared with normal marrow CD34⁺ cells. (D) Heat may showing the 30 upregulated genes shared among blasts from pDC-AML, blasts from AML with *RUNX1* mutations or pDC expansion. (E) Gene ontology analysis showing upregulated pathways in blasts from pDC-AML compared with normal marrow CD34⁺ cells. (F) Expression levels of a subset of IFN-related genes upregulated in pDC-AML. These genes are also among the subset of genes depicted in panel D. ***P* < .01.



Figure 6. Tagraxofusp treatment results in significant reduction of leukemic blasts and pDCs in vivo. (A-C) BM cells harvested from primary NSG mice were injected IV into secondary NSG mice. The mice were treated by intraperitoneal injection for 1 cycle with PBS, tagraxofusp 0.1 mg/kg per day, or 0.2 mg/kg per day. (A) One week after the last dose, hCD45⁺ cells were examined in peripheral blood. (B) Leukemic blasts and pDC proportions are shown in hCD45⁺ compartments. (C) CD34⁺ leukemic blast proportions in total white blood cells (WBC) were also evaluated. Data are mean \pm SD. (D-F) Representative flow plots show leukemic blasts and pDCs after treatment with PBS (D), tagraxofusp 0.1 mg/kg per day (E), or tagraxofusp 0.2 mg/kg per day (F). (G) BM cells were harvested from mice treated with PBS or tagraxofusp 0.1 mg/kg per day. CD34⁺ leukemic blasts and pDCs were enumerated. **P < .01.

The presence of pDCs in patients with AML raises important diagnostic challenges. Very few patients in our cohort had a history of CMML, and the mutational profiles are largely different from that of CMML with or without pDC expansion,^{26,54} arguing against a progression from CMML in the majority of patients. On the other hand, a minority of patients had >20% pDCs in the marrow with immature pDC morphology, raising the differential

diagnosis of BPDCN. However, they all had *RUNX1* mutations, and their pDCs expressed CD34, but not CD56, which is distinct from BPDCN. There are several lines of evidence supporting pDC-AML as an entity distinct from BPDCN.⁵⁵ First, skin presentation is relatively infrequent (only 14%) in pDC-AML but is typical (70-100%) in BPDCN.⁴² Second, myeloid blasts are >20% in all patients with pDC-AML. Third, *RUNX1* mutations are present in >70% of pDC-AML but are rare (<5%) in BPDCN.^{56,57} Fourth, pDCs from pDC-AML alone are not leukemogenic, in contrast to studies showing that pDCs from BPDCN can maintain/propagate the diseases.⁵³ These data suggest that pDC-AML is a distinct entity from BPDCN or CMML with pDC expansion and that important shared features can confound diagnostic evaluation and may inform shared insights into disease pathogenesis and therapy. Although the immunophenotype of CD4, CD123, and CD56 coexpression is used to define neoplastic pDCs, it is of paramount importance to note that this immunophenotype is also commonly seen in myeloid blasts with monocytic differentiation, but in the setting of \geq 1-log lower expression level of CD123 than pDCs. In this regard, the reports of BPDCN-like AML with NPM-1 mutation (which is exceedingly rare in pDC-AML and BPDCN) may merely represent AML with monocytic differentiation,⁵⁸ although rare cases of pDC-AML or BPDCN cannot be entirely ruled out based on the reports.

Despite the recognition of BPDCN as a distinct entity,^{30,42} the role of pDCs in myeloid transformation remains poorly understood. A recent study demonstrated that CMML patients with pDC expansion have a higher transformation risk to AML and that pDCs share genetic mutations with the CMML clone.²⁶ However, the role of pDCs in leukemic transformation and maintenance, as well as in the interaction with malignant cells in different cancer contexts, has not been extensively studied. Our studies suggest that pDCs in AML are derived from the leukemic clone and that there may be interactions between leukemic cells and pDC progeny. Studies have shown that direct contact with E-cadherin and PD-L1-expressing myeloma cells could convert pDCs into tumor-promoting cells by suppressing pDC IFN-a production.^{59,60} pDCs can also induce regulatory T cells,⁶¹ thereby inhibiting the antitumor response mediated by other effector T cells.⁶²⁻⁶⁴ Moreover, leukemia-derived pDCs may have aberrant functionalities, including with respect to cytokine signaling and secretion, despite the largely preserved type I IFN program that we and other investigators have observed.26 Second, pDCs may be a surrogate marker for disease aggressiveness. Because pDCs are markedly depleted in the majority of cases of AML and other high-grade myeloid neoplasms,^{25,65} increased pDC differentiation may implicate an early hematopoietic stem cell or multipotent progenitor as the cell of origin of the myeloid neoplasms, which may correlate with poor outcome.66,67 Consistent with this hypothesis, crosslineage marker expression is much more common in pDC-AML than in AML as a whole. Alternatively, pDC differentiation may be indicative of the presence of adverse risk somatic alterations, such as RUNX1 mutations, which may or may not relate to the cell of origin or the differentiation stage of leukemic blasts.

Our data indicate that leukemic blasts from pDC-AML upregulate an IFN-driven pDC transcriptional program, potentially contributing to the increased pDC output. The enrichment of *RUNX1* mutations in pDC-AML implies a crucial role for this genetic event and downstream altered gene regulatory networks in pDC differentiation. Prior studies have shown that pDC-related transcription factors are highly expressed in *RUNX1*-mutated AML,⁶⁸⁻⁷⁰ which is further supported by our data. Our study also provides direct evidence that the leukemic blasts from *RUNX1*-mutated AML are primed to differentiate into pDCs. *VAV-iCre*-inducible *Runx1*-deficient mice have markedly decreased Flt3⁺ lymphoid progenitors, including the entire dendritic cell compartment,⁷¹ consistent with the indispensable role of *Runx1* in lymphoid development.⁷² Future elucidation of RUNX1 function in dendritic cell development requires novel mouse models that have inducible mutant *RUNX1* alleles seen in pDC-AML under the control of pan-hematopoietic–specific vs DC-specific promoters. In addition, whether and how the BM microenvironment, including cytokines and infection responses, contributes to pDC differentiation/expansion remains to be studied.

Based on the clonal origin and possible proleukemic activities of pDCs in AML, we reasoned that targeting pDCs may have therapeutic benefit in pDC-AML. The proof-of-principle experiments demonstrated efficacy of tagraxofusp in eliminating pDCs, consistent with the clinical efficacy in patients with BPDCN.⁵⁰ Tagraxofusp has previously shown objective responses as a single therapeutic agent in a small subset of patients with AML.73 Consistently, this agent led to a marked reduction in BM cellularity, including CD34⁺ blasts. Approximately half of the mice obtained complete responses with tagraxofusp monotherapy in our preclinical model of pDC-AML, suggesting that it will be important to combine CD123-targeted therapies with other antileukemic targets to increase therapeutic efficacy in this subset of AML, as is being investigated in a broader suite of tagraxofusp-refractory myeloid neoplasms^{52,53} (also see ClinicalTrials.gov Identifier: NCT03113643).

In summary, our study identifies a subset of AML patients with pDC expansion, a high frequency of *RUNX1* mutations, unique transcriptional profile, and increased CD123 expression that may represent a therapeutic target in this high-risk AML subset. Future studies are warranted to elucidate the role of pDCs in pDC-AML initiation and maintenance and to develop effective combination therapeutic approaches for this unique AML subset with poor clinical outcome.

Acknowledgments

This work was supported by the Center for Hematologic Malignancies at MSKCC and, in part, through National Institutes of Health/National Cancer Institute Cancer Center Support grant P30 CA008748. It was also supported by National Institutes of Health/National Cancer Institute grants R01CA173636 and R35197594 (R.L.L.), and by a Leukemia and Lymphoma Society Specialized Center of Research grant. M.S. received funding from the MSKCC Clinical Scholars T32 Program under award 2T32 CA009512-31.

Authorship

Contribution: W.X., M.R., and R.L.L. conceived the study, collected and analyzed data, and wrote the manuscript; A.C. annotated the clinical and flow data; M.R.W., R.L.B., and R.P.K. performed the RNA sequencing analysis; T.M., S.F.C., I.S.C., and N.L.D. performed PDX experiments; Y.L., J.Y., and M.E.A. collected and annotated the mutational data; A. Derkach performed statistical analyses; Q.G. helped with cell sorting; J.B., S.Y., C.A.F., and M.P. coordinated the project; M.S., R.K.R., M.S.T., and A.D.G. provided clinical data; Y.Z. reviewed cytogenetic data; A. Dogan analyzed data; and all authors approved the final version of the manuscript.

Conflict-of-interest disclosure: W.X. has received research support from Stemline Therapeutics. S.F.C. is a consultant for Imago Biosciences and has received honoraria from DAVA Oncology. A.D.G. served on advisory boards or as a consultant for AbbVie, Aptose, Celgene, Daiichi Sanyko, and Genentech; received research funding from AbbVie, ADC Therapeutics, Aprea, AROG, Daiichi Sanyko, and Pfizer; and received honoraria from Dava Oncology. R.K.R has received consulting fees from Constellation, Incyte, Celgene, Promedior, CTI, Jazz Pharmaceuticals, Blueprint, and Stemline Therapeutics and has received research funding from Incyte, Constellation, and Stemline Therapeutics. M.S.T. has received research funding from AbbVie, Cellerant, Orsenix, ADC Therapeutics, Biosight, Glycomimetics, Rafael Pharmaceuticals and Amgen; has served on advisory Boards for AbbVie, BioLineRx, Daiichi-Sankyo, Orsenix, KAHR, Rigel, Nohla, Delta Fly Pharma, Tetraphase, Oncolyze, Jazz Pharma, Roche, Biosight, and Novartis; and has received royalties from UpToDate. A. Dogan has received personal fees from Roche, Corvus Pharmaceuticals, Physicians' Education Resource, Seattle Genetics, Peerview Institute, Oncology Specialty Group, Takeda, and EUSA Pharma and research grants from Roche. R.L.L. is on the supervisory board of QIAGEN and is a scientific advisor to Loxo (until 2019), Auron, Ajax, Mission Bio, Imago, C4 Therapeutics, and Isoplexis, which each include an equity interest; received research support from and consulted for Celgene and Roche; received research support from Prelude Therapeutics; consulted for Incyte, Novartis, and Janssen; and received honoraria from Lilly and Amgen for invited lectures and from Gilead for grant reviews. The remaining authors declare no competing financial interests.

ORCID profiles: W.X., 0000-0001-8586-8500; A.C., 0000-0001-5884-8424; Y.L., 0000-0002-9167-3205; S.F.C., 0000-0002-2708-887X; A. Derkach, 0000-0003-2178-8493; S.Y., 0000-0003-1518-7210; A. Dogan, 0000-0001-6576-5256; A.D.G., 0000-0002-2892-2643.

REFERENCES

- Grouard G, Rissoan M-C, Filgueira L, Durand I, Banchereau J, Liu Y-J. The enigmatic plasmacytoid T cells develop into dendritic cells with interleukin (IL)-3 and CD40-ligand. J Exp Med. 1997;185(6):1101-1111.
- Cella M, Jarrossay D, Facchetti F, et al. Plasmacytoid monocytes migrate to inflamed lymph nodes and produce large amounts of type l interferon. Nat Med. 1999;5(8):919-923.
- Dzionek A, Fuchs A, Schmidt P, et al. BDCA-2, BDCA-3, and BDCA-4: three markers for distinct subsets of dendritic cells in human peripheral blood. *J Immunol.* 2000;165(11): 6037-6046.
- Sathe P, Vremec D, Wu L, Corcoran L, Shortman K. Convergent differentiation: myeloid and lymphoid pathways to murine plasmacytoid dendritic cells. *Blood*. 2013; 121(1):11-19.
- Shigematsu H, Reizis B, Iwasaki H, et al. Plasmacytoid dendritic cells activate lymphoid-specific genetic programs irrespective of their cellular origin. *Immunity*. 2004;21(1):43-53.
- Reizis B. Plasmacytoid dendritic cells: development, regulation, and function. *Immunity*. 2019;50(1):37-50.
- Cisse B, Caton ML, Lehner M, et al. Transcription factor E2-2 is an essential and specific regulator of plasmacytoid dendritic cell development. *Cell.* 2008;135(1):37-48.
- Nagasawa M, Schmidlin H, Hazekamp MG, Schotte R, Blom B. Development of human plasmacytoid dendritic cells depends on the combined action of the basic helix-loop-helix factor E2-2 and the Ets factor Spi-B. Eur J Immunol. 2008;38(9):2389-2400.
- Ghosh HS, Ceribelli M, Matos I, et al. ETO family protein Mtg16 regulates the balance of dendritic cell subsets by repressing Id2. J Exp Med. 2014;211(8):1623-1635.

- Wu X, Satpathy AT, Kc W, Liu P, Murphy TL, Murphy KM. Bcl11a controls Flt3 expression in early hematopoietic progenitors and is required for pDC development in vivo. PLoS One. 2013;8(5):e64800.
- Honda K, Yanai H, Negishi H, et al. IRF-7 is the master regulator of type-l interferondependent immune responses. *Nature*. 2005; 434(7034):772-777.
- Bao M, Wang Y, Liu Y, et al. NFATC3 promotes IRF7 transcriptional activity in plasmacytoid dendritic cells. J Exp Med. 2016; 213(11):2383-2398.
- Sasaki I, Hoshino K, Sugiyama T, et al. Spi-B is critical for plasmacytoid dendritic cell function and development. *Blood*. 2012;120(24): 4733-4743.
- Chopin M, Preston SP, Lun ATL, et al. RUNX2 mediates plasmacytoid dendritic cell egress from the bone marrow and controls viral immunity. *Cell Rep.* 2016;15(4):866-878.
- Ma S, Wan X, Deng Z, et al. Epigenetic regulator CXXC5 recruits DNA demethylase Tet2 to regulate TLR7/9-elicited IFN response in pDCs. J Exp Med. 2017;214(5):1471-1491.
- Díaz-Rodríguez Y, Cordeiro P, Belounis A, Herblot S, Duval M. In vitro differentiated plasmacytoid dendritic cells as a tool to induce anti-leukemia activity of natural killer cells. Cancer Immunol Immunother. 2017; 66(10):1307-1320.
- Cordeau M, Belounis A, Lelaidier M, et al. Efficient killing of high risk neuroblastoma using natural killer cells activated by plasmacytoid dendritic cells. *PLoS One*. 2016;11(10): e0164401.
- Tel J, Smits EL, Anguille S, Joshi RN, Figdor CG, de Vries IJ. Human plasmacytoid dendritic cells are equipped with antigenpresenting and tumoricidal capacities. *Blood*. 2012;120(19):3936-3944.
- 19. Stary G, Bangert C, Tauber M, Strohal R, Kopp T, Stingl G. Tumoricidal activity of TLR7/8-

Correspondence: Wenbin Xiao, Memorial Sloan Kettering Cancer Center, 1275 York Ave, New York, NY 10065; e-mail: xiaow@mskcc.org; or Ross L. Levine, Memorial Sloan Kettering Cancer Center, 1275 York Ave, New York, NY 10065; e-mail: leviner@mskcc.org.

Footnotes

Submitted 26 June 2020; accepted 18 August 2020; prepublished online on *Blood* First Edition 1 September 2020. DOI 10.1182/blood. 2020007897.

Presented in abstract form at the 61st annual meeting of the American Society of Hematology, San Diego, CA, 2 December 2018.

Data sharing requests should be sent to Wenbin Xiao (xiaow@mskcc.org) or Ross L. Levine (leviner@mskcc.org).

The online version of this article contains a data supplement.

There is a Blood Commentary on this article in this issue.

The publication costs of this article were defrayed in part by page charge payment. Therefore, and solely to indicate this fact, this article is hereby marked "advertisement" in accordance with 18 USC section 1734.

activated inflammatory dendritic cells. J Exp Med. 2007;204(6):1441-1451.

- Drobits B, Holcmann M, Amberg N, et al. Imiquimod clears tumors in mice independent of adaptive immunity by converting pDCs into tumor-killing effector cells. J Clin Invest. 2012; 122(2):575-585.
- Chauhan D, Singh AV, Brahmandam M, et al. Functional interaction of plasmacytoid dendritic cells with multiple myeloma cells: a therapeutic target. *Cancer Cell.* 2009;16(4): 309-323.
- 22. Sawant A, Hensel JA, Chanda D, et al. Depletion of plasmacytoid dendritic cells inhibits tumor growth and prevents bone metastasis of breast cancer cells. *J Immunol.* 2012;189(9):4258-4265.
- Sisirak V, Faget J, Gobert M, et al. Impaired IFN-α production by plasmacytoid dendritic cells favors regulatory T-cell expansion that may contribute to breast cancer progression. *Cancer Res.* 2012;72(20):5188-5197.
- Beird HC, Khan M, Wang F, et al. Features of non-activation dendritic state and immune deficiency in blastic plasmacytoid dendritic cell neoplasm (BPDCN). *Blood Cancer J.* 2019;9:99.
- 25. Xiao W, Goldberg AD, Famulare CA, et al. Loss of plasmacytoid dendritic cell differentiation is highly predictive for post-induction measurable residual disease and inferior outcomes in acute myeloid leukemia. *Haematologica*. 2019;104(7):1378-1387.
- Lucas N, Duchmann M, Rameau P, et al. Biology and prognostic impact of clonal plasmacytoid dendritic cells in chronic myelomonocytic leukemia. *Leukemia*. 2019; 33(10):2466-2480.
- Derolf AR, Laane E, Björklund E, Saft L, Björkholm M, Porwit A. Dendritic cells in bone marrow at diagnosis and after chemotherapy in adult patients with acute myeloid leukaemia. Scand J Immunol. 2014;80(6):424-431.

- Chaperot L, Bendriss N, Manches O, et al. Identification of a leukemic counterpart of the plasmacytoid dendritic cells. *Blood*. 2001; 97(10):3210-3217.
- Jacob MC, Chaperot L, Mossuz P, et al. CD4+ CD56+ lineage negative malignancies: a new entity developed from malignant early plasmacytoid dendritic cells. *Haematologica*. 2003;88(8):941-955.
- Vermi W, Facchetti F, Rosati S, et al. Nodal and extranodal tumor-forming accumulation of plasmacytoid monocytes/interferonproducing cells associated with myeloid disorders. Am J Surg Pathol. 2004;28(5):585-595.
- Dargent JL, Delannoy A, Pieron P, Husson B, Debecker C, Petrella T. Cutaneous accumulation of plasmacytoid dendritic cells associated with acute myeloid leukemia: a rare condition distinct from blastic plasmacytoid dendritic cell neoplasm. J Cutan Pathol. 2011; 38(11):893-898.
- Song HL, Huang WY, Chen YP, Chang KC. Tumorous proliferations of plasmacytoid dendritic cells and Langerhans cells associated with acute myeloid leukaemia. *Histopathology*. 2012;61(5):974-983.
- 33. Dargent JL, Henne S, Pranger D, et al. Tumorforming plasmacytoid dendritic cells associated with myeloid neoplasms. Report of a peculiar case with histopathologic features masquerading as lupus erythematosus. J Cutan Pathol. 2016;43(3):280-286.
- Wang P, Feng Y, Deng X, et al. Tumor-forming plasmacytoid dendritic cells in acute myelocytic leukemia: a report of three cases and literature review. Int J Clin Exp Pathol. 2017; 10(7):7285-7291.
- Wang M, Chen Y-J, Wang L-R, Wang Y-Z, Lu J. Plasmacytoid dendritic cells proliferation coexisted with acute myeloid leukemia. *Chin Med J* (Engl). 2018;131(15):1866-1867.
- Hamadeh F, Awadallah A, Meyerson HJ, Beck RC. Flow cytometry identifies a spectrum of maturation in myeloid neoplasms having plasmacytoid dendritic cell differentiation. *Cytometry B Clin Cytom.* 2020;98(1):43-51.
- Huang Y, Wang Y, Chang Y, et al. Myeloid neoplasms with elevated plasmacytoid dendritic cell differentiation reflect the maturation process of dendritic cells. *Cytometry A*. 2020; 97(1):61-69.
- Rickmann M, Krauter J, Stamer K, et al. Elevated frequencies of leukemic myeloid and plasmacytoid dendritic cells in acute myeloid leukemia with the FLT3 internal tandem duplication. Ann Hematol. 2011;90(9): 1047-1058.
- Rickmann M, Macke L, Sundarasetty BS, et al. Monitoring dendritic cell and cytokine biomarkers during remission prior to relapse in patients with FLT3-ITD acute myeloid leukemia. Ann Hematol. 2013;92(8):1079-1090.
- 40. Xiao W, Goldberg AD, Famulare C, et al. Acute myeloid leukemia with plasmacytoid dendritic cell differentiation: predominantly secondary AML, enriched for RUNX1 mutations, frequent cross-lineage antigen expression and poor prognosis [abstract]. *Blood.* 2018;132(suppl 1). Abstract 2789.

- 41. Wang W, Khoury JD, Miranda RN, et al. Immunophenotypic characterization of reactive and neoplastic plasmacytoid dendritic cells permits establishment of a 10-color flow cytometric panel for initial workup and residual disease evaluation of blastic plasmacytoid dendritic cell neoplasm [published online ahead of print 2 Apr 2020]. *Haematologica.* doi:10.3324/haematol.2020. 247569.
- Arber DA, Orazi A, Hasserjian R, et al. The 2016 revision to the World Health Organization classification of myeloid neoplasms and acute leukemia [published correction appears in *Blood*. 2016;128(3):462-463]. *Blood*. 2016; 127(20):2391-2405.
- Boitano AE, Wang J, Romeo R, et al. Aryl hydrocarbon receptor antagonists promote the expansion of human hematopoietic stem cells. *Science*. 2010;329(5997):1345-1348.
- 44. Ito Y, Nakamura S, Sugimoto N, et al. Turbulence activates platelet biogenesis to enable clinical scale ex vivo production. *Cell.* 2018;174(3):636-648.e18.
- 45. Thordardottir S, Hangalapura BN, Hutten T, et al. The aryl hydrocarbon receptor antagonist StemRegenin 1 promotes human plasmacytoid and myeloid dendritic cell development from CD34+ hematopoietic progenitor cells. Stem Cells Dev. 2014;23(9): 955-967.
- 46. Laustsen A, Bak RO, Krapp C, et al. Interferon priming is essential for human CD34+ cellderived plasmacytoid dendritic cell maturation and function. *Nat Commun.* 2018;9(1): 3525.
- Wang K, Sanchez-Martin M, Wang X, et al. Patient-derived xenotransplants can recapitulate the genetic driver landscape of acute leukemias. *Leukemia*. 2017;31(1): 151-158.
- See P, Dutertre C-A, Chen J, et al. Mapping the human DC lineage through the integration of high-dimensional techniques. *Science*. 2017;356(6342):eaag3009.
- Villani AC, Satija R, Reynolds G, et al. Singlecell RNA-seq reveals new types of human blood dendritic cells, monocytes, and progenitors. *Science*. 2017;356(6335):eaah4573.
- Pemmaraju N, Lane AA, Sweet KL, et al. Tagraxofusp in blastic plasmacytoid dendriticcell neoplasm. N Engl J Med. 2019;380(17): 1628-1637.
- Frankel AE, Woo JH, Ahn C, et al. Activity of SL-401, a targeted therapy directed to interleukin-3 receptor, in blastic plasmacytoid dendritic cell neoplasm patients. *Blood.* 2014; 124(3):385-392.
- Montero J, Stephansky J, Cai T, et al. Blastic plasmacytoid dendritic cell neoplasm is dependent on BCL2 and sensitive to venetoclax. *Cancer Discov.* 2017;7(2):156-164.
- Togami K, Pastika T, Stephansky J, et al. DNA methyltransferase inhibition overcomes diphthamide pathway deficiencies underlying CD123-targeted treatment resistance. J Clin Invest. 2019;129(11):5005-5019.
- 54. Elena C, Gallì A, Such E, et al. Integrating clinical features and genetic lesions in the risk assessment of patients with chronic

myelomonocytic leukemia. *Blood*. 2016; 128(10):1408-1417.

- 55. Garnache-Ottou F, Vidal C, Biichlé S, et al. How should we diagnose and treat blastic plasmacytoid dendritic cell neoplasm patients? *Blood Adv.* 2019;3(24):4238-4251.
- Menezes J, Acquadro F, Wiseman M, et al. Exome sequencing reveals novel and recurrent mutations with clinical impact in blastic plasmacytoid dendritic cell neoplasm. *Leukemia*. 2014;28(4):823-829.
- Stenzinger A, Endris V, Pfarr N, et al. Targeted ultra-deep sequencing reveals recurrent and mutually exclusive mutations of cancer genes in blastic plasmacytoid dendritic cell neoplasm. Oncotarget. 2014;5(15):6404-6413.
- Minetto P, Guolo F, Clavio M, et al. A blastic plasmacytoid dendritic cell neoplasm-like phenotype identifies a subgroup of npm1mutated acute myeloid leukemia patients with worse prognosis. *Am J Hematol.* 2018;93(2): E33-E35.
- 59. Bi E, Li R, Bover LC, et al. E-cadherin expression on multiple myeloma cells activates tumor-promoting properties in plasmacytoid DCs. J Clin Invest. 2018;128(11):4821-4831.
- 60. Ray A, Das DS, Song Y, et al. Targeting PD1-PDL1 immune checkpoint in plasmacytoid dendritic cell interactions with T cells, natural killer cells and multiple myeloma cells. *Leukemia*. 2015;29(6):1441-1444.
- Lynch JP, Werder RB, Loh Z, et al. Plasmacytoid dendritic cells protect from viral bronchiolitis and asthma through semaphorin 4a-mediated T reg expansion. J Exp Med. 2018;215(2):537-557.
- 62. Takagi H, Fukaya T, Eizumi K, et al. Plasmacytoid dendritic cells are crucial for the initiation of inflammation and T cell immunity in vivo. *Immunity*. 2011;35(6):958-971.
- Cervantes-Barragan L, Lewis KL, Firner S, et al. Plasmacytoid dendritic cells control T-cell response to chronic viral infection. Proc Natl Acad Sci USA. 2012;109(8):3012-3017.
- Rogers GL, Shirley JL, Zolotukhin I, et al. Plasmacytoid and conventional dendritic cells cooperate in crosspriming AAV capsidspecific CD8⁺ T cells. *Blood*. 2017;129(24): 3184-3195.
- 65. Chan A, Zhang Y, Devlin SM, et al. Plasmacytoid dendritic cell proportion is predictive of risk and outcomes in myelodysplastic syndromes [abstract]. *Blood.* 2019; 134(suppl 1). Abstract 5439.
- Eppert K, Takenaka K, Lechman ER, et al. Stem cell gene expression programs influence clinical outcome in human leukemia. *Nat Med.* 2011;17(9):1086-1093.
- 67. Roshal M, Chien S, Othus M, et al. The proportion of CD34(+)CD38(low or neg) myeloblasts, but not side population frequency, predicts initial response to induction therapy in patients with newly diagnosed acute myeloid leukemia. *Leukemia*. 2013;27(3):728-731.
- 68. Simon L, Lavallée VP, Bordeleau ME, et al. Chemogenomic landscape of RUNX1mutated AML reveals importance of RUNX1 allele dosage in genetics and glucocorticoid sensitivity. Clin Cancer Res. 2017;23(22): 6969-6981.

- Gerritsen M, Yi G, Tijchon E, et al. RUNX1 mutations enhance self-renewal and block granulocytic differentiation in human in vitro models and primary AMLs. *Blood Adv.* 2019; 3(3):320-332.
- 70. In 't Hout FEM, Gerritsen M, Bullinger L, et al. Transcription factor 4 (TCF4) expression predicts clinical outcome in RUNX1 mutated and translocated acute myeloid leukemia [published online ahead of print 19 Dec 2019].

Haematologica. doi:org/10.3324/haematol. 2019.232827.

- 71. Satpathy AT, Briseño CG, Cai X, et al. Runx1 and Cbfβ regulate the development of Flt3+ dendritic cell progenitors and restrict myeloproliferative disorder. *Blood*. 2014; 123(19):2968-2977.
- 72. Ichikawa M, Asai T, Saito T, et al. AML-1 is required for megakaryocytic maturation and

lymphocytic differentiation, but not for maintenance of hematopoietic stem cells in adult hematopoiesis [published correction appears in Nat Med. 2005; 11(1):102]. *Nat Med.* 2004;10(3):299-304.

 Frankel A, Liu J-S, Rizzieri D, Hogge D. Phase I clinical study of diphtheria toxin-interleukin 3 fusion protein in patients with acute myeloid leukemia and myelodysplasia. *Leuk Lymphoma*. 2008;49(3):543-553.