MYELOID NEOPLASIA

Ivosidenib or enasidenib combined with intensive chemotherapy in patients with newly diagnosed AML: a phase 1 study

Eytan M. Stein,^{1,*} Courtney D. DiNardo,^{2,*} Amir T. Fathi,³ Alice S. Mims,⁴ Keith W. Pratz,⁵ Michael R. Savona,⁶ Anthony S. Stein,⁷ Richard M. Stone,⁸ Eric S. Winer,⁸ Christopher S. Seet,⁹ Hartmut Döhner,¹⁰ Daniel A. Pollyea,¹¹ James K. McCloskey,¹² Olatoyosi Odenike,¹³ Bob Löwenberg,¹⁴ Gert J. Ossenkoppele,¹⁵ Prapti A. Patel,¹⁶ Mikhail Roshal,¹⁷ Mark G. Frattini,¹⁸ Frederik Lersch,¹⁹ Aleksandra Franovic,²⁰ Salah Nabhan,²¹ Bin Fan,²¹ Sung Choe,²¹ Hongfang Wang,²¹ Bin Wu,²¹ Lei Hua,²¹ Caroline Almon,²¹ Michael Cooper,²¹ Hagop M. Kantarjian,^{2,†} and Martin S. Tallman^{1,†}

¹Department of Medicine, Memorial Sloan Kettering Cancer Center, New York, NY; ²Department of Leukemia, University of Texas MD Anderson Cancer Center, Houston, TX; ³Dana-Faber/Harvard Cancer Center, Massachusetts General Hospital, Boston, MA; ⁴Ohio State University Wexner Medical Center, Columbus, OH; ⁵Sidney Kimmel Comprehensive Cancer Center, Johns Hopkins University, Baltimore, MD; ⁶Vanderbilt-Ingram Center, Vanderbilt University School of Medicine, Nashville, TN; ⁷Department of Hematology & Hematopoietic Cell Transplantation, City of Hope Medical Center, Duarte, CA; ⁸Department of Medical Oncology, Dana-Farber Cancer Institute, Boston, MA; ⁹Department of Medicine, University of California Los Angeles Medical Center, Los Angeles, CA; ¹⁰Department of Internal Medicine III, University Hospital Ulm, Ulm, Germany; ¹¹Department of Medicine, University of Colorado School of Medicine, Aurora, CO; ¹²John Theurer Cancer Center, Hackensack University Medical Center, Hackensack, NJ; ¹³University of Chicago Medicine Comprehensive Cancer Center, Chicago, IL; ¹⁴Department of Hematology, Erasmus University Medical Centre, Rotterdam, The Netherlands; ¹⁵Department of Internal Medicine, University of Texas Southwestern Medical Center, Dallas, TX; ¹⁷Department of Pathology, Memorial Sloan Kettering Cancer Center, New York, NY; ¹⁸Bristol-Myers Squibb, Summit, NJ; ¹⁹Celgene International, Boudry, Switzerland; ²⁰Bristol-Myers Squibb, San Francisco, CA; and ²¹Agios Pharmaceuticals, Inc., Cambridge, MA

KEY POINTS

- Ivosidenib or enasidenib combined with induction and consolidation chemotherapy were both well tolerated in newly diagnosed mIDH1/2 AML.
- CR/CRi/CRp rates: 77% (ivosidenib) and 74% (enasidenib); 39% and 23% of patients had mIDH1/2 clearance by digital polymerase chain reaction.

Ivosidenib (AG-120) and enasidenib (AG-221) are targeted oral inhibitors of the mutant isocitrate dehydrogenase (mIDH) 1 and 2 enzymes, respectively. Given their effectiveness as single agents in mIDH1/2 relapsed or refractory acute myeloid leukemia (AML), this phase 1 study evaluated the safety and efficacy of ivosidenib or enasidenib combined with intensive chemotherapy in patients with newly diagnosed mIDH1/2 AML. Ivosidenib 500 mg once daily and enasidenib 100 mg once daily were well tolerated in this setting, with safety profiles generally consistent with those of induction and consolidation chemotherapy alone. The frequency of IDH differentiation syndrome was low, as expected given the concurrent administration of cytotoxic chemotherapy. In patients receiving ivosidenib, the frequency and grades of QT interval prolongation were similar to those observed with ivosidenib monotherapy. Increases in total bilirubin were more frequently observed in patients treated with enasidenib, consistent with this inhibitor's known potential to inhibit UGT1A1, but did not appear to have significant clinical consequences. In patients receiving ivosidenib (n = 60) or enasidenib (n = 91), end-of-induction complete remission (CR) rates were 55% and 47%, respectively, and CR/CR with incomplete neutrophil or platelet recovery (CR/CRi/CRp) rates were 72% and 63%, respectively. In pa-

tients with a best overall response of CR/CRi/CRp, 16/41 (39%) receiving ivosidenib had *IDH1* mutation clearance and 15/64 (23%) receiving enasidenib had *IDH2* mutation clearance by digital polymerase chain reaction; furthermore, 16/20 (80%) and 10/16 (63%), respectively, became negative for measurable residual disease by multiparameter flow cytometry. This trial was registered at www.clinicaltrials.gov as #NCT02632708. (*Blood*. 2021;137(13):1792-1803)

Introduction

Intensive induction chemotherapy with cytarabine and an anthracycline ("7 + 3") remains the most effective treatment for adults with newly diagnosed acute myeloid leukemia (AML) who can withstand its toxicities. Recent modifications to this backbone have improved event-free survival and overall survival in defined subsets of patients.¹⁻³ Mutations in the isocitrate dehydrogenase (*IDH*) 1 or 2 genes are seen in ~20% of patients with AML.⁴⁻⁷ Mutant IDH (mIDH) proteins catalyze the production of D-2-hydroxyglutarate (2-HG),^{4,8} resulting in DNA and histone hypermethylation, with consequent changes in gene expression and impaired cellular differentiation.⁹⁻¹¹ Ivosidenib and enasidenib are targeted, oral, small-molecule inhibitors of the

mIDH1 and mIDH2 enzymes, respectively, approved by the US Food and Drug Administration as monotherapies for adults with relapsed or refractory (R/R) AML and a susceptible *IDH1* or *IDH2* mutation.^{12,13} In a phase 1 study, ivosidenib 500 mg once daily resulted in an overall response rate of 41.6% and a median response duration of 6.5 months in patients with m*IDH1* R/R AML.¹⁴ Similarly, the overall response rate in patients with m*IDH2* R/R AML receiving enasidenib 100 mg once daily was 40.3%, with a median response duration of 5.8 months.¹⁵ We hypothesized that combining ivosidenib or enasidenib with intensive induction and consolidation chemotherapy would improve outcomes for patients with newly diagnosed m*IDH1/2* AML.

Methods

Study design

This phase 1, multicenter, open-label study enrolled patients with m*IDH1* or m*IDH2* newly diagnosed AML. Induction therapy consisted of continuous ivosidenib 500 mg once daily (m*IDH1*) or enasidenib 100 mg once daily (m*IDH2*) in combination with cytarabine (200 mg/m² per day for 7 days) and either daunorubicin (60 mg/m² per day for 3 days) or idarubicin (12 mg/m² per day for 3 days). A second cycle of induction was permitted according to institutional practice.

Six patients were initially enrolled to each of the 4 induction treatment arms. If ≤ 2 of the 6 patients initially enrolled in a treatment arm experienced a dose-limiting toxicity (DLT) at the standard dose of either mIDH inhibitor, then an additional 6 patients were to be evaluated at that dose. If \leq 3 of the expanded cohort of 12 patients experienced a DLT, then that dose was to be declared suitable for further evaluation. However, if \geq 3 of the initial 6 patients or \geq 4 of the first 12 patients experienced a DLT, a lower dose was to be administered. After the safety of the combination regimens had been determined for each of the 4 induction therapy cohorts, \sim 30 patients in total were to be enrolled into each cohort to better characterize the safety profile of these regimens. This larger number of patients in each induction cohort also ensured that an adequate number would proceed to the consolidation phase of treatment, so that the safety profile of the mIDH inhibitors in combination with consolidation therapy could be evaluated. Owing to an early concern regarding delayed recovery of blood counts in patients receiving enasidenib with induction chemotherapy, 2 additional cohorts (for the daunorubicin- and idarubicin-based regimens) were treated with enasidenib 100 mg once daily beginning on day 8 (instead of day 1) of the first cycle of induction therapy.

Patients achieving at least a partial remission at the end of induction could receive consolidation therapy (up to 4 cycles of intermediate- or high-dose cytarabine, or 1 cycle of mitoxantrone/etoposide), while continuing to receive ivosidenib or enasidenib continuously once daily. The safety of each consolidation therapy regimen was evaluated after the first 6 evaluable patients had completed ≥28 days of consolidation treatment or discontinued owing to toxicity.

Patients remaining in remission at the end of consolidation could receive maintenance treatment with ivosidenib or enasidenib monotherapy daily until relapse, development of an unacceptable toxicity, or allogeneic hematopoietic stem cell transplantation (HSCT). Patients appropriate for HSCT could proceed to HSCT at any point; resumption of mIDH inhibitor therapy was not allowed after HSCT. See supplemental Appendix on the *Blood* Web site for further details on treatment.

The study was conducted in accordance with the principles of the Declaration of Helsinki and International Conference on Harmonization Guidelines for Good Clinical Practice, and the protocol was approved by human investigation committees at participating sites. Written informed consent was provided by all patients before screening and enrollment.

The study sponsor analyzed the data and conducted the statistical analyses. All authors had access to the primary clinical trial data on request.

Patients

Patients \geq 18 years of age with newly diagnosed AML (de novo or secondary) were eligible if they had an Eastern Cooperative Oncology Group performance status score of 0 to 2 and a documented *IDH1* and/or *IDH2* mutation by local laboratory testing. Secondary AML was defined as AML arising after myelodysplastic syndrome or another hematologic disorder, or AML arising after exposure to genotoxic injury (ie, radiation and/ or chemotherapy). Patients were ineligible if they had received any prior chemotherapy for AML except hydroxyurea, but previous treatment of myelodysplastic syndrome or another antecedent hematologic disorder, including hypomethylating agents, was permitted if the last dose was \geq 14 days before study treatment initiation.

Assessments

The primary objective was to assess the safety and tolerability of the combination regimens. Treatment-emergent adverse events were graded according to the National Cancer Institute Common Terminology Criteria for Adverse Events, version 4.03. Analyses of time to hematologic recovery were conducted for the induction phase (supplemental Appendix).

Secondary objectives included investigator assessment of clinical responses using the modified 2003 International Working Group response criteria for AML, and overall survival (further details in supplemental Appendix).

Pharmacokinetics and pharmacodynamics

Analyses were conducted as reported in the supplemental Appendix.

Translational analyses

Retrospective confirmation of *IDH* mutation and analysis of cooccurring mutations at baseline were performed by nextgeneration sequencing (NGS) using the 95-gene Rapid Heme Panel (detection sensitivity, 5%).¹⁶ Samples from baseline and specified on-study time points were analyzed for mutation evolution using the Personalis ACE Cancer Panel (Menlo Park, CA)¹⁷ for ivosidenib-treated patients, and the Archer VariantPlex Core Myeloid Panel (Boulder, CO) for enasidenib-treated patients; analyses were limited to the 33 genes represented on both platforms (supplemental Table 1), and a 2% limit of detection for variant allele frequency (VAF) was applied to both datasets. Heatmaps were produced in R, version 3.6.0. Longitudinal m*IDH1/2* VAF was assessed using the OncoBEAM

Table 1. Baseline characteristics

Characteristic	lvosidenib 500 mg + chemotherapy, n = 60	Enasidenib 100 mg + chemotherapy, n = 93		
Age, median (range), y	62.5 (24-76)	63.0 (27-77)		
Age category, n (%), y <60 ≥60	21 (35) 39 (65)	34 (37) 59 (63)		
Men, n (%)	30 (50)	52 (56)		
Type of AML, n (%) De novo Secondary	42 (70) 18 (30)	58 (62) 35 (38)		
Prior hypomethylating agent, n (%)*	4 (22)	17 (49)		
IDH1 mutation type, n (%)† R132 Other or unknown	58 (97) 2 (3)	2 (2) 1 (1)		
IDH2 mutation type, n (%)† R140 R172 Other or unknown	1 (2) 1 (2) 0	66 (71) 25 (27) 2 (2)		
Cytogenetic risk status by investigator, n (%) Favorable Intermediate Poor Unknown	0 42 (70) 13 (22) 5 (8)	2 (2) 64 (69) 20 (22) 7 (8)		

*For patients with secondary AML only.

[†]Patients with dual *IDH1* and *IDH2* mutations were assigned to ivosidenib or enasidenib on the basis of the *IDH* mutation with the higher allele burden.

BEAMing digital polymerase chain reaction (dPCR) assay (lower limit of detection, 0.02% to 0.04%; Sysmex Inostics Inc., Hamburg, Germany).¹⁸ Mutation clearance was defined as a reduction in *mIDH1/2* VAF that was below the limit of detection for \geq 1 ontreatment time point on or after day 21 of induction therapy in patients who had detectable *mIDH1/2* at baseline. Measurable residual disease (MRD) was assessed centrally by multiparameter flow cytometry for a subset of patients. Further detail on all exploratory assessments is provided in the supplemental Appendix.

Statistical analysis

The data cutoff date was 13 December 2018. The safety sets included all patients who received at least 1 dose of ivosidenib, enasidenib, or chemotherapy during the relevant study period (ie, induction, consolidation, and maintenance). All efficacy analyses were performed using the full analysis set (FAS), comprising all enrolled patients who received at least 1 dose of ivosidenib or enasidenib. Only response assessments occurring on or after day 21 were used to determine the best response.

The proportion of patients in each best overall response category was evaluated using the Clopper-Pearson method for the overall treatment period (induction, consolidation, and maintenance; excludes post-HSCT response) and for the de novo AML and secondary AML subgroups. Median overall survival and 12-month overall survival rates were estimated using the Kaplan-Meier method, along with associated 95% confidence intervals based on log-log transformation. See supplemental Appendix for additional statistical methods.

Results

Patient characteristics and disposition

Sixty patients were dosed in the ivosidenib plus induction chemotherapy cohorts and 93 in the enasidenib plus induction chemotherapy cohorts from January 2016 to July 2018. Baseline characteristics are provided in Table 1 and patient disposition in Figure 1.

In the ivosidenib cohorts, the median age was 62.5 years and 70% of patients had de novo AML. Of the 60 patients in the ivosidenib induction cohorts, 9 (15%) received 2 induction cycles, 35 (58%) went on to receive consolidation, and 19 (32%) received ivosidenib maintenance therapy. At the data cutoff date, 47 (78%) ivosidenib-treated patients had discontinued study treatment, most commonly for HSCT (28 patients), whereas 13 patients were continuing to receive ivosidenib as maintenance therapy.

In the enasidenib cohorts, the median age was 63.0 years and 62% of patients had de novo AML. Of the 93 patients in the enasidenib induction cohorts, 2 who were assigned to receive enasidenib starting on day 8 had an ongoing adverse event or died on day 8 and, as a result, never received a dose of enasidenib. These 2 patients are included in the induction safety analysis set but not in the FAS used for efficacy analyses. Twenty-two patients (24%) received 2 induction cycles, 46 patients (49%) received consolidation, and 24 (26%) received enasidenib maintenance therapy. At the data cutoff date, 81 enasidenib-treated patients (87%) had discontinued study treatment, most commonly for HSCT (43 patients), whereas 12 patients continued to receive enasidenib as maintenance therapy.

Safety

During the initial safety evaluation of the combination therapies, the sole DLT occurred during induction in a 64-year-old man with de novo AML in the enasidenib plus daunorubicin/cytarabine cohort. He experienced persistent grade 4 thrombocytopenia in the absence of residual leukemia on day 42 of induction cycle 1. No patients in the ivosidenib cohorts experienced a DLT. Ivosidenib and enasidenib were well tolerated at their starting doses of 500 and 100 mg each day; therefore, these doses were chosen for evaluation in additional patients.

The toxicities observed during induction and consolidation therapy with the combinations were similar to the toxicities seen with "7 + 3" and cytarabine alone (Table 2; supplemental Tables 2 and 3). Only 2 patients in the enasidenib cohorts and none in the ivosidenib cohorts received mitoxantrone/etoposide as consolidation therapy; hence, the safety of this combination consolidation regimen could not be formally evaluated.



Figure 1. Disposition of the study population. ARA-C, cytarabine; DNR, daunorubicin; IDR, idarubicin.

The 30- and 60-day mortality rates were 5% and 10%, respectively, in the ivosidenib cohorts, and 5% and 9%, respectively, in the enasidenib cohorts. There were 13 on-study deaths (including the survival follow-up period) in the ivosidenibtreated cohorts, of which 7 (54%) occurred during treatment (occurring within 28 days of the last dose). Of 31 on-study deaths (including survival follow-up) in the enasidenib-treated cohort, 13 (42%) occurred during treatment. None of the deaths were related to ivosidenib or enasidenib; the majority were attributed to disease progression or complications of the underlying disease, such as respiratory failure, lung infection, intracranial hemorrhage, or sepsis.

The use of ivosidenib and enasidenib in combination with induction chemotherapy did not affect the time to recovery of the absolute neutrophil count (ANC) or platelet count. In the ivosidenib cohorts, the median times to recovery of the ANC (>500/ μ L) and platelet count (>50 000/ μ L) during induction were both 28 days (supplemental Table 4). Early in the study there was concern regarding delayed recovery of platelets and neutrophils in patients who received enasidenib with induction chemotherapy. Therefore, a cohort of 25 patients received an alternative dosing schedule in which enasidenib dosing started on day 8, after completion of the administration of "7 + 3," instead of at the beginning. Although a comparison of the time to blood count recovery between the cohorts starting enasidenib on day 1 vs day 8 was limited by the small number of patients who started enasidenib on day 8, the time to count recovery was generally similar between the 2 groups (supplemental Table 5). Overall, when combining the enasidenib day 1 and day 8 cohorts, the median times to ANC and platelet count recovery during the induction phase were 34 and 29 days, respectively (supplemental Table 4).

IDH differentiation syndrome (IDH-DS) was reported in 2 patients (3.3%) receiving ivosidenib and 2 patients (2.2%) receiving enasidenib. All 4 cases occurred during induction, starting between days 29 and 48. Three of the 4 cases were grade \geq 3 in severity. Ultimately, IDH-DS was reported as resolved in 3 of the 4 patients and all 3 went on to achieve a complete remission (CR) or CR with incomplete platelet recovery (CRp). The fourth patient died of a lung infection and IDH-DS was reported as ongoing at the time of death.

In the ivosidenib cohorts, QT interval prolongation was observed in 16 patients (26.7%) during induction and was grade \geq 3 in 6 of these patients (10%). During the consolidation phase, QT prolongation was observed in 3 patients (8.6%) in the ivosidenib cohorts and was grade \geq 3 in 1 patient (2.9%). In the enasidenib cohorts, QT prolongation during induction was reported in 11 patients (11.8%) and was grade \geq 3 in 7 patients (7.5%), whereas during the consolidation period, QT prolongation was reported in 7 patients (15.2%) and was grade \geq 3 in 3 patients (6.5%).

In patients treated with enasidenib, increased blood bilirubin was reported during induction in 46 patients (49.5%) and was grade \geq 3 in 15 patients (16%). During consolidation, increased blood bilirubin was reported in 13 patients (28.3%) and was

Table 2. Nonhematologic TEAEs of any grade reported in >20% of patients in any treatment group, and the corresponding frequencies of grade ≥3 events, during the induction and consolidation periods, regardless of attribution

	Induction period				Consolidation period			
	Ivosidenib 500 mg + chemotherapy, n = 60		Enasidenib 100 mg + chemotherapy, n = 93		Ivosidenib 500 mg + chemotherapy, n = 35		Enasidenib 100 mg + chemotherapy, n = 46	
TEAE, n (%)	All grades	Grade ≥3						
Any TEAE	60 (100.0)	58 (96.7)	92 (98.9)	87 (93.5)	35 (100.0)	34 (97.1)	45 (97.8)	41 (89.1)
Diarrhea	43 (71.7)	1 (1.7)	55 (59.1)	5 (5.4)	7 (20.0)	0	17 (37.0)	0
Nausea	33 (55.0)	0	50 (53.8)	2 (2.2)	11 (31.4)	0	15 (32.6)	1 (2.2)
Rash*	33 (55.0)	3 (5.0)	51 (54.8)	13 (14.0)	12 (34.3)	1 (2.9)	13 (28.3)	1 (2.2)
Decreased appetite	32 (53.3)	5 (8.3)	31 (33.3)	3 (3.2)	4 (11.4)	0	11 (23.9)	1 (2.2)
Vomiting	21 (35.0)	0	31 (33.3)	1 (1.1)	9 (25.7)	1 (2.9)	10 (21.7)	0
Stomatitis	20 (33.3)	3 (5.0)	23 (24.7)	4 (4.3)	4 (11.4)	0	9 (19.6)	4 (8.7)
Fatigue	19 (31.7)	0	24 (25.8)	2 (2.2)	5 (14.3)	0	9 (19.6)	2 (4.3)
Hypokalemia	17 (28.3)	7 (11.7)	29 (31.2)	9 (9.7)	3 (8.6)	1 (2.9)	11 (23.9)	4 (8.7)
Pyrexia	16 (26.7)	4 (6.7)	31 (33.3)	2 (2.2)	8 (22.9)	1 (2.9)	13 (28.3)	0
Constipation	14 (23.3)	0	25 (26.9)	0	6 (17.1)	0	13 (28.3)	0
Hypophosphatemia	14 (23.3)	10 (16.7)	20 (21.5)	12 (12.9)	0	0	7 (15.2)	4 (8.7)
Edema peripheral	14 (23.3)	0	37 (39.8)	0	2 (5.7)	0	13 (28.3)	0
Abdominal pain	13 (21.7)	2 (3.3)	19 (20.4)	0	4 (11.4)	0	8 (17.4)	1 (2.2)
Aspartate aminotransferase increased	13 (21.7)	4 (6.7)	18 (19.4)	4 (4.3)	1 (2.9)	0	6 (13.0)	2 (4.3)
Electrocardiogram QT prolonged†	16 (26.7)	6 (10.0)	11 (11.8)	7 (7.5)	3 (8.6)	1 (2.9)	7 (15.2)	3 (6.5)
Alanine aminotransferase increased	11 (18.3)	4 (6.7)	22 (23.7)	5 (5.4)	2 (5.7)	1 (2.9)	6 (13.0)	2 (4.3)
Cough	11 (18.3)	0	22 (23.7)	1 (1.1)	3 (8.6)	0	9 (19.6)	0
Headache	10 (16.7)	0	29 (31.2)	0	6 (17.1)	1 (2.9)	8 (17.4)	0
Hypoalbuminemia	10 (16.7)	3 (5.0)	21 (22.6)	5 (5.4)	1 (2.9)	1 (2.9)	3 (6.5)	0
Blood bilirubin increased‡	11 (18.3)	4 (6.7)	46 (49.5)	15 (16.1)	2 (5.7)	1 (2.9)	13 (28.3)	5 (10.9)
Hypocalcemia	9 (15.0)	3 (5.0)	24 (25.8)	6 (6.5)	2 (5.7)	1 (2.9)	6 (13.0)	2 (4.3)

TEAE, treatment-emergent adverse event.

*Rash includes preferred terms rash, rash maculopapular, rash pruritic, rash erythematous, rash macular, dermatitis, dermatitis acneiform, dermatitis allergic, dermatitis bullous, exfoliative rash, skin ulcer, drug eruption, and urticaria.

†Electrocardiogram QT prolonged includes ventricular tachycardia, ventricular arrhythmia, cardiac arrest, cardiorespiratory arrest, electrocardiogram QT prolonged, multiple organ dysfunction syndrome, and syncope.

‡Blood bilirubin increased includes preferred terms of increased blood bilirubin and hyperbilirubinemia.

grade \geq 3 in 5 patients (11%). In patients treated with ivosidenib, increased blood bilirubin was reported during induction in 11 patients (18.3%) and was grade \geq 3 in 4 patients (6.7%). During consolidation, increased blood bilirubin was reported in 2 patients (5.7%) and was grade \geq 3 in 1 patient (2.9%). Further analysis of these data showed that in patients treated with enasidenib, grade 1 and 2 increases in blood bilirubin were primarily due to increases in unconjugated bilirubin. When grade \geq 3 increases in blood bilirubin occurred in enasidenib-treated patients, they consisted of increases in both the conjugated and unconjugated forms, but with a trend toward greater proportional increases in unconjugated bilirubin than in patients treated with ivosidenib. This is consistent with enasidenib's inhibition of UGT1A1.

Table 3. Best overall responses at any t	time during the study in the FAS
--	----------------------------------

	Ivoside	nib 500 mg + chen	notherapy, n (%)	Enasidenib 100 mg + chemotherapy, n (%)			
Response category	All, N = 60	De novo AML, n = 42	Secondary AML, n = 18	All, N = 91*	De novo AML, n = 56	Secondary AML, n = 35	
CR/CRi/CRp	46 (77)	37 (88)	9 (50)	67 (74)	45 (80)	22 (63)	
CR	41 (68)	32 (76)	9 (50)	50 (55)	36 (64)	14 (40)	
CRi/CRp	5 (8)	5 (12)	—	17 (19)	9 (16)	8 (23)	
MLFS	4 (7)	3 (7)	1 (6)	10 (11)	5 (9)	5 (14)	
PR	2 (3)	—	2 (11)	2 (2)	1 (2)	1 (3)	
Treatment failure†	8 (13)	2 (5)	6 (33)	12 (13)	5 (9)	7 (20)	

MLFS, morphologic leukemia-free state

*Two patients assigned to receive enasidenib starting on day 8 had an ongoing adverse event or died on day 8 and thus never received enasidenib; these 2 patients were not included in the FAS used for efficacy analyses.

+Treatment failure = stable disease + progressive disease + discontinuation before response assessment on or after induction day 21 + discontinuation with best response of not evaluable.

Efficacy

Response In the ivosidenib-treated cohorts, rates of CR and the combined measure of CR plus CR with incomplete neutrophil recovery (CRi) or CRp (CR/CRi/CRp) at the end of the induction period were 55% and 72%, respectively, and were 68% and 77%, respectively, when evaluated at any time on study. In the enasidenib-treated cohorts, CR and CR/CRi/CRp rates were 47% and 63%, respectively, at the end of the induction period, and were 55% and 74%, respectively, when evaluated at any time on study (supplemental Figure 1).

In the ivosidenib-treated cohorts, CR and CR/CRi/CRp rates at any time during the study were higher in patients with de novo AML than in those with secondary AML (Table 3). Similar findings were observed in the enasidenib-treated cohorts. Post hoc analyses of best overall response in patients with de novo AML with or without myelodysplastic syndrome-related cytogenetic abnormalities, in secondary AML with or without prior HMA treatment, and by *IDH2* mutation type were conducted but proved to be inconclusive because of the limited number of patients in these subgroups (supplemental Tables 6-8). **Overall survival** After a median follow-up period of 9.3 months (range, 0.4-32.1 months), median overall survival was not reached in the ivosidenib-treated cohorts; the 12-month survival probability after induction day 1 was 78% (Figure 2). When patients were censored at the time of HSCT, the 12-month survival probability was 74%. In the enasidenib-treated cohorts, median overall survival was 25.6 months (95% confidence intervals, 25.5 and not calculable) after a median follow-up period of 14.5 months (range, 0.5-31.8), and the 12-month survival probability was 76% (Figure 2). When patients were censored at time of HSCT, the 12-month survival probability as 76% (Figure 2). When patients were censored at time of HSCT, the 12-month survival probability was 76% (Figure 2). When patients were censored at time of HSCT, the 12-month survival probability was 76% (Figure 2). When patients were censored at time of HSCT, the 12-month survival probability was 67%. Event-free survival analyses were conducted but were limited by the relatively high number of patients who were censored when they discontinued treatment to proceed to HSCT (supplemental Appendix).

Pharmacokinetic and pharmacodynamic data Pharmacokinetic/ pharmacodynamic parameters observed in this study were consistent with findings from previous studies in which ivosidenib and enasidenib were administered as single agents^{19,20} and appeared to be similar across daunorubicin and idarubicin induction cohorts. Ivosidenib and enasidenib were rapidly absorbed, with peak plasma concentrations at 4 hours following single (data not shown)



Figure 2. Overall survival in the FAS set. Patients not censored at the time of HSCT.



Downloaded from http://ashpublications.net/blood/article-pdf/137/13/1792/1803916/bloodbld2020007233.pdf by guest on 08 June 2024

Figure 3. Baseline mutational landscape and best overall clinical responses. Each column represents an individual patient, organized by best overall response. Genes (rows) are grouped by biological pathway. A blue box indicates the detection of a known or likely oncogenic variant in at least 1 sample type (peripheral blood and/or bone marrow). NA, not assessed; NE, not evaluable; PD, progressive disease; RTK, receptor tyrosine kinase; SD, stable disease.

and multiple doses (supplemental Table 9). Exposure at steady state following 14 days of daily dosing was higher than after a single dose, with moderate accumulation (ranging from 1.7- to 2.4-fold) for ivosidenib and high accumulation (ranging from 6.3- to 8.3-fold) for enasidenib.

Plasma 2-HG concentrations were elevated at baseline and decreased after single and multiple doses of ivosidenib or enasidenib (supplemental Table 9). After multiple doses, mean trough plasma 2-HG concentrations were reduced by up to 99%, to levels observed in healthy participants, and were maintained throughout continued ivosidenib or enasidenib dosing. However, in contrast to the decreases in 2-HG observed in single-agent studies of these mIDH inhibitors, the decreases observed

in this study likely also reflect the antileukemic activity of the standard chemotherapeutic agents.

Baseline mutation profiling and clinical response In the ivosidenib-treated cohorts, the most frequent baseline comutations were DNMT3A (41%), NPM1 (34%), ASXL1 (20%), and BCOR (14%), and in the enasidenib cohorts were DNMT3A (39%), SRSF2 (25%), ASXL1 (23%), and RUNX1 (20%) (supplemental Figure 2). Figure 3 summarizes the known or likely oncogenic variants detected at baseline, with patients organized by best overall response. In the ivosidenib-treated cohort, there were no co-mutations associated with a higher likelihood of CR/CRi/CRp. In the enasidenib-treated cohort, mutations in ASXL1, NRAS, U2AF1, and TP53 were associated with a lack of

Treatment	n	IDH mutation clearance, n (%)	n	MRD⁻, n (%)
lvosidenib + chemotherapy	41	16 (39)	20	16 (80)
Enasidenib + chemotherapy				
Total	64	15 (23)	16	10 (62)
R140	47	11 (23)	12	8 (67)
R172	17	4 (24)	4	2 (50)

CR/CRi/CRp, whereas *DNMT3A* mutations were marginally associated with CR/CRi/CRp (supplemental Table 10).

IDH1/2 mutation clearance and MRD assessment In ivosidenibtreated patients with a best response of CR/CRi/CRp and who had samples available for analysis, 16 of 41 (39%) had m/DH1 clearance. In enasidenib-treated patients with a best response of CR/CRi/CRp and who had samples available for analysis, 15 of 64 (23%) had m/DH2 clearance from bone marrow mononuclear cells by dPCR (Table 4).

MRD negativity was assessed using multiparameter flow cytometry in a subset of patients that was not identical to that in which m*IDH* clearance was assessed. Among ivosidenib-treated patients with a best response of CR/CRi/CRp, 16 of 20 (80%) were MRD⁻, whereas among enasidenib-treated patients with a best response of CR/CRi/CRp, 10 of 16 (63%) were MRD⁻ (Table 4). Supplemental Table 11 shows mutation clearance and MRD⁻ rates for all response categories and by *IDH2* mutation type. Longitudinal mutation clearance at the per-patient level is shown in supplemental Figures 3 and 4.

In the subset of patients with a best overall response of CR/CRi/ CRp who had data for both mIDH1/2 clearance by dPCR and MRD by multiparameter flow cytometry, 6 of 6 (100%) ivosidenibtreated patients and 1 of 2 (50%) enasidenib-treated patients who cleared their IDH1/2 mutation also achieved MRD⁻. However, 9 of 15 (60%) ivosidenib-treated patients and 9 of 10 (90%) enasidenibtreated patients who achieved elimination of MRD had detectable mIDH1/2, suggesting that dPCR for mIDH clearance is more sensitive for determining the presence of underlying disease than assessment of MRD using multiparameter flow cytometry (supplemental Table 12).

Co-mutation clearance at end of induction We performed targeted NGS profiling at baseline and the end of induction in patients with a best of response of CR/CRi/CRp (ivosidenib n = 31, enasidenib n = 28) to monitor co-occurring mutation clearance at the 2% VAF level. Figure 4 shows a per-gene overview of the number of patients with a detectable mutation at baseline vs the end of induction. The frequency of mutation clearance at the end of induction varied considerably by gene. Mutations in *DNMT3A*, *TET2*, and/or *ASXL1* (DTA mutations) are common in patients with age-related clonal hematopoiesis, and DTA mutations have been

shown frequently to persist after chemotherapy, with no significant impact on prognosis.^{21,22} We observed that aside from DTA mutations, most co-occurring mutations were cleared by the combination of ivosidenib or enasidenib and induction chemotherapy.

Supplemental Table 13 summarizes mutation clearance (NGS; sensitivity 2% VAF) at the end of induction. In ivosidenib-treated patients with a response of CR/CRi/CRp, the rates of clearance of m/DH1 and non-DTA co-mutations were 67% and 58%, respectively. In enasidenib-treated patients with a response of CR/CRi/CRp, the rate of clearance for m/DH2-R140 was 75%, whereas for m/DH2-R172 it was 43%. The rate of non-DTA mutation clearance was the same in patients with m/DH2-R140 (43%) and those with m/DH2-R172 (43%).

Mutation profiling at relapse Of the 4 ivosidenib-treated and 8 enasidenib-treated patients who relapsed, mutational profiling at relapse was available for 3 patients from the ivosidenib cohorts and 1 patient from the enasidenib cohorts. Co-mutations at relapse were the same as those at screening, except in 1 ivosidenib-treated patient in whom a *GNAS* mutation (L46del) emerged, and in another ivosidenib-treated patient in whom a *KDM5C* F1376fs mutation emerged and m*IDH1* was not detectable.

Discussion

Because ivosidenib and enasidenib have activity as single agents in R/R mIDH1/2 AML,14,15 we assessed their combination with intensive induction and consolidation chemotherapy in patients with newly diagnosed mIDH1/2 AML. Ivosidenib 500 mg once daily and enasidenib 100 mg once daily were well tolerated when combined with induction and consolidation therapy, with safety profiles similar to those seen with induction and consolidation chemotherapy alone. The addition of ivosidenib or enasidenib, regardless of start date, did not affect the time needed for hematologic recovery after induction chemotherapy when compared with historical references in relatively similar patient populations.^{23,24} Therefore, future studies will initiate treatment with the IDH inhibitors concurrently with the start of induction therapy. Thirty- and 60-day mortality rates in this older patient population were comparable with those observed in patients with newly diagnosed AML given intensive chemotherapy. IDH-DS has been observed in ${\sim}10\%$ of patients treated with ivosidenib or enasidenib as single agents.^{14,15} Because ivosidenib and enasidenib were administered in combination with cytotoxic chemotherapy in this study, rates of IDH-DS were low, as expected. QT interval prolongation has been associated with ivosidenib monotherapy,¹⁴ and in the present study the frequency and grades of QT prolongation were similar to those observed with ivosidenib monotherapy in R/R AML.¹⁴ However, ivosidenib's independent contribution to QT prolongation in this study could not be determined, given that the use of supportive medications known to prolong the QT interval (eg, quinolones, azole antifungals, 5-HT3 antagonists) was permitted. Increased blood bilirubin was more frequently observed in the enasidenib cohorts, consistent with enasidenib's known inhibition of the UGT1A1 enzyme,¹⁵ but it did not appear to have significant clinical consequences.

In previous studies, CR rates in patients with newly diagnosed or previously treated mIDH1/2 AML who received induction and



Figure 4. Paired sample analysis of mutations by NGS at screening and end of induction. (A) lvosidenib-treated patients with a best response of CR/CRi/CRp only, n = 31, Personalis ACE Cancer Panel. (B) Enasidenib-treated patients with a best response of CR/CRi/CRp only, n = 28, Archer VariantPlex Core Myeloid panel. Values under each bar denote the number of patients with a mutation detected at baseline and end of induction, and the mutation clearance rate (%) for each gene.

consolidation chemotherapy ranged from 38% to 65%, although definitions of CR varied across the studies.^{5,25,26} The end-ofinduction CR and CR/CRi/CRp rates observed in our study were 55% and 72%, respectively, in the ivosidenib-treated patients, and 47% and 63%, respectively, in the enasidenib-treated patients. The best overall CR and CR/CRi/CRp rates were 68% and 77%, respectively, in ivosidenib-treated patients, and 55% and 74%, respectively, in enasidenib-treated patients. These response rates are encouraging, particularly in this group of patients, which included a substantial proportion with secondary AML (~30%) and a majority \geq 60 years of age. However, because this was a phase 1 study with no comparator group, and owing to the limited data available for older patients with newly diagnosed m*IDH* AML treated with intensive induction and consolidation therapy, the interpretation of our remission rates is challenging. A randomized, controlled trial is needed to

adequately compare response rates for each mIDH inhibitor combination with intensive chemotherapy alone.

To better understand the depth of clinical responses in the present study, mIDH clearance (by highly sensitive dPCR) and the elimination of MRD (by multiparameter flow cytometry) were assessed. These analyses were limited in that samples were not regularly collected after the end of induction and thus are not time-matched across patients. Additionally, the subset of patients with samples available for mIDH clearance analysis was not the same as the subset with samples available for MRD analysis. Nevertheless, among patients who achieved CR/CRi/CRp, ivosidenib given with intensive induction and consolidation chemotherapy was associated with mIDH1 clearance in 39% of patients (16 of 41) and with elimination of MRD in 80% of patients (16 of 20). Enasidenib given with the same chemotherapy was associated with mIDH2 clearance in 23% of patients (15 of 64) and with the elimination of MRD in 63% of patients (10 of 16). In the small group of patients for whom results were available both for mIDH1/2 clearance and for MRD, clearance of mIDH1/2 was often associated with elimination of MRD. However, a substantial proportion of patients who demonstrated persistence of mIDH1/ 2 achieved elimination of MRD, suggesting that mIDH clearance as assessed by dPCR is a more sensitive measure for detecting the presence of the underlying disease than assessing MRD using multiparameter flow cytometry. Comparisons of mutation clearance rates with those of other studies cannot be made because of the limited number of samples collected for this assessment in our study and the irregular timing of sample collection after cycle 1 day 28; moreover, the cutoff used for determination of mutation clearance varies between studies.

In patients with mIDH R/R AML, the presence of co-mutations in receptor tyrosine kinase pathway genes such as NRAS, KRAS, and FLT3 has been associated with resistance to single-agent therapy with inhibitors of mIDH.^{14,27,28} Our longitudinal molecular profiling in a subset of patients using NGS (cutoff 2%) of mutations at baseline, and at the time of best response, showed that mutations in FLT3 and RAS were cleared after induction chemotherapy, suggesting that the combination of an mIDH inhibitor with chemotherapy in the first-line setting may overcome one of the most common mechanisms of resistance to single-agent inhibitors of mIDH1/2. Although mutational profiling data were available for only 4 of the patients who relapsed, the data suggest that relapse can occur both with and without the emergence of new mutations.

In this study, ivosidenib-treated and enasidenib-treated patients had 12-month survival probabilities of >75%, which is promising given that survival rates have historically been low, especially for older patients with AML,^{29,30} although improvements have been noted in the past few decades, particularly in younger patients.³¹ Survival follow-up is ongoing to determine whether these encouraging rates will be maintained over a longer period.

In summary, the combination of ivosidenib or enasidenib with intensive induction and consolidation therapy was well tolerated in patients with newly diagnosed m*IDH1* or m*IDH2* AML, and the initial clinical activity was encouraging. The benefit of adding ivosidenib or enasidenib to induction and consolidation chemotherapy followed by single-agent maintenance therapy for

patients with newly diagnosed mIDH AML is being further evaluated in an ongoing randomized phase 3 trial.

Acknowledgments

Editorial assistance was provided to the authors by Helen Varley, CMPP, Excel Medical Affairs, Horsham, United Kingdom, and supported by Agios.

This work was supported by Agios Pharmaceuticals, Inc., in collaboration with Bristol-Myers Squibb. Agios provided financial support for the study and participated in the design, study conduct, and analysis and interpretation of data, as well as the writing, review, and approval of the manuscript.

Authorship

Contribution: E.M.S., C.D.D., A.T.F., M.R.S., R.M.S., H.D., B.L., G.J.O., S.N., L.H., C.A., M.C., and M.S.T. designed the study; E.M.S., C.D.D., A.T.F., A.S.M., K.W.P., M.R.S., A.S.S., R.M.S., E.S.W., C.S.S., H.D., D.A.P., J.K.M., O.O., B.L., G.J.O., P.A.P., M.R., H.M.K., and M.S.T. participated in patient enrollment and care of patients; E.M.S., C.D.D., A.T.F., A.S.M., K.W.P., M.R.S., A.S.S., R.M.S., E.S.W., C.S.S., H.D., D.A.P., J.K.M., OO., B.L., G.J.O., P.A.P., M.R., H.G.F., F.L., A.F., S.N., B.F., S.C., H.W., B.W., L.H., C.A., M.C., H.M.K., and M.S.T. participated in data collection, analysis, and interpretation; E.M.S., C.D.D., B.F., S.C., H.W., L.H., C.A., and M.C. oversaw drafting of the manuscript; and all authors participated in manuscript development and final approval of the submitted version.

Conflict-of-interest disclosure: E.M.S. is a stockholder in/has ownership of Auron Therapeutics; is a consultant/advisor to AbbVie, Agios, Astellas, Bayer, BioLineRx, Celgene, Daiichi Sankyo, Genentech, Novartis, Pfizer, PTC Therapeutics, and Syros; received research funding from Agios, Amgen, Bayer, Celgene, and Syros; and received travel expenses from AbbVie, Astellas, BioTheryX, Celgene, Daiichi Sankyo, Novartis, Syndax, and Syros. C.D.D. is a consultant/advisor to AbbVie, Agios, Celgene, Daiichi Sankyo, Jazz, MedImmune, and Notable Labs; and received research funding from AbbVie, Agios, Calithera, Celgene, and Daiichi Sankyo. A.T.F. is a consultant/advisor to Agios, AbbVie, Amgen, Astellas, Boston Biomedical, Celgene (Bristol-Myers Squibb), Daiichi Sankyo, Jazz, Takeda, and Trovagene and received research funding from Agios and Celgene (Bristol-Myers Squibb). A.S.M. is a consultant/advisor to AbbVie, Agios, Astellas, Jazz, and PTC Therapeutics. K.W.P. is an advisor to Agios, AbbVie, Astellas, Boston Biomedical, and Celgene/Bristol-Myers Squibb and received institutional research funding from AbbVie, Agios, Astellas, and Takeda. M.R.S. received research support from Astex, Incyte, Sunesis, Takeda, and TG Therapeutics; is a stockholder in/has ownership of Karyopharm; is a consultant/advisor to AbbVie, Bristol-Myers Squibb, Celgene, Merck, Ryvu, Sierra Oncology, and TG Therapeutics; and has patents with/received royalties from Boehringer Ingelheim. A.S.S. is on a speakers' bureau for Amgen, Celgene, and Stemline and is an advisor to Ämgen. R.M.S. is on a steering committee for Celgene; is an advisor to AbbVie, Actinium, Agios, Amgen, Astellas, BioLineRx, Celgene, Daiichi Sankyo, ElevateBio, GEMoaB, Janssen, MacroGenics, Novartis, Syndax, Takeda, and Trovagene; received clinical research funding from AbbVie, Agios, and Syndax; and is on data safety monitoring boards for Argenx, Celgene, and Takeda. E.S.W. is an advisor to Jazz, Pfizer, and Shionogi. H.D. is a consultant to/received honoraria from AbbVie, Agios, Amgen, Astellas, Astex, Celgene, Helsinn, Janssen, Jazz, Novartis, Oxford Biomedicals, and Roche and received institutional clinical research funding from Amgen, AROG, Bristol-Myers Squibb, Celgene, Jazz, Pfizer, and Sunesis. D.A.P. is a consultant to AbbVie, Agios, Amgen, Celgene, Celyad, Daiichi Sankyo, Forty Seven, Gilead, Pfizer, and Takeda; and received research funding from AbbVie and Pfizer. J.K.M. is on a speakers' bureau for Amgen, Celgene, Jazz, and Takeda; received honoraria from Celgene; is a consultant to Jazz, Pfizer, and Takeda; and is a stockholder in/has ownership in COTA. O.O. is an advisor to AbbVie, Celgene, and Impact Biomedicines and received institutional clinical research funding from AbbVie, Agios, AstraZeneca, Celgene, Daiichi Sankyo, Incyte, NS Pharma, and OncoTherapy Science. B.L. is an advisor to AbbVie, Agios, AIMM Therapeutics, Astellas, Astex, Celgene, Clear Creek Bio, F. Hoffman-La Roche, Frame, GEMoaB, and Oxford Biomedical. G.J.O. is a consultant to AbbVie, Agios, Amgen, Astellas,

Celgene, Daiichi Sankyo, Jazz, Novartis, Pfizer, and Roche and received research funding from BD, Celgene, Genentech, Johnson & Johnson, and Novartis. P.A.P. received honoraria from DAVA Oncology and France Foundation and is a member of an entity's board of directors/advisory committees/speakers' bureau for Celgene. M.R. is involved with the provision of services for Auron Therapeutics, Celgene, and Physicians' Education Resource and is a stockholder in/has ownership of Auron Therapeutics. M.G.F. is an employee of and stockholder in/has ownership of Bristol-Myers Squibb. F.L. is an employee of and stockholder in/ has ownership of Celgene International. A.F. is an employee of Bristol-Myers Squibb. S.N., S.C., C.A., and M.C. are employees of and stockholders in/have ownership of Agios. B.F., H.W., and L.H. were employees of and stockholders in/had ownership of Agios at the time of the study. B.W. is an employee of, stockholder in/has ownership of, and received patents/royalties from Agios. H.M.K. received research funding from AbbVie, Agios, Amgen, ARIAD, Astex, Bristol-Myers Squibb, Cyclacel, Daiichi Sankyo, ImmunoGen, Jazz, Novartis, and Pfizer and received honoraria from AbbVie, Actinium, Agios, Amgen, Pfizer, and Takeda. M.S.T. received patents/royalties from UpToDate; is a consultant/ member of an entity's board of directors/advisor to AbbVie, BioLineRx, Daiichi Sankyo, Delta-Fly Pharma, Jazz, KAHR, Nohla, Oncolyze, Orsenix, Rigel, Roche, and Tetraphase; and received research funding from AbbVie, ADC Therapeutics, BioSight, Cellerant, and Orsenix. C.S.S. declares no competing financial interests.

The current affiliation for K.W.P. is Penn Medicine, Philadelphia, PA.

ORCID profiles: C.D.D., 0000-0001-9003-0390; A.T.F., 0000-0003-4907-6447; A.S.M., 0000-0002-8971-2199; M.R.S., 0000-0003-3763-5504; E.S.W., 0000-0002-4515-8245; D.A.P., 0000-0001-6519-4860; B.L.,

REFERENCES

- Lancet JE, Uy GL, Cortes JE, et al. CPX-351 (cytarabine and daunorubicin) liposome for injection versus conventional cytarabine plus daunorubicin in older patients with newly diagnosed secondary acute myeloid leukemia. *J Clin Oncol.* 2018;36(26):2684-2692.
- Hills RK, Castaigne S, Appelbaum FR, et al. Addition of gemtuzumab ozogamicin to induction chemotherapy in adult patients with acute myeloid leukaemia: a meta-analysis of individual patient data from randomised controlled trials. *Lancet Oncol.* 2014;15(9): 986-996.
- Stone RM, Mandrekar SJ, Sanford BL, et al. Midostaurin plus chemotherapy for acute myeloid leukemia with a FLT3 mutation. N Engl J Med. 2017;377(5):454-464.
- Ward PS, Patel J, Wise DR, et al. The common feature of leukemia-associated IDH1 and IDH2 mutations is a neomorphic enzyme activity converting alpha-ketoglutarate to 2hydroxyglutarate. *Cancer Cell*. 2010;17(3): 225-234.
- DiNardo CD, Ravandi F, Agresta S, et al. Characteristics, clinical outcome, and prognostic significance of IDH mutations in AML. *Am J Hematol.* 2015;90(8):732-736.
- Paschka P, Schlenk RF, Gaidzik VI, et al. IDH1 and IDH2 mutations are frequent genetic alterations in acute myeloid leukemia and confer adverse prognosis in cytogenetically normal acute myeloid leukemia with NPM1 mutation without FLT3 internal tandem duplication. J Clin Oncol. 2010;28(22): 3636-3643.
- Medeiros BC, Fathi AT, DiNardo CD, Pollyea DA, Chan SM, Swords R. Isocitrate dehydrogenase mutations in myeloid malignancies. *Leukemia*. 2017;31(2):272-281.

- Dang L, White DW, Gross S, et al. Cancerassociated IDH1 mutations produce 2hydroxyglutarate. *Nature*. 2009;462(7274): 739-744.
- Lu C, Ward PS, Kapoor GS, et al. IDH mutation impairs histone demethylation and results in a block to cell differentiation. *Nature*. 2012; 483(7390):474-478.
- Saha SK, Parachoniak CA, Ghanta KS, et al. Mutant IDH inhibits HNF-4α to block hepatocyte differentiation and promote biliary cancer [published correction appears in Nature. 2015;528(7580):152]. Nature. 2014; 513(7516):110-114.
- Xu W, Yang H, Liu Y, et al. Oncometabolite 2hydroxyglutarate is a competitive inhibitor of α-ketoglutarate-dependent dioxygenases. *Cancer Cell*. 2011;19(1):17-30.
- TIBSOVO. (ivosidenib tablets), for oral use. Cambridge, MA: Agios Pharmaceuticals, Inc; 2019 [prescribing information].
- 13. IDHIFA. (enasidenib) tablets, for oral use. Summit, NJ: Celgene Corporation; 2019 [prescribing information].
- DiNardo CD, Stein EM, de Botton S, et al. Durable remissions with ivosidenib in IDH1mutated relapsed or refractory AML. N Engl J Med. 2018;378(25):2386-2398.
- Stein EM, DiNardo CD, Pollyea DA, et al. Enasidenib in mutant *IDH2* relapsed or refractory acute myeloid leukemia. *Blood*. 2017; 130(6):722-731.
- Kluk MJ, Lindsley RC, Aster JC, et al. Validation and implementation of a custom next-generation sequencing clinical assay for hematologic malignancies. J Mol Diagn. 2016;18(4):507-515.
- 17. Patwardhan A, Harris J, Leng N, et al. Achieving high-sensitivity for clinical

0000-0001-8982-5217; B.W., 0000-0001-9232-4456; H.M.K., 0000-0002-1908-3307.

Correspondence: Eytan M. Stein, Leukemia Service, Department of Medicine, Memorial Sloan Kettering Cancer Center, 530 East 74th St, New York, NY 10021; e-mail: steine@mskcc.org.

Footnotes

Submitted 2 June 2020; accepted 25 September 2020; prepublished online on *Blood* First Edition 5 October 2020. DOI 10.1182/ blood.2020007233.

*E.M.S. and C.D.D. are joint lead authors.

†H.M.K. and M.S.T. are joint lead senior authors.

Investigators interested in data sharing and collaboration should contact datasharing@agios.com.

The online version of this article contains a data supplement.

There is a Blood Commentary on this article in this issue.

The publication costs of this article were defrayed in part by page charge payment. Therefore, and solely to indicate this fact, this article is hereby marked "advertisement" in accordance with 18 USC section 1734.

applications using augmented exome sequencing. *Genome Med.* 2015;7(1):71.

- Dressman D, Yan H, Traverso G, Kinzler KW, Vogelstein B. Transforming single DNA molecules into fluorescent magnetic particles for detection and enumeration of genetic variations. Proc Natl Acad Sci USA. 2003;100(15): 8817-8822.
- Fan B, Dai D, DiNardo CD, et al. Clinical pharmacokinetics and pharmacodynamics of ivosidenib in patients with advanced hematologic malignancies with an IDH1 mutation. *Cancer Chemother Pharmacol.* 2020;85(5): 959-968.
- Li Y, Liu L, Gomez D, et al. Pharmacokinetics and safety of enasidenib following single oral doses in Japanese and Caucasian subjects. *Pharmacol Res Perspect*. 2018;6(6):e00436.
- Klco JM, Miller CA, Griffith M, et al. Association between mutation clearance after induction therapy and outcomes in acute myeloid leukemia. JAMA. 2015;314(8): 811-822.
- Jongen-Lavrencic M, Grob T, Hanekamp D, et al. Molecular minimal residual disease in acute myeloid leukemia. N Engl J Med. 2018; 378(13):1189-1199.
- 23. Löwenberg B, Ossenkoppele GJ, van Putten W, et al; for the Dutch-Belgian Cooperative Trial Group for Hemato-Oncology (HOVON), German AML Study Group (AMLSG), and Swiss Group for Clinical Cancer Research (SAKK) Collaborative Group. High-dose daunorubicin in older patients with acute myeloid leukemia [published correction appears in N Engl J Med. 2010;362(12):1155]. N Engl J Med. 2009;361(13):1235-1248.
- Lancet JE, Cortes JE, Hogge DE, et al. Phase 2 trial of CPX-351, a fixed 5:1 molar ratio of cytarabine/daunorubicin, vs

cytarabine/daunorubicin in older adults with untreated AML. *Blood*. 2014;123(21): 3239-3246.

- 25. Mims AS, Kohlschmidt J, Borate U, et al. A precision medicine hierarchical classification developed using variant allele frequency (VAF) for treatment of older patients (pts) with acute myeloid leukemia (AML): Alliance Clinical Trials in Oncology (Alliance) Historical Patient Control [abstract]. *Blood*. 2018;132(suppl 1). Abstract 617.
- 26. Paschka P, Schlenk RF, Weber D, et al. Outcome of patients with refractory or relapsed AML with IDH1 and IDH2 mutations

after conventional salvage therapy: a study of the German-Austrian AML study group (AMLSG) [abstract]. *Haematologica*. 2016; 101(suppl 1). Abstract S809.

- Choe S, Wang H, DiNardo CD, et al. Molecular mechanisms mediating relapse following ivosidenib monotherapy in *IDH1*-mutant relapsed or refractory AML. *Blood Adv.* 2020; 4(9):1894-1905.
- Amatangelo MD, Quek L, Shih A, et al. Enasidenib induces acute myeloid leukemia cell differentiation to promote clinical response. *Blood.* 2017;130(6):732-741.
- 29. Appelbaum FR, Gundacker H, Head DR, et al. Age and acute myeloid leukemia. *Blood*. 2006;107(9):3481-3485.
- 30. Kantarjian H, O'brien S, Cortes J, et al. Results of intensive chemotherapy in 998 patients age 65 years or older with acute myeloid leukemia or high-risk myelodysplastic syndrome: predictive prognostic models for outcome. *Cancer.* 2006;106(5):1090-1098.
- 31. Thein MS, Ershler WB, Jemal A, Yates JW, Baer MR. Outcome of older patients with acute myeloid leukemia: an analysis of SEER data over 3 decades. *Cancer*. 2013;119(15): 2720-2727.