IMMUNOBIOLOGY AND IMMUNOTHERAPY

CME Article

HLA associations, somatic loss of HLA expression, and clinical outcomes in immune aplastic anemia

Yoshitaka Zaimoku,^{1,*} Bhavisha A. Patel,^{1,*} Sharon D. Adams,² Ruba Shalhoub,³ Emma M. Groarke,¹ Audrey Ai Chin Lee,² Sachiko Kajigaya,¹ Xingmin Feng,¹ Olga Julia Rios,¹ Holly Eager,¹ Lemlem Alemu,¹ Diego Quinones Raffo,¹ Colin O. Wu,³ Willy A. Flegel,² and Neal S. Young¹

¹Hematology Branch, National Heart, Lung, and Blood Institute, National Institutes of Health, Bethesda, MD; ²Department of Transfusion Medicine, NIH Clinical Center, National Institutes of Health, Bethesda, MD; and ³Office of Biostatistics Research, National Heart, Lung, and Blood Institute, National Institutes of Health, Bethesda, MD

KEY POINTS

- In patients with AA, somatic mutations in HLA genes correlated with functionality, as measured by cell surface expression.
- Loss of HLA expression and specific HLA genotypes correlated with clinical manifestations, especially age of onset and clonal evolution.

Immune aplastic anemia (AA) features somatic loss of HLA class I allele expression on bone marrow cells, consistent with a mechanism of escape from T-cell-mediated destruction of hematopoietic stem and progenitor cells. The clinical significance of HLA abnormalities has not been well characterized. We examined the somatic loss of HLA class I alleles and correlated HLA loss and mutation-associated HLA genotypes with clinical presentation and outcomes after immunosuppressive therapy in 544 AA patients. HLA class I allele loss was detected in 92 (22%) of the 412 patients tested, in whom there were 393 somatic HLA gene mutations and 40 instances of loss of heterozygosity. Most frequently affected was HLA-B*14:02, followed by HLA-A*02:01, HLA-B*40:02, HLA-B*08:01, and HLA-B*07:02. HLA-B*14:02, HLA-B*40:02, and HLA-B*07:02 were also overrepresented in AA. High-risk clonal evolution was correlated with HLA loss, HLA-B*14:02 genotype, and older age, which yielded a valid prediction model. In 2 patients, we traced monosomy 7 clonal evolution from preexisting clones harboring somatic mutations in HLA-A*02:01 and HLA-B*40:02. Loss of HLA-B*40:02 correlated

with higher blood counts. *HLA-B*07:02* and *HLA-B*40:01* genotypes and their loss correlated with late-onset of AA. Our results suggest the presence of specific immune mechanisms of molecular pathogenesis with clinical implications. HLA genotyping and screening for HLA loss may be of value in the management of immune AA. This study was registered at clinicaltrials.gov as NCT00001964, NCT00061360, NCT00195624, NCT00260689, NCT00944749, NCT01193283, and NCT01623167.



Medscape Continuing Medical Education online

In support of improving patient care, this activity has been planned and implemented by Medscape, LLC and the American Society of Hematology. Medscape, LLC is jointly accredited by the Accreditation Council for Continuing Medical Education (ACCME), the Accreditation Council for Pharmacy Education (ACPE), and the American Nurses Credentialing Center (ANCC), to provide continuing education for the healthcare team.

Medscape, LLC designates this Journal-based CME activity for a maximum of 1.00 AMA PRA Category 1 Credit(s)TM. Physicians should claim only the credit commensurate with the extent of their participation in the activity.

Successful completion of this CME activity, which includes participation in the evaluation component, enables the participant to earn up to 1.0 MOC points in the American Board of Internal Medicine's (ABIM) Maintenance of Certification (MOC) program. Participants will earn MOC points equivalent to the amount of CME credits claimed for the activity. It is the CME activity provider's responsibility to submit participant completion information to ACCME for the purpose of granting ABIM MOC credit.

All other clinicians completing this activity will be issued a certificate of participation. To participate in this journal CME activity: (1) review the learning objectives and author disclosures; (2) study the education content; (3) take the post-test with a 75%

minimum passing score and complete the evaluation at **http://www.medscape.org/journal/blood**; and (4) view/print certificate. For CME questions, see page 2891.

Disclosures

Associate Editor Jeanne Hendrickson, CME questions author Laurie Barclay, freelance writer and reviewer, Medscape, LLC, and the authors declare no competing financial interests.

Learning objectives

Upon completion of this activity, participants will:

- 1. Determine somatic loss of HLA class I alleles in a study of 544 patients with immune aplastic anemia (AA)
- 2. Describe allele frequencies of HLA-A and HLA-B associated with somatic loss in a study of 544 patients with immune AA
- 3. Identify correlations of HLA alleles and HLA loss with clinical presentation and outcome after IST in a study of 544 patients with immune AA

Release date: December 30, 2021; Expiration date: December 30, 2022

Introduction

Immune aplastic anemia (AA) is caused by T cells that destroy hematopoietic stem cells (HSCs), and marrow failure is successfully treated with hematopoietic cell transplantation (HCT) or immunosuppressive therapy (IST).¹ Eltrombopag (EPAG) combined with IST yielded higher hematologic responses and survival compared with IST alone,² but long-term outcomes such as relapse and clonal evolution remain clinically problematic and biologically not well understood.

The immune pathophysiology of AA has been, in part, inferred from frequent somatic loss of HLA class I alleles. Increased frequency of some HLA alleles has been reported in AA patients of various ethnicities,^{3–14} and has been confirmed in a recent genome-wide association study.¹⁵ Somatic loss of HLA class I alleles may result from copy-neutral chromosome 6p loss of heterozygosity (6p LOH)^{11,16,17} or acquired inactivating HLA gene mutations.^{18,19} A limited set of *HLA-A* and *HLA-B* alleles are more likely to acquire somatic mutations^{18,19}; a T-cell line specific for a missing HLA class I allele has been isolated from an AA patient.²⁰ Loss in a recurrently mutated HLA allele may characterize a specific immune pathogenesis and associated clinical manifestations. Previous studies suggested better IST responses

and survival in patients with HLA loss^{11,18,21} and poor outcomes, including frequent clonal evolution, in patients who harbored HLA alleles related to somatic mutations, irrespective of somatic loss of an HLA allele.¹⁹

Clonality is common in AA.²² Paroxysmal nocturnal hemoglobinuria (PNH)-clones, cells deficient in glycosylphosphatidylinositol (GPI)-anchored proteins due to acquired *PIGA* mutations, are most frequent. PNH is closely related to immune marrow failure, and the GPI anchor itself may be a target of immune attack.^{23,24} Somatic mutations are also present in genes recurrently mutated in myelodysplastic syndromes or acute myeloid leukemia (AML), especially *DNMT3A*, *ASXL1*, and *BCOR*.¹⁷ However, the allelic burden of these clones in AA are usually small, clones remain stable for years, and affected cells infrequently drive evolution to myeloid neoplasms, which are usually characterized by complete or partial loss of chromosome 7.^{1,25,26} Of all clonal associations, the mechanism of HLA loss is most clearly related to escape from immune cell destruction.

Our aim was to clarify and enlarge on the clinical significance of HLA class I allele loss and recurrently mutated genotypes in a large cohort of patients with immune AA.



Figure 1. Patients and clinical parameters studied. Correlations of clinical parameters with HLA genotype and HLA loss were tested in subgroups based on the availability of data.

Table 1. Clinical characteristics of patients

	n	Value, n (%)
Age, y	544	29 (18-54)
Range		2-82
Male sex	544	304 (56)
Ethnicity	544	
White		275 (51)
African American		120 (22)
Hispanic		109 (20)
Asian		35 (6)
Other		5 (1)
Disease status	544	
Treatment naïve severe AA		515 (95)
Relapsed or refractory AA		29 (5)
Pretreatment blood values		
Neutrophil count, ×10 ⁹ /L	515	0.29 (0.09-0.51)
Lymphocyte count, ×10 ⁹ /L	515	1.27 (0.92-1.63)
Reticulocyte count, ×10 ⁹ /L	515	15.3 (6.5-32.0)
Platelet count, ×10 ⁹ /L	515	9 (5-13)
Thrombopoietin, ng/mL	140	2610 (2220-3080)
PNH clone ≥1%	470	177 (38)
hATG-based IST	416	
hATG and CsA		102 (25)
hATG, CsA, and MMF		103 (25)
hATG, CsA, and rapamycin		35 (8)
hATG, CsA, and EPAG		176 (42)
Response to IST	416	
Overall response		293 (70)
Complete response		101 (24)
Clonal evolution	416	
High-risk clonal evolution		31 (7)
Low-risk clonal evolution		26 (6)

Values are n (%) or median (IQR).

CsA, cyclosporine; MMF, mycophenolate mofetil.

Methods

Study design and participants

A total of 544 patients with AA, aged 2 years or older, treated at the NIH Clinical Center, Bethesda, MD, between May 1999 and August 2019 were enrolled in this retrospective study. Patients were enrolled in various IST protocols approved by the Institutional Review Board of the National Heart, Lung, and Blood Institute (clinicaltrials.gov: NCT00001964, NCT00061360, NCT00195624, NCT00260689, NCT00944749, NCT01193283, and NCT01623167), which entailed informed consent. All horse antithymocyte globulin (hATG)-based IST protocols have been published^{2,27–29} and briefly described in the supplemental Methods (available on the *Blood* Web site).

Definitions

All definitions have been consistent across protocols. Severe AA was diagnosed when at least 2 of the following 3 criteria were met: a neutrophil count <0.5 × 10⁹/L, a reticulocyte count <60 × 10⁹/L, and a platelet count <20 × 10⁹/L. Hematologic responses were assessed 6 months after institution of hATG: an overall response was defined as blood counts no longer meeting the above stated criteria for severe AA; a complete response required a neutrophil count of at least 1.0 × 10⁹ cells/L, a hemoglobin level of at least 10 g/dL, and a platelet count of at least 100 × 10⁹/L. High-risk clonal evolution was defined as the acquisition of either chromosome 7 abnormalities, complex cytogenetics, myelodysplastic syndrome, or AML, and low-risk clonal evolution was the other cytogenetic change, as previously published.³⁰

Detection of HLA class I loss

Detailed methods for the detection of HLA loss are described in the supplemental Methods. In brief, HLA loss was assessed by either HLA flow cytometry plus deep nucleotide sequencing or deep sequencing alone. HLA flow cytometry was performed using cryopreserved peripheral blood mononuclear cells (PBMCs) stained with HLA-A and HLA-B allele-specific monoclonal antibodies and assessed in CD33^{hi} monocytes. Deep sequencing of HLA class I genes was performed using sorted peripheral blood cell subpopulations, including HLA allelelacking and expressing monocytes or using whole blood for the deep sequencing-only samples. HLA class I genes were enriched by locus-specific long-range PCR, as previously described.³¹ To precisely detect low-frequency mutations, we used hla-mapper (version 2.3),³² and used thresholds defined by base-position error rates in T-cell samples, as previously published for non-HLA genes.^{33,34} We inferred 6p LOH using read counts of allelespecific single-nucleotide polymorphisms in HLA genes.

Statistics

Statistical analyses were performed using R (version 4.0.3) with EZR package (version 1.54)³⁵; individual statistical methods are described in the supplemental Methods. HLA allele frequencies in patients were compared with those in in the National Marrow Donor Program (NMDP) datasets according to 4 wide ethnic groups including Whites, African Americans, Hispanics, and Asians,³⁶ and the results were combined by fixed-effect meta-analysis to elucidate consistent impacts across ethnic groups.

Data sharing statement

All somatic HLA gene mutation information is available in the supplemental $\ensuremath{\mathsf{Data}}$.

Results

Patients

All 544 patients had HLA allele genotyping performed during evaluation on clinical protocols; 515 of the 544 were treatmentnaïve severe AA, and 416 of 515 were initially treated with hATG-based standard IST (Figure 1). Clinical characteristics of the patients are summarized in Table 1. The median time to clonal evolution and observation period after hATG treatment was 25 (interquartile range [IQR], 6 to 41) months, and 41 (IQR, 22 to 61) months, respectively. Of the 544 total subjects, 412 were tested for the presence of HLA allele loss.



Figure 2. HLA class I allele loss. (A) Number of patients with HLA loss due to HLA gene mutations, 6p LOH, or both in a total of 412 subjects tested. (B) The number and proportion of patients who lacked individual HLA alleles (left); the total and mean number of clones with HLA gene mutations and 6p LOH among patients who lost respective HLA alleles (right). The number of patients tested for the presence of HLA loss is shown in parenthesis. (C) Combined clone size of cells with HLA loss (upper) and the clone size of individual HLA gene mutations and 6p LOH among cells with HLA loss (lower). (D) Positions and types of somatic inactivating mutations in HLA-A (n = 123) and HLA-B (n = 267). Hotspot mutations are noted on the figure.



Downloaded from http://ashpublications.net/blood/article-pdf/138/26/2799/1890463/bloodbld2021012895.pdf by guest on 08 June 2024

Figure 3. High-risk clonal evolution. Cumulative incidence of high-risk clonal evolution, defined as the acquisition of either chromosome 7 abnormalities, complex cytogenetics, myelodysplastic syndrome, or AML after institution of hATG-based IST, are shown according to the presence or absence of *HLA-B*14:02* genotype (A), the presence or absence of HLA loss (B), age groups (C), and 3 risk groups of the prediction model for high-risk clonal evolution incorporating *HLA-B*14:02* genotype, HLA loss, and age (D): a high-risk group, any HLA risk (*HLA-B*14:02* genotype or HLA loss) present and aged 40 years or older; a low-risk group, no HLA risk present and aged less than 40 years; and an intermediate-risk group, not meeting the criteria for groups of high-risk nor low-risk.

Detection of HLA class I gene mutations

Our sequencing method allowed us to identify variant allele frequency as low as 0.3% in more than 50% of base positions of single nucleoide variants and 90% of indels (supplemental Figure 1). The accuracy of the method was supported by the following observations and experiments: HLA class I gene mutations were not detected from additional 16 control T-cell samples nor from monocytes of patients expressing respective HLA alleles as determined by flow cytometry (except for missense mutations and 1 intronic mutation); somatic mutations detected from patient samples were generally predicted to inactivate affected HLA alleles, irrespective of variant allele frequencies and samples (supplemental Figure 2), in contrast to pseudomutations created by PCR errors (supplemental Figure 3); serial mixing of HLA-lacking monocytes and HLA-expressing T cells showed a linear correlation between variant allele frequency and mixing ratio (supplemental Figure 4). Low-frequency mutations were confirmed by droplet digital PCR (supplemental Figure 5), and functional HLA loss was demonstrated in the large cohort of samples also studied by flow cytometry. Small cell populations deficient in surface HLA-A or HLA-B antigens were not detected in 41 healthy individuals by flow cytometry at a threshold of 0.1%, consistent with the very infrequent spontaneous inactivating mutations in leukocytes of healthy individuals.^{37,38} Deep



Figure 4. Clonal evolution of monosomy 7 from HLA class I allele-lacking clones. (A) A 56-year-old female at the time of severe AA diagnosis, who received hATG, CsA, and mycophenolate mofetil (MMF), and multiple rounds of salvage IST, including rabbit ATG, daclizumab and danazol, and alemtuzumab, without adequate response, had clonal evolution to monosomy 7 approximately 7.5 years after her initial diagnosis of AA and died 6 months later. Her blood, sampled 4 years after the institution of initial IST, contained a PNH clone and HLA-A0201-lacking cells (A0201⁺), consistent with the diagnosis of immune AA. Multiple clones with *HLA-A*02:01* deactivating mutations constituted sorted A0201⁻ cells (multiple clones with missense mutations also did A0201⁺ cells), from which chromosome 7 deletion was not detected by fluorescent in situ hybridization (FISH). After 3.5 years, virtually all hematopoiesis was replaced by a preexisting A0201⁻ clone with an intron 1 mutation (intron 1^{mut}; c.74-19A>G) that had acquired monosomy 7. (B) A 72-year-old man at the time of severe AA diagnosis was treated with hATG, CsA, and EPAG with a partial response at 6 months but had declining cytopenia which responded to reinitiation of CsA. About 2.5 years from the initial diagnosis and IST treatment, BM cytogenetics revealed monosomy 7, and myelodysplastic syndrome was diagnosed. The patient received symptomatic treatment (infrequent blood transfusions twice a year) and died of pneumonia 3.5 years later. HLA flow cytometry of cryopreserved cells revealed a gradual expansion of a clone partially lacking HLA-B4002 (B4002^{dim}), which was attributed to an *HLA-B*40:02* intron 1 mutation (c.74-9C-A). A small monosomy 7 clone that was not visible by conventional BM cytogenetics by Giersa-banding was detected by FISH from the sorted B4002^{dim} cells at least 2 years before clinical diagnosis of monosomy 7. NA, not assessed (due to deficient cryopreserved cells for analysis). See also supplemental Tables 8 and 9.

sequencing could detect 6p LOH at a clone size of 15% (supplemental Figure 6).

Loss of HLA class I allele

HLA class I allele loss was detected in 92 (22%) of 412 patients (Figure 2A) by flow cytometry plus deep sequencing of HLA allele-expressing and -deficient cells (51 of 201) (supplemental Figures 7 and 8) or only whole blood deep sequencing (41 of 211). A total of 393 somatic HLA gene mutations (supplemental Data) were identified from 92 patients, in *HLA-A* (n = 124), *HLA-B* (n = 267), and *HLA-C* (n = 2); 29 of the 92 patients also had 6p LOH and a further 11 patients had 6p LOH only. Multiple mutant clones (median, 5 [range, 2 to 20]) with HLA gene mutations or with 6p LOH coexisted in 62 of the 92 patients, almost always restricted to a single *HLA-A* or *HLA-B* allele in individuals. *HLA-B*14:02, *HLA-A*02:01, *HLA-B*40:02, *HLA*-

B*07:02, and HLA-B*08:01 were frequently absent (Figure 2B). The median proportion of cells with HLA loss (combined size of all clones with HLA gene mutations and 6p LOH) was 16.3% (range, 0.2% to 100%; IQR, 2.4% to 62%) (Figure 2C).

Several "hotspots" were observed (Figure 2D; supplemental Figure 9): nonsense mutation R7X (n = 55) and frameshift mutation R7fs (n = 28) in both HLA-A and HLA-B; and start codon mutation (n = 15) and missense mutation T118P (n = 11) in HLA-B. Mutations in the start codon were detected only for HLA- $B^{*}40:02$, HLA- $B^{*}27:05$, and HLA- $B^{*}13:01$, which lack the second start signal in the fourth codon (supplemental Figure 10).

When assessed by flow cytometry, all frameshift mutations (n = 76), nonsense mutations (n = 72), intronic mutations (n = 24), and even synonymous mutations reduced cell surface expression

Odds ratio of allele frequency (95% CI)		2.44 (1.91-3.12)	1.88 (1.36-2.61)	1.42 (1.19-1.70)	0.99 (0.86-1.15)	0.92 (0.72-1.17)	
Allele frequency, % (phenotype frequency, %) White African American Hispanic Asian	Asian	Control n = 568 597	0.2 (0.3)	2.4 (4.8)	2.9 (5.8)	8.4 (16.2)	1.8 (3.5)
		AA n = 35	0.0 (0)	11.4 (20.0)	1.4 (2.9)	7.1 (11.4)	0 (0)
	anic	Control n = 712 764	4.2 (8.3)	5.0 (9.7)	5.8 (11.2)	21.0 (37.5)	4.2 (8.2)
	Hisp	AA n = 109	7.8 (14.7)	7.3 (13.8)	8.3 (16.5)	17.0 (29.4)	4.6 (9.2)
	African American	Control n = 505 250	2.2 (4.3)	0.3 (0.7)	7.3 (14.0)	12.2 (23.0)	3.7 (7.2)
		AA n = 120	5.0 (10.0)	0.4 (0.8)	8.8 (15.8)	12.9 (24.2)	3.8 (7.5)
	ite Control	Control n = 3 912 440	2.7 (5.4)	1.3 (2.7)	12.5 (23.4)	27.2 (46.9)	10.6 (20.0)
	Wh	AA n = 275	7.3 (12.7)	2.4 (4.7)	17.6 (32.4)	28.4 (50.5)	9.5 (17.8)
HLA allele		HLA-B*14:02	HLA-B*40:02	HLA-B*07:02	HLA-A*02:01	HLA-B*08:01	

Table 2. HLA allele frequencies in patients with AA and healthy individuals in the NMDP dataset

of affected HLA alleles (except for 1 mutation in HLA-A intron 7). Missense mutations were inconsistently associated with loss of expression: 21 of 69 expressed HLA antigens on the cell surface, and for 48, there was a loss of expression (supplemental Figure 11A); 6 of the 21 missense mutations that were expressed on cell surface located to the peptide-binding groove (S37F [HLA-B*08:01], F57L [HLA-B*40:01], L102Q [HLA-B*40:01], H117Y [HLA-A*02:01], T118P [HLA-B*40:02], and D126Y [HLA-B*14:02]) (supplemental Figure 11B), where they might interact with antigens presented by HLA alleles; another 13 missense mutations were located in the α -3 domain where they could associate with CD8 molecules on cytotoxic T cells (supplemental Figure 11C).¹⁸ Eleven missense mutations and 2 synonymous mutations were predicted to create alternative splice sites and generate truncated proteins (supplemental Table 1). Of note, 12 of 24 functionally confirmed intronic mutations and 7 unconfirmed mutations did not involve donor 5'-GT nor acceptor AG-3' canonical splice sites (supplemental Table 2): 8 of these mutations proximate to the splice sites have been predicted to reduce splicing efficiency; 6 mutations at 19 to 21 nucleotides from the 3' splice site were found to impair the consensus branchpoint sequences, CTGAC, CTCAC, or CTCAG, also critical for splicing³⁹; the remaining 5 mutations in the middle of HLA-B intron 3 (c.619 + 123A>G and c.619 + 133C>G) have been reported to create alternative 5' splice sites.¹⁸ One mutation in intron 7 of HLA-A and 2 mutations in cytoplasmic tail (exon 7) of HLA-C (S357A and E359del) were assumed to be passenger mutations, as these clones were small (1.0%, 3.0%, and 0.6%) and multiple clones with HLA-B mutations (14, 7, and 6) were also detected from the 3 patients.

The proportions of HLA allele-lacking cells in total monocytes were reduced in response to IST plus EPAG in all 18 patients sequentially tested, irrespective of hematological responses, and in 2 patients, HLA allele-lacking cells had disappeared at 6 months; they were thereafter stable over many years in 13 patients (supplemental Figure 12), as has been reported.¹⁸

HLA loss and other hematopoietic clones

Allele frequency is copy number/2n. The phenotype frequency in the NMDP donors was estimated by the equation of (phenotype frequency) = $1 - (1 - [allele frequency))^2$

HLA loss was relatively infrequent among patients with other types of clonal hematopoiesis compared with patients without clonal hematopoiesis, including PNH clones (18% [24 of 136] vs 24% [54 of 225], P = .19) (supplemental Figure 13A; supplemental Table 4) and somatic myeloid neoplasm gene (20% [4 of 20] vs 28% [33 of 119], P = .25) (supplemental Figure 14). When PNH clones and HLA loss were simultaneously analyzed by sensitive flow cytometry, PNH clones at a median frequency of 4.6% (range, 0.1% to 39%) in total monocytes were present in 27 of the 51 patients (53%) with concomitant HLA loss; such PNH clones always retained cell surface expression of HLA class I (supplemental Figure 13B-C). Neither PNH clone nor clones mutated in myeloid malignancy-related genes correlated with loss of specific HLA alleles.

Clinical significance of HLA class I allele loss and individual alleles

Recurrently inactivated class I alleles *HLA-B*14:02*, *HLA-B*40:02*, and *HLA-B*07:02* were significantly overrepresented in patients with AA (Table 2; supplemental Table 3).

	Univariate model			Multivariate model 1			Multivariate model 2		
	HR	95% CI	Р	HR	95% CI	Р	HR	95% CI	Р
Age ≥40 y	4.00	1.90-8.40	.00026	4.87	2.16-11.0	.00014	4.73	2.10-10.7	.00018
HLA-B*14:02 genotype	2.83	1.32-6.10	.0077	2.47	1.01-6.05	.048			
HLA loss	3.68	1.77-7.66	.00050	2.70	1.20-6.08	.017			
Combined HLA risk†	4.44	2.04-9.68	.00018				4.26	1.02-9.48	.00038

Table 3. Fine-Gray proportional hazard regression for high-risk clonal evolution

†Presence of HLA loss or HLA-B*14:02 genotype.

Patient age did not correlate with HLA loss in general, but did correlate with 2 mutation-associated genotypes HLA-B*07:02 (median age [IQR], 46 [22-64] years vs 27 [18-49] years, P < .0001) (supplemental Table 4) and HLA-B*40:01 (median age [IQR], 46 [27-65] years vs 28 [18-53] years, P = .0022), and therefore, HLA-B*40:01 was exclusively overrepresented among patients 40 years or older compared with healthy individuals (odds ratio 1.56 [95% CI, 1.06-2.52]). Moreover, 7 and 2 patients who lost HLA-B*07:02 and HLA-B*40:01 were also older, with a median age of 65 and 68 years, respectively (supplemental Figure 15; supplemental Table 5). These findings suggested an age-associated immune pathogenesis involving specific HLA-B alleles.

Correlations of HLA loss and HLA genotypes with pretreatment blood values and hematologic responses to IST were weak (supplemental Table 4). However, loss of HLA-B*40:02 and certain minor alleles was related to higher blood counts, and all patients who lost these alleles responded to IST (supplemental Figure 16; supplemental Table 5). These relationships are consistent with the predictive value of reticulocytes for IST response,^{40,41} and the generally high response rate to IST in Japanese patients with HLA loss, in whom HLA-B*40:02 was most frequently inactivated,^{18,42} but HLA-B*14:02 and HLA-B*08:01 genotypes are virtually absent.43

High-risk clonal evolution was correlated with the most frequently inactivated allele HLA-B*14:02 (P = .0092) (Figure 3A), as previously suggested by Babushok et al,¹⁹ and with HLA loss (P = .00040) (Figure 3B), especially with the loss of HLA-A*02:01 and HLA-B*14:02 (supplemental Figure 17A; supplemental Table 5). In particular, HLA-B*14:02 genotype was strongly correlated among patients without HLA loss (P = .0072), while none of 43 patients who did not carry any of the mutation-associated HLA genotypes exhibited highrisk clonal evolution (supplemental Figure 17B-C); this suggested that the presence of specific HLA class I-mediated immunity, rather than HLA loss itself, predisposed to highrisk clonal evolution. A combined HLA risk, defined as the presence of HLA loss or HLA-B*14:02 genotype (irrespective of loss), yielded a better prediction factor for high-risk clonal evolution than did either HLA loss alone or HLA-B*14:02 alone (supplemental Figure 17D). The HLA associations with high-risk clonal evolution were independent of patient age (Table 3; supplemental Table 6), a known predictor of secondary myeloid malignancies (Figure 3C).^{44,45} A prediction model for high-risk clonal evolution incorporating the combined HLA risk and age 40 years or older clearly divided patients into 3 risk groups (Figure 3D; supplemental Figure 17E), which was validated in various subgroups of patients (supplemental Figure 18).

Origin of monosomy 7

We further studied somatic HLA class I gene mutations in monosomy 7 clones evolved in 11 patients with immune AA, using cryopreserved PBMCs; 4 of the patients had HLA allele lacking-leukocytes present before clonal evolution, and 3 did not, and in the other 4 cases, samples prior to the evolution event were not available. In 2 patients with known HLA allele loss, monosomy 7 clonal evolution was associated with the expansion of clones with a preexisting mutation within intron 1 of HLA-A*02:01 (c.74-19A>G) and HLA-B*40:02 (c.74-9C>A); before clonal evolution, their blood was supported by 15 and 19 clones with HLA gene mutations including the intron 1 mutant clone and a PNH clone (Figure 4; supplemental Tables 7-9). The clone with an HLA-B*40:02 intron 1 mutation could be isolated by flow cytometry, as this mutation partially expressed HLA-B4002 (B4002^{dim}); fluorescent in situ hybridization (FISH) confirmed monosomy 7 was a subclone of this B4002^{dim} intron 1 mutant clone. A monosomy 7 clone accounting for 3% of total monocytes and <1% of total mononuclear cells was detected by FISH from the sorted B4002^{dim} clone (11% in B4002^{dim} cells) at least 2 years before clinical diagnosis of monosomy 7, although bone marrow (BM) cytogenetics at this time and 1 year later were normal. In the other 9 patients, somatic mutations were not detected from HLA-A, HLA-B, and HLA-C genes in the monosomy 7 clones. Similar intron 1 mutations in HLA class I genes were observed in 2 other patients with AA, and 1 of the 2 acquired 7p deletion and t(7;17)(g22;g11.2), implying a possible relationship of this intron 1 mutation with chromosome 7 abnormalities. No PNH clones in 9 patients expanded after monosomy 7 clonal evolution (supplemental Figure 13D), consistent with previous observations.²⁵

Discussion

Our retrospective study of HLA genes in many hundreds of patients with immune AA demonstrated novel findings regarding HLA allele frequency, loss of HLA class I alleles, and important clinical correlations.

We confirmed the high prevalence of HLA class I alleles recurrently inactivated by somatic mutations across 4 major ethnic groups in the United States, which included previously reported *HLA-B*14:02* and *HLA-B*40:02*,^{18,19,46} and an unreported allele *HLA-B*07:02*. Further, correlations of HLA loss with patient age at disease onset, peripheral blood counts, IST-response, and high-risk clonal evolution suggest specific pathogeneses related to certain HLA class I alleles.

In cancer generally, HLA allele loss is frequent and represents an escape from immune surveillance, 47,48 which clinically correlates with poor outcomes after treatment with immune checkpoint inhibitors⁴⁹ and relapse of AML after HCT.⁵⁰ While HLA loss in immune AA is protective of the stem cell under self-destructive immune attack, it also should allow escape from immune surveillance and therefore of malignant clonal evolution. Indeed, we observed this association and tracked it in 2 patients. Nevertheless, high-risk clonal evolution is more likely consequent to HLA class I-mediated destruction of BM cells than due to immune escape, as HLA-B*14:02, the most frequently mutated and overrepresented class I allele in AA, was associated with high-risk clonal evolution irrespective of HLA mutation status, as previously suggested.¹⁹ Most (but not all) monosomy 7 clonal evolution arose from cells with intact HLA genes; high-risk evolution was not observed in any patient without an HLA allele prone to loss. A simple prediction model incorporating HLA loss and HLA-B*14:02, as a convincing indicator of class I-mediated pathogenesis, in addition to age, stratified the incidence of high-risk clonal evolution, even within subgroups. Monitoring for clonal evolution after IST is desirable in patients with HLA allele loss or HLA-B*14:02; early HCT could be considered for the high-risk group patients aged 40 years or older rather than IST as generally recommended for the aged individuals.⁵¹

The underlying mechanisms responsible for the predominant development of monosomy 7 in immune AA (and other marrow failure syndromes) remain elusive.⁵² We detected a small monosomy 7 clone not visible by conventional BM cytogenetics at least 2 years before the clinical diagnosis of clonal evolution in 1 patient; extended time periods may be necessary for aneuploid clones (and somatically mutated clones⁵³) to expand and become clinically evident, either by accumulating additional genetic aberrations or altered environmental selection. Relatively infrequent mutations in non-HLA genes among patients with HLA loss argue against HLA gene mutations arising due to an intrinsically higher somatic mutation rate or from genetic drift in a paucicellular stem cell pool.²⁴ More sensitive methods would benefit both the monitoring and deeper understanding of secondary myeloid malignancies in AA.

PNH clones and HLA allele-lacking clones are both assumed to represent immune escape of HSCs from immune cell attack and destruction, manifested by the recurrent presence of multiple clones with somatic mutations in *PIGA*⁵⁴ or HLA class I genes.¹⁸ PNH and HLA absent clones were always mutually exclusive, as in the 27 patients who had both types of clones. Loss of surface

GPI-anchored proteins may confer no additional survival advantage on HLA allele loss and the reverse. A remarkable difference between the 2 types of escape clones was their relationship to clonal evolution, which occurred regardless of the presence of PNH clones but much more frequently in patients with HLA allele loss. Malignant clones always evolved from non-PNH phenotype cells as previously reported,²⁵ but they did arise from clones with HLA allele loss.

Our deep sequencing method for HLA class I genes allowed detection of somatic mutations at an allele frequency as low as 0.3%. Many coexisting small clones with somatic mutations in HLA-A and HLA-B alleles were identified, most of which functionally confirmed by flow cytometry. Even intronic mutations not involving canonical splice sites and a synonymous mutation were found to reduce cell surface expression of HLA class I alleles; such mutations likely have been underreported in studies using exome or RNA sequencing data and in those lacking functional validation. A limitation of our sequencing method was the lower sensitivity for 6p LOH detection, the result of unbalanced amplification of heterozygous HLA alleles, and the use of whole blood for some patients. Underestimation of 6p LOH would not be predicted to affect the overall detection of HLA loss, as most 6p LOH clones coexist with clones with inactivating mutations in an HLA allele, but detection of such small clones may be of value if they predicted secondary myeloid malignancies.

In summary, our study suggested the presence of distinct pathogenic pathways related to HLA genotype and loss in immune AA. Most important in the clinic was the relationship between high-risk malignant clonal evolution and class I-mediated immunity, as for the most frequently inactivated *HLA-B*14:02* genotype and HLA loss in general, which provide a simple prediction tool for this serious complication.

Acknowledgments

This work was supported by the Intramural Research Program of the National Heart, Lung, and Blood Institute (NIH), and the NIH Clinical Center (ZIC CL002128). NCT01623167 is funded by Novartis and previously by GSK through a cooperative research and development agreement (CRADA). The authors acknowledge the support of the National Heart, Lung, and Blood Institute core facilities and the CCR Genomics Core facility: cell sorting was performed at the Flow Cytometry Core Facility, deep sequencing was performed at the DNA Sequencing and Genomics Core Facility, bioinformatics analysis was performed at the Bioinformatics and Computational Biology Core Facility, and droplet digital PCR was performed at the CCR Genomics Core facility.

Authorship

Contribution: Y.Z., S.K., and N.S.Y. designed research; B.A.P., O.J.R., H.E., E.M.G., and N.S.Y. provided clinical care and collected clinical data; Y.Z., S.D.A., A.A.C.L., S.K., X.F., L.A., and D.Q.R. performed research; Y.Z. analyzed and interpreted data; Y.Z., R.S., and C.O.W. performed statistical analysis; and Y.Z., B.A.P., S.K., W.A.F., and N.S.Y. wrote the manuscript.

Conflict-of-interest disclosure: The authors declare no competing financial interests.

ORCID profiles: Y.Z., 0000-0002-4108-5245; B.A.P., 0000-0002-2974-7701; E.M.G., 0000-0002-4648-5926.

Correspondence: Yoshitaka Zaimoku, Hematology Branch, National Heart, Lung, and Blood Institute, National Institutes of Health, Bethesda, MD, 20892; e-mail: zaimokuyoshitaka@gmail.com.

Footnotes

Submitted 14 June 2021; accepted 18 October 2021; prepublished online on *Blood* First Edition 1 November 2021. DOI 10.1182/ blood.2021012895.

*Y.Z. and B.A.P. contributed equally to this study.

REFERENCES

- Young NS. Aplastic Anemia. N Engl J Med. 2018;379(17):1643-1656.
- Townsley DM, Scheinberg P, Winkler T, et al. Eltrombopag added to standard immunosuppression for aplastic anemia. N Engl J Med. 2017;376(16):1540-1550.
- D'amaro J, van Rood JJ, Rimm AA, Bortin MM. HLA associations in Italian and non-Italian Caucasoid aplastic anaemia patients. *Tissue Antigens.* 2008;21(3):184-191.
- Chapuis B, Von Fliedner VE, Jeannet M, et al. Increased frequency of DR2 in patients with aplastic anaemia and increased DR sharing in their parents. Br J Haematol. 1986;63(1):51-57.
- Nimer SD, Ireland P, Meshkinpour A, Frane M. An increased HLA DR2 frequency is seen in aplastic anemia patients. *Blood*. 1994; 84(3):923-927.
- Nakao S, Takamatsu H, Chuhjo T, et al. Identification of a specific HLA class II haplotype strongly associated with susceptibility to cyclosporine-dependent aplastic anemia. *Blood*. 1994;84(12): 4257-4261.
- Shichishima T, Noji H, Ikeda K, Akutsu K, Maruyama Y. The frequency of HLA class I alleles in Japanese patients with bone marrow failure. *Haematologica*. 2006;91(6): 856-857.
- Sugimori C, Yamazaki H, Feng X, et al. Roles of DRB1 *1501 and DRB1 *1502 in the pathogenesis of aplastic anemia. *Exp Hematol.* 2007;35(1):13-20.
- Song EY, Park S, Lee DS, Cho HI, Park MH. Association of human leukocyte antigen-DRB1 alleles with disease susceptibility and severity of aplastic anemia in Korean patients. *Hum Immunol.* 2008;69(6):354-359.
- Rehman S, Saba N, Khalilullah, Munir S, Ahmed P, Mehmood T. The frequency of HLA class I and II alleles in Pakistani patients with aplastic anemia. *Immunol Invest.* 2009; 38(8):812-819.
- Katagiri T, Sato-Otsubo A, Kashiwase K, et al; Japan Marrow Donor Program. Frequent loss of HLA alleles associated with copy number-neutral 6pLOH in acquired aplastic anemia. *Blood.* 2011;118(25): 6601-6609.
- Dhaliwal JS, Wong L, Kamaluddin MA, Yin LY, Murad S. Susceptibility to aplastic anemia is associated with HLA-DRB1*1501

in an aboriginal population in Sabah, Malaysia. *Hum Immunol.* 2011;72(10):889-892.

- Chen C, Lu S, Luo M, Zhang B, Xiao L. Correlations between HLA-A, HLA-B and HLA-DRB1 allele polymorphisms and childhood susceptibility to acquired aplastic anemia. Acta Haematol. 2012;128(1):23-27.
- 14. Qi J, Wang TJ, Li HX, et al. Association of HLA class II (-DRB1,-DQB1,-DPB1) alleles and haplotypes on susceptibility to aplastic anemia in northern Chinese Han. *Hum Immunol.* 2020;81(12):685-691.
- Savage SA, Viard M, O'hUigin C, et al. Genome-wide association study identifies HLA-DPB1 as a significant risk factor for severe aplastic anemia. Am J Hum Genet. 2020;106(2):264-271.
- Afable MG II, Wlodarski M, Makishima H, et al. SNP array-based karyotyping: differences and similarities between aplastic anemia and hypocellular myelodysplastic syndromes. *Blood.* 2011;117(25):6876-6884.
- Yoshizato T, Dumitriu B, Hosokawa K, et al. Somatic mutations and clonal hematopoiesis in aplastic anemia. N Engl J Med. 2015; 373(1):35-47.
- Zaimoku Y, Takamatsu H, Hosomichi K, et al. Identification of an HLA class I allele closely involved in the autoantigen presentation in acquired aplastic anemia. *Blood.* 2017; 129(21):2908-2916.
- Babushok DV, Duke JL, Xie HM, et al. Somatic HLA mutations expose the role of class I-mediated autoimmunity in aplastic anemia and its clonal complications. *Blood Adv.* 2017;1(22):1900-1910.
- Espinoza JL, Elbadry MI, Chonabayashi K, et al. Hematopoiesis by iPSC-derived hematopoietic stem cells of aplastic anemia that escape cytotoxic T-cell attack. *Blood Adv.* 2018;2(4):390-400.
- Maruyama H, Katagiri T, Kashiwase K, et al. Clinical significance and origin of leukocytes that lack HLA-A allele expression in patients with acquired aplastic anemia. *Exp Hematol.* 2016;44(10):931-939.e3.
- Mustjoki S, Young NS. Somatic mutations in "benign" disease. N Engl J Med. 2021; 384(21):2039-2052.
- Gargiulo L, Zaimoku Y, Scappini B, et al. Glycosylphosphatidylinositol-specific T cells, IFN-γ-producing T cells, and pathogenesis of idiopathic aplastic anemia. *Blood.* 2017; 129(3):388-392.

Presented at the sixty-first annual meeting of the American Society of Hematology, Orlando, FL, 9 December 2019.

There is a *Blood* Commentary on this article in this issue.

The online version of this article contains a data supplement.

The publication costs of this article were defrayed in part by page charge payment. Therefore, and solely to indicate this fact, this article is hereby marked "advertisement" in accordance with 18 USC section 1734.

- Luzzatto L, Risitano AM. Advances in understanding the pathogenesis of acquired aplastic anaemia. *Br J Haematol.* 2018; 182(6):758-776.
- Sun L, Babushok DV. Secondary myelodysplastic syndrome and leukemia in acquired aplastic anemia and paroxysmal nocturnal hemoglobinuria. *Blood.* 2020; 136(1):36-49.
- Negoro E, Nagata Y, Clemente MJ, et al. Origins of myelodysplastic syndromes after aplastic anemia. *Blood.* 2017;130(17): 1953-1957.
- Scheinberg P, Nunez O, Wu C, Young NS. Treatment of severe aplastic anaemia with combined immunosuppression: antithymocyte globulin, ciclosporin and mycophenolate mofetil. *Br J Haematol.* 2006; 133(6):606-611.
- Scheinberg P, Wu CO, Nunez O, et al. Treatment of severe aplastic anemia with a combination of horse antithymocyte globulin and cyclosporine, with or without sirolimus: a prospective randomized study. *Haematologica*. 2009;94(3):348-354.
- Scheinberg P, Nunez O, Weinstein B, et al. Horse versus rabbit antithymocyte globulin in acquired aplastic anemia. N Engl J Med. 2011;365(5):430-438.
- Scheinberg P, Young NS. How I treat acquired aplastic anemia. *Blood*. 2012; 120(6):1185-1196.
- Hosomichi K, Jinam TA, Mitsunaga S, Nakaoka H, Inoue I. Phase-defined complete sequencing of the HLA genes by nextgeneration sequencing. *BMC Genomics*. 2013;14(1):355.
- Castelli EC, Paz MA, Souza AS, Ramalho J, Mendes-Junior CT. Hla-mapper: an application to optimize the mapping of HLA sequences produced by massively parallel sequencing procedures. *Hum Immunol.* 2018;79(9):678-684.
- Pécuchet N, Rozenholc Y, Zonta E, et al. Analysis of base-position error rate of nextgeneration sequencing to detect tumor mutations in circulating DNA. *Clin Chem.* 2016;62(11):1492-1503.
- 34. Pécuchet N, Zonta E, Didelot A, et al. Baseposition error rate analysis of nextgeneration sequencing applied to circulating tumor DNA in non-small cell lung cancer: a prospective study. *PLoS Med.* 2016;13(12): e1002199.
- 35. Kanda Y. Investigation of the freely available easy-to-use software 'EZR' for medical

statistics. Bone Marrow Transplant. 2012; 48(3):452-458.

- Gragert L, Madbouly A, Freeman J, Maiers M. Six-locus high resolution HLA haplotype frequencies derived from mixed-resolution DNA typing for the entire US donor registry. *Hum Immunol.* 2013;74(10):1313-1320.
- Rondelli T, Berardi M, Peruzzi B, et al. The frequency of granulocytes with spontaneous somatic mutations: a wide distribution in a normal human population. *PLoS One*. 2013; 8(1):e54046.
- Watson CJ, Papula AL, Poon GYP, et al. The evolutionary dynamics and fitness landscape of clonal hematopoiesis. *Science*. 2020; 367(6485):1449-1454.
- Mercer TR, Clark MB, Andersen SB, et al. Genome-wide discovery of human splicing branchpoints. *Genome Res.* 2015;25(2): 290-303.
- Scheinberg P, Wu CO, Nunez O, Young NS. Predicting response to immunosuppressive therapy and survival in severe aplastic anaemia. Br J Haematol. 2009;144(2): 206-216.
- Zaimoku Y, Patel BA, Shalhoub R, et al. Predicting response of severe aplastic anemia to immunosuppression combined with eltrombopag. *Haematologica*. In press.

- Mizumaki H, Hosomichi K, Hosokawa K, et al. A frequent nonsense mutation in exon 1 across certain HLA-A and -B alleles in leukocytes of patients with acquired aplastic anemia. *Haematologica*. 2021;106(6):1581–1590.
- Ikeda N, Kojima H, Nishikawa M, et al. Determination of HLA-A, -C, -B, -DRB1 allele and haplotype frequency in Japanese population based on family study. *Tissue Antigens.* 2015;85(4):252-259.
- 44. Li Y, Li X, Ge M, et al. Long-term follow-up of clonal evolutions in 802 aplastic anemia patients: a single-center experience. Ann Hematol. 2011;90(5):529-537.
- 45. Patel BA, Groarke EM, Lotter J, et al. Longterm outcomes in severe aplastic anemia patients treated with immunosuppression and eltrombopag: a phase 2 study. *Blood.* 2021;blood.2021012130.
- 46. Katagiri T, Kawamoto H, Nakakuki T, et al. Individual hematopoietic stem cells in human bone marrow of patients with aplastic anemia or myelodysplastic syndrome stably give rise to limited cell lineages. Stem Cells. 2013;31(3): 536-546.
- 47. Shukla SA, Rooney MS, Rajasagi M, et al. Comprehensive analysis of cancer-associated

somatic mutations in class I HLA genes. Nat Biotechnol. 2015;33(11):1152-1158.

- McGranahan N, Rosenthal R, Hiley CT, et al; TRACERx Consortium. Allele-specific HLA loss and immune escape in lung cancer evolution. *Cell*. 2017;171(6):1259-1271.e11.
- Chowell D, Morris LGT, Grigg CM, et al. Patient HLA class I genotype influences cancer response to checkpoint blockade immunotherapy. *Science*. 2018;359(6375): 582-587.
- Vago L, Perna SK, Zanussi M, et al. Loss of mismatched HLA in leukemia after stem-cell transplantation. N Engl J Med. 2009;361(5): 478-488.
- 51. Bacigalupo A. How I treat acquired aplastic anemia. *Blood.* 2017;129(11):1428-1436.
- Zhao X, Gao S, Wu Z, et al. Single-cell RNAseq reveals a distinct transcriptome signature of aneuploid hematopoietic cells. *Blood*. 2017;130(25):2762-2773.
- Patel BA, Ghannam J, Groarke EM, et al. Detectable mutations precede late myeloid neoplasia in aplastic anemia. *Haematologica*. 2020;106(2):647-650.
- Clemente MJ, Przychodzen B, Hirsch CM, et al. Clonal PIGA mosaicism and dynamics in paroxysmal nocturnal hemoglobinuria. *Leukemia*. 2018;32(11):2507-2511.