## Remission, treatment failure, and relapse in pediatric ALL: an international consensus of the Ponte-di-Legno Consortium

Swantje Buchmann,<sup>1,\*</sup> Martin Schrappe,<sup>1,\*</sup> Andre Baruchel,<sup>2,3</sup> Andrea Biondi,<sup>4</sup> Michael Borowitz,<sup>5,6</sup> Myriam Campbell,<sup>6</sup> Gunnar Cario,<sup>1</sup> Giovanni Cazzaniga,<sup>4</sup> Gabriele Escherich,<sup>7</sup> Christine J. Harrison,<sup>8</sup> Mats Heyman,<sup>9</sup> Stephen P. Hunger,<sup>10</sup> Csongor Kiss,<sup>11</sup> Hsi-Che Liu,<sup>12</sup> Franco Locatelli,<sup>13</sup> Mignon L. Loh,<sup>14,15</sup> Atsushi Manabe,<sup>16</sup> Georg Mann,<sup>17</sup> Rob Pieters,<sup>18</sup> Ching-Hon Pui,<sup>19</sup> Susana Rives,<sup>20</sup> Kjeld Schmiegelow,<sup>21</sup> Lewis B. Silverman,<sup>22</sup> Jan Stary,<sup>23</sup> Ajay Vora,<sup>24</sup> and Patrick Brown,<sup>25</sup> on behalf of the Ponte-di-Legno Consortium

<sup>1</sup>Department of Pediatrics, University Medical Center Schleswig-Holstein, Kiel, Germany; <sup>2</sup>Pediatric Hematology-Immunology Department, University Hospital Robert Debré Assistance Publique-Hôpitaux de Paris (AP-HP), Université de Paris, Paris; <sup>3</sup>Société Française de Lutte contre les Cancers et Leucémies de l'Enfant et de l'Adolescent (SFCE), Paris, France; <sup>4</sup>Department of Pediatrics and Tettamanti Research Center, Fondazione MBBM (Monza e Brianza per il Bambino e la sua Mamma)/Ospedale San Gerardo, University of Milano-Bicocca, Monza, Italy; <sup>5</sup>Department of Pediatrics and Pediatric Surgery, Hospital de Niños Roberto del Río, Facultad de Medicina, Universidad de Chile, Santiago, Chile; <sup>6</sup>Chilean National Pediatric Oncology Group (PINDA), Santiago, Chile; <sup>7</sup>Clinic of Pediatric Hematology and Oncology, University Medical Center Hamburg-Eppendorf, Hamburg, Germany;<sup>8</sup>Translational and Clinical Research Institute, Newcastle University Centre for Cancer, Newcastle-upon-Tyne, United Kingdom; <sup>9</sup>Childhood Cancer Research Unit, Karolinska Institutet–Astrid Lindgren's Children's Hospital, Karolinska University Hospital, Stockholm, Sweden; <sup>10</sup>Department of Pediatrics, Center for Childhood Cancer Research, Children's Hospital of Philadelphia, Philadelphia, PA; <sup>11</sup>Department of Pediatric Hematology and Oncology, Institute of Pediatrics, Faculty of Medicine, University of Debrecen, Debrecen, Hungary; <sup>12</sup>Division of Pediatric Hematology-Oncology, MacKay Memorial Hospital–MacKay Children's Hospital, Taipei, Taiwan; <sup>13</sup>Department of Pediatric Hematology and Oncology, Istituto di Ricovero e Cura a Carattere Scientifico (IRCCS) Ospedale Pediatrico Bambino Gesù, Sapienza, Università di Roma, Rome, Italy; 14 Department of Pediatrics, Hokkaido University Graduate School of Medicine, Sapporo, Japan;<sup>15</sup> Japan Children's Cancer Group Japan (JCCG), Sapporo, Japan;<sup>16</sup> Department of Pediatrics, Hokkaido University Graduate School of Medicine, Sapporo, Japan; <sup>17</sup>Children's Cancer Research Institute–St Anna Children's Hospital, Department of Pediatrics, Medical University of Vienna, Vienna, Austria; <sup>18</sup> Princess Maxima Center for Pediatric Oncology, Utrecht, The Netherlands; <sup>19</sup> Department of Oncology, St Jude Children's Research Hospital, Memphis, TN; <sup>20</sup>Pediatric Hematology and Oncology Department, Hospital Sant Joan de Déu de Barcelona-Institut de Recerca Sant Joan de Déu, Barcelona, Spain; <sup>21</sup>Department of Pediatrics and Adolescent Medicine, University Hospital Rigshospitalet–Institute of Clinical Medicine, Faculty of Medicine, University of Copenhagen, Copenhagen, Denmark;<sup>22</sup>Department of Pediatric Oncology, Dana-Farber Cancer Institute, Boston, MA; <sup>23</sup>University Hospital Motol–Second Faculty of Medicine, Charles University, Prague, Czech Republic; <sup>24</sup>Great Ormond Street Hospital, London, United Kingdom; and <sup>25</sup>Johns Hopkins Kimmel Cancer Center, Baltimore, MD

Comparison of treatment strategies in de novo pediatric acute lymphoblastic leukemia (ALL) requires standardized measures of efficacy. Key parameters that define diseaserelated events, including complete remission (CR), treatment failure (TF; not achieving CR), and relapse (loss of CR) require an updated consensus incorporating modern diagnostics. We collected the definitions of CR, TF, and relapse from recent and current pediatric clinical trials for the treatment of ALL, including the key components of response evaluation (timing, anatomic sites, detection methods, and thresholds) and found significant heterogeneity, most notably in the definition of TF. Representatives of the major international ALL clinical trial groups convened to establish consensus definitions. CR should be defined at a time point no earlier than at the end of induction and should include the reduction of blasts below a specific threshold in bone marrow and extramedullary sites, incorporating minimal residual disease (MRD) techniques for marrow evaluations. TF should be defined as failure to achieve CR by a prespecified time point in therapy. Relapse can only be defined in patients who have achieved CR and must include a specific threshold of leukemic cells in the bone marrow confirmed by MRD, the detection of central nervous system leukemia, or documentation of extramedullary disease. Definitions of TF and relapse should harmonize with eligibility criteria for clinical trials in relapsed/refractory ALL. These consensus definitions will enhance the ability to compare outcomes across pediatric ALL trials and facilitate development of future international collaborative trials.

## Introduction

Children with newly diagnosed acute lymphoblastic leukemia (ALL) in high-income countries are usually treated in clinical trials encompassing different therapeutic approaches. Treatment strategies and overall results are quite comparable among national and international study groups despite treatment variations.<sup>1-3</sup> There is no uniform consensus on the basic definitions of response criteria, including complete remission (CR), treatment failure (TF, sometimes also referred to as non-CR, primary treatment failure, induction failure, or refractory disease), and relapse in pediatric ALL. These definitions are highly relevant to treatment decisions; for example, when to conclude that the frontline therapy is only partially effective (prompting stratification into a more intensive treatment arm) or largely ineffective (prompting use of alternative or salvage therapy). Up to approximately the year 2000, morphologic evaluation (cytology) was the primary method to establish the diagnosis of ALL and to assess the treatment response. Twenty-five percent or more lymphoblasts in the bone marrow (BM) defined the diagnosis of ALL and BM relapse.<sup>4,5</sup> Since 2000, new diagnostic methods, able to detect leukemic cells with greater sensitivity and specificity compared with morphology, have emerged. Molecular and immunophenotypic techniques can measure minimal residual disease (MRD) reliably at the threshold of 1 in 10000 to 100000 cells. Response assessment by MRD detection has become the most important independent predictor of treatment outcome in many studies.<sup>6-8</sup>

The objective of all pediatric ALL study groups is to increase the event-free survival (EFS) and overall survival (OS) rates and improve short- and long-term quality of life. Because of the numerous new therapeutic options being investigated by various study groups worldwide, it is crucial to be able to compare results between different protocols. Thus, standardized definitions of CR and TF after the initial therapy (assessed after a comparable treatment period) and standardized definitions of relapse after achieving CR are required. Common definitions of TF and relapse may provide more uniform eligibility criteria for relapsed/refractory studies. In pediatric oncology, there are other well-established examples of uniform consensus definitions of treatment response and failure being used routinely in clinical trials (eg, response evaluation criteria in solid tumors [RECIST], solid tumors; Cheson, acute myeloid leukemia; Lugano, Hodgkin disease).<sup>9-11</sup>

A key question is which parameters in the evaluation of outcome are most relevant in upfront ALL trials. OS (time-to-death) is important, but it provides incomplete information. The single most widely accepted parameter is EFS. The main advantage of EFS over OS is the incorporation of clinically important outcomes in addition to death, such as the inability to achieve CR (TF) or relapse (both of which reflect unsatisfactory response to frontline treatment requiring salvage therapy), therapy-related mortality (induction death and death in CR, caused by resistant disease or treatment-related toxicity), and second malignancies. Furthermore, OS in a clinical trial is influenced by the efficacy of salvage therapy and thus may not inform about the efficacy of frontline therapy.

For calculation of time-to-event from diagnosis, the following parameters must be defined consistently: (1) CR (first CR; because relapse can only occur if CR has been achieved); (2) primary treatment failure (TF; not achieving CR at a predefined time point; eg, end of induction or end of consolidation); and (3) relapse (re-emergence of leukemia after CR has been achieved).

To address these issues, the Ponte-di-Legno (PDL) group representatives endeavored first to collect and summarize the current definitions of CR, TF, relapse, and other events in the pediatric ALL study groups and then to create consensus definitions to be used in future clinical trials.

### Methods

This collaborative project of the PDL group comprises the following study groups with their recent protocols: Associazone Italiana Ematologia Oncologia Pediatrica (AIEOP), Berlin-Frankfurt-Münster group (BFM) (AIEOP-BFM ALL 2017); Intercontinental BFM group for ALL (ALL-IC BFM) (ALL-IC BFM 2009); ALLtogether (ALLTogether1); Co-operative Study Group (COALL) (CoALL 08-09); Children's Oncology Group (COG) (AALL1731, AALL1732, AALL1721); Dutch Childhood Oncology Group (DCOG) (DCOG-ALL-11); Dana Farber Cancer Institute (DFCI) ALL Consortium (DFCI 16-001); European Study Group for Philadelphia positive ALL (EsPhALL)/COG (EsPhALL2017/COG (AALL1631); Japan Children's Cancer Group (JCCG) (JPLSG-ALL-B12), Nordic Society of Paediatric Haematology and Oncology (NOPHO) (NOPHOALL 2008); Spanish Society of Paediatric Haematology Oncology (SEHOP Group) and Programa para el Tratamiento de Hemopatías Malignas (PETHEMA) (ALL/SEHOP-PETHEMA 2013); Société Française de lutte contre les Cancers de l'Enfant et de l'Adolescent (SFCE) (CAALL F01); St. Jude Children's Research Hospital (SJCRH) (TOT-XVI); Taiwan Pediatric Oncology Group (TPOG) (TPOG-ALL2013); and United Kingdom Acute Lymphoblastic leukaemia (UKALL) (UKALL 2011 Trial).<sup>12-27</sup>

Initially, recent protocols were reviewed for the definitions of CR, TF, relapse, and events. A summary was provided to all participant study groups to review the definitions derived from these study protocols. In preparation for a face-to-face meeting of the PDL group in Zbiroh, Czech Republic, on 19 May 2019, an overview of all definitions was prepared, and the first classifications were made.

#### Terminology

**BM and central nervous system involvement** The following morphologic classifications of BM and central nervous system (CNS) involvement with ALL were developed decades ago and will be referred to in the presentation of the results.<sup>5,28,29</sup>

BM classification by cytomorphology is defined: (1) M1 marrow, <5% blasts; (2) M2 marrow,  $\geq$ 5 to <25% blasts; (3) M3 marrow,  $\geq$ 25% blasts. Of note, COG has historically defined M3 marrow as >25% blasts rather than  $\geq$ 25% blasts.<sup>28</sup>

CNS classification by cytomorphology and/or clinical findings (±imaging and biopsy):<sup>30</sup> (1) CNS1, absence of blasts on cytospin preparation in cerebrospinal fluid (CSF) and no clinical or imaging findings of CNS leukemia; (2) CNS2,  $\leq$ 5/µL nucleated cells in CSF, cytospin preparation positive for blasts, and no clinical or imaging findings of CNS disease; (3) CNS3: >5/µL nucleated cells in CSF and cytospin preparation positive for blasts or clinical or imaging findings of CNS disease. Of note, COG, EsPhALL/COG, DFCI, NOPHO, SFCE, SJCRH, and TPOG defined CNS3 as  $\geq$ 5/µL nucleated cells in CSF.

**Treatment phases** In most protocols, induction chemotherapy (phase Ia in BFM-type regimens) lasts about 4 to 6 weeks and is followed by a second phase, frequently known as consolidation phase (phase Ib in BFM-type regimens), then a third intensification phase often targeting the extramedullary space, and finally a fourth reintensification phase (delayed intensification) with or without CNS irradiation, depending on risk stratification and treatment strategy. In a small minority of cases, the latter parts of the protocol sequence are replaced by allogeneic hematopoietic stem cell transplantation.<sup>32</sup> 
 Table 1. Definition, methods, and time points for remission assessment in current treatment protocols for pediatric

 ALL (remission requires fulfillment of all sites)

Anatomic sites	Methods	Study group	Time point, d	Exceptions
<b>BM</b> <5% blasts in BM	Cytomorphology	AIEOP-BFM ALL-IC BFM CoALL DCOG DFCI JCCG TPOG	33, EOI 33, EOI 29, EOI 33, EOI 32, EOI 33, EOI 33, EOI 35-42, EOI	
<5% blasts in BM	Cytomorphology plus confirmation by FCM-/ PCR-MRD/genetics	NOPHO SEHOP-PETHEMA SFCE SJCRH	29, EOI EOI 35-42, EOI 38-42, EOI	SJCRH mainly uses FCM-MRD
<5% blasts in BM	PCR-MRD (±cytomorphology)	UKALL	29, EOI	
<1% blasts in BM	FCM-/PCR-MRD (±cytomorphology)	ALLTogether COG	EOI as the earliest EOI for early CR, EOC for late CR End of Cons. block 3	
CNS 1	Cytomorphology, imaging, clinical examination	Every participating study group	See above for respective study group	
CNS 1	Cytomorphology plus confirmation by FCM-MRD	ALLTogether	See above for respective study group	If suspect cells are seen; evaluation at EOC (d 71)
EM				
Resolution of leukemic infiltrates	Clinical examination, imaging, histology	ALL-IC COG DCOG JCCG EsPhALL/COG	See above for respective study group	DCOG: retesting of testicular involvement after Prot. M NOPHO: retesting EM involvement at d 85
Reduction of initial leukemic mass to 1/3	Clinical examination, imaging, histology	AIEOP-BFM ALLTogether CoALL DFCI SEHOP-PETHEMA SFCE UKALL	See above for respective study group	UKALL: retesting of testicular involvement at wk 8

BM: bone marrow, CNS: central nervous system, d: day, EM: extramedullary sites, EOC: end of consolidation, EOI: end of induction, FCM-MRD: flow cytometric minimal residual disease, MRD: minimal residual disease, PCR-MRD: polymerase chain reaction minimal residual disease.

## Results

#### **Complete remission**

Large differences among groups were found in definitions of CR, reflecting differences in timing of CR assessment, in threshold levels of leukemic cells to define CR, and the detection method (Table 1; supplemental Table 1, available on the *Blood* Web site).

#### Time point(s) of remission assessment

All study groups agreed that the earliest assessment of CR should occur no sooner than the completion of induction chemotherapy. Remission assessment at the end of induction is used for risk stratification in every study group except EsPhALL/COG. In some protocols (AIEOP-BFM ALL 2017, ALL-IC BFM 2009, ALLTogether1, AALL1731, CoALL 08-09, DCOG-ALL-11, NOPHO-ALL 2008, DFCI 16-001), remission status is also explicitly re-examined at later therapy time points for patients not in CR (eg, after high-intensity intensification blocks).

#### Anatomic sites included in remission assessment

All study groups evaluate CR in BM and CNS. All groups except SJCRH and TPOG also evaluate CR in other extramedullary sites.

#### Thresholds of lymphoblasts in BM

COG and ALLTogether currently use a threshold of <1% lymphoblasts in BM as the criterion for CR. This is a new MRD-based addition to current definitions and was not used in earlier trials. All other study groups continue to use a historical threshold of <5% lymphoblasts in BM as a criterion for CR.

#### Table 2. Definitions of treatment failure events in current treatment protocols

Event	Methods	Study group
Induction failure (EOI) ≥5% blasts in BM	Cytomorphology and/or FCM-/PCR-MRD or genetics	JCCG, SEHOP-PETHEMA, SFCE, SJCRH, TPOG
Induction failure (EOI) ≥25% blasts in BM	See above	DFCI, UKALL
No induction failure events; treatment failure is an event if CR has not been achieved at later timepoints	Different combinations of methods	AIEOP-BFM, ALLTogether, ALL-IC-BFM, CoALL, COG, DCOG, EsPhALL-COG, NOPHO

BM: bone marrow, EOI: end of induction, FCM-MRD: flow cytometric minimal residual disease, MRD: minimal residual disease, PCR-MRD: polymerase chain reaction minimal residual disease.

#### **Detection methods for response assessment**

BM Methods for detection of leukemic cells have been described in the study protocols, particularly for BM assessment, and as MRD techniques have evolved, the variability has increased. CR assessment is currently based solely (without incorporating MRD methodology) on cytomorphology in AIEOP-BFM, ALL-IC BFM, CoALL, DCOG, DFCI, JCCG, and TPOG, and thus, the morphologic classification of the BM in M1, M2, and M3 applies.<sup>28</sup> In unclear cases, alternative methods are applied. NOPHO and SEHOP-PETHEMA initially assess the therapy response cytomorphologically but require confirmation of a result with <5% lymphoblasts by flow cytometry (FCM). If the results differ, FCM takes precedence.<sup>33</sup> SFCE also examines the bone marrow cytomorphologically and then uses polymerase chain reaction-MRD (PCR-MRD) as a confirmation method. SJCRH mainly uses FCM for remission assessment. COG, ALLTogether, and UKALL consider FCM- and/or PCR-MRD as the primary source of information and only consider cytomorphological results as a backup in the unlikely situation of failure of MRD technologies. In EsPhALL/ COG, CR is primarily determined by PCR-MRD, but backup by FCM-MRD is permitted to assess CR if PCR-MRD was unavailable, and lack of CR can be confirmed by cytomorphology (M2/M3) or by an additional method (PCR-MRD, FCM-MRD, fluorescence in situ hybridization [FISH]).33-36

**CSF** Cytomorphology is used by all study groups to determine CR status in CSF. The presence of CNS leukemia is classified according to published criteria.<sup>29,31,37</sup> SJCRH uses indirect nuclear terminal deoxynucleotidyl transferase immunofluorescence assay to confirm the leukemic nature of blasts in CSF. ALLTogether and CoALL recommend a differentiation of malignant and nonmalignant contamination with FCM-MRD for abnormal CSF findings.<sup>38</sup> In addition, all groups do clinical examinations and imaging in case of neurologic symptoms. In exceptional cases where imaging is equivocal, biopsy may be required.

Other extramedullary sites In all study groups except SJCRH and TPOG, non-CNS extramedullary (EM) sites are examined clinically by imaging, and/or biopsy to assess reduction of tumor mass or to assess a viability of residual tumor tissue histologically. ALL-IC, COG, DCOG, JCCG, and EsPhALL/COG require a resolution of EM leukemia. AIEOP-BFM, ALLTogether, CoALL, DFCI, SEHOP-PETHEMA, SFCE, and UKALL require a reduction of initial tumor mass (especially mediastinal mass) to at least one-third of the initial size. NOPHO evaluates EM leukemia after consolidation therapy for the first time. Some study groups (UKALL, DCOG, and NOPHO) do not incorporate resolution of testicular disease at the end of induction for assessment of CR; they assess testicular response at later therapy time points for the first time.

#### TF

Significant differences among groups were found in definitions of TF as events (Table 2; supplemental Table 1). Although there is a broad consensus that TF should be defined as failure to achieve a CR before a prespecified time point, there are significant differences in the prespecified time point.

#### Defining treatment time points

A uniform definition of the time points end of induction (EOI; approximately 4 weeks from start of treatment) and end of consolidation (EOC; approximately 8-12 weeks from start of treatment) appears feasible. Later time points, such as end of intensification or end of reintensification are also potentially useful, because the results of assessment at these time points may be used as criteria for intensification of therapy or pursuit of alternative rescue therapies.

#### Defining TF based on treatment time points

Some study groups consider failure to achieve CR at EOI as a TF event for EFS analysis, whereas others do not: instead, they use the result to stratify the patient to more intensive postinduction therapy, and a TF event only occurs if CR is not achieved at a later time point. This is an important limitation when trying to compare the results of different trials, because identical end-of-induction evaluations could lead to different EFS results for different protocols. For example, JCCG, SFCE, SEHOP-PETHEMA, SJCRH, and TPOG consider M2 or more blasts in BM at EOI as a TF event. In COG trials, on the other hand, a TF event occurs only if CR is not achieved after consolidation therapy (EOC). For UKALL and DFCI, M3 marrow at EOI is considered a TF event. In AIEOP-BFM ALL, ALL-IC BFM, CoALL, DCOG, and ALLTogether, lack of CR at EOI has an impact on subsequent risk stratification and intensification of postinduction treatment; however, this does not define a TF event. In these groups, a TF event is defined as lack of CR at various later time points.

#### Consensus for CR and TF

 Each study group should define the decisive treatment points EOI and EOC (and, if useful for the individual protocol, end of intensification phase) for their protocols.

- Several methods are available to assess MRD in ALL, including quantitative polymerase chain reaction of rearranged immunoglobulin/T-cell receptor genes using either standard allele-specific oligonucleotide PCR (ASO-PCR) or next-generation sequencing (NGS) technologies, reverse transcriptase-PCR of fusion transcripts, and FCM. All these methods are acceptable as long as standardized evaluation criteria (quality standards) have been established.
- Definition of CR:
  - CR is to be assessed no earlier than EOI
    - For CR, the following is required:
    - BM: MRD <1% and/or M1 cytomorphology
    - CNS: CNS1
    - Testes: normalization of clinical examination, or a negative biopsy if clinical examination is not considered normal.
    - EM: no evidence of leukemic infiltrates as evaluated clinically and by imaging; a preexisting leukemic mass (mediastinal mass included) must have decreased at least to one-third of the initial tumor volume
- CR assessment of BM should be performed by a standardized MRD method and cytomorphology. The consensus was that MRD measurement is now the 'gold standard' to assess CR in addition to its established role in risk stratification at the end of induction. If MRD is not available, the response is assessed by cytomorphology. If both methods are available, MRD generally takes precedence over cytomorphology. Complementary methods including genetic analysis may be used to verify CR achievement. CNS status should be based on CSF cytomorphology (other methods such as FCM or genetic analysis may be used in unclear cases), and clinical neurological examination, and CNS imaging in case of neurological clinical findings. Physical examination, imaging or histologic examination of a tissue biopsy should be used for the evaluation of non-CNS EM disease.
- We propose that every study group report CR status on early time points such as end of induction or end of consolidation to improve comparability between trials.
- Definition of TF:
  - Failure to achieve CR at a clearly predefined time point (EOI, EOC, or other time points during intensification) should be considered as a TF event, and this time point should be specified at the onset of the clinical trial.
  - There was progress toward a consensus that a TF event is to be defined no earlier than the EOC, because this would allow a patient not achieving CR by EOI to be offered a consolidation therapy with agents not given during induction in an effort to overcome blast resistance and potentially achieve CR. If such a patient still does not achieve CR after this risk-adapted consolidation phase, this could reasonably define a TF event.

#### Relapse

An overview and classification of current relapse definitions was prepared (supplemental Table 2). Obviously, a patient can only have a relapse event after a CR has previously been achieved. Relapse should be subclassified by anatomic sites involved at the time of relapse: isolated BM relapse, isolated CNS relapse, isolated testicular relapse, isolated other EM relapse involving other anatomical sites, and various combinations of these sites.

#### Historical definitions of relapse

Traditionally, isolated BM relapse has been defined as a M3 marrow occurring after CR with no other sites involved, with cytomorphology as the gold standard for defining a BM relapse. Combined relapse has been defined as reappearance of lymphoblasts in the BM with a threshold of  $\geq$ 5%, with the involvement of one or more EM sites confirmed to be leukemic by cytology/histology or imaging. Isolated CNS relapse is defined as CNS3 status (with a M1 marrow) emerging after CR. A testicular relapse is defined as uni- or bilateral testicular enlargement with leukemic infiltration confirmed by biopsy (required by most groups) or imaging. Defining relapse in other EM anatomic sites (one and more) requires biopsy (supplemental Table 3).

# Change in the threshold of leukemic blasts in BM with a suspected isolated BM relapse

Highly sensitive methods to detect leukemic cells have prompted all study groups except JCCG and SEHOP-PETHEMA to lower the threshold of morphologically defined blast percentage required for BM relapse if additional conditions are fulfilled. Thus, AIEOP-BFM, ALL-Together, CoALL, COG, DCOG, DFCI, EsPhALL/ COG, NOPHO, SFCE, SJCRH, and UKALL define BM relapse as ≥5% lymphoblasts if confirmed to be leukemic by 1 (ALLTogether, CoALL, COG, DCOG, DFCI, EsPhALL/COG, NOPHO, SJCRH, UKALL) or 2 (AIEOP-BFM, SFCE) other detection methods such as PCR-MRD, FCM-MRD, NGS-MRD, or genetics. ALLTogether and COG define BM relapse as <5% (but >1%) blasts by morphology in BM if 2 additional methods indicate the re-emergence of leukemia. If only 1 sensitive diagnostic test is available, a repeat BM evaluation is required, with additional conditions to be met for this second BM assessment. It is important to emphasize that all groups agreed that relapse in the absence of a M3 morphologic result must be confirmed by specified validated diagnostic methods.

#### Special features of CNS relapse

All study groups accept a single lumbar puncture (LP) with CNS3 status as a relapse event. COG, DCOG, DFCI, SJCRH, and UKALL also incorporate a definition of CNS relapse that does not require CNS3. Patients with a single LP with CNS2 status after attaining CR require a second LP to be performed 1 to 4 weeks later. If this LP again shows CNS2, COG, DCOG, SJCRH, and UKALL define this to be a CNS relapse if the presence of leukemic blasts is confirmed by FCM-MRD, FISH, and/ or indirect nuclear terminal deoxynucleotidyl transferase immunofluorescence assay. DFCI does not require a confirmatory test. ALLTogether and CoALL require a confirmatory test also when CNS3 is detected.

#### **Consensus for relapse**

The following consensus was reached (Tables 3-5):

- The diagnosis of relapse can only be made if CR has been previously achieved.
- BM:
  - For the purposes of defining bone marrow relapse, the percentage of blasts in any marrow evaluation should be determined using a validated MRD technique if available, with cytomorphology used only as a backup.
  - To define bone marrow relapse, there must be clear evidence that the patient has ≥1% leukemic blasts in the

#### Table 3. Consensus criteria for relapse definition in pediatric ALL: BM relapse (MRD available)

BM 1		BM 2*
MRD	Others	MRD
≥25%	Not necessary to define relapse	Not necessary to define relapse
Five to <25%	One other test <sup><math>\ddagger</math></sup> with $\geq$ 1% blasts	Not necessary to define relapse
Five to <25%	None	Two tests <sup>†</sup> with with $\geq$ 1% blasts
One to <5%	Two other tests <sup>†</sup> with with $\geq$ 1% blasts	Not necessary to define relapse
One to <5%	Zero or 1 other test <sup>‡</sup> with $\geq$ 1% blasts	Two tests <sup>†</sup> with $\geq$ 1% blasts

\*Second bone marrow evaluation ≥1 wk later.

<sup>†</sup>FCM/PCR/NGS-based MRD or FISH/karyotype/PCR demonstrating leukemia-specific marker or M2/M3 morphology.

#### Table 4. Consensus criteria for relapse definition in pediatric ALL: BM relapse (MRD unavailable)

BM 1		BM 2*
Cytomorphology	Others	Cytomorphology
М3	Not necessary to define relapse	Not necessary to define relapse
M2	One other test <sup>‡</sup> with $\geq$ 1% blasts	Not necessary to define relapse
M2	None	M2
M1	Two other tests <sup>†</sup> with $\geq$ 1% blasts	Not necessary to define relapse

\*Second bone marrow evaluation ≥1 wk later.

<sup>†</sup>FISH/karyotype/PCR demonstrating leukemia-specific marker.

bone marrow. The specific conditions that must be met to provide clear evidence of  $\geq 1\%$  leukemic blasts in the bone marrow include the following:

- If MRD testing is available, then a single marrow is sufficient to define relapse if MRD is ≥25%; or MRD is ≥5% with 1 additional sensitive diagnostic test demonstrating ≥1% blasts (FCM-/PCR-/NGS-MRD or FISH or cytogenetics or RT-PCR of leukemic specific marker or M2/M3 morphology); or MRD is ≥1% with 2 additional sensitive diagnostic tests demonstrating ≥1% blasts.
  - If the MRD is ≥1% but without the required confirmatory tests, then a relapse can still be defined if a consecutive marrow evaluation separated by ≥1

week demonstrates  $\geq$  1% using 2 sensitive diagnostic tests.

- If MRD testing is unavailable, then a single marrow is sufficient to define relapse if M3 morphology or M2 morphology with 1 additional sensitive diagnostic test demonstrating ≥1% blasts (FISH or cytogenetics or RT-PCR of leukemic specific marker) or M1 morphology with 2 additional sensitive diagnostic tests demonstrating ≥1% blasts is present.
  - If the morphology is M2 with no confirmatory tests, then a relapse can still be defined if a consecutive marrow evaluation separated by ≥1 week demonstrates M2 morphology.

#### Table 5. Consensus criteria for relapse definition in pediatric ALL: CNS relapse

CSF 1	CSI	CSF 2*		
Cytomorphology	Cytomorphology	Others		
CNS3 <sup>†</sup>	Not necessary to define relapse	Not necessary to define relapse		
CNS2	CNS2	One other positive test		

\*Second CSF evaluation  $\geq 1$  wk later.

<sup>†</sup>May be defined by cytomorphology, imaging, or biopsy.



Figure 1. Differences and consequences of current vs new consensus relapse criteria: a case study. HSCT, hematopoietic stem cell transplantation; pB-ALL, precursor B-acute lymphoblastic leukemia; TCF3-R, transcription factor 3 rearrangement.

- CNS:
  - CNS3 (by cytomorphology or imaging or biopsy)
  - CNS2 on 2 consecutive cytomorphologic evaluations at least 1 week apart with 1 additional consistent and sensitive diagnostic test present in ≥1 of the evaluations
- Other EM sites:
  - EM tumor burden, confirmed by imaging and/or biopsy. Imaging could be diagnostic if combined with other sites, but a positive biopsy is needed to define relapse if it is the only potential relapse site
- Combined relapse:
  - any BM relapse combined with any non-BM relapse (with each defined above)

The immediate impact of this newly agreed on consensus is illustrated by an example in Figure 1.

## Discussion

Establishing common criteria for response and relapse assessment in pediatric ALL is urgently needed, and the PDL group believes that these criteria should be adopted by all study groups. The implementation of new consensus criteria will entail significant additional effort for most study groups. Amendments to the respective protocols might be required, or it may be necessary to wait until the next clinical trial is implemented. Obviously, any prioritization techniques to determine slow response, treatment failure, and/or relapse will depend on the availability of such diagnostic tests. Currently, crucial end points in clinical trials are poorly defined. This inhibits consistency not only in the definition of EFS but also in eligibility criteria to enroll patients into trials for relapsed or refractory ALL. This creates major challenges in comparing the efficacy of the treatments in such trials. Uniform eligibility for trials for relapsed or refractory ALL may facilitate seamless transition from upfront to second-line trials. It may also reduce the proportion of patients who are taken off protocol as they previously did not qualify for a subsequent clinical trial.

It appears no longer appropriate to require identification of  $\geq$ 25% BM blasts by morphology to define relapse, when an impending relapse can be detected using sensitive, objective, highly reproducible, and well-standardized MRD methods.<sup>7,39,40</sup> The results of trials that do not adopt these response definitions will be confounded if patients are being removed from protocol therapy without meeting a protocol-specified event, which complicates EFS comparisons. For example, individual clinicians, patients, and their families may be unwilling to continue frontline therapy in the face of clear evidence of recurrent disease that is not defined as relapse in the treatment protocol. Such patients may reasonably pursue alternative therapies (eg, chimeric antigen receptor T cells or other immunotherapeutic approaches) before morphologic relapse, especially because recent data suggest that such therapies are more likely to be safe and effective in the MRD setting than following an overt morphologic relapse.<sup>41,42</sup> Indeed, patients may benefit from more timely initiation of alternative salvage therapy. Particularly in the context of clinical trials in relapsed or refractory ALL, uniform eligibility criteria are crucial to compare the efficacy of alternative salvage therapies. Clear

Downloaded from http://ashpublications.net/blood/article-pdf/139/12/1785/1882168/bloodbld2021012328.pdf by guest on 29 May 202-

definitions for CR, TF, relapse, and EFS will be also useful for regulatory agencies in evaluating applications for approval of new drugs or therapies or by expansion of the indication for an approved drug or therapy.

During the work for this project, it was repeatedly emphasized that an actionable threshold for each diagnostic method should be carefully considered, and although the threshold may differ among various methods, the results of FCM- and PCR-MRD are very consistent.<sup>43</sup> This aspect is important because the sensitivity for detection and quantification depends on the method itself, and availability of each method may vary between study groups. For example, for most probes, the sensitivity of FISH testing ranges from 1% to 5%, whereas for FCM-MRD, PCR-MRD, and NGS-MRD, much lower thresholds of disease detection are possible. Even within a given methodology, however, it is extremely difficult to set uniform threshold levels for diagnostic tests in all study groups because individual assays may have different validated sensitivity thresholds. We provide a supplemental Appendix comprising suggestions how thresholds may be defined for different scenarios of test availability.

The authors acknowledge that the MRD technologies used to define response and relapse may not be widely available in lowand middle-income countries. In this case, standard morphologic definitions should be used. However, there is also an urgent need to establish standardized and affordable MRD technologies to more sensitively detect patient responses and improve treatment outcomes.

In summary, the efforts of the PDL group to establish uniform definitions of CR, TF, and relapse among the major international ALL consortia, provided insights into the adaptations that individual groups have made with the ever-advancing technologies to detect microscopic disease in the era of molecular diagnostics. Although it was not possible for the groups to agree on every aspect, important progress was made to come to consensus on major aspects of these definitions. This will facilitate a more uniform interpretation of treatment outcomes in newly diagnosed patients, and perhaps even more importantly, about the benefit of novel therapeutic interventions in patients for whom upfront therapies fail.

### Acknowledgments

The authors thank Julia Alten, Isabel Badell, Claus R. Bartram, Giuseppe Basso, Monika Brüggemann, Mireia Camós, Hélène Cavé, Valentino

#### REFERENCES

- 1. Smith M, Arthur D, Camitta B, et al. Uniform approach to risk classification and treatment assignment for children with acute lymphoblastic leukemia. *J Clin Oncol.* 1996; 14(1):18-24.
- Pui CH, Yang JJ, Hunger SP, et al. Childhood acute lymphoblastic leukemia: progress through collaboration. J Clin Oncol. 2015;33(27): 2938-2948.
- Schrappe M, Nachman J, Hunger S, et al. 'Educational symposium on long-term results of large prospective clinical trials for

Conter, José Luis Dapena, Michael Dworzak, Eva Fronkova, Hester de Groot-Kruseman, Renate Panzer-Grümayer, Oskar Haas, Valerie de Haas, Jochen Harbott, Ondrej Hrusak, Janos Kappelmayer, Der-Cherg Liang, Wolf-Dieter Ludwig, Eva Mejstrikova, Anja Möricke, Manuel Ramírez, Jose M. Ribera, Brigitte Schlegelberger, Martin Stanulla, Maria Grazia Valsecchi, Jan Trka, Marketa Zaliova, Martin Zimmermann, and Jan Zuna for their valuable contribution and input.

This project was supported by Deutsche Krebshilfe grants 108588, 70112517, and 70112958, the Italian Association for Cancer Research (AIRC IG 2017: 20564, AIRC IG 2018: 21724 and IG2015:17593, CRUK/AIRC/FC AECC 22791, AIRC Molecular Clinical Oncology 5 per mille 21147), the French Hospital Program for Clinical Research (PHRC 10-02-05), the Danish Childhood Cancer Foundation (2019-5934), the Danish Cancer Society (R257-A14720), US National Cancer Institute (CA021765), American Lebanese Syrian Associated Charities, the Swedish Childhood Cancer Fund (TJ2017-0076), Japan Agency for Medical Research and Development (AMED) (21ck0106612h0002), Blood Cancer UK, NCTN Operations Center and Statistics & Data Center (SDC) grants (U10CA180886 and U10CA180899, respectively) and the St Baldrick's Foundation (COG).

## Authorship

Contribution: M.S. and P.B. initiated the topic; S.B. and M.S. wrote the paper; and A. Baruchel, A. Biondi, M.B., M.C., G. Cario, G. Cazzaniga, G.E., C.J.H., M.H., S.P.H., C.K., H.-C.L., F.L., M.L.L., A.M., G.M., R.P., C.-H.P., S.R., K.S., L.B.S., J.S., A.V., and P.B. discussed the consensus and reviewed the paper.

Conflict-of-interest disclosure: The authors declare no competing financial interests.

ORCID profiles: M.S., 0000-0003-0034-5845; A. Baruchel, 0000-0003-0120-6089; A. Biondi, 0000-0002-6757-6173; G. Cazzaniga, 0000-0003-2955-4528; M.H., 0000-0001-7637-1949; S.P.H., 0000-0002-5492-3957; H.-C.L., 0000-0001-9721-167X; M.L.L., 0000-0003-4099-4700; A.M., 0000-0002-6698-2348; G.M., 0000-0002-2258-0822; C.-H.P., 0000-0003-0303-5658; S.R., 0000-0002-5658-1831; K.S., 0000-0002-0829-4993; P.B., 0000-0002-8653-1069.

Correspondence: Martin Schrappe, Department of Pediatrics, University Medical Center Schleswig-Holstein, Campus Kiel, Arnold-Heller-Str. 3, Bldg C, 24105 Kiel, Germany; email: m.schrappe@pediatrics.uni-kiel.de.

## Footnotes

childhood acute lymphoblastic leukemia

4. Arber DA, Orazi A, Hasserjian R, et al. The

Organization classification of myeloid

neoplasms and acute leukemia [correction

published in Blood. 2016;128(3):462-463].

5. van der Does-van den Berg A, Bartram CR,

Basso G, et al. Minimal requirements for the

diagnosis, classification, and evaluation of the

treatment of childhood acute lymphoblastic

leukemia (ALL) in the "BFM Family" Cooper-

ative Group. Med Pediatr Oncol. 1992;20(6):

2016 revision to the World Health

Blood. 2016;127(20):2391-2405.

497-505

(1985-2000). Leukemia. 2010;24(2):253-254.

Submitted 19 April 2021; accepted 22 June 2021; prepublished online on *Blood* First Edition 30 June 2021; DOI 10.1182/blood.2021012328.

\*S.B. and M.S. contributed equally to this study.

The online version of this article contains a data supplement.

- Coustan-Smith E, Sancho J, Hancock ML, et al. Clinical importance of minimal residual disease in childhood acute lymphoblastic leukemia. *Blood.* 2000;96(8): 2691-2696.
- Szczepański T, Orfão A, van der Velden VH, San Miguel JF, van Dongen JJ. Minimal residual disease in leukaemia patients. *Lancet* Oncol. 2001;2(7):409-417.
- Nyvold C, Madsen HO, Ryder LP, et al; Nordic Society for Pediatric Hematology and Oncology. Precise quantification of minimal residual disease at day 29 allows identification of children with acute lymphoblastic leukemia

and an excellent outcome. *Blood*. 2002;99(4): 1253-1258.

- Cheson BD, Fisher RI, Barrington SF, et al; United Kingdom National Cancer Research Institute. Recommendations for initial evaluation, staging, and response assessment of Hodgkin and non-Hodgkin lymphoma: the Lugano classification. J Clin Oncol. 2014; 32(27):3059-3068.
- Cheson BD, Cassileth PA, Head DR, et al. Report of the National Cancer Institutesponsored workshop on definitions of diagnosis and response in acute myeloid leukemia. J Clin Oncol. 1990;8(5):813-819.
- Eisenhauer EA, Therasse P, Bogaerts J, et al. New response evaluation criteria in solid tumours: revised RECIST guideline (version 1.1). Eur J Cancer. 2009;45(2):228-247.
- Schrappe M. Treatment protocol for children and adolescents with acute lymphoblastic leukemia—AIEOP-BFM ALL. Available at: https://ClinicalTrials.gov/show/ NCT03643276.
- Schmiegelow K. ALL2008 protocol for childhood acute lymphoblastic leukemia intermittent versus continuous PEG asparaginase. Available at: https:// ClinicalTrials.gov/show/NCT00819351.
- Mc Neer JL. Inotuzumab ozogamicin and post-induction chemotherapy in treating patients with high-risk B-ALL, mixed phenotype acute leukemia, and B-LLy. Available at: https://ClinicalTrials.gov/show/ NCT03959085.
- NCI. A study to investigate blinatumomab in combination with chemotherapy in patients with newly diagnosed B-lymphoblastic leukemia. Available at: https://ClinicalTrials.gov/ show/NCT03914625.
- Baruchel A, Bertrand Y. A French protocol for the treatment of acute lymphoblastic leukemia (ALL) in children and adolescents. Available at: https://ClinicalTrials.gov/show/ NCT02716233.
- Biondi A, Silverman L. Imatinib mesylate and combination chemotherapy in treating patients with newly diagnosed Philadelphia chromosome positive acute lymphoblastic leukemia. Available at: https://ClinicalTrials. gov/show/NCT03007147.
- Koh K, Kato M, Saito AM, et al. Phase II/III study in children and adolescents with newly diagnosed B-cell precursor acute lymphoblastic leukemia: protocol for a nationwide multicenter trial in Japan. *Jpn J Clin Oncol.* 2018;48(7):684-691.
- UKALL. United Kingdom Trial for children and young adults with Acute lymphoblastic Leukaemia and Lymphoma 2011; 2012.

- 20. SEHOP/PETHEMA. Tratamiento de la Leucemia Aguda Linfoblástica de Nuevo Diagnóstico; 2014.
- DCOG. Protocol ALL-11: Treatment study protocol of the Dutch Childhood Oncology Group for children and adolescents (1-19 year) with newly diagnosed acute 2012.
- Heyman M. A treatment study protocol for participants 1-45 years with acute lymphoblastic leukaemia. Available at: https:// ClinicalTrials.gov/show/NCT04307576.
- Biondi A, Silverman L. International phase 3 trial in Philadelphia chromosome-positive acute lymphoblastic leukemia (Ph+ ALL) testing imatinib in combination with two different cytotoxic chemotherapy backbones. Available at: https://www.clinicaltrialsregister. eu/ctr-search/search?query=2017-000705-20.
- 24. Liang D-C. TPOG-ALL-2013. 2012.
- Cario G, Escherich G, Möricke A, et al. New strategies for treatment of children and adolescents with acute lymphoblastic leukemia. *Monatsschr Kinderheilkd*. 2021; 169(1):20-29.
- Silverman LB. Treatment of newly diagnosed acute lymphoblastic leukemia in children and adolescents. Available at: https:// ClinicalTrials.gov/show/NCT03020030.
- Jeha S, Pei D, Choi J, et al. Improved CNS control of childhood acute lymphoblastic leukemia without cranial irradiation: St Jude total therapy study 16. J Clin Oncol. 2019; 37(35):3377-3391.
- Bennett JM, Catovsky D, Daniel MT, et al. Proposals for the classification of the acute leukaemias. French-American-British (FAB) co-operative group. Br J Haematol. 1976; 33(4):451-458.
- 29. Bürger B, Zimmermann M, Mann G, et al. Diagnostic cerebrospinal fluid examination in children with acute lymphoblastic leukemia: significance of low leukocyte counts with blasts or traumatic lumbar puncture. *J Clin Oncol.* 2003;21(2):184-188.
- 30. Mastrangelo R, Poplack D, Bleyer A, Riccardi R, Sather H, D'Angio G. Report and recommendations of the Rome workshop concerning poor-prognosis acute lymphoblastic leukemia in children: biologic bases for staging, stratification, and treatment. Med Pediatr Oncol. 1986;14(3):191-194.
- Mahmoud HH, Rivera GK, Hancock ML, et al. Low leukocyte counts with blast cells in cerebrospinal fluid of children with newly diagnosed acute lymphoblastic leukemia. N Engl J Med. 1993;329(5):314-319.
- 32. Vora A, Andreano A, Pui CH, et al. Influence of cranial radiotherapy on outcome in children with acute lymphoblastic leukemia treated

with contemporary therapy. J Clin Oncol. 2016;34(9):919-926.

- Dworzak MN, Panzer-Grümayer ER. Flow cytometric detection of minimal residual disease in acute lymphoblastic leukemia. *Leuk Lymphoma*. 2003;44(9):1445-1455.
- 34. van der Velden VH, Cazzaniga G, Schrauder A, et al; European Study Group on MRD detection in ALL (ESG-MRD-ALL). Analysis of minimal residual disease by Ig/TCR gene rearrangements: guidelines for interpretation of real-time quantitative PCR data. *Leukemia*. 2007;21(4):604-611.
- Armstrong SA, Look AT. Molecular genetics of acute lymphoblastic leukemia. J Clin Oncol. 2005;23(26):6306-6315.
- Shago M. Recurrent cytogenetic abnormalities in acute lymphoblastic leukemia. *Methods Mol Biol.* 2017;1541: 257-278.
- Gajjar A, Harrison PL, Sandlund JT, et al. Traumatic lumbar puncture at diagnosis adversely affects outcome in childhood acute lymphoblastic leukemia. *Blood.* 2000;96(10): 3381-3384.
- Thastrup M, Marquart HV, Levinsen M, et al; Nordic Society of Pediatric Hematology and Oncology (NOPHO). Flow cytometric detection of leukemic blasts in cerebrospinal fluid predicts risk of relapse in childhood acute lymphoblastic leukemia: a Nordic Society of Pediatric Hematology and Oncology study [correction published in Leukemia. 2020;34:2822]. Leukemia. 2020; 34(2):336-346.
- Pui CH, Campana D. New definition of remission in childhood acute lymphoblastic leukemia. *Leukemia*. 2000;14(5):783-785.
- 40. van Dongen JJ, van der Velden VH, Brüggemann M, Orfao A. Minimal residual disease diagnostics in acute lymphoblastic leukemia: need for sensitive, fast, and standardized technologies. *Blood*. 2015; 125(26):3996-4009.
- Brown P. Blinatumomab for MRD<sup>+</sup> B-ALL: the evidence strengthens. *Blood*. 2018;131(14): 1497-1498.
- Locatelli F, Whitlock JA, Peters C, et al. Blinatumomab versus historical standard therapy in pediatric patients with relapsed/ refractory Ph-negative B-cell precursor acute lymphoblastic leukemia. *Leukemia*. 2020; 34(9):2473-2478.
- Neale GA, Coustan-Smith E, Stow P, et al. Comparative analysis of flow cytometry and polymerase chain reaction for the detection of minimal residual disease in childhood acute lymphoblastic leukemia. *Leukemia*. 2004; 18(5):934-938.

© 2022 by The American Society of Hematology