



Lulla PD. CAR T cells and autologous transplantation can coexist for DLBCL. *Blood*. 2021;139(9):1266-1267.

Page 1266: In the paragraph that begins with “Results from,” the passage “In practice, however, we will still see patients who have received 1 to 2 cycles of salvage chemotherapy (not allowed in ZUMA7 or TRANSFORM) before consideration of a cell therapy, who do not fit the early chemotherapy failure criteria in those trials (ie, refractory to or relapsed within 12 months of first-line chemotherapy), or who cannot readily access a CAR T-cell center. Indeed, the EFS was not different between the standard of care and CAR T cells in the BELINDA trial, which had a design similar to those of ZUMA-7 and TRANSFORM except that it permitted salvage chemotherapy prior to administration of another potent CD19 CAR T-cell product, tisagenlecleucel.⁸” should read, “In practice, however, we will still see patients who have received salvage chemotherapy (not allowed in ZUMA7 and only 1 cycle in TRANSFORM, which is not typical salvage) before consideration of a cell therapy, who do not fit the early chemotherapy failure criteria in those trials (ie, refractory to or relapsed within 12 months of first-line chemotherapy), or who cannot readily access a CAR T-cell center. Indeed, the EFS was not different between the standard of care and CAR T cells in the BELINDA trial, which had a design similar to those of ZUMA-7 and TRANSFORM except that it permitted standard salvage chemotherapy (~48% of patients received 2 or more salvage cycles) prior to administration of another potent CD19 CAR T-cell product, tisagenlecleucel.⁸” The errors have been corrected in the online version of the article.

<https://doi.org/10.1182/blood.2022017891>

© 2022 by The American Society of Hematology

Sangaletti S, Tripodo C, Chiodoni C, et al. Neutrophil extracellular traps mediate transfer of cytoplasmic neutrophil antigens to myeloid dendritic cells toward ANCA induction and associated autoimmunity. *Blood*. 2012;120(15):3007-3018.

Page 3011: In the left column of Figure 2C, the image for necrotic polymorphonuclear leukocytes (PMNs) stained with PKH-26 was inadvertently duplicated for apoptotic PMNs. In the legend to Figure 2, the sentence “Scale bars represent 5 μm ” should not appear in the description of panel A; the sentence “Scale bars represent 30 μm ” should not appear in the description of D and E, and the sentence “The scale bar represents 30 μm ” should be added to the description of panel F. The corrected Figure 2 and legend are shown below.