chemotherapy-refractory acute lymphoblastic leukemia. *Sci Transl Med.* 2013;5(177):177ra38.

- Weber EW, Parker KR, Sotillo E, et al. Transient rest restores functionality in exhausted CAR-T cells through epigenetic remodeling. *Science*. 2021;372(6537): eaba1786.
- Locke FL, Rossi JM, Neelapu SS, et al. Tumor burden, inflammation, and product attributes determine outcomes of axicabtagene ciloleucel in large B-cell lymphoma. *Blood* Adv. 2020;4(19):4898-4911.
- 7. Blank CU, Haining WN, Held W, et al. Defining 'T cell exhaustion'. Nat Rev Immunol. 2019;19(11):665-674.

### LYMPHOID NEOPLASIA

Comment on Largeot et al, page 3166

# Targeting mRNA translation in CLL

**Ralf Küppers** | University of Duisburg-Essen

In this issue of *Blood*, Largeot et al<sup>1</sup> show that flavagline FL3-mediated inhibition of enhanced messenger RNA (mRNA) translation in chronic lymphocytic leukemia (CLL) cells has multiple effects, including blocking proliferation, rewiring MYC-driven metabolism, and controlling CLL growth in a murine model.

CLL is the most frequent leukemia in older people in the Western world. The disease now may often be controlled for years or sometimes may not even require treatment for extended periods. However, CLL still can not be cured, and aggressive forms, such as a transformation into a diffuse large B-cell lymphoma (Richter transformation) or the immunodeficiency caused by CLL, can be rapidly life threatening. Thus, there is a continuing search for cellular vulnerabilities that can be therapeutically exploited. Largeot et al focused on the prior observation that CLL cells have an enhanced protein translation rate compared with normal B cells<sup>2</sup> and on indications that inhibiting mRNA translation may be toxic for CLL cells.<sup>3</sup> The authors first validated an increased translation rate in human CLL cells and then showed that this is also a feature of murine CLL cells in the TCL1-driven murine CLL model. Triggering B-cell receptor (BCR) or Toll-like receptor pathways further increased translation in CLL cells. A known translation inhibitor and modifier, the synthetic flavagline

compound FL3, inhibited translation in CLL cells. FL3 was active at a low concentration, at which most other immune cells were not affected. Thus, CLL cells seem particularly sensitive to translation inhibition, opening a promising therapeutic window. Treatment of activated CLL cells with FL3 modified several pathways, including a reduction of MYC activity. Importantly, FL3 treatment also decreased proliferation of CLL cells, rewired multiple metabolic pathways, and induced apoptosis in these cells. MYC seems to play a major role in the metabolic rewiring, as direct MYC inhibition showed similar effects as FL3 treatment. This may be linked to a positive feedback loop, as strong translation initiation promotes MYC activity, and MYC itself promotes translation.

8. Sen DR, Kaminski J, Barnitz RA, et al. The

Science. 2016;354(6316):1165-1169.

9. Melenhorst JJ, Chen GM, Wang M, et al.

2022;612(7941):E22.

576(7786):293-300.

Decade-long leukaemia remissions with

10. Lynn RC, Weber EW, Sotillo E, et al. c-Jun

overexpression in CAR T cells induces

exhaustion resistance. Nature. 2019;

https://doi.org/10.1182/blood.2023020935

© 2023 by The American Society of Hematology

persistence of CD4(+) CAR T cells. Nature.

epigenetic landscape of T cell exhaustion.

The striking effects of FL3 on CLL cells prompted the authors to investigate its mechanism of action in more detail. An unexpected finding from these studies was that the primary effect of FL3 on its prohibitin targets was not the known impairment of the RAS/RAF pathway.

Rather, in CLL cells prohibitins bind directly to the eukaryotic translation initiation factor complex involving eIF4A, eIF4G, and eIF4F, and this binding is inhibited by FL3 (see figure). Thus, a novel mechanism of FL3-mediated inhibition of translation initiation was identified for CLL cells. The important role of prohibitins for translation initiation in CLL cells was confirmed by showing that silencing of prohibitin expression had similar inhibitory effects on translation as FL3 treatment. Finally, in the CLL mouse model, the authors showed that FL3 treatment of transplanted CLL cells slowed CLL growth and prolonged survival of the mice. Notably, FL3 treatment in this model not only inhibited translation in CLL cells (while not affecting normal B cells) but also in regulatory T cells, which also have a relatively high translation rate. As regulatory T cells promote CLL development,<sup>4</sup> the additional effect of FL3 on these cells may be of added value for FL3 treatment of CLL.

The obvious question is whether these in vitro findings are relevant in vivo. Indeed, most CLL proliferation takes place in proliferation centers in lymph nodes, where CLL cells are in contact with T helper cells and presumably receive autoantigen triggering of their often autoreactive BCRs. In these proliferation centers, a fraction of the CLL cells shows MYC expression.<sup>5</sup> Hence, FL3 may specifically target these proliferating and metabolically active tumor cells, thereby attacking the root of CLL tumor clone expansion.

Several aspects deserve further considerations and study. First, CLL is presumably derived from mature CD5<sup>+</sup> B cells, and MYC expression and activity were reported for a subset of these normal CD5<sup>+</sup> B cells.<sup>6</sup> Hence, a moderate propensity for increased translation may already be an inherent property of the normal counterparts of CLL cells. Second, CLL is usually preceded by the premalignant condition monoclonal B-cell lymphocytosis, in which expanding B-cell clones already carry genetic lesions typical for CLL cells. It would be interesting to know whether these premalignant cells also already show increased translation initiation and metabolic rewiring. If this is the case, would treatment with a translation initiation inhibitor such as FL3 (assuming it



Prohibitins directly binding to the eukaryotic translation initiation complex (shown here are the factors elF4G and elF4F) promote strong translation initiation, which results in CLL cell proliferation, metabolic rewiring, MYC activity, and other effects (not shown). FL3 directly binds to prohibitins, thereby replacing them from the translation initiation factors. As a consequence, translation initiation is reduced, causing reduced proliferation, metabolic rewiring, and MYC activity in CLL cells.

has minimal adverse effects) prevent evolution to CLL? Third, there are 2 forms of CLL, 1 with mutated BCR variable region genes and 1 with unmutated BCR. The latter has an unfavorable clinical behavior, and it shows stronger BCR signaling upon activation and also MYC activation.<sup>7,8</sup> ZAP70-mediated Hence, are unmutated CLL cells more vulnerable to translation inhibition than BCR-mutated CLL, so that such a therapy may be useful only for unmutated CLL? Fourth, Richter transformation of CLL, which has a particularly poor prognosis, is often associated with genetic lesions of MYC causing increased MYC activity.9 This points to an even more pronounced translational dysregulation and metabolic rewiring in this transformed form of CLL. Even though translation inhibition with FL3 alone is unlikely to control this aggressive lymphoma, it is certainly worthwhile to study whether Richter transformation is particularly sensitive to inhibition of translation. In that case, this could become a valuable component of a multimodality treatment of Richter transformation.

Conflict-of-interest disclosure: The author declares no competing financial interests.

### REFERENCES

- Largeot A, Klapp V, Viry E, et al. Inhibition of MYC translation through targeting of the newly identified PHB-eIF4F complex as therapeutic strategy in CLL. *Blood*. 2023;141(26):3166-3183.
- Yeomans A, Thirdborough SM, Valle-Argos B, et al. Engagement of the B-cell receptor of chronic lymphocytic leukemia cells drives global and MYC-specific mRNA translation. *Blood.* 2016;127(4):449-457.
- Willimott S, Beck D, Ahearne MJ, Adams VC, Wagner SD. Cap-translation inhibitor,

4EGI-1, restores sensitivity to ABT-737 apoptosis through cap-dependent and -independent mechanisms in chronic lymphocytic leukemia. *Clin Cancer Res.* 2013;19(12):3212-3223.

 Wierz M, Pierson S, Guyonnet L, et al. Dual PD1/LAG3 immune checkpoint blockade limits tumor development in a murine model of

## MYELOID NEOPLASIA

Comment on Xie et al, page 3184

## IL-34: a novel differentiation therapy for AML?

Nainita Roy and Christopher Y. Park | New York University Grossman School of Medicine

In this issue of *Blood*, Xie et al<sup>1</sup> describe a novel role for interleukin 34 (IL-34) in promoting hematopoietic progenitor and myeloid leukemia differentiation by binding to a previously unrecognized receptor for IL-34, triggering receptor expressed on myeloid cells 2 (TREM2).

Extending their prior observation that osteoclast-specific deletion of tuberous sclerosis 1 (Tsc1) results in increased numbers of osteoclasts,<sup>2</sup> the authors evaluated the hematopoietic system and observed increased granulocytic differentiation and enhanced osteoclast production of IL-34, a cytokine previously shown to promote myeloid maturation.<sup>3</sup> The authors investigated whether IL-34 may promote maturation of acute myeloid leukemia (AML) blasts and found that IL-34 induced differentiation of human AML cell lines and primary mouse and human blasts by binding to TREM2 (see figure). In addition, AML cells engrafted into osteoclast-specific Tsc1-deleted mice resulted in reduced disease aggressiveness. Impressively, IL-34 treatment of mice engrafted with primary human AML cells or human and mouse AML cell lines resulted in enhanced blast differentiation, reduced disease burden, and improved survival.

Although IL-34 has been reported to bind to receptors including colony stimulating factor 1 receptor (CSF1R), protein tyrosine phosphatase receptor type Z1 (PTPz), and syndecan 1 (CD138), deletion of these receptors did not attenuate the differentiation activity of IL-34. Using a combination of approaches, the authors identified TREM2 as a novel receptor required for IL-34's differentiation activity and found that IL-34 binding to TREM2 inhibits

- chronic lymphocytic leukemia. *Blood*. 2018; 131(14):1617-1621.
- Krysov S, Dias S, Paterson A, et al. Surface IgM stimulation induces MEK1/2-dependent MYC expression in chronic lymphocytic leukemia cells. *Blood*. 2012;119(1):170-179.
- Seifert M, Sellmann L, Bloehdorn J, et al. Cellular origin and pathophysiology of chronic lymphocytic leukemia. J Exp Med. 2012; 209(12):2183-2198.
- 7. Chen J, Sathiaseelan V, Moore A, et al. ZAP-70 constitutively regulates gene expression and protein synthesis in chronic lymphocytic leukemia. *Blood*. 2021;137(26): 3629-3640.
- Stevenson FK, Forconi F, Kipps TJ. Exploring the pathways to chronic lymphocytic leukemia. *Blood.* 2021;138(10):827-835.
- Rossi D, Spina V, Deambrogi C, et al. The genetics of Richter syndrome reveals disease heterogeneity and predicts survival after transformation. *Blood*. 2011;117(12): 3391-3401.

https://doi.org/10.1182/blood.2023020420 © 2023 by The American Society of Hematology