


## Follicular lymphoma involving the synovium in rheumatoid arthritis

Yanna Ding and Elaine S. Jaffe, National Cancer Institute



A 79-year-old woman with a prior follicular lymphoma (FL) diagnosed in 2007 underwent knee replacement for long-standing rheumatoid arthritis (RA). At the time of surgery the patient was in remission, with negative imaging studies. The synovial lining showed striking involvement by FL (panel A, original magnification  $\times 1.25$ , hematoxylin and eosin stain). Expanded villous structures were infiltrated by abnormal follicles composed of centrocytes (panel A inset, original magnification  $\times 40$ ). The atypical follicles were strongly positive for CD20 (panel B, original magnification  $\times 5$ , immunohistochemical staining), CD10 (panel C, original magnification  $\times 5$ , immunohistochemical staining), and BCL2 (panel D, original magnification  $\times 5$ , immunohistochemical staining) with surrounding interfollicular CD3<sup>+</sup> T cells (panel E, original magnification  $\times 5$ , immunohistochemical staining).

The neoplastic follicles contained CD21<sup>+</sup> follicular dendritic cell meshworks (not shown). The bone marrow space was negative for lymphoma.

Synovial tissues in RA can show prominent lymphoid aggregates with reactive germinal centers, seen in  $\sim 25\%$  of patients with well-established disease. The reactive follicles express activation-induced cytidine deaminase. FL B cells show homing to germinal centers throughout the lymphoid system, but involvement of the synovium in autoimmune conditions has not been well described. The synovial involvement in this patient was clinically silent, with the symptoms related to long-standing joint damage. Local chronic inflammatory stimulation likely played a role in the unique pattern of dissemination seen in this case.