Check for updates





Blood 142 (2023) 1800–1802

The 65th ASH Annual Meeting Abstracts

# POSTER ABSTRACTS

## 632.CHRONIC MYELOID LEUKEMIA: CLINICAL AND EPIDEMIOLOGICAL

## Characteristics, Outcomes, Tfr Rates in Young Adults with Chronic Myeloid Leukemia in the Tyrosine Kinase Inhibitor Era: A French Observational Study

Laurence Legros, MD PhD<sup>1,2,3</sup>, Francoise Huguet, MD<sup>4</sup>, Pascale Cony-Makhoul, MD<sup>5</sup>, Irit Touitou, PhD<sup>6</sup>, Franck E. Nicolini, MDPhD<sup>7,8</sup>, Delphine Rea, MD PhD<sup>9,3</sup>, Marc G. Berger, MD PhD<sup>10,11</sup>, Madalina Uzunov, MD<sup>12</sup>, Lydia Roy, MD<sup>3,13</sup>, Martine Escoffre Barbe, MD<sup>14</sup>, Agnes Guerci Bresler, MD<sup>15,11</sup>, Gabriel Etienne, MD PhD<sup>3</sup>, Elisabeth Daguenet, MD<sup>16</sup>, Amelie Penot, MD<sup>17</sup>, Martine Gardembas Pain, MD<sup>11,18</sup>, Shanti Ame, MD<sup>19,11</sup>, Eric Jourdan, MD PhD<sup>20</sup>, Hyacinthe A. Johnson-Ansah, MD<sup>3,21</sup>, Valerie Coiteux, MD<sup>3</sup>, Antoine Machet, MD<sup>22</sup>, Eric Lippert, MDPhD<sup>23</sup>, Anne Vekoff<sup>24</sup>, Jean Michel Karsenti, MD<sup>25</sup>, Stéphane Morisset, Biostat<sup>26</sup>, Francois Mahon, MD PhD<sup>11</sup>

<sup>1</sup>Hematology Department, AP-HP, University of Paris Saclay, Bicêtre Hospital, Paris, France

<sup>2</sup>INSERM U1197, Villejuif, France

<sup>3</sup>Centre Léon Bérard, Fi-LMC, Lyon, France

<sup>4</sup>IUCT-Oncopole, Toulouse, France

<sup>5</sup>Department of Hematology, Annecy-Genevois Hospital Centre, Pringy, France

<sup>6</sup>CHU de Nice, Nice, France

<sup>7</sup> Fi-LMC group, Pessac, France

<sup>8</sup>Hématologie Clinique, Centre Léon Bérard, Lyon, France

<sup>9</sup>Hopital Saint Louis AP-HP, Paris, FRA

- <sup>10</sup>Laboratoire d'Hematologie Biologique CHU ESTAING, Clermont-Ferrand, FRA
- <sup>11</sup> FILMC group, Lyon, France

<sup>12</sup>Clinical hematology, Pitie-Salpetriere Hopstal, APHP, Sorbonne Universite, Paris, FRA

<sup>13</sup>Hematology Department, Henri Mondor University Hospital, UPEC, IMRB, Assistance Publique Hopitaux de Paris,

Créteil, France

<sup>14</sup>Department of hematology, Rennes, France

<sup>15</sup> HOPITAUX DE BRABOIS, VANDOEUVRE LES NANCY CEDEX, FRA

<sup>16</sup>Institut de Cancerologie Lucien Neuwirth, CHU Saint Etienne, Saint Etienne, FRA

- <sup>17</sup> Department of hematology, Centre Hospitalier Universitaire Dupuytren, Limoges, France
- <sup>18</sup>Department of hematology, Angers, FRA
- <sup>19</sup>Department of hematology, Strasbourg, FRA

<sup>20</sup>CHU Nimes, Nimes, FRA

<sup>21</sup>Caen University Hospital, Normandy, Hematology Institute, Caen, France

<sup>22</sup> depatment of hematology, Tours, France

<sup>23</sup>Department of hematology, Brest, FRA

<sup>24</sup>APHP-Hopital Saint Antoine, Paris, France

<sup>25</sup>Department of hematology, NICE, FRA

<sup>26</sup>Centre Léon Bérard, Lyon, France

**Introduction**. Chronic myeloid leukemia (CML) is a myeloproliferative disorder with a median age of approximately 60 years. Our study emphasizes the fact that CML can present at younger ages. These patients have significantly different life challenges compared with older patients. The 2020 update of the ELN recommendations mentioned that" TFR may be the main goal for any patient, irrespective of age, but it is clear that the younger the patient the stronger the case for achieving TFR". We report here the main characteristics and TFR data in CML young adults.

**Methods**. The French CML group (FI-LMC) initiated an observational study (CCTIRS no. 14.745) to describe the characteristics of CML in young adult patients in terms of characteristics at diagnosis, treatment choices, tolerance, adherence, therapeutic response, evolution and survival. Patients ranging from 18 to < 30 years with newly diagnosed CML in chronic phase (CP) or accelerated phase (AP) treated with a tyrosine kinase inhibitor (TKI) in first line were eligible for enrolment.

### POSTER ABSTRACTS

#### Session 632

**Results**. The observatory included 253 unselected young adults from 22 centers with newly diagnosed CML in CP (n=246) or in AP (n=7) among which 51.6% were treated within a first line trial. At the time of CML diagnosis, the median patient age was 24 years (range, 18-29 years); 165 patients were males (63%). Median (range) values were: WBC 154.3 G/L (6.5; 841); platelets 374 G/L (51; 1799), hemoglobin 11.3 g/dL (4.1; 18.3), blast cells 1 % (0; 20), spleen below costal margin 4 cm (0; 30). In CP patients, Sokal score was low in 132 (53.6%) patients, intermediate in 46 (18.7%), high in 37 (15.1%) and unknown in 31 (12.6%) and ELTS score was low in 142 (57.7%) patients, intermediate in 48 (19.5%), high in 22 (9%) and unknown in 34 (13.8%). Initial TKI was imatinib alone (52.4%) or in combination with IFN (2.4%) or Ara-C (0.8%), nilotinib alone (24.6%) or asciminib (1.2%). Median follow-up is 106.2 months (range: 0.03; 279.2). Ninety-nine (39.3%) patients remained under first-line TKI at latest follow-up and 21 pts (8.3%) underwent bone marrow transplantation (BMT) for resistance or transformation. Median overall survival at 9 years is 96.9% [95% CI: 94.5- 99.4]. Event free survival without death, transformation, or BMT at 9 years is 93.1% [95% CI: 89.7-96.7]. Six patients died, 1 from sarcoma, 5 from CML including 1 before any therapy (brain hemorrhage). We investigated the TFR strategy. In 169 patients treated for at least 5 years, 56 (33.1%) attempted TKI discontinuation. TKI free survival at 24 months is 57.9% [95% CI: 45.8- 73.2] (Figure 1). Complete analysis comparing characteristics and outcome between the 2 groups of patients (TFR vs no TFR), will be available for ASH presentation.

**Conclusion**. Despite aggressive features including prominent splenomegaly, high WBC counts and high blast cell percentage, TFR seems achievable in young adults with CML.

Disclosures Legros: Correspondances en Hématologie: Consultancy, Honoraria, Speakers Bureau; BMS: Honoraria; Incyte Biosciences: Honoraria; NOVARTIS: Honoraria, Other; PFIZER: Honoraria; AMGEN: Honoraria. Huguet: NOVARTIS: Honoraria; INCYTE: Honoraria; PFIZER: Honoraria. Nicolini: INCYTE BIOSCIENCES: Honoraria, Membership on an entity's Board of Directors or advisory committees, Research Funding; Novartis: Honoraria, Membership on an entity's Board of Directors or advisory committees, Research Funding, Speakers Bureau; SUN pharma: Honoraria, Membership on an entity's Board of Directors or advisory committees; Pfizer: Honoraria, Membership on an entity's Board of Directors or advisory committees; BMS: Honoraria, Membership on an entity's Board of Directors or advisory committees. Rea: Pfizer: Honoraria, Membership on an entity's Board of Directors or advisory committees, Speakers Bureau; INCYTE BIOSCIENCES: Consultancy, Honoraria, Speakers Bureau; Novartis: Consultancy, Honoraria, Membership on an entity's Board of Directors or advisory committees, Speakers Bureau. Berger: NOVARTIS: Research Funding; PFIZER: Consultancy, Research Funding; NOVARTIS: Consultancy, Research Funding; INCYTE: Research Funding. Roy: Incyte biosciences: Honoraria; Novartis: Honoraria, Research Funding, Speakers Bureau; Pfizer: Honoraria; BMS: Honoraria, Research Funding. Guerci Bresler: PFIZER: Honoraria. Etienne: Pfizer: Honoraria; Incyte biosciences: Honoraria; Novartis: Consultancy, Honoraria, Research Funding; BMS: Honoraria. Jourdan: Novartis: Honoraria, Membership on an entity's Board of Directors or advisory committees; Abbvie: Honoraria, Membership on an entity's Board of Directors or advisory committees; BMS: Honoraria, Membership on an entity's Board of Directors or advisory committees; GSK: Honoraria, Membership on an entity's Board of Directors or advisory committees. Johnson-Ansah: PFIZER: Consultancy; GILEAD: Consultancy; NOVARTIS: Consultancy, Honoraria. Coiteux: Pfizer: Honoraria; Incyte Biosciences: Honoraria, Speakers Bureau; Novartis: Honoraria. Machet: novartis: Honoraria; pfizer: Honoraria.





Figure 1

https://doi.org/10.1182/blood-2023-173132