



The 65th ASH Annual Meeting Abstracts

POSTER ABSTRACTS

632. CHRONIC MYELOID LEUKEMIA: CLINICAL AND EPIDEMIOLOGICAL

Mutation of Epigenetic Regulators at Diagnosis Is an Independent Predictor of Tyrosine Kinase Inhibitor Treatment Failure: A Report from the Residiag Study

Hippolyte Guerineau, PharmD¹, Jean-Michel Cayuela, PharmD, PhD^{2,3}, Stéphanie Dulucq, PharmD, PhD^{3,4,5}, Violaine Tran Quang, PharmD¹, Sihem Tarfi, PharmD¹, Quentin Barathon⁶, Corine Joy⁶, Orianne Wagner-Ballon, MDPH¹, Stéphane Morisset, Biostat^{7,3}, Franck E. Nicolini, MDPH^{8,3}, Lydia Roy, MD^{3,9}, Sébastien Maury⁹, Gabriel Etienne, MD PhD^{10,3}, Delphine Rea, MD PhD^{11,3}, Ivan Sloma, PhDPharmD^{1,3}

¹Hematology and Immunology Department, Henri Mondor University Hospital, UPEC, IMRB, Assistance Publique Hopitaux de Paris, Créteil, France

²University Hospital Saint-Louis, AHP and Paris DIDEROT, Paris, FRA

³Centre Léon Bérard, Fi-LMC, Lyon, France

⁴Laboratoire Hématopoïèse Leucémique Et Cible Th, Bordeaux Cedex, FRA

⁵Laboratory of Hematology, Hôpital Haut Lévêque, Centre Hospitalier Universitaire de Bordeaux, Pessac, France

⁶Hematology and Immunology Department, Henri Mondor University Hospital, Assistance Publique-Hôpitaux de Paris, Créteil, France

⁷Centre Léon Bérard, Lyon, France

⁸Hematology department & CRCL INSERM U1052, Centre Léon Bérard, Lyon, France

⁹Hematology Department, Henri Mondor University Hospital, UPEC, IMRB, Assistance Publique Hopitaux de Paris, Créteil, France

¹⁰Hematology department, Institut Bergonié, Bordeaux, France

¹¹Hematology Department, Hôpital Saint-Louis, Paris, France

Introduction

Additional mutations at chronic myeloid leukemia (CML) diagnosis have been shown to variably affect tyrosine kinase inhibitor (TKI) response and were inconstantly detected at loss of response. Contradictory observations may have resulted from difficulties in reliably inferring CML clonal architecture from mutations quantified by NGS, *BCR::ABL1* by *qRT-PCR*, *ABL1-TK* by RNA-Seq. The RESIDIAG study was designed to identify at diagnosis molecular markers predictive of TKI treatment failure and analyze the clonal dynamics using asymmetric capture sequencing (aCAP-Seq).

Methods

DNA samples from 60 TKI responders (median follow-up: 7.1 years) were sequenced at diagnosis by aCAP-Seq, allowing quantification and breakpoint sequencing of genomic *BCR::ABL1* fusion together with 43 myeloid genes, including *ABL1* with a limit of detection of ~0.1% for *BCR::ABL1* and 0.5-1% for SNVs and Indels. In parallel, CML patients who experienced treatment failure (ELN 2020) were sequentially analyzed at diagnosis and at failure (n=53), at diagnosis only (n=1), and at failure only (n=3). Small deletions and *BCR::ABL1* breakpoints were analyzed to search for microhomology-mediated end-joining (MMEJ) repair signatures. Mutation analysis followed strict medical standards.

Results

The proportion of patients receiving imatinib at diagnosis wasn't statistically different (p=0.08; Pearson's Chi-square test) between responders (70%) and the failure group (56%). At diagnosis, the number of mutations was higher (p<0.001, t-test with Welch correction) in the failure group (0.9±0.1) as compared with responders (0.18±0.05). *ASXL1*, *DNMT3A*, and *TET2* were more frequently mutated at diagnosis in the failure group (38.6%, p<0.001; 10.5%, p=0.01, and 4.3%, p=0.025 respectively; Pearson's Chi² test) as compared with responders (6.7%, 0% and 0% respectively) and the presence of a mutation in one of these genes was highly associated with a reduced failure-free survival (p<0.001, Log-ranked test, Figure 1). An MMEJ signature at *BCR::ABL1* breakpoints was more frequent (p=0.014, Pearson's Chi² test) in the failure group (29.1%) than in the responder group (9.1%), and was significantly associated with the ELTS score (p=0.005, Pearson's Chi² test) but not the Sokal score (p=0.017, Pearson's Chi² test). At first failure, 69% of patients had additional mutations in *ASXL1* (39%), *ABL1-TK* (38%), *DNMT3A* (18%), or *TET2* (15%), while other genes were rarely (<5%) mutated. *ASXL1* mutations found at diagnosis were

still detectable in 18/22 patients at failure and VAF suggested that double mutant *BCR::ABL1/ASXL1* clones were driving TKI failure with or without *BCR::ABL1* tyrosine kinase domain mutation (TKD). *ASXL1* mutations disappearing upon treatment were mostly at low frequency <5% at diagnosis and thus possibly not clonally related to CML. Furthermore, a stepwise selection of variables followed by multivariate analysis with the Fine-Gray model identified the age of patients (sdHR [95% C.I.] = 0.99 [0.97-1.00]; p=0.08) and the presence of a mutation in epigenetics regulators at diagnosis as factors affecting the onset of TKD mutations (sdHR = 3.01 [1.76-5.15]; p<0.001). Intriguingly, co-occurring mutations in *DNMT3A* and *TET2* were found in five patients at TKI failure. Their allelic frequency kinetics suggested that these belonged to the same clone driving CML relapse in 4/5 CML patients, while one patient also had a TKD mutation likely involved in CML relapse. Finally, multivariate COX regression analysis identified two independent predictors of TKI failure at diagnosis: a high-risk ELTS score (p=0.001; hazard ratio=3.42 [1.62-7.22] with low-risk as baseline) and a mutation in *ASXL1*, *DNMT3A* or *TET2* (p=0.002; hazard ratio=2.87 [1.49-5.51]).

Conclusion

Altogether these results strongly point to the contribution of epigenetic regulator mutations in the emergence of TKI failure in CML and warrant further biological studies to better understand the underlying mechanisms

Disclosures Dulucq: Novartis, Incyte Biosciences: Honoraria, Speakers Bureau. **Wagner-Ballon:** Novartis: Honoraria; Alexion: Consultancy, Honoraria. **Nicolini:** Pfizer: Honoraria, Membership on an entity's Board of Directors or advisory committees; SUN pharma: Honoraria, Membership on an entity's Board of Directors or advisory committees; INCYTE BIOSCIENCES: Honoraria, Membership on an entity's Board of Directors or advisory committees, Research Funding; Novartis: Honoraria, Membership on an entity's Board of Directors or advisory committees, Research Funding, Speakers Bureau; BMS: Honoraria, Membership on an entity's Board of Directors or advisory committees. **Roy:** Novartis: Honoraria, Research Funding, Speakers Bureau; Pfizer: Honoraria; BMS: Honoraria, Research Funding; Incyte biosciences: Honoraria. **Etienne:** Novartis: Consultancy, Honoraria, Research Funding; BMS: Honoraria; Pfizer: Honoraria; Incyte biosciences: Honoraria. **Rea:** Pfizer: Honoraria, Membership on an entity's Board of Directors or advisory committees, Speakers Bureau; INCYTE BIOSCIENCES: Consultancy, Honoraria, Speakers Bureau; Novartis: Consultancy, Honoraria, Membership on an entity's Board of Directors or advisory committees, Speakers Bureau. **Sloma:** Novartis: Honoraria, Speakers Bureau; Incyte Biosciences: Honoraria, Speakers Bureau; Agilent Technologies: Patents & Royalties.




Figure 1

<https://doi.org/10.1182/blood-2023-186271>

Downloaded from http://ashpublications.net/blood/article-pdf/142/Supplement_1/4540/2186631/blood-141-main.pdf by guest on 18 May 2024