



## The 65th ASH Annual Meeting Abstracts

## ONLINE PUBLICATION ONLY

## 501. HEMATOPOIETIC STEM AND PROGENITOR CELLS AND HEMATOPOIESIS: BASIC AND TRANSLATIONAL

**Interleukin-17A Directly Signals to Bone Marrow-Derived Hematopoietic Stem and Progenitor Cells to Promote Their Expansion and Differentiation in Vitro**

Cristobal González Losada, MD<sup>1,2</sup>, William Wai Lam Poon, M.Sc.<sup>1,2</sup>, Matthew Salaciak, M.Sc.<sup>1,2</sup>, Samy Amghar<sup>1,2</sup>, Nathalie A. Johnson, MDPhD<sup>2,1</sup>, Bahram Sharif-Askari, PhD<sup>2</sup>, Francois E. Mercier, MD<sup>2,1</sup>

<sup>1</sup> McGill University, Montreal, Canada

<sup>2</sup> Lady Davis Institute for Medical Research, Jewish General Hospital, Montreal, Canada

**Rationale:** IL-17 is a family of six pro-inflammatory cytokines (named from A to F) with distinct patterns of cellular expression. Among these, IL-17A is secreted by lymphoid subsets such as T-helper 17 cells and innate lymphoid cells type 3. Through its action on many tissues, IL-17A is recognized as a key mediator of response to infection and pathogenic states such as cancer and auto-immunity. Until now, the main effect of IL-17A on hematopoiesis has been attributed to an indirect loop through IL-17RA signaling on bone marrow stromal cells, which stimulates secretion of hematopoietic factors such as G-CSF. However, we hypothesized that IL-17A can directly signal to hematopoietic stem and progenitor cells (HSPCs) to determine their self-renewal and differentiation.

**Methodology:** We analyzed the expression of the IL-17 signaling pathway in publicly available human and mouse hematopoietic single-cell RNAseq datasets. Mouse HSPCs were isolated using FACS to perform clonogenic and differentiation assays in Methocult (StemCell Technologies) and StemPro-34 (Gibco). Spectral flow cytometry (SONY ID7000) was used for analysing IL-17RA expression and differentiation of mouse HSPCs. IL-17A levels were measured in blood and bone marrow serum using ELISA (Invitrogen).

**Results:** IL-17RA is expressed at the surface of mouse HSPCs, with greatest levels in common lymphoid progenitors and granulocyte-monocyte progenitors. mRNA expression of IL-17RA signaling pathway is also detectable in human HSPCs, with greatest levels in young individuals. IL-17A supplementation leads to an increase in clonogenic capacity of mouse HSPCs in methylcellulose colony-formation assays. In liquid culture, supplementation with IL-17A leads to a greater expansion of sorted Lin<sup>-</sup>/s-kit<sup>+</sup>/Sca-1<sup>+</sup> progenitors, accompanied by an increased proportion of myeloid-committed cells. IL-17A is also detected in the bone marrow serum, with possible production from subsets of cells expressing ROR $\gamma$ t, a master regulator in development of T helper 17 (Th17) cells.

**Conclusions:** Our work suggests the ability of IL-17A to directly promote clonogenic potential of HSPCs and drive a myeloid-biased lineage commitment in progenitor subsets. Additional studies are required to characterize the response of HSPCs to direct IL-17A signaling in vivo. Considering the important role of IL-17A as a mediator of immune response, and tissue repair, further studies are warranted to elucidate whether modulation of this pathway may be therapeutic in the context of hematopoietic failure or malignancy.

**Disclosures Poon:** Strand Therapeutics: Current Employment. **Johnson:** Roche: Consultancy, Honoraria; Merck: Consultancy, Honoraria; Abbvie: Consultancy; Gilead: Consultancy.

<https://doi.org/10.1182/blood-2023-190975>