Augmented Intracellular Glutathione Inhibits Fas-Triggered Apoptosis of Activated Human Neutrophils

By R. William G. Watson, Ori D. Rotstein, Maria Jimenez, Jean Parodo, and John C. Marshall

Agonist signals delivered through cell surface Fas induce apoptosis. However, the apoptotic program can be modulated by signals from the environment, and in particular, by signals delivered through adhesion molecules. Because neutrophil functional activity in inflammation is contingent on cell survival, and because circulating neutrophils normally die rapidly through a constitutively expressed apoptotic program, we evaluated Fas-mediated apoptosis in resting and inflammatory human neutrophils. We show that normal neutrophils respond to Fas engagement with accelerated rates of apoptosis, but cross-linking of β_2 integrins or priming with bacterial lipopolysaccharide (LPS) prevents this increase. Adhesion molecule cross-linking results in increased intracellular glutathione (GSH). Augmentation of in-

F AS (CD95/APO-1), A MEMBER of the tumor necrosis factor (TNF)/nerve growth factor receptor family, is a type 1 transmembrane receptor that is expressed constitutively on a variety of cell types.¹ Ligation of Fas by agonistic antibody or by its physiologic ligand, Fas ligand (FasL), transduces a signal leading to apoptotic cell death.² Although Fas itself lacks a discrete intracellular signaling domain, it aggregates with a family of membrane-associated molecules³ that, in turn, generate a death signal either directly,^{4,5} or through the downstream activation of proteases of the ICE/ CED3 family.⁶

Fas-mediated signaling for apoptotic cell death is centrally involved in the deletion of autoreactive T cell clones,⁷ and is the mechanism for elimination of transformed cells by natural killer cells.⁸ Following their release from bone marrow stores, neutrophils are induced to die a rapid death by apoptosis, with the result that the entire circulating population is eliminated within 72 hours.⁹ Since neutrophils express both Fas and its physiologic ligand, it has been proposed that autocrine Fas/FasL interactions mediate the constitutive apoptosis of this cell population.¹⁰

Rapid expression of programmed cell death in the neutrophil may serve to limit nonspecific tissue injury resulting from the release of proteases and oxygen intermediates by activated neutrophils. The apoptotic program in the neutrophil is exquisitely sensitive to the influence of factors in the local inflammatory environment. Neutrophil apoptosis is delayed, for example, by bacterial lipopolysaccharide (LPS),¹¹ or host-derived cytokines including granulocyte colony-stimulating factor, granulocyte-macrophage colonystimulating factor, interleukin-8 (IL-8), IL-1, and interferon- γ .¹¹⁻¹³ Its expression is similarly delayed by the process of transmigration into an inflammatory focus, an effect mediated in part by cross-linking of the β_2 integrins, CD11a and CD11b.¹⁴ Conversely, the phagocytosis of *Escherichia coli*,¹⁵ and cytokines such as IL-4, transforming growth factor- β (TGF- β), and TNF- α^{16-18} accelerate expression of the apoptotic program.

Fas engagement signals for the programmed cell death of a number of hematopoietic cells, including lymphocytes,¹⁹ eosinophils,²⁰ monocytes,²¹ and neutrophils.⁹ Fas-triggered signaling in the monocyte can be modulated by intracellular tracellular GSH with exogenous GSH or N-acetylcysteine is sufficient to reduce the Fas-triggered increase in apoptotic rates. Prevention of the activation induced GSH increase by buthionine sulfoximine, a cell permeable inhibitor of GSH biosynthesis, restored Fas responsiveness in activated neutrophils, an effect that could be blocked with exogenous GSH. Taken together, these data show that Fas-induced signaling for neutrophil apoptosis is blocked in a redox sensitive manner by costimulatory signals delivered through β_2 integrins or activation by LPS, and provide a biologic explanation for sustained neutrophil survival in the inflammatory environment.

© 1997 by The American Society of Hematology.

reactive oxygen intermediates.²¹ Therefore, we sought to evaluate the role of Fas-triggered signaling for apoptosis in the neutrophil, and to determine its susceptibility to modulation during neutrophil activation. We show that resting neutrophils respond to Fas engagement with accelerated rates of apoptosis, but that activated neutrophils are refractory to signals delivered through Fas. Moreover, the studies suggest that this state of nonresponsiveness to Fas-mediated signals results from elevated levels of intracellular thiols.

MATERIALS AND METHODS

Reagents. Dulbecco's modified Eagle's medium (DMEM), penicillin, and streptomycin solution, L-glutamine, phosphate-buffered saline, and fetal calf serum (FCS) were purchased from GIBCO Life Technologies Ltd (Burlington, Ontario, Canada). Dextran T500 and Ficoll were purchased from Pharmacia (Baie D'Urfe, Quebec, Canada). The endothelial cell line, ECV-304, was supplied by American Type Culture Collection (ATCC, Rockville, MD). LPS from *E coli* 0111:B4 was purchased from Difco Laboratories (Detroit, MI) and dihydrorhodamine 123 from Molecular Probes (Eugene, OR). All other chemicals, unless otherwise stated, were purchased from Sigma Chemical Co (San Diego, CA). E-Lyse was supplied by Cardinal Associates Inc (Santa Fe, NM).

Antibodies. Murine antibodies to human CD11b (Bear-1) and Fas (CH-11), a mouse IgM negative control, fluorescein isothiocyanate (FITC)-conjugated murine antihuman Fas (UB2), FITC-labeled goat antimouse IgG, and $F(ab')_2$ were purchased from Immunotech

From the Department of Surgery, The Toronto Hospital, University of Toronto, Toronto, Canada.

Submitted November 8, 1996; accepted January 21, 1997.

Supported in part by a Grant from the Canadian Liver Foundation (Toronto, Ontario), the Surgical ICU Research Group, the Toronto Hospital, and the Medical Research Council (Ottawa, Ontario, Canada).

Address reprint requests to John C. Marshall, MD, Eaton N 9-234, Toronto Hospital, General Division, 200 Elizabeth St, Toronto, Ontario, Canada M5G 2C4.

The publication costs of this article were defrayed in part by page charge payment. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. section 1734 solely to indicate this fact.

© 1997 by The American Society of Hematology. 0006-4971/97/8911-0016\$3.00/0





Fig 1. Effects of anti-Fas IgM ligation on neutrophil apoptosis. Human neutrophils were incubated with increasing concentrations of anti-Fas IgM, CH-11, and assessed for apoptosis after 18 hours. Apoptosis was measured by flow cytometry using PI DNA staining with representative histograms shown in (A) and the percentage of cells with apoptotic DNA plotted in Fig 1B. (*P < .05 v ng/mL anti-Fas). The high-intensity peak at a relative fluorescence intensity of 500 Ln represents normal DNA, whereas the lower peak between 10 and 100 Ln intensity indicates apoptotic or fragmented DNA with reduced uptake of propidium iodide. Results are the mean \pm SD for n = 5 experiments with three separate donors of neutrophils.

Table 1. Neutrophil Respiratory Burst Activity, Adhesion Receptor Expression and Fas Expression in Response to Activating Factors

	Control	LPS (1 µg/ mL)	Migrated	Cross- linked
Reactive oxygen				
intermediates				
(LnMCF)	14.4 ± 12.0	34.9 ± 3.6*	22.1 ± 3.6*	$25.6 \pm 4.0*$
CD11b (LnMCF)	$\textbf{88.3} \pm \textbf{9.4}$	$144.3 \pm 11.5^{*}$	$120.0 \pm 7.3^{*}$	—
L-selectin (LnMCF)	12.3 ± 2.7	$3.9 \pm 0.6^{*}$	6.1 ± 2.0*	_
Fas (LnMCF)	$\textbf{8.4} \pm \textbf{3.9}$	$\textbf{8.8} \pm \textbf{2.6}$	$\textbf{9.1} \pm \textbf{3.1}$	$\textbf{7.9} \pm \textbf{1.8}$

Neutrophils were incubated with LPS (1 μ g/mL), migrated or cross-linked with anti-CD11b and then assessed for reactive oxygen intermediates 5, 10, and 8 minutes, respectively. Surface receptor expression of CD11b and L-selectin were assessed in LPS and migrated neutrophils. Results are the mean \pm SD of n = 3 experiments.

*P < .05 (not performed because of cross-linking interface to antibodies).

(Burlington, Ontario, Canada). Murine antihuman CD11b (Bear-1) and L-selectin (Dreg-56) FITC-conjugated antibodies were purchased from Becton Dickinson (Mountain View, CA).

Isolation of human neutrophils. Blood from 10 healthy volunteers was collected in heparinized tubes. Neutrophils (PMN) were isolated by dextran sedimentation and centrifugation through a discontinuous ficoll gradient; red blood cells were lysed using E-Lyse.¹⁰ The PMN pellet was resuspended in DMEM supplemented with 10% FCS, 1% glutamine, and 1% penicillin/streptomycin solution at a concentration of 1×10^6 cells/mL. Cells were incubated in polypropylene tubes to prevent adherence. PMN purity was greater than 95% as assessed by size and granularity on flow cytometry.

Quantification of PMN apoptosis. Neutrophil apoptosis was quantitated using the technique of Nicoletti et al.²² Briefly, PMN cell suspensions were centrifuged at 200g for 10 minutes. The cell pellets were gently resuspended in 1 mL of hypotonic fluorochrome solution (50 μ g/mL PI, 3.4 mmol/L sodium citrate, 1 mmol/L Tris, 0.1 mmol/L EDTA, 0.1% Triton X-100 [Sigma]) and stored in the dark at 4°C for 3 to 4 hours before they were analyzed using a Coulter EPICS XL-MCL cytofluorometer (Miami, FL). A minimum of 5,000 events were collected and analyzed. Apoptotic nuclei were distinguished from normal nuclei by their hypodiploid DNA; debris was excluded from analysis by raising the forward threshold. All measurements were performed under the same instrument settings.

Neutrophil migration. Neutrophil transmigration was performed as previously described.¹⁴ Briefly, transwell chambers, 6-well size (Costar, Cambridge, MA), with polycarbonate membranes of 3.0 μ m pore size were coated for 1 hour with fibronectin (50 μ g/mL), then seeded with the endothelial cell line ECV-304 (1 × 10⁶ /mL). After 24-hours incubation at 37°C and 5% CO₂, plates were washed three times in assay medium. Confluence of ECV-304 monolayer with tight cell junctions was confirmed by the ability of the monolayer to prevent diffusion of radiolabeled ¹²⁵I-labeled albumin diffusion. Control PMN, and PMN incubated with LPS (1 μ g/mL for 1 hour) (1 × 10⁶ /mL) were added to the top chamber and allowed to migrate in response to FMLP (10⁻⁷ mol/L), for 45 minutes at 37°C and 5% CO₂. Migrated PMN were collected from the bottom chamber, washed, and resuspended at 1×10⁶ PMN/mL, a concentration that necessitated pooling cells from 2- to 3-bottom chambers.

Cross-linking. To cross-link β_2 -integrin receptors, PMN (2 × 10⁶/200 μ L) were incubated at 4°C for 20 minutes with differing concentrations of monoclonal antibody, then washed twice in cold Hanks' balanced salt solution (HBSS) without Mg²⁺/Ca²⁺. Cells were then incubated with F(ab')₂ fragments for 20 minutes at 4°C, to cross-link the primary antibody. Washed PMN were resuspended at 1 × 10⁶/mL and incubated at 37°C before study. To confirm cross-linking, FITC-conjugated F(ab')₂ fragment was used to cap the primary antibodies, then intensity of binding was analyzed by flow cytometry.¹⁴

Respiratory burst activity. Generation of reactive oxygen intermediates was assessed by the technique of Smith et al.²³ Briefly, neutrophils (5×10^5 /mL) were incubated with 1 μ mol/L dihydrorhodamine 123 for 5 minutes at 37°C and then stimulated with PMA (5 nmol/L) for 10 minutes. Neutrophil fluorescence intensity was assessed by flow cytometry and expressed as Ln mean channel fluorescence (LnMCF).

Chemiluminescence. Whole cell chemiluminescence was assessed by the method of Allen et al.²⁴ Briefly, 20 μ L of cells at a concentration of 5×10⁶ /mL was added to 400 μ L of luminol (5-amino-2,3-dihydro-1,4-phthalazinedione) balanced salt solution. PMA (5 nmol/L) stimulated chemiluminescence was followed over a 60-minute period using an Automat LB 953 (Wildbad, Germany). Total chemiluminescence during this interval was integrated using software provided by the manufacturer.

Assay of intracellular glutathione (GSH). GSH was assessed using a DTNB (5,5'-dithiobis ([2-nitrobenzoic acid])-based assay as described by Jocelyn et al.²⁵ Isolated PMN (5×10^6 /mL) were sonicated for 30 seconds in 300 µL of 5% 5-sulfosalicylic acid, and centrifuged for 10 minutes at 10,000g. The resultant acid thiol extract was assayed for nonprotein sulfhydryls by quantitating the reduction of DTNB through its conversion to TNB (5-thiol-2 nitrobenzoic acid) at 412 nm using a spectrophotometer. Sample values were then calculated from a standard curve generated using known amounts of GSH and expressed as nmol/10⁷ PMN.

Statistics. Statistical analysis was performed using ANOVA with Scheffe's correction; significance was assumed for values of P < .05.

RESULTS

Ligation of Fas induces neutrophil apoptosis. Incubation of resting neutrophils with an agonistic anti-Fas antibody (anti-Fas IgM), CH-11, caused a dose-dependent increase in rates of spontaneous apoptosis in vitro (Fig 1A and B). Neutrophils incubated with anti-Fas IgM (CH-11, 100 ng/mL) and washed after 1 hour also showed the characteristic morphologic features of apoptosis when examined by light microscopy (data not shown).

Surface expression of Fas on the neutrophil was evaluated using an FITC-conjugated murine anti-Fas IgM. Resting neutrophils express Fas with a relative fluorescence intensity of 8.2 ± 0.3 LnMCF, significantly higher than the binding of an isotype-matched mouse IgM negative control (data not shown).

Activated neutrophils are resistant to Fas-triggered apoptosis. We have previously shown that rates of spontaneous apoptosis are retarded in neutrophils that have migrated into an inflammatory focus; this refractory state can be reproduced in vitro by exposure to LPS, by transmigration across an endothelial cell monolayer in response to fMLP, or by cross-linking of cell surface β_2 integrins.¹⁴ Each of these maneuvers resulted in functional and phenotypic changes of activation, evidenced by increased levels of intracellular reactive oxygen intermediates, upregulation of cell surface CD11b, and shedding of L-selectin; none, however, altered Fas expression (Table 1). The increased levels of reactive oxygen intermediates, although statistically significant, were biologically modest: PMA treatment of resting neutrophils (5 nmol/L for 10 minutes) generated levels of reactive oxygen intermediates of 495 \pm 48 LnMCF, or 14fold higher than those resulting from LPS stimulation.

Neutrophils incubated with LPS (1 μ g/mL) for 1 hour showed a significant delay in rates of spontaneous apoptosis. Unlike resting neutrophils, LPS primed neutrophils were refractory to the proapoptotic effects of incubation with anti-Fas IgM at doses of up to 200 ng/mL. Similarly transmigrated neutrophils and neutrophils in which the β_2 -integrin CD11b had been cross-linked failed to respond to anti-Fas IgM engagement with increased rates of apoptosis (Fig 2).

The refractoriness of activated neutrophils to anti-Fas IgM-triggered apoptosis was a transient, reversible phenomenon. By 4 hours after exposure to LPS or cross-linking of CD11b, cultured cells again responded to anti-Fas IgM engagement with an accelerated apoptotic rate (Fig 3). The restoration of susceptibility to anti-Fas antibody required that the LPS and CD11b antibody be washed out of the culture plates.

Fas engagement does not alter levels of intracellular reactive oxygen intermediates or thiols. Depletion of endogenous antioxidant thiols induces apoptosis of resting neutrophils²⁶; conversely, superoxide anion has been shown to inhibit anti-Fas IgM-induced apoptosis in a TNFR p55-Fas transfected M14 cell line.²⁷ Activated neutrophils manifest a rapid but transient increase in levels of intracellular reactive oxygen intermediates and GSH (Table 1 and Fig 4). Therefore, we hypothesized that altered responsiveness to anti-Fas IgM in activated neutrophils might occur through a redoxsensitive mechanism.

Engagement of Fas had no effect on levels of intracellular reactive oxygen intermediates as measured by oxidation of dihydrorhodamine or chemiluminescent response to PMA (5 nmol/L) (Table 2); similarly, Fas engagement alone had no effect on GSH levels (Table 2).

Increased levels of GSH inhibit Fas-induced apoptosis. Although Fas induced apoptosis did not require changes in the intracellular redox state, refractoriness to Fas-induced apoptosis in activated neutrophils was associated with increased levels of reactive oxygen intermediates and GSH. We have previously showed that depletion of GSH induces neutrophil apoptosis, an effect that can be reversed by the antioxidant, NAC (N-acetylcysteine). Therefore, we hypothesized that elevated GSH levels following neutrophil activation might block Fas-mediated apoptotic signaling.

The antioxidant NAC is rapidly deacylated and converted to glutathione after its uptake by the cell.²⁸ Augmentation of intracellular GSH levels by incubation of neutrophils with either authentic GSH (25 mmol/L) or NAC (25 mmol/L) raised intracellular concentrations of GSH to levels observed in activated neutrophils (Fig 5A), and reduced the percentage of cells undergoing apoptosis in response to anti-Fas IgM (Fig 5B). Although apoptotic rates were reduced, they did not return to baseline levels. Having shown that elevation of intracellular GSH was sufficient to inhibit Fas-triggered apoptosis of resting neutrophils, we then sought to determine whether elevation of GSH was necessary to Fas-triggered apoptosis of activated cells.

In resting cells, Fas ligation triggered apoptosis without a corresponding increase in intracellular GSH; LPS induced both a rise in concentrations of intracellular GSH and a reduction in apoptotic rates that was not altered by ligation



Fig 2. Effects of anti-Fas IgM on apoptotic rates of activated neutrophils. Neutrophils $(1 \times 10^6/mL)$ incubated with LPS $(1 \mu g/mL)$, migrated through an endothelial cell/fibronectin monolayer, or cross-linked with antibodies to CD11b, were incubated with anti-Fas IgM (100 or 200 ng/mL) for 1 hour before being washed. Apoptosis was assessed after 18-hour incubation in medium alone by flow cytometry. *P < .05 versus control, \$P < .05 versus medium alone. Results are the mean \pm SD for n = 5 experiments with four separate donors of neutrophils.

of Fas (Fig 6A). Buthionine sulfoximine (BSO) is a specific inhibitor of GSH biosynthesis, but has no effect on extant levels of preformed protein.²⁹ BSO pretreatment did not alter GSH levels or Fas-induced apoptosis in resting neutrophils (data not shown). Pretreatment of neutrophils with 50 mmol/ L BSO before LPS exposure prevented the LPS-induced increase in GSH (Fig 6A), but not the LPS-induced reduction in apoptosis (Fig 6b). However, blockade of the LPS-triggered increase in GSH resulted in restoration of normal responsiveness to Fas (Fig 6B). Restoration of GSH levels after BSO and LPS treatment with exogenous GSH (25 mmol/ L) was associated again with a resistance to Fas-induced apoptosis.

Together these studies show that augmentation of intracellular GSH concentrations is both necessary and sufficient to prevent Fas-mediated apoptosis of activated neutrophils.

DISCUSSION

Engagement of Fas or the p55 component of the receptor for TNF (TNFR1) results in rapid cell death through the controlled process of apoptosis.¹ Such a response has clear adaptive benefit to the host in facilitating the elimination of transformed cells,⁸ or terminating an activated inflammatory response.9 Yet the interaction of TNF with its cellular receptor can also activate NF- κ B,³⁰ leading to enhanced proinflammatory gene transcription, a response that supports antimicrobial activity in the inflammatory microenvironment. Fas engagement has similarly been reported to activate NF- κ B leading to IL-6 synthesis in an SV80 fibroblast cell line,³¹ although not in an L929 cell line stably transfected with Fas cDNA.32 Thus, a single signal delivered through a common receptor can produce fundamentally divergent responses: cellular activation or cellular death; the mechanism underlying the preferential expression of one response or other is unknown.

A variety of biologic functions are altered as a consequence of the adherence of neutrophils to biologic substrata,³³ and cell surface adhesion molecules are now recognized to function as signaling molecules, altering gene expression in response to stimuli originating external to the cell,³⁴ and, in particular, modulating the expression of programmed cell death.³⁵ We have shown that engagement of adhesion molecules involved in the delivery of neutrophils to an inflammatory focus alters the temporal expression of spontaneous neutrophil apoptosis, L-selectin engagement accelerating rates and β_2 -integrin engagement retarding them.¹⁴ The present study sought to determine whether cell activation or adhesion molecule engagement could alter apoptotic signals delivered through the death receptor Fas.

We confirmed that quiescent neutrophils that die in vitro by spontaneous apoptosis are induced to die more rapidly when Fas is engaged with an agonist monoclonal antibody.³⁶ It has recently been shown that neutrophils express Fas ligand constitutively, and further, that they can release soluble Fas ligand in tissue culture.³⁷ The autocrine interaction of neutrophil Fas with soluble or cell-associated Fas ligand provides an attractive model to explain the rapid process of spontaneous apoptosis that occurs in normal neutrophils.¹⁰ Our data show that this process is susceptible to modulation during the process of neutrophil activation.

The development of a state of refractoriness to Fas-induced death signals was a transient, reversible phenomenon. It could not be explained by alterations in cell surface expression of Fas, since expression was similar in both activated and nonactivated cells. Neutrophil activation by LPS, endothelial transmigration, or CD11b cross-linking led to a rapid increase in levels of reactive oxygen intermediates and a later increase in GSH levels. Clement et al²⁷ showed that increased superoxide levels inhibit the Fas-mediated death of melanoma cells, indicating that alterations in the redox potential of the cell influence the ability of Fas to induce apoptosis.

We have previously reported that the depletion of intracellular thiols by diethylmaleate results in increased rates of



Fig 3. Temporal course of responsiveness to anti-Fas IgM ligation. Neutrophils ($1 \times 10^6/mL$) were incubated with LPS ($1 \mu g/mL$) (A) or cross-linked with antibodies to CD11b ($8 \mu g/mL$) (B) for 0, 1, or 4 hours, washed, and then incubated with anti-Fas IgM (100 ng/mL for 1 hour). Apoptosis was assessed by flow cytometry at 18 hours after incubation with LPS or CD11b ligation. **P* < .05 versus control, § *P* < .05 versus Fas (100 ng/mL). Results are the mean ± SD for n = 5 experiments with four separate donors of neutrophils.

spontaneous neutrophil apoptosis, a phenomenon that can be reversed by the anti-oxidants GSH and NAC,²⁶ although these exert no intrinsic effect on rates of spontaneous apoptosis. In the present study we found that these same antioxidants reduced the Fas-induced apoptosis of resting neutrophils. This finding is in keeping with the observation that mutant T cells expressing increased levels of intracellular glutathione as a result of augmented activity of γ -glutamylcysteine synthetase acquire resistance to Fas-mediated death signals,³⁸ and suggests a role for intracellular thiols in cellular resistance to apoptotic stimuli. Consistent with the hypothesis that augmented levels of GSH are necessary for the inhibition of Fas-mediated apoptosis in activated neutro-



Fig 4. Neutrophil intracellular glutathione levels. Neutrophils (10 × 10⁶/5 mL) were incubated with LPS (1 µg/mL) (\Box) or cross-linked with antibodies to CD11b (8 µg/mL) (**I**). **P* < .05 versus control (\bigcirc). Results are the mean ± SD for n = 3 experiments with three separate donors of neutrophils.

phils, we found that prevention of GSH synthesis in response to LPS by BSO permitted normal expression of Fas-induced signaling. Taken together, these observations show that augmentation of intracellular GSH is both necessary and sufficient to block Fas-mediated death signals in activated human neutrophils. Conversely, it can be hypothesized that depletion of GSH following neutrophil activation may again render the cell susceptible to Fas-induced apoptosis, and favor the resolution of an inflammatory response once the inciting stimulus has been removed. In keeping with the hypothesis we have previously shown that neutrophil ingestion of *E coli* results in the induction of apoptosis through an oxygen dependent mechanism associated with a reduction in intracellular thiol levels induced by the oxidant stress, since the effect is inhibited by the antioxidants GSH and NAC.¹⁵

Reconciliation of the divergent imperatives of cell growth and activation on the one hand, and programmed cell death and quiescence on the other is critical to the maintenance of normal immunologic homeostasis. Mice of the MRL/lpr strain, lacking functional Fas, develop an autoimmune disor-

 Table 2. Neutrophil Respiratory Burst Activity and Intracellular

 GSH After Fas Incubation

	Anti-Fas IgM (100 ng/mL)		
Control	30 Min	60 Min	120 Min
15.0 . 11.0			45.0.07
15.2 ± 11.0	14.9 ± 4.9	12.1 ± 5.0	15.6 ± 8.7
$\begin{array}{c} 18.3\pm8.2\\ 8.3\pm0.5\end{array}$	24.1 ± 10.6	$\begin{array}{c} 20.3\pm9.1\\ 9.1\pm1.2 \end{array}$	19.0 ± 10.0 —
	Control 15.2 ± 11.0 18.3 ± 8.2 8.3 ± 0.5	$\begin{tabular}{ c c c c c } \hline Anti-F & \\ \hline Control & 30 Min \\ \hline 15.2 \pm 11.0 & 14.9 \pm 4.9 \\ \hline 18.3 \pm 8.2 & 24.1 \pm 10.6 \\ \hline 8.3 \pm 0.5 & - \\ \hline \end{tabular}$	$\begin{tabular}{ c c c c c c } \hline Anti-Fas IgM (100 r \\ \hline Control & 30 Min & 60 Min \\ \hline 15.2 \pm 11.0 & 14.9 \pm 4.9 & 12.1 \pm 5.0 \\ \hline 18.3 \pm 8.2 & 24.1 \pm 10.6 & 20.3 \pm 9.1 \\ \hline 8.3 \pm 0.5 & - & 9.1 \pm 1.2 \\ \hline \end{tabular}$

Neutrophils were incubated with anti-Fas IgM (CH-11, 100 ng/mL) and then assessed for reactive oxygen intermediates after 0, 30, 60, and 120 minutes by flow cytometry using dihydrorhodamine 123 or neutrophil induced chemiluminescence. Intracellular GSH was measured at 60 minutes. Results represent the mean \pm SD of n = 3 experiments.

der characterized by nephritis and arthritis,³⁹ and are particularly sensitive to septic shock resulting from *Staphylococcal enterotoxin B*.⁴⁰ In contrast, administration of an agonistic antibody to Fas induces lethal hepatitis and neutropenia in mice.⁴¹ An optimal response to an acute infectious challenge necessitates the sequential expression of cell activation and cell quiescence. The observations reported here suggest that the cell surface molecules whose upregulation accompanies neutrophil activation serve a costimulatory function, and block the apoptotic response to death signals delivered by Fas. Because this costimulatory role is dependent on augmentation of intracellular thiols, the removal of an inflammatory stimulus, may permit the return of normal apoptosis in response to Fas, and so permit the normal resolution of inflammation.



Fig 5. Effects of the antioxidants GSH and NAC on neutrophil GSH levels (A) and Fas induced apoptosis (B). (A) Neutrophils (10×10^6 /mL) were incubated for 1 hour with GSH (25 mmol/L) or NAC (25 mmol/L), washed extensively, then assayed for intracellular GSH levels as described in Materials and Methods. (B) Neutrophils (1×10^6 /mL) were incubated for 1 hour with GSH (25 mmol/L) or NAC (25 mmol/L), then incubated with anti-Fas lgM (100 ng/mL) for 1 hour before washing. Apoptosis was assessed by flow cytometry 18 hours after the addition of anti-Fas IgM. *P < .05 versus control; \$P < .05 versus control; \$P < .05 versus control PMN and medium alone. Results are the mean \pm SD for n = 4 experiments with four separate donors of neutrophils.



Fig 6. Effects of BSO on (A) LPS induced increases in GSH and (B) LPS-induced blockade of anti-Fas IgM induced neutrophil apoptosis. Neutrophils (1 × 10⁶/mL) were preincubated with or without BSO (50 mmol/L) and then incubated with LPS (1 µg/mL) or LPS (1 µg/mL) + GSH (25 mmol/L), all groups were incubated after 1 hour with or without Fas (100 ng/mL). Intracellular GSH and apoptosis were assessed at 1 hour and 18 hours, respectively. **P* < .05 versus control. Results are the mean ± SD for n = 3 experiments with three separate donors of neutrophils.

REFERENCES

1. Nagata S, Golstein P: The Fas death factor. Science 267:1449, 1995

2. Suda T, Nagata S: Purification and characterization of the Fasligand that induces apoptosis. J Exp Med 179:873, 1994

3. Chinnaiyan AM, O'Rourke K, Tewari M, Dixit VM: FADD, a novel death domain-containing protein, interacts with the death domain of Fas and initiates apoptosis. Cell 81:505, 1995

4. Boldin MP, Goncharov TM, Goltsev YV, Wallach D: Involvement of MACH, a novel MOR1/FADD-interacting protease, in Fas/ APO-1- and TNF receptor-induced cell death. Cell 85:803, 1996

5. Muzio M, Chinnaiyan AM, Kischkel FC, O'Rourke K, Shevchenko A, Ni J, Scaffidi C, Bretz JD, Zhang M, Gentz R, Mann M, Krammer PH, Peter ME, Dixit VM: FLICE, a novel FADDhomologous ICE/CED-3-like protease, is recruited to the CD95 (Fas/ APO-1) death-inducing signaling complex. Cell 85:817, 1996

6. Schlegel J, Peters I, Orrenius S, Miller DK, Thornberry NA, Yamin T, Nicholson DW: CPP32/apopain is a key interleukin 1 beta converting enzyme-like protease involved in Fas-mediated apoptosis. J Biol Chem 271:1841, 1996

7. Griffith TS, Brunner T, Fletcher SM, Green DR, Ferguson TA:

Fas ligand-induced apoptosis as a mechanism of immune privilege. Science 270:1189, 1995

8. Kägi D, Vignaux F, Ledermann B, Bürki K, Depraetere V, Nagata S, Hengartner H, Golstein P: Fas and perforin pathways as major mechanisms of T cell-mediated cytotoxicity. Science 265:528, 1994

9. Haslett C: Resolution of acute inflammation and the role of apoptosis in the tissue fate of granulocytes. Clin Sci 83:639, 1992

10. Liles WC, Klebanoff SJ: Regulation of apoptosis in neutrophils - Fas Track to Death? J Immunol 155:3289, 1995

11. Lee A, Whyte MKB, Haslett C: Inhibition of apoptosis and prolongation of neutrophil functional longevity by inflammatory mediators. J Leuk Biol 54:283, 1993

12. Yamamoto C, Yoshida S, Taniguchi H, Qin MH, Miyamoto H, Mizuguchi Y: Lipopolysaccharide and granulocyte colony stimulating factor delay neutrophil apoptosis and ingestion by guinea pig macrophages. Infect Immun 61:1972, 1993

13. Armant M, Delespesse G, Sarfati M: IL-2 and IL-7 but not IL-12 protect natural killer cells from death by apoptosis and up-regulate bcl-2 expression. Immunology 85:331, 1995

14. Watson RWG, Rotstein OD, Nathens AB, Marshall JC: Neutrophil apoptosis is modulated by endothelial transmigration and adhesion molecule engagement. J Immunol 158:945, 1997

15. Watson RWG, Redmond HP, Wang JH, Condron C, Bouchier-Hays D: Neutrophils undergo apoptosis following ingestion of *Escherichia coli*. J Immunol 156:3986, 1996

16. Mangan EF, Robertson B, Wahl SM: IL-4 enhances programmed cell death (apoptosis) in stimulated human monocytes. J Immunol 148:1812, 1992

17. Bursch W, Oberhammer F, Jirtle RL, Askari M, Sedivy R, Grasl-Kraupp B, Purchio AF, Schulte-Hermann R: Transforming growth factor-1 as a signal for induction of cell death by apoptosis. Br J Cancer 67:531, 1993

18. Woods KM, Chapes SK: Three distinct cell phenotypes of induced-TNF cytotoxicity and their relationship to apoptosis. J Leuk Biol 53:37, 1993

19. Crispe IN: Fatal interactions: Fas-induced apoptosis of mature T cells. Immunity 1:347, 1994.

20. Druilhe A, Cai Z, Haile S, Chouaib S, Pretolani M: Fasmediated apoptosis in cultured human eosinophils. Blood 87:2822, 1996

21. Um H, Orenstein JM, Wahl SM: Fas mediates apoptosis in human monocytes by a reactive oxygen intermediate dependent pathway. J Immunol 156:3469, 1996

22. Nicoletti I, Migliorati G, Pagliacci MC, Grignani F, Riccardi C: A rapid and simple method for measuring thymocyte apoptosis by propidium iodide staining and flow cytometry. J Immunol Methods 139:271, 1991

23. Smith JA, Weidemann MJ: Further characterization of neutrophil oxidative burst by flow cytometry. J Immunol Methods 162:261, 1993

24. Allen RC: Phagocytic leukocyte oxygenation activities and

chemiluminescence: A kinetic approach to analysis. Methods Enzymol 133:449, 1986

25. Jocelyn PC: Spectrophotometric assay of thiols. Methods Enzymol 143:44, 1987

26. Watson RWG, Rotstein OD, Nathens AB, Dackiw APB, Marshall JC: Thiol-mediated redox regulation of neutrophil apoptosis. Surgery 120:150, 1996

27. Clement M, Stamenkovic I: Superoxide anion is a natural inhibitor of Fas-mediated cell death. EMBO J 15:216, 1996

28. Droge W, Eck HP, Mihms: HIV-induced cysteine deficiency and T-cell dysfunction-a rational for treatment for N-acetylcysteine. Immunol Today 13:472, 1992

29. Griffith OW: Mechanisms of action, metabolism, and toxicity of buthionine sulfoximine and its higher homologs, potent inhibitors of glutathione biosynthesis. J Biol Chem 257:13704, 1982

30. Hsu H, Xiong J, Goeddel DV: The TNF receptor 1-associated protein TRADD signals cell death and NF-kappaB activation. Cell 81:495, 1995

31. Rensing-Ehl A, Hess S, Ziegler-Heitbrock HW, Riethmuller G, Engelmann H: Fas/APO-1 activates nuclear factor kappa B and enhances interleukin-6 production. J Inflamm 45:161, 1995

32. Schulze-Osthoff K, Krammer PH, Droge W: Divergent signalling via APO-1/Fas and the TNF receptor, two homologous molecules involved in physiological cell death. EMBO J 13:4587, 1994.

33. Carlos TM, Harlan JM: Leukocyte-endothelial adhesion molecules. Blood 84:2068, 1995

34. Rosales C, Juliano RL: Signal transduction by cell adhesion receptors in leukocytes. J Leuk Biol 57:189, 1995

35. Boudreau N, Sympson CJ, Werb Z, Bissell MJ: Suppression of ICE and apoptosis in mammary epithelial cells by extracellular matrix. Science 267:891, 1995

36. Iwai K, Miyawaki T, Takizawa T, Konno A, Ohta K, Yachie A, Seki H, Taniguchi N: Differential expression of bcl-2 and susceptibility to anti-Fas-mediated cell death in peripheral blood lymphocytes, monocytes, and neutrophils. Blood 84:1201, 1994

37. Liles WC, Kiener PA, Ledbetter JA, Aruffo A, Klebanoff SJ: Differential expression of Fas (CD95) and Fas ligand on normal human phagocytes: Implications for the regulation of apoptosis in neutrophils. J Exp Med 184:429, 1996

38. Kohno T, Yamada Y, Hata T, Mori H, Yamamura M, Tomonaga M, Urata Y, Goto S, Kondo T: Relation of oxidative stress and glutathione synthesis to CD95 (Fas/APO-1)-mediated apoptosis of adult T cell leukemia cells. J Immunol 154:4722, 1995

39. Nagata S, Suda T: Fas and Fas ligand: lpr and gld mutations. Immunol Today 16:39, 1995

40. Mountz JD, Baker TJ, Borcherding DR, Bluethmann H, Zhou T, Edwards CK: Increased susceptibility of fas mutant MRL-lpr/ lpr mice to Staphylococcal enterotoxin B-induced septic shock. J Immunol 155:4829, 1995

41. Ogasawara J, Watanabe-Fukunaga R, Adachi M, Matsuzawa A, Kasugai T, Kitamura Y, Itoh N, Suda T, Nagata S: Lethal effect of the anti-Fas antibody in mice. Nature 364:806, 1993