

Molecular Remission in PML/RAR α -Positive Acute Promyelocytic Leukemia by Combined All-*trans* Retinoic Acid and Idarubicin (AIDA) Therapy

By Franco Mandelli, Daniela Diverio, Giuseppe Avvisati, Anna Luciano, Tiziano Barbui, Carlo Bernasconi, Giorgio Broccia, Raffaella Cerri, Michele Falda, Giuseppe Fioritoni, Franco Leoni, Vincenzo Liso, Maria Concetta Petti, Francesco Rodeghiero, Giuseppe Saglio, Maria Luce Vegna, Giuseppe Visani, Ulrich Jehn, Roel Willemze, Petra Muus, Pier Giuseppe Pelicci, Andrea Biondi, and Francesco Lo Coco for the Gruppo Italiano Malattie Ematologiche Maligne dell' Adulto and Associazione Italiana di Ematologia ed Oncologia Pediatrica Cooperative Groups

Two hundred fifty-three patients with newly diagnosed acute promyelocytic leukemia (APL) were eligible to enter the multicentric GIMEMA-AIEOP "AIDA" trial during the period July 1993 to February 1996. As a mandatory prerequisite for eligibility, all patients had genetic evidence of the specific t(15;17) lesion in their leukemic cells confirmed by karyotyping or by reverse transcription-polymerase chain reaction (RT-PCR) of the PML/RAR α fusion gene (the latter available in 247 cases). Median age was 37.8 years (range, 2.2 to 73.9). Induction treatment consisted of oral all-*trans* retinoic acid (ATRA), 45 mg/m²/d until complete remission (CR), given with intravenous idarubicin, 12 mg/m²/d on days 2, 4, 6, and 8. Three polychemotherapy cycles were given as consolidation. Hematologic and molecular response by RT-PCR was assessed after induction and after consolidation. At the time of analysis, 240 of the 253 eligible patients were evaluable

for induction. Of these, 11 (5%) died of early complications and 229 (95%) achieved hematologic remission. No cases of resistant leukemia were observed. Of 139 cases studied by RT-PCR after induction, 84 (60.5%) were PCR-negative and 55 (39.5%) PCR-positive. One hundred sixty-two patients were evaluable by RT-PCR at the end of consolidation. Of these, 159 (98%) tested PCR-negative and 3 (2%), PCR-positive. After a median follow up of 12 months (range, 0 to 33), the estimated actuarial event-free survival for the whole series of 253 eligible patients was 83% \pm 2.6% and 79% \pm 3.2% at 1 and 2 years, respectively. This study indicates that the AIDA protocol is a well-tolerated regimen that induces molecular remission in almost all patients with PML/RAR α -positive APL. Preliminary survival data suggest that a remarkable cure rate can be obtained with this treatment.

© 1997 by The American Society of Hematology.

ACUTE PROMYELOCYTIC leukemia (APL) is a particular type of acute myeloid leukemia (AML) with characteristic biological and clinical features, which include the presence of a specific t(15;17) chromosome translocation in

the leukemic blasts, the frequent association at diagnosis of a severe hemorrhagic diathesis, and in vitro and in vivo sensitivity to the differentiating agent all-*trans* retinoic acid (ATRA).¹⁻⁴

The t(15;17) involves the retinoic acid receptor α (RAR α) gene on chromosome 17 and the PML (for promyelocytic) gene on chromosome 15 and generates a chimeric gene, which is transcribed into the PML/RAR α fusion transcript. The resulting PML/RAR α chimeric protein is crucial to the pathogenesis of the disease.^{3,4}

The PML/RAR α hybrid mRNA is readily detectable by the reverse transcription-polymerase chain reaction (RT-PCR) assay.^{5,6} As reported by several groups, including ours, RT-PCR studies of the PML/RAR α have proved extremely useful in the clinical management of the disease.⁵⁻¹¹ Detection of PML/RAR α at diagnosis predicts response to ATRA, whereas cases with morphological features of APL, which are PML/RAR α negative, are refractory to differentiative treatment.⁶ Moreover, when they are applied to evaluating residual disease during remission, RT-PCR tests can identify patients at risk of relapse and in need of additional therapy.⁷⁻¹¹

Anthracycline-based chemotherapy (CHT) and ATRA are the mainstay of APL treatment.²⁻⁴ Anthracyclines alone or in association with cytosine arabinoside induce complete remission (CR) in a high percentage of cases (75% to 80%), but this approach is associated with a high mortality rate (up to 20%) during the early phases of treatment.^{2-4,12-17} Differentiative treatment with ATRA results in greater than 90% CRs, but patients remain invariably PCR positive after induction, and all relapse if no consolidation CHT is added.^{2-4,18-21} Furthermore, ATRA treatment has been associated with the occurrence, in a sizable fraction of patients, of life-threatening complications. These include a severe respiratory distress due to pulmonary infiltrates (ATRA syndrome) usually (but

From Ematologia, Dipartimento di Biotecnologie Cellulari ed Ematologia, University "La Sapienza," Rome; Clinica Pediatrica, Ospedale S. Gerardo, University of Milano-Monza; Divisione di Ematologia, Ospedali Riuniti di Bergamo; Divisione di Ematologia, Policlinico S. Matteo, University of Pavia; Divisione di Ematologia, Ospedale A. Businco, Cagliari; Divisione di Ematologia, Ospedale S. Martino, Genova; Divisione di Ematologia, Ospedale S. Giovanni Battista, Torino; Divisione di Ematologia, Ospedale di Pescara; Divisione di Ematologia, Policlinico Careggi, Firenze; Servizio di Ematologia, University of Bari; Divisione di Ematologia, Ospedale S. Bortolo, Vicenza; Patologia Medica, Ospedale S. Luigi Gonzaga, Orbassano, Torino; Istituto di Ematologia L. e A. Seragnoli, University of Bologna; Clinica Medica I, University of Perugia, Italy; Medizinische Klinik III, University of Munich, Munich, Germany; Department of Hematology, University Hospital, Nijmegen; Department of Hematology, University of Leiden, The Netherlands.

Submitted November 21, 1996; accepted April 1, 1997.

Supported by grants from CNR (Consiglio Nazionale delle Ricerche) contratti ACRO no. 95.00414.PF39, 95.02140, and 95.00440.PF39, ROMAIL (Associazione Italiana contro le Leucemie, Sezione di Roma), Fondazione Tettamanti, Monza, and AIRC (Associazione Italiana per la Ricerca sul Cancro), Milano.

Presented in part at the 37th Annual Meeting of the American Society of Hematology, Seattle, WA, December 1-5, 1995.

Address reprint requests to Francesco Lo Coco, MD, Hematology, Department of Human Biopathology, University La Sapienza, Via Benevento 6, 00161 Roma, Italy.

The publication costs of this article were defrayed in part by page charge payment. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. section 1734 solely to indicate this fact.

© 1997 by The American Society of Hematology.

0006-4971/97/9003-0014\$3.00/0

not uniformly) correlated to a rapid increase of white blood cell counts^{21,22} and pseudotumor cerebri, which is more frequently observed in younger patients.²³

Preliminary clinical studies have been performed with various combinations of ATRA and CHT in an attempt to obtain more durable remissions and reduced ATRA-related toxicity. The combinations, as demonstrated in a randomized trial, seem to improve disease-free survival over that achieved with chemotherapy alone.²⁴ Since 1983, the Italian cooperative group, Gruppo Italiano Malattie Ematologiche Maligne dell' Adulto (GIMEMA), adopted disease-tailored protocols for APL, showing that anthracyclines alone are as effective in inducing remission as polychemotherapy schemes.²⁵ In 1993, aiming at improving the efficacy of APL-targeted treatment, a new protocol combining ATRA and Idarubicin (AIDA protocol) was designed by the GIMEMA, first as a pilot,²⁶ then in association with the Associazione Italiana di Ematologia ed Oncologia Pediatrica (AIEOP), as a larger multicentric study. Prospective RT-PCR analyses were performed on all patients at presentation to assess study eligibility and after treatment to better evaluate response and minimal residual disease.

MATERIALS AND METHODS

Patients. Two hundred seventy-four consecutive patients with newly diagnosed APL were registered during the period July 1993 to February 1996. All cases were diagnosed as AML French-American-British (FAB) M3 or M3 variant (M3v) by conventional morphocytological criteria²⁷ in 71 participating institutions (see Appendix). The majority of these centers belong to the Italian cooperative groups GIMEMA and AIEOP, while five belong to other European countries. Eligibility criteria to be enrolled in the AIDA protocol included: age between 1 and 75 years, karyotypic and/or molecular evidence of the t(15;17) in leukemic blasts at presentation, serum creatinine <2.5 mg/dL, serum alkaline phosphatase, bilirubin, and aspartate aminotransferase (AST) <3 times the upper normal limit, no cardiac contraindications to anthracycline chemotherapy, negative pregnancy test, and informed consent. Cases presenting with hyperleukocytosis were also included in the study.

AIDA protocol. Induction therapy consisted of 45 mg/m²/d ATRA administered orally until the achievement of CR or for a maximum of 90 days, and four 12 mg/m² doses of Idarubicin given intravenously (IV) on days 2, 4, 6, and 8. In patients less than 20 years old, ATRA doses were adjusted to 25 mg/m². After the achievement of hematologic remission, patients received three consolidation polychemotherapy courses as reported.²⁶ Each consolidation course was administered at recovery from the previous course, when polymorphonuclear cells and platelet counts were >1,500/ μ L and >100,000/ μ L, respectively.

At the end of consolidation, residual disease was estimated by RT-PCR before deciding further therapy. PCR-negative patients were randomized into four arms including chemotherapy alone with 6-mercaptopurine and methotrexate (arm 1), ATRA alone (arm 2), alternating chemotherapy and ATRA (arm 3), or no further therapy (arm 4). Patients who were PCR-positive underwent, if eligible, allogeneic bone marrow transplantation in first CR.

ATRA syndrome was defined as "definitely present" or "overt" in the presence of the following five signs and symptoms: fever, dyspnea, pleural and/or pericardial effusion, pulmonary infiltrates on chest radiograph, and unexplained weight gain greater than 5 kg, according to the original report by Frankel et al.²² We defined as "indeterminate" ATRA syndrome a combination of two to four

signs and symptoms of the above mentioned \pm lower extremities edema and hypotension. In all other patients, the ATRA syndrome was defined as "definitely absent."²⁸ Rules for supportive therapy, treatment of the ATRA syndrome, and prevention and control of the coagulopathy have been reported in detail elsewhere.²⁶

Collection and preparation of samples for molecular studies. Samples for RT-PCR analyses were requested at diagnosis, at the time of CR after induction, and at the end of consolidation. With the exception of some cases with hyperleukocytosis in which peripheral blood (PB) samples were studied at presentation, all other diagnostic and all remission samples were obtained from the bone marrow (BM). Mononuclear cells, obtained by centrifuging BM or PB specimens on a Ficoll-Hypaque density gradient, were washed twice in sterile phosphate-buffered saline (PBS), suspended in a 4-mol/L guanidium thiocyanate (GTC) solution and stored at -20°C . These procedures were performed in each GIMEMA and AIEOP center using RNAase-free disposable materials. The GTC solution was previously prepared by the referral molecular biology laboratories and distributed to all peripheral centers. Samples cryopreserved in GTC were then sent in dry ice to two referral molecular biology laboratories for RT-PCR studies (Hematology, University "La Sapienza" of Rome and Clinica Pediatrica, University of Milano, Monza).

RT-PCR of PML/RAR α . Total RNA was extracted by the method of Chomczynsky and Sacchi.²⁹ The integrity of RNAs was assessed in all diagnostic and remission samples by electrophoretic run through a formaldehyde minigel. The protocol and the primers used to amplify the PML/RAR α hybrid gene have been reported elsewhere.⁷ To assess the efficiency of the RT step and to further verify RNA integrity, a cDNA fragment containing RAR α exons 2 and 3 was coamplified in each analyzed sample (internal control). A positive control (amplification of RNA extracted from the promyelocytic cell line NB4) and a negative control (all reagents plus water with no RNA) was included in each experiment. To assess the sensitivity of our RT-PCR method, total RNA isolated from a diagnostic sample with 100% blastic infiltration was serially diluted by mixing it with the t(15;17) negative myeloid cell line GF-D8 RNA.⁵ Our assay allowed us to detect the PML/RAR α transcript in the presence of 0.1 ng total RNA, that is a final dilution of 10^{-4} .

Evaluation of response. CR was defined as the reconstitution of normal BM cellularity with less than 5% leukemic promyelocytes, together with PB cell counts of polymorphonuclear leukocytes (PMN) > 1,500/ μ L and platelets >100,000/ μ L. Molecular remission was defined as the disappearance, on ethidium bromide-stained electrophoresis gel, of the specific PML/RAR α amplification band identified at diagnosis, in the presence of RNA integrity as evaluated by minigel visualization and successful amplification of the internal control. Overall survival (OS) and event-free survival (EFS) duration was calculated by the method of Kaplan and Meier. The median observation was 12 months (range, 0 to 33).

RESULTS

Patient characteristics at diagnosis. Of the 274 patients registered, 21 (7.7%) were considered ineligible. Exclusion criteria for these 21 patients are reported in Table 1. These 21 patients were withdrawn from the study and treated differently.

The clinical and biological characteristics of the 253 eligible patients are shown in Table 2. Median age was 37.8 years (range, 2.2 to 73.9). Thirty-two cases (12.6%) were classified as microgranular or variant (M3v) APL on morphologic examination. The median white blood cell (WBC) count was $2.5 \times 10^9/\text{L}$ (range, 0.2 to 140.0). Bleeding symptoms were present at diagnosis in 67% of patients; of these,

Table 1. Reasons, Other Than Age, for the Exclusion From Entering the AIDA Protocol in 21 Patients Registered as AML-FAB M3

Exclusion Criteria	No. of Cases
Karyotype NA*/PML-RAR α -negative	7
Karyotype NA/PML-RAR α NE†	3
Karyotype NE‡/PML-RAR α -negative	2
Absence of t(15;17)/PML-RAR α -negative	2
Absence of t(15;17)/PML-RAR α NE	2
Karyotype NA/specimen for RT-PCR not sent	1
t(11;17)/PML-RAR α -negative	1
HIV-positive	1
Poor performance status/active serious infection	1
Other concomitant malignancy	1

Abbreviations: NA, not available; HIV, human immunodeficiency virus.

* Karyotype, not available.

† PML-RAR α NE, poor quality RNA.

‡ Karyotype NE, no mitosis or poor quality metaphases.

94% also had laboratory evidence of coagulopathy defined as fibrinogen <150 mg/dL and fibrinogen degradation products (FDP) >40 μ g/mL or D-dimers (XDP) >400 μ g/mL.

The type of PML/RAR α transcript as detected by RT-PCR (available in 247 cases) was bcr1-2 (or long type) in 152 (61.5%) patients, and bcr3 (or short type) in 95 (38.5%) patients. In the remaining six eligible patients, the t(15;17) abnormality was detected by karyotypic examination only and no material was available for molecular studies at diagnosis. The type of PML breakpoint was not significantly correlated with age, sex, WBC count, or presence of coagulopathy at presentation. However, the incidence of bcr3 was significantly higher in patients with a diagnosis of M3v. In fact, a bcr3 breakpoint was observed in 20 of 32 (62%) M3v cases and in 75 of 215 (35%) hypergranular APL cases ($P = .005$).

Hematological and molecular response to induction therapy. Of the 253 eligible patients, three died of intracranial hemorrhage before the start treatment, one was lost to follow up, two were crossed to chemotherapy at days +16 and +20 due to ATRA toxicity and were considered as protocol violations, and it was too early for response to induction to be evaluated in seven patients.

Of the remaining 240 evaluable patients, 229 (95%) com-

Table 3. Response to Induction Therapy

Eligible	253
Pretherapy deaths	3
Lost to follow-up	1
Violation	2
Too early	7
Evaluable	240
CR	229 (95%)
Induction deaths	11 (5%)
Resistant disease	0

pleted induction therapy and achieved hematologic CR, and 11 patients (5%) died during induction. No cases of resistant disease were documented. Of the 11 early deaths, eight were due to intracranial hemorrhage and occurred at days +1, +1, +3, +7, +8, +8, +13, and +36, respectively; one to ATRA syndrome (day +17), one to cerebral thrombosis (day +13), and one to myocardial infarction (day +20). The median time of ATRA therapy was 31 days (range, 3 to 90). The median time for polymorphonuclear cells to recover to $>1.0 \times 10^9/L$ and platelets to $>100 \times 10^9/L$ was 27 days (range, 4 to 48) and 28 days (range, 9 to 49), respectively. Table 3 shows clinical results after induction.

RT-PCR analysis for the evaluation of the PML/RAR α hybrid at the end of induction and before consolidation therapy was performed in 139 cases. As to the other patients who achieved CR, four cases were not evaluable due to poor quality RNA and no samples were sent in for the remaining cases. Fifty-five (39.5%) of the 139 patients analyzed in CR after induction were PCR-positive and 84 (60.5%) PCR-negative. There was no significant correlation between the type of PML breakpoint and molecular response to induction therapy.

Molecular response after consolidation and follow-up analysis. Of the 229 patients who entered hematologic CR, 5 died of complications during consolidation, 1 was lost to follow up, 1 relapsed after the second cycle, and 60 have yet to complete consolidation therapy. Of the remaining 162 tested by RT-PCR after consolidation, 159 (98%) were PCR-negative and three (2%) PCR-positive. The three PCR-positive patients had all been positive at the end of induction. The remaining 52 patients who had been PCR-positive at the end of induction converted to PCR negativity after consolidation.

PCR assays between cycles of consolidation were performed in a limited number of cases in this study. Of 23 patients PCR-positive postinduction, 16 (69.5%) converted to PCR-negative and 7 (30.5%) remained PCR-positive after the first consolidation. Of these seven PCR-positive, 2 were still PCR-positive and 2 converted to PCR-negative following the second consolidation (no analyses were available at this time in the other three cases. However, all three converted to PCR-negative after the third cycle). In summary, of 20 postinduction PCR-positive cases tested either after one or two consolidation cycles, 18 (90%) achieved molecular remission before receiving the third chemotherapy consolidation course.

Sixteen (9.8%) of the 162 patients evaluable after consolidation did not complete the three courses because of therapy-related toxicity. Clinical and molecular data available in this

Table 2. Clinical and Biological Features of Eligible Patients at Diagnosis

Sex	
M	138 (54.5%)
F	115 (45.5%)
Age median	37.8
(range)	2.2-73.9
FAB M3	221 (86.4%)
M3v	32 (12.6%)
WBC $\times 10^9/L$ median	2.5
(range)	0.2-140.0
Type of PML/RAR α transcript	
Long (bcr 1-2)	152 (61.5%)
Short (bcr 3)	95 (38.5%)
NA	6*

Abbreviation: NA, not available.

* All had the t(15;17).

Table 4. Clinical and Molecular Outcome of 16 Patients Withdrawn From Study Due To Therapy-Related Toxicity

UPN	Age	No. of Consolid. Cycles Received	Type of Toxicity	Further Therapy Postconsolid.	Outcome	Follow-up RT-PCR Analyses
28024	65	0*	Cardiac	ATRA/MTX/6MP	CCR 14+†	Negative (2 tests)
28030	60	0	Infectious	ATRA/MTX/6MP	CCR 10+	Negative (2 tests)
04003	63	1	Infectious	IFN	CCR 33+	Negative (3 tests)
10004	62	1	Infectious	None	Relapsed at 6 mo/dead	Negative (2 tests)
35003	55	2	Neurologic	None	CCR 31+	Negative (5 tests)
04002	33	2	Infectious	ATRA/MTX/6MP	CCR 32+	Negative (3 tests)
36001	59	2	Infectious	IFN	CCR 32+	Negative (4 tests)
28015	63	2	Prolonged BM hypoplasia	MTX/6MP	CCR 33+	Negative (6 tests)
D6004	9	2	Severe mucositis	None	CCR 29+	Negative (6 tests)
43002	66	2	Prolonged BM hypoplasia	None	CCR 29+	Negative (4 tests)
37005	62	2	Prolonged BM hypoplasia	None	CCR 30+	Negative (5 tests)
55001	43	2	Infectious	None	Relapsed at 7 mo/dead	Positive after ind.‡
07006	63	2	Infectious	None	Dead at 2 mo§	Negative (1 test)
43005	66	2	Neurologic	None	CCR 16+	Negative (3 tests)
28021	23	2	Infectious	MTX/6MP	CCR 17+	Negative 5 tests, 1 +ve
08005	31	2	Infectious	None	CCR 10+	Negative (3 tests)

Abbreviations: MTX, methotrexate; 6MP, 6-mercaptopurine; IFN, alpha 2b interferon.

* Patients 1 and 2 received induction therapy only.

† Months in continuous complete remission.

‡ Tested after induction therapy only.

§ Died of septic shock while in CR.

|| Tested PCR-positive at the last molecular follow-up analysis.

series are shown in Table 4. Median age was 61 years. In the majority of cases, treatment withdrawal was due to infection and/or prolonged BM hypoplasia. These patients were not randomized for maintenance; 7 patients received heterogeneous therapy as maintenance, whereas 9 had no further treatment. The median remission duration of patients in this group is 23 months (range, 2 to 33).

The results of the RT-PCR studies performed at diagnosis, after induction, and after consolidation in the whole series are shown in Table 5. The three patients who tested PCR-positive at the end of consolidation underwent allogeneic BMT from an HLA-identical sibling. Of the 159 who tested PCR-negative, 14 went off the study for protocol violation (9 cases), treatment-related toxicity (4 cases), and refusal to continue the therapeutic program (1 case). The remaining 145 patients were randomized for maintenance and are currently under evaluation for the effect of the different treatment approaches on relapse-free survival.

As of March 1996, 17 patients relapsed at 5 to 20 months from the achievement of hematologic remission. Among these, two had not completed all cycles of consolidation chemotherapy. Fifteen had typical hypergranular morphology at presentation, whereas two were classified as M3v. The median WBC count in these series of 17 patients was

$7.2 \times 10^9/L$ (0.9 to 54.9). The type of breakpoint within the PML gene, available in 16 cases, was bcr 1-2 in nine patients, and bcr 3 in seven patients.

The estimated 12 and 24 months OS for all of the 253 eligible patients was $90\% \pm 1\%$ and $87\% \pm 2\%$, whereas the EFS was calculated at $83\% \pm 2.6\%$ and $79\% \pm 3.2\%$, respectively (Fig 1). The comparison of EFS (calculated from the achievement of CR) between patients who were PCR-positive and PCR-negative at the end of induction showed no statistically significant difference.




Fig 1. Kaplan-Meier estimates of overall survival and event-free survival (EFS).

Table 5. PCR Monitoring of PML/RAR α

	Evaluable	PCR+(%)	PCR-(%)
At diagnosis	247	247 (100%)	
After induction	139	55 (39.5%)	84 (60.5%)
After consolidation	162	3 (2%)	159 (98%)

Table 6. Clinical Features of Patients With Overt or “Definitely Present” ATRA Syndrome

UPN	Age/Sex	FAB	WBC Count ($\times 10^9/L$)		Therapy	Outcome
			Initial	Peak (day of treatment)		
07006	63/F	M3	0.5	5.9 (6)	ATRA discontin./Dexamet.	CR
24009	46/M	M3v	54.2	101.0 (2)	ATRA discontin./Dexamet.	CR
24010	36/F	M3	7.2	18.2 (4)	ATRA discontin./Dexamet.	CR
35005	45/M	M3	11.6	25.8 (7)	ATRA discontin./Dexamet.	CR
55003	40/F	M3	0.6	1.6 (16)	ATRA discontin./Dexamet.	CR
24001	31/F	M3	1.5	5.1 (4)	ATRA discontin./Dexamet.	Died

Abbreviation: discontin./Dexamet., discontinuation/Dexamethasone.

ATRA syndrome and other therapy-related toxicity. Of 240 patients evaluable for induction therapy, 6 (2.5%) developed an overt or “definitely present” ATRA syndrome and one patient died of it. Clinical features including WBC initial count and peak, specific treatment for the syndrome, and outcome of these patients are reported in Table 6. In 11 additional cases, a combination of signs and symptoms was observed, which allowed us to classify them as having “indefinite” ATRA syndrome. Finally, ATRA syndrome was considered “definitely absent” in 223 of 240 patients (93%).

Other ATRA-related adverse reactions included mucosal and skin dryness (29% of cases), hypotension (7%), headache (13%), severe bone pain (6%), pseudotumor cerebri (2%), and hypercholesterolemia (6%). Scrotal ulcerations were observed in two patients (1%).

As to the other therapy-related toxicity, mainly due to anthracycline treatment, these were limited to stomatitis (6%), hemorrhages (15%), nausea and vomiting (5%), diarrhea (2%), cardiac (2.5%), hepatic (3%), and renal (0.5%) toxicities considering together only World Health Organization (WHO) grades 3 and 4.

Of the 5 deaths occurring after the end of induction, 3 were recorded during the first consolidation course and 2 during the second course. Three deaths were due to bacterial infection, one to hepatitis, and one to an ATRA syndrome developed in CR.

DISCUSSION

Of all acute leukemias, APL is at present the form to which molecular studies contribute the most clinically relevant information. Three lines of evidence support this assumption, ie, (1) the PML/RAR α genetic lesion is absolutely APL-specific and found in virtually 100% of cases²⁻⁴; (2) the disease responds specifically to a distinct treatment approach that is also effective against the life-threatening coagulopathy²⁻⁴; and (3) PML/RAR α detection in the leukemic blasts predicts response to ATRA.⁶ A similar combination of circumstances is not observed in any other human leukemia.

The mandatory prerequisite for entry into this study was molecular or cytogenetic evidence of the t(15;17) in leukemia cells at diagnosis. The characterization of the PML/RAR α junction type, available in 247 of 253 (98%) eligible patients, also provided an ideal specific marker for the sensitive assessment of response to therapy. Based on past experi-

ence gained as a referral laboratory for RT-PCR analysis in a multicentric study,²⁶ we put special emphasis on the logistic organization of the molecular studies. In particular, we recommended that rapid isolation and storage of the mononuclear (MNC) cell fraction of marrow aspirates in an RNAase inhibitor (guanidium isothiocyanate), be performed locally in each participating center before shipment. This procedure turned out to be extremely important for the extraction of good quality RNA in the vast majority of cases.

The results of induction therapy with ATRA and Idarubicin show that all PML/RAR α positive patients are responsive to this treatment. As a genetic diagnosis of APL was not required in any previously published study, it might seem inappropriate to compare our data with results obtained using other protocols, whether they included ATRA or not. However, it is conceivable that the lower CR rate reported in other studies could have been due, at least in part, to PML/RAR α negative patients being enrolled.^{12-25,28,30} Several new acute leukemia entities, which morphologically resemble typical APL but manifest distinct genetic and/or immunophenotypic features, have been described recently. They include cases with variant translocation that involve RAR α on 17q with partner chromosomes other than 15, for example t(11;17) and t(5;17) associated APL,^{31,32} as well as cases with a myeloid-natural killer (CD56⁺) immunophenotype.³³ Because these cases do not express the PML/RAR α fusion gene and fail to respond to ATRA, patients with no cytogenetic or molecular proof of the t(15;17) in their leukemia blasts at presentation were excluded from our series (Table 1).

The AIDA induction regimen appears to be well tolerated. In fact, the incidence and the severity of complications, including hemorrhagic deaths, ATRA syndrome, and other therapy-related toxicity were lower in our series than in other induction protocols irrespective of whether or not they included ATRA.^{12-25,28,30} The combination of a differentiating agent and a cytotoxic drug administered in the early phase of therapy very likely provided a counteraction against either the bleeding diathesis due to the coagulopathy or the occurrence of the severe pulmonary symptoms associated with the ATRA syndrome. Several investigators have shown that ATRA treatment significantly improves the APL-associated coagulopathy,³⁴⁻³⁶ while cytotoxic chemotherapy is known to prevent the onset of the ATRA syndrome, being often anticipated in patients considered at risk because of hyper-

leukocytosis at presentation.²¹ These two severe complications, which so far represented the major causes of induction death in APL, were recorded in a minority of cases in this study. Overall, 11 of 240 patients (5%) died during induction treatment.

Following the guidelines of the New York Group (Frankel et al²⁸ and previous personal communication by Dr R.P. Warrell Jr, April 1993), we recommended to all physicians of our Cooperative Group to promptly administer IV dexamethasone as prophylactic treatment at the earliest clinical sign of dyspnea. It is conceivable that, in conjunction with early chemotherapy administration, this policy contributed to prevent the onset of a full-blown and life-threatening ATRA syndrome in the vast majority of our patients.

Concerning consolidation, our clinical and molecular data suggest that the administration of less postremission treatment might be considered in the future, at least for elderly patients. In fact, as shown in Table 4, the median age of patients withdrawn from the study because of therapy-related toxicity was 61 years. Furthermore, the preliminary follow-up analysis in this series showed a short-term outcome comparable to that of patients receiving the three consolidation courses.

PCR results between consolidation cycles, although only available in a fraction of cases, indicate that the vast majority of patients achieve molecular remission after two cycles. A prolonged follow-up analysis is needed to establish the prognostic significance of these early achieved molecular remissions and to verify whether less postremission treatment could be effective in APL, regardless of age.

Molecular remission was achieved in 60% of our cases after induction and 98% after consolidation. We are not sure whether the PCR positivity detected at the end of induction reflects persistence of resistant blasts or delayed ATRA-induced maturation. We are also unable to assess whether the different kinetics of molecular response and leukemia-cell clearance have prognostic value. However, the attainment of a remission status corresponding to less than 10⁻⁴ PML/RAR α -positive cells (according to the sensitivity of our assay) in almost all patients is further testimony for the therapeutic efficacy of the AIDA regimen.

While PCR positivity during hematologic remission in APL is generally considered a strong predictor of clinical relapse, the achievement of a PCR-negative status is no guarantee of cure, as indicated by the 17 relapses occurring in patients who had previously tested PCR-negative. This probably reflects the limited sensitivity of the RT-PCR assay for PML/RAR α .⁴ At the time of the present analysis, due to the low number of adverse events, no predictive features could be found that were associated with a higher probability of relapse. In particular, no statistically significant association was found between treatment outcome and type of PML breakpoint, WBC count, or PCR status after induction. However, it is interesting to observe that patients who relapsed had higher median WBC counts at diagnosis. Future analysis of these parameters in a larger series and on a longer follow-up, might provide relevant information as to the possibility of identifying patients at higher risk of relapse.

Because of the uncertainty of the fate of patients who

achieve PCR negativity, it is too early to predict the long-term outcome in the present series. However, the actuarial 1- and 2-year EFS estimates (83% \pm 2.6% and 79% \pm 3.2%, respectively) are promising figures with respect to long-term outcome. In addition, APL patients who relapse are easily rescued into second remission and become long-term survivors more often than patients with other myeloid leukemias.³⁷

In conclusion, our results indicate that a specific treatment approach targeted to a molecular abnormality might dramatically change the natural history of APL, turning this once rapidly fatal disease into one of the most frequently curable acute leukemias. In addition, APL offers a model that should trigger interest to identify other genetico-clinical subsets of acute leukemia so that currently available therapy and future advancements can be more precisely directed at treating these diseases.

APPENDIX

The following clinical departments participated in the AIDA 0493 trial: Dipartimento di Biotecnologie Cellulari ed Ematologia, Università "La Sapienza," Roma, F. Mandelli, G. Avvisati, D. Diverio, F. Lo Coco, M.C. Petti, M.L. Vegna; Divisione di Ematologia, Ospedale S. Martino, Genova, E. Damasio, R. Cerri; Istituto di Ematologia L. e A. Seragnoli, Università, Bologna, S. Tura, G. Visani, G. Martinelli; Divisione di Ematologia, Ospedale S. Bortolo, Vicenza, F. Rodeghiero, E. Di Bona; Divisione di Ematologia, Policlinico S. Matteo, Pavia, C. Bernasconi, M. Lazzarino; Divisione di Medicina E, Ospedale S. Giovanni, Torino, L. Resegotti, M. Falda; Divisione di Ematologia, Policlinico Careggi, Firenze, P. Rossi Ferrini, F. Leoni; Divisione di Ematologia, Ospedali Riuniti, Bergamo, T. Barbui, A. Rambaldi; Divisione di Ematologia, Ospedale Civile, Pescara, G. Fioritoni, A. Recchia; Servizio di Ematologia, Policlinico, Bari, V. Liso, G. Specchia; Divisione di Ematologia, Ospedale A. Businco, Cagliari, G. Brocchia, W. Deplano; Servizio di Ematologia, Ospedale Civile, Avellino, E. Volpe, N. Cantore; Divisione di Ematologia, Ospedale A. Pugliese, Catanzaro, A. Peta, F. Iuliano; Divisione di Ematologia, Ospedale S. Gerardo, Monza, E. Pogliani, G. Corneo; Ematologia, Ospedale Generale e Regionale, Bolzano, P. Coser, P. Fabris; Sezione di Ematologia Spedali Civili, Brescia, T. Izzi, G. Rossi; Cattedra di Ematologia, Università, Catania, E. Cacciola, F. Di Raimondo; Cattedra di Ematologia, Università, Parma, V. Rizzoli, C. Almici; Cattedra di Ematologia, Università, Verona, G. Perona, D. Veneri; Cattedra di Ematologia, Università, Genova, M. Gobbi, M. Clavio; Divisione di Ematologia, Ospedale Cardarelli, Napoli, R. Cimino, F. Ferrara; Divisione di Ematologia, Osp. Nuovo Pellegrini, Napoli, R. De Biasi, E. Miraglia; Divisione di Ematologia, T.E.R.E., Napoli, L. De Rosa, V. Mettivier; Cattedra di Ematologia, Università Tor Vergata, Roma, S. Amadori, G. Aronica; Clinica Pediatrica, Ospedale S. Gerardo, Monza, G. Masera, A. Biondi, A. Luciano; Divisione di Ematologia, Università Cattolica, Roma, G. Leone, S. Sica; Divisione di Ematologia, Ospedali Riuniti, Reggio Calabria, F. Nobile, B. Martino; Sezione di Ematologia, Ospedale S. Croce, Cuneo, E. Gallo, A. Gallamini; Divisione di Ematologia, Ospedale S. Maria Goretti, Latina, L. Deriu, A. Cherichini; Sezione di Ematologia, CTMO, Cremona, A. Porcellini, S. Morandi; Divisione di Ematologia, Nuovo Policlinico, Napoli, B. Rotoli, C. Selleri; Cattedra di Ematologia, Università, Perugia, M.F. Martelli, A. Tabilio; Clinica Medica, Università, Palermo, A. Caizzo, M. Musso; Divisione di Ematologia, Ospedale V. Cervello, Palermo, F. Caronia, S. Mirto, A. Santoro; Divisione di Ematologia, Ospedale B. Gesù, Roma, G. De Rossi, M. Caniggia; Istituto di Ematologia, Nuovo Ospedale Torrette, Ancona, P. Leoni, M. Montillo;

Centro di Riferimento Oncologico, Aviano, S. Monfardini, V. Zagonel; Patologia Medica, Università, Genova, R. Ghio, E. Balleari; Clinica Medica, Policlinico S. Matteo, Pavia, E. Ascari, R. Invernizzi; Divisione di Ematologia, Università, Pisa, B. Grassi, M. Petrini; Ematologia, Ospedale S.S. Annunziata, Taranto, P. Mazza, G. Lazzari; Cattedra di Ematologia, Università, Udine, M. Baccarani, A. Candoni; Ematologia Pediatrica, Università, Catania, G. Schilirò, A.M. Ippolito; Ematologia, IV Divisione Pediatrica, Genova, L. Massimo, C. Micalizzi; Clinica Pediatrica, Università, Pavia, F. Severi, F. Locatelli; Ematologia, Ospedale Regionale A. Di Summa, Brindisi, G. Quarta, A. Melpignano; Cattedra di Ematologia, Università, Ferrara, G. Castoldi, F. Lanza; Semeiotica Medica, Università, Genova, F. Patrone, M. Sessarego; Divisione di Ematologia, Ospedale Niguarda, Milano, E. Morra, A.M. Nosari; Ematologia, Ospedale S. Raffaele, Milano, C. Bordignon, L. Camba; Ematologia ed Autotrapianto Ospedale S. Martino, Genova, A. M. Carella, F. Frassoni; Sezione di Ematologia, Ospedale S. Francesco, Nuoro, A. Gabbas, G. Latte; Cattedra di Ematologia, Policlinico, Palermo, P. Citarella, S. Grisanti; Divisione di Ematologia, Ospedale S. Salvatore, Pesaro, G. Lucarelli, G. Sparaventi; Sezione di Ematologia, Ospedale S. Carlo, Potenza, F. Ricciuti, M. Pizzuti; Divisione di Ematologia, Ospedale S. Camillo, Roma, A. De Laurenzi, L. Pacilli; Divisione di Ematologia, Casa Sollievo della Sofferenza, S.G. Rotondo, M. Carotenuto, S. Ladogana; Divisione di Ematologia, Ospedale A. Scavo, Siena, E. Dispensa, A. Bucalossi; Clinica Pediatrica, Ospedale G. Salesi, Ancona, P. Giorgi, L. Felici; Clinica Pediatrica I, Policlinico, Bari, F. Schettini, N. Santoro; Onco-Ematologia Pediatrica, Ospedale Regionale, Cagliari, P. Biddau; II Divisione Pediatrica, Ospedale Pausilipon, Napoli, V. Poggi, M.F. Pintà; Clinica Pediatrica I, Università, Napoli, M.T. Di Tullio, M. Giuliano; Clinica Pediatrica II, Università, Padova, L. Zanesco, M. Pilon; Clinica Pediatrica III, Università, Pisa, P. Macchia, C. Favre; Clinica Pediatrica, Università, Torino, E. Madon, R. Miniero; Department of Hematology, University of Nijmegen (NL) T. de Witte, P. Muus; Medizinische Klinik III, University of Munich (D), U. Jehn; Department of Hematology, University of Leiden (NL), R. Willemze; Department of Hematology, University of Ankara (TK), M. Beksac; and Az Middelheim, Afdeling Hemato-Oncologie, Antwerpen (B), R. De Bock.

REFERENCES

1. Stone R, Mayer RJ: The unique aspects of acute promyelocytic leukemia. *J Clin Oncol* 8:1913, 1990
2. Avvisati G, ten Cate JW, Mandelli F: Acute promyelocytic leukemia. *Br J Haematol* 81:315, 1992
3. Warrell RP Jr, de Thé H, Wang Z-Y, Degos L: Acute promyelocytic leukemia. *N Engl J Med* 329:177, 1993
4. Grignani F, Fagioli M, Alcalay M, Longo L, Pandolfi PP, Dotti E, Biondi A, Lo Coco F, Grignani F, Pelicci PG: Acute promyelocytic leukemia: From genetics to treatment. *Blood* 83:10, 1994
5. Biondi A, Rambaldi A, Pandolfi PP, Rossi V, Giudici G, Alcalay M, Lo Coco F, Diverio D, Pogliani EM, Lanzi EM, Mandelli F, Masera G, Barbui G, Pelicci PG: Molecular monitoring of the myl/RAR α fusion gene in acute promyelocytic leukemia by polymerase chain reaction. *Blood* 80:49, 1992
6. Miller WH, Kakizuka A, Frankel SR, Warrell RP Jr, DeBlasio A, Levine K, Evans RM, Dmitrovsky E: Reverse transcription-polymerase chain reaction for the rearranged retinoic acid receptor clarifies diagnosis and detects minimal residual disease in acute promyelocytic leukemia. *Proc Natl Acad Sci USA* 89:2694, 1992
7. Lo Coco F, Diverio D, Pandolfi PP, Biondi A, Rossi V, Avvisati G, Rambaldi A, Arcese W, Petti MC, Meloni G, Mandelli F, Grignani F, Masera G, Barbui T, Pelicci PG: Molecular evaluation of residual disease as a predictor of relapse in acute promyelocytic leukemia. *Lancet* 340:1437, 1992
8. Huang W, Sun G-L, Li X-S, Cao Q, Lu Y, Jang G-S, Zhang F-Q, Chai J-R, Wang ZY, Waxman S, Chen Z, Chen S-J: Acute promyelocytic leukemia: Clinical relevance of two major PML/RAR α isoforms and detection of minimal residual disease by reverse transcription-polymerase chain reaction to predict relapse. *Blood* 82:1264, 1993
9. Miller WH, Levine K, DeBlasio A, Frankel SR, Dmitrovsky E, Warrell RP Jr: Detection of minimal residual disease in acute promyelocytic leukemia by a reverse transcription-polymerase chain reaction assay for the PML/RAR α fusion mRNA. *Blood* 6:1689, 1993
10. Diverio D, Pandolfi PP, Rossi V, Biondi A, Pelicci PG, Lo Coco F: Monitoring of treatment outcome in acute promyelocytic leukemia by reverse transcription-polymerase chain reaction. *Leukemia* 12:327, 1994
11. Fukutani H, Naoe T, Ohno R, Yoshida H, Kiyoi H, Miyawaki S, Morishita H, Sano F, Kamibayashi H, Matsue K, Miyake T, Hasegawa S, Ueda Y, Kato Y, Kobayashi H, Shimazaki C, Kobayashi M, Kurane R, Sakota H, Masaki K, Wakayama T, Tohyama K, Nonaka Y, Natori H: Prognostic significance of the RT-PCR assay of PML/RAR α transcripts in acute promyelocytic leukemia. *Leukemia* 9:588, 1995
12. Bernard J, Weil M, Boiron M, Jacquillat C, Flandrin G, Gemon MF: Acute promyelocytic leukemia. Results of treatment with daunorubicin. *Blood* 41:489, 1973
13. Marty M, Ganem G, Fischer J, Flandrin G, Berger R, Schaison G, Degos L, Boiron M: Leucémie aigue promyélocitaire: étude retrospective de 119 malades traités par daunorubicine. *Nouv Rev Fr Hematol* 26:371, 1984
14. Petti MC, Avvisati G, Amadori S, Baccarani M, Guarini AR, Papa G, Rosti GA, Tura S, Mandelli F: Acute promyelocytic leukemia: Clinical aspects and results of treatment in 62 patients. *Haematologica* 72:151, 1987
15. Sanz MA, Jarque I, Martin G, Lorenzo I, Martinez J, Rafecas J, Pastor E, Sayas MJ, Sanz G, Gomis F: Acute promyelocytic leukemia. Therapy results and prognostic factors. *Cancer* 61:7, 1988
16. Avvisati G, Mandelli F, Petti MC, Vegna ML, Spadea A, Liso V, Specchia G, Bernasconi C, Alessandrino EP, Piatti C, Carella AM: Idarubicin (4-Demethoxidaunorubicin) as single agent for remission induction of previously untreated acute promyelocytic leukemia. A pilot study of the Italian cooperative group GIMEMA. *Eur J Haematol* 44:257, 1990
17. Head D, Kopecky KJ, Weick J, Files JC, Ryan D, Foucar K, Montiel M, Bickers J, Fishleder A, Miller M, Spier C, Hanson C, Bitter M, Brazier R, Mills G, Welborn J, Williams W, Hewlett J, Willman C, Appelbaum FR: Effect of aggressive daunomycin therapy on survival in acute promyelocytic leukemia. *Blood* 86:1717, 1995
18. Huang ME, Ye HC, Chen SR, Chai JR, Lu HX, Zoha L, Gu LJ, Wang ZY: Use of all-trans retinoic acid in the treatment of acute promyelocytic leukemia. *Blood* 72:567, 1988
19. Castaigne S, Chomienne C, Daniel MT, Ballerini P, Berger R, Fenaux P, Degos L: All-trans retinoic acid as a differentiation therapy for acute promyelocytic leukemia. I Clinical results. *Blood* 76:1704, 1990
20. Warrell RP Jr, Frankel SR, Miller WH Jr, Scheinberg DA, Itri LM, Hittelman WN, Vyas R, Andreeff M, Tafuri A, Jakubowski A, Gabrilove J, Gordon MS, Dmitrovsky E: Differentiation therapy for acute promyelocytic leukemia with tretinoin (all-trans retinoic acid). *N Engl J Med* 324:1385, 1991
21. Degos L, Dombret H, Chomienne C, Daniel MT, Miclea JM, Chastang C, Castaigne S, Fenaux P: All-trans retinoic acid as a differentiating agent in the treatment of acute promyelocytic leukemia. *Blood* 85:2643, 1995
22. Frankel SR, Eardley A, Lauwers G, Weiss M, Warrell RP Jr:

The 'retinoic acid syndrome' in acute promyelocytic leukemia. *Ann Intern Med* 117:292, 1992

23. Mahmoud HH, Hurwitz CA, Roberts WM, Santana VM, Ribeiro RC, Krance RA: Tretinoin toxicity in children with acute promyelocytic leukemia. *Lancet* 342:1394, 1993

24. Fenaux P, Le Deley MC, Castaigne S, Archimbaud E, Chomienne C, Link H, Guerci A, Duarte M, Daniel MT, Bowen D, Huembner G, Bauters F, Fegueux N, Fei M, Sanz MA, Lowenberg B, Maloisel F, Auzanneau G, Sadoun A, Gardin C, Bastion Y, Ganser A, Jacky E, Dombret H, Chastang C, Degos L: Effect of all-trans retinoic acid in newly diagnosed acute promyelocytic leukemia. Results of a multicenter randomised trial. *Blood* 82:3241, 1993

25. Avvisati G: Treatment of newly diagnosed acute promyelocytic leukemia: The experience of the Italian Cooperative Group GIMEMA during 12 years (1983-1995). *Sangre (Barc)* 40:136, 1995 (suppl 3)

26. Avvisati G, Lo Coco F, Diverio D, Falda M, Ferrara F, Lazarino M, Russo D, Petti MC, Mandelli F: AIDA (All-trans retinoic acid plus Idarubicin) in newly diagnosed acute promyelocytic leukemia: A GIMEMA pilot study. *Blood* 88:1390, 1996

27. Bennett JM, Catovsky D, Daniel MT, Flandrin G, Galton DAG, Gralnick HR, Sultan C: Proposed revised criteria for the classification of acute leukemias: A report of the French-American-British cooperative group. *Ann Intern Med* 103:629, 1985

28. Frankel SR, Eardley A, Heller G, Berman E, Miller WH Jr, Dmitrovsky E, Warrell RP Jr: All-trans retinoic acid for acute promyelocytic leukemia. Results of the New York study. *Ann Intern Med* 120:278, 1994

29. Chomczynsky P, Sacchi N: Single step method of RNA isolation by acid guanidium thiocyanate - phenol chloroform extraction. *Anal Biochem* 162:156, 1987

30. Kanamuru A, Takemoto Y, Tanimoto M, Murakami H, Asou N, Kobayashi T, Kuriyama K, Ohmoto E, Sakamaki H, Tsubaki K, Hiraoka A, Yamada O, Oh H, Saito K, Matsuda S, Minato K, Ohno R: All-trans retinoic acid for the treatment of newly diagnosed acute promyelocytic leukemia. *Blood* 85:1202, 1995

31. Licht JD, Chomienne C, Goy A, Chen A, Scott AA, Head DR, Michaux JL, Wu Y, DeBlasio A, Miller WH Jr, Zelentz AD, Willman CL, Chen Z, Chen SJ, Zelent A, Macintyre E, Veil A, Cortes J, Kantarjian H, Waxman S: Clinical and molecular characterization of a rare syndrome of acute promyelocytic leukemia associated with translocation t(11;17). *Blood* 85:1083, 1995

32. Redner RL, Rush EA, Faas S, Rudert WA, Corey SJ: The t(5;17) variant of acute promyelocytic leukemia expresses a nucleophosmin-retinoic acid receptor. *Blood* 87:882, 1996

33. Scott AA, Head DR, Kopecky KJ, Appelbaum FR, Theil KS, Grever MR, Chen IM, Whittaker MH, Grift BB, Licht JD, Waxman S, Whalen MM, Bankhurst AD, Richter LC, Groogan TM, Willman CL: HLA-DR-, CD33+, CD56+, CD16- myeloid/natural killer acute leukemia: A previously unrecognized form of acute leukemia potentially misdiagnosed as French-American-British acute myeloid leukemia. *Blood* 84:244, 1994

34. Dombret H, Scrobohaci ML, Ghorra P, Zini JM, Daniel MT, Castaigne S, Degos L: Coagulation disorder in acute promyelocytic leukemia: Corrective effect of all-trans retinoic acid treatment. *Leukemia* 7:2, 1993

35. Falanga A, Iacoviello L, Evangelista V, Bellotti D, Consonni R, D'Orazio A, Robba L, Donati MB, Barbui T: Loss of blasts cell procoagulant activity and improvement of hemostatic variables in patients with acute promyelocytic on all-trans retinoic acid. *Blood* 86:1072, 1995

36. Avvisati G, Dragoni F, Chistolini A, Mazzucconi MG, Pallotta A, Latagliata R, Petti MC, Mandelli F: Rapid amelioration of the coagulopathy in acute promyelocytic leukemia (APL) treated with a combination of all-trans retinoic acid (ATRA) plus idarubicin (AIDA protocol). *Thromb Haemost* 73:1432, 1995 (abstr 2034)

37. Mandelli F, Labopin M, Granena A, Iriando A, Prentice G, Bacigalupo A, Sierra J, Meloni G, Frassoni F, Goldman J, Gratwohl A, Gorin NC, for the EBMT: European survey of bone marrow transplantation in acute promyelocytic leukemia (M3). *Bone Marrow Transplant* 14:293, 1994