

## REVIEW ARTICLE

# Biology and Treatment of Childhood T-Lineage Acute Lymphoblastic Leukemia

By Fatih M. Uckun, Martha G. Sensel, Lei Sun, Peter G. Steinherz, Michael E. Trigg, Nyla A. Heerema, Harland N. Sather, Gregory H. Reaman, and Paul S. Gaynon

**A**CUTE LYMPHOBLASTIC LEUKEMIA (ALL) is the most prevalent type of cancer, as well as the most common form of leukemia in children.<sup>1</sup> This lymphoid malignancy, manifested by the proliferation of lymphopoietic blast cells, represents a heterogeneous group of diseases that vary with respect to morphological, cytogenetic, and immunologic features of the transformed cells. Technical improvements in immunofluorescence staining and flow cytometry together with the availability of numerous monoclonal antibodies (MoAbs) that recognize lineage-associated membrane molecules have illuminated the immunophenotypic heterogeneity in ALL. We now know that leukemia cells from patients with ALL may express various combinations of surface antigens that are found normally on lymphocyte precursors at discrete stages of maturation.<sup>2,3</sup> Thus, the malignant clones in patients with ALL are thought to originate from normal lymphoid progenitor cells arrested at early stages of B- or T-lymphocyte ontogeny. Although cells from the majority ( $\approx 85\%$ ) of pediatric patients express B-lineage-associated antigens, those from approximately 15% of patients express the T-lineage-associated antigens CD1, CD2, CD3, CD4, CD5, CD7, or CD8.<sup>4,6</sup> T-lineage ALL in children is associated with numerous unfavorable presenting features, thus it is not surprising that children with T-lineage ALL frequently have been reported to have a worse prognosis than children with B-lineage ALL.<sup>4,5,7-10</sup> However, a number of encouraging reports from recent clinical studies using contemporary risk-adjusted multiagent chemotherapy programs have documented remarkably improved outcomes for patients with T-lineage ALL.<sup>6,10-14</sup> Moreover, advanced preclinical studies have triggered much optimism that new agent discovery programs may lead to further improvements in outcome in the near future. In this review, we discuss current concepts regarding the etiology, biological characteristics, clinical features, and treatment of pediatric T-lineage ALL.

### ETIOLOGY

The role of numerous epidemiological factors, including maternal and paternal exposure to radiation, history of maternal fetal loss or fertility problems, higher birthweight at diagnosis, and use of exogenous growth hormone, remains controversial in the cause of pediatric ALL.<sup>15-17</sup> A recent comprehensive review found no relationship between exposure to electromagnetic field (EMF) radiation and incidence of childhood ALL.<sup>18</sup> The reported space-time clustering of ALL cases, which might suggest an etiologic agent such as a virus, is also controversial.<sup>19-23</sup> Human T-cell leukemia virus-I and II may be associated with adult, but not pediatric T-lineage leukemia or lymphoma,<sup>24,25</sup> and Epstein-Barr virus infection has been linked to a limited number of cases of T-cell lymphoma, but not T-lineage ALL, in children.<sup>26</sup>

The autosomal recessive disorder ataxia telangiectasia (AT) appears to be a true etiologic factor because patients with AT have an increased risk of developing lymphoid malignancies,

including T-lineage ALL.<sup>27</sup> Translocations involving the T-cell receptor (TCR) loci are reported in approximately 10% of the T cells from patients with AT,<sup>28</sup> but interestingly, the most frequent of these translocations appear to involve different regions within the TCR loci compared with those observed in patients with T-lineage ALL without AT.<sup>29-31</sup> The molecular basis for these effects as well as other genetic abnormalities that may play a role in T-lineage leukemia will be discussed below. Taken together, these data suggest that multiple factors may be involved in the origin of T-lineage ALL.

### BIOLOGICAL FEATURES OF T-LINEAGE ALL

Because leukemic cells are thought to originate from normal T-lymphocyte precursors arrested at early stages of ontogeny,<sup>2,32</sup> every pathway that ensures homeostasis of a functional immune system is a potential target for disruption. Still, the fundamental issue of how many different mutations are required for malignant transformation to the leukemic state remains to be delineated. Nevertheless, clear associations have been identified between the occurrence of nonrandom translocations or other gene mutations and the development of T-lineage ALL. Below, we describe the specific molecular defects found in T-lineage leukemias and discuss altered signal transduction pathways that may contribute to the malignancy.

*Chromosomal translocations.* An array of nonrandom translocations that are specific to T-lineage ALL have been identified; all appear to occur preferentially in the TCR loci on chromosomes 14 and 7.<sup>33</sup> The breakpoints in many cases resemble TCR recombination signals, implying that the aberration arose during TCR rearrangement.<sup>34-39</sup> Translocations involv-

---

*From the Children's Cancer Group (CCG) ALL Biology Reference Laboratory, and Wayne Hughes Institute, St Paul, MN; Group Operations Center, Children's Cancer Group, Arcadia, CA; the Department of Pediatrics, Sloan-Kettering Cancer Center, New York, NY; the Department of Hematology-Oncology, University of Iowa Hospitals and Clinic, Iowa City; the Department of Medical and Molecular Genetics, Indiana University School of Medicine, Indianapolis; the Department of Preventive Medicine, University of Southern California, Los Angeles, and Group Operations Center, Children's Cancer Group, Arcadia, CA; the Department of Hematology-Oncology, Children's National Medical Center, and the George Washington University, Washington, DC; and the Department of Pediatric Hematology-Oncology, University of Wisconsin, Madison.*

*Submitted May 21, 1997; accepted October 9, 1997.*

*Supported in part by research grants including CCG Chairman's Grants No. CA-13539, CA-51425, CA-42633, CA-42111, CA-60437, and CA-27137 from the National Cancer Institute, National Institutes of Health. F.M.U. is a Stohlman Scholar of the Leukemia Society of America, New York, NY.*

*Address reprint requests to Fatih M. Uckun, MD, PhD, Children's Cancer Group ALL Biology Reference Laboratory and Wayne Hughes Institute, 2665 Long Lake Rd, St Paul, MN 55113.*

*© 1998 by The American Society of Hematology.*

*0006-4971/98/9103-0037\$3.00/0*

ing chromosomes 1 and 14, such as t(1;14)(p33;q11) and t(1;14)(p32;q11), have been estimated to occur in approximately 3% of T-lineage ALL cases.<sup>40</sup> In such rearrangements, the SCL/TCL5/TAL-1 gene from chromosome 1 and the TCR $\delta$  gene on chromosome 14<sup>36,41,42</sup> are juxtaposed, resulting in deregulation of normal TAL-1 expression.<sup>41,43</sup> TAL-1 was predicted to encode a protein containing a helix-loop-helix DNA binding motif,<sup>42,43</sup> suggesting that the t(1;14) translocations could contribute to leukemogenesis by inducing aberrant expression of novel or TAL-1-regulated genes.

A distinct TAL-1 disruption occurs via an interstitial deletion between a locus called SIL (SCL interrupting locus) and the 5'UTR of SCL, resulting in a fusion transcript SIL/SCL, and is estimated to occur with a frequency of 16% to 26% in T-lineage ALL.<sup>44-46</sup> Presence of a TAL-1 disruption was correlated with high white blood cell (WBC) count, high hemoglobin level, and CD2<sup>+</sup>/CD10<sup>-</sup> immunophenotypes, and interestingly, 4-year event-free survival (EFS) was higher for patients with TAL-1 disruption compared to those without TAL-1 alterations (59%  $\pm$  11% v 44%  $\pm$  7%, respectively), although this difference did not reach conventional significance.<sup>46</sup> Although TAL-1 is required for development of all hematopoietic lineages in mice,<sup>47</sup> the gene is not expressed in B- or T-lineage cells,<sup>41,48</sup> and interestingly, SCL-transfected, v-ABL-transformed cells appear to be oncogenic in mice.<sup>49</sup> Taken together, these data suggest that disruption of normal TAL-1 expression may contribute to the transformation of T-cell precursors into leukemic blasts.

The t(10;14)(q24;q11) translocation, first identified in T-cell neoplasms including T-lineage ALL, involves the TCR $\alpha$ /TCR $\delta$  locus on chromosome 14<sup>34,35,50</sup> and the TCL3 locus on chromosome 10.<sup>34,35</sup> An open reading frame within TCL3 encodes a novel homeobox protein, HOX-11, whose expression is deregulated as a result of the translocation.<sup>51-53</sup> Moreover, like TAL-1, HOX-11 is capable of DNA binding and transcriptional activation of reporter genes, suggesting a role for this gene in leukemic transformation.<sup>54</sup> Additional studies showed that whereas HOX-11 was expressed in leukemic cell lines and leukemic blasts, it was not expressed in normal T lymphocytes,<sup>52,53,55</sup> but was required for normal spleen development.<sup>56</sup> Reverse transcriptase-polymerase chain reaction (RT-PCR) assays have suggested that HOX-11 alterations may occur with high frequency in patients with T-lineage ALL.<sup>57</sup> Thus, deregulation of HOX-11 is likely to be a biologically significant factor in development of T-lineage ALL.

Translocations t(11;14)(p13;q11) and t(11;14)(p15;q11) also are observed frequently in T-lineage ALL<sup>58-60</sup>; both involve breakpoints within diversity or J segments the region the TCR $\alpha$  or TCR $\delta$  genes on chromosome 14.<sup>37,61,62</sup> McGuire et al<sup>61</sup> described multiple open reading frames near the chromosome 11 breakpoints and identified one at 11p15 as the open reading frame of the TTG-1 gene. Similarly, Boehm et al<sup>63</sup> identified the involved region of 11p15 as the rhombotin gene. Both genes encode proteins characterized by duplicate cysteine-rich zinc-finger protein binding homology domains.<sup>61,63</sup> A related gene, rhombotin-2/TTG-2, was shown to be deregulated in cases involving 11p13.<sup>63,64</sup> Consistent with the predicted structure of the rhombotins, a recent report described the identification of an ets family transcription factor, ELF-2, that contains rhom-

botin-2 binding domains, suggesting a transcriptional regulatory role for rhombotin-2.<sup>65</sup> Clinically, several investigators have associated t(11;14) translocations with an immature stage of thymocyte development,<sup>37,59,60</sup> but the overall prognostic significance of this translocation remains unclear.

Although translocations involving chromosome 7 occur in both B-precursor and T-lineage ALL, those involving the TCR- $\beta$  locus at 7q32-36 are specific for T-lineage ALL.<sup>66</sup> One such translocation, t(7;19), truncates the lyl-1 gene on chromosome 19,<sup>67</sup> presumably resulting in altered DNA-binding ability for lyl-1.<sup>68</sup> Another case, t(7;9), results in truncation of the TAN-1 gene on chromosome 9.<sup>69</sup> The mouse homologue of TAN-1 is expressed ubiquitously, but is most abundant in lymphoid tissues, suggesting that normal expression of TAN-1 is disrupted in t(7;9)<sup>+</sup> ALL.<sup>69</sup>

The distinct translocation t(1;7)(p34;q34) was shown to juxtapose the TCR- $\beta$  constant region enhancer upstream of the LCK gene, which encodes an SRC family protein tyrosine kinase that is involved in signal transduction through CD4.<sup>70,71</sup> Notably, overexpression of LCK in transgenic mice causes thymomas or both thymomas and peripheral lymphoid malignancies,<sup>72,73</sup> suggesting a role for deregulated LCK expression in leukemogenesis. The c-myc locus on chromosome 8 defines yet another class of translocations associated with T-lineage ALL. In t(8;14)(q24;q11), c-myc is translocated with the TCR $\alpha$  loci on chromosome 14, resulting in deregulation of myc expression.<sup>74,75</sup> In t(2;8), a fusion protein is produced that consists of c-myc and the product of an unidentified locus on chromosome 2.<sup>76</sup> The frequency and significance of these translocations are unclear at present.

We have recently determined the frequency and clinical significance of chromosomal abnormalities in a large cohort of patients with T-lineage ALL enrolled on contemporary CCG studies (Heerema N., et al, submitted for publication). Translocations involving 14q11 and 7q32-q36 were among the most frequent abnormalities, but non-TCR loci, including 9p, 6q, 11q23, and 14q32, also were frequently altered. Notably, none of these abnormalities had prognostic significance in the context of the intensive therapies used in contemporary CCG studies. Nevertheless, the array of chromosomal rearrangements described above are a hallmark of the biological diversity of T-lineage ALL and are likely to result from alterations in underlying cellular control mechanisms. Indeed, recent advances in our understanding of cell signaling and cell cycle control suggest that defective cell surveillance mechanisms are likely to be the major factors leading both to unrestrained proliferation of leukemic cells and to the development of chromosomal abnormalities, including translocations, pseudodiploidy, and hyperdiploidy, that are associated with leukemic cells.<sup>77-80</sup> Alterations in such control mechanisms are discussed below.

*Mutation or loss of cell cycle control genes.* Mutations present in malignant cells allow them to circumnavigate regulators that control proliferation and differentiation. The retinoblastoma (Rb) gene was originally identified as a tumor suppressor gene because of its inactivation in cases of retinoblastoma; prostate, breast, and lung cancers; and leukemias.<sup>81</sup> Notably, the telomeric Rb1 gene is located on the long arm of chromosome

13 (13q14), which is inactivated or deleted in approximately 6% of T-lineage ALL cases.<sup>82,83</sup>

In addition to Rb, other proteins that affect cell cycle progression include the cyclin-dependent kinase inhibitors p21, p27, and p57, as well as the inhibitors of Cdk4 (Ink4): p15<sup>Ink4b</sup>, p16<sup>Ink4a</sup>, p18<sup>Ink4c</sup>, and p19<sup>Ink4d</sup>.<sup>84-89</sup> Among the Ink4 family of inhibitors, p15<sup>Ink4b</sup> and p16<sup>Ink4a</sup> have been implicated for a role in the biology of T-lineage ALL.<sup>90-95</sup> Both genes map to 9p21, a region on the short arm of chromosome 9 previously shown to be deleted frequently in T-lineage ALL.<sup>33,96-98</sup> In addition, Batova et al<sup>95</sup> recently reported that the 5' promoter region of the p15 gene is preferentially hypermethylated, presumably resulting in loss of transcriptional expression in 38% of newly diagnosed T-lineage ALL.

Another critical regulator of cell cycle progression, the p53 gene, is the most frequently mutated gene in human cancers.<sup>81</sup> The major function of p53 is to ensure that cells arrest and attempt to repair genotoxic damage before replicating DNA and entering mitosis.<sup>99</sup> In p53-deficient mice, the most common tumor that arises is a T-lineage lymphoid malignancy.<sup>100</sup> Although p53 mutations are infrequently observed at diagnosis, they are associated with relapse in pediatric T-lineage ALL.<sup>101,102</sup>

Another sensor for cell damage appears to be the ATM gene product, which is mutated in patients with AT.<sup>103</sup> After insult with agents that induce sublethal DNA damage, cells from patients with AT fail to block DNA synthesis and thereby fail to repair the damaged DNA.<sup>104</sup> These effects are apparently caused by a failure of the mutated ATM gene to regulate p53.<sup>105</sup> ATM-deficient mice develop an aggressive form of T-lineage leukemia/lymphoma,<sup>106,107</sup> and, as described above, children with AT frequently develop T-lineage ALL,<sup>27,108,109</sup> implicating ATM in leukemogenesis.

Other genes implicated in the malignant transformation of leukemic cells are Ets-1 and IKAROS. The Ets-1 T-lymphocyte transcription factor is thought to be important for normal thymic development and for prevention of cell death in normal mature T cells. A mutation in the DNA binding domain of the Ets-1 was reported in a case of T-lineage ALL,<sup>110</sup> but the clinical significance of this finding remains to be proven. The IKAROS gene encodes a zinc finger DNA binding protein that is required for lymphoid cell differentiation.<sup>111</sup> Heterozygous transgenic mice harboring a defective IKAROS gene develop a very aggressive form of T-cell leukemia, suggesting that IKAROS may serve as a suppressor of leukemic transformation.<sup>112</sup>

Leukemic cells also appear to be altered in their responses to various stimuli that induce apoptosis. Debatin et al<sup>113,114</sup> reported that primary leukemic cells and cell lines from adult patients with T-cell leukemia were sensitive to FasL-induced cell killing in vitro, whereas leukemic cells from pediatric patients with T-lineage ALL were resistant. Resistance was unrelated to the quantity of Fas on the cell surface, but was reversed by treatment with the protein synthesis inhibitor cycloheximide, suggesting that short-lived proteins were required for maintenance of the resistant phenotype. In vivo treatment of a human T-lineage ALL-engrafted severe combined immunodeficiency (SCID) mouse with an anti-Fas antibody resulted in prolonged survival, but did not eradicate the disease, supporting the existence of Fas sensitive and insensitive leukemic cells.<sup>115</sup> These data suggest that altered responses

to apoptotic stimuli or regulatory factors may contribute to the ability of leukemic cells to escape killing by either immune surveillance or cytotoxic agents.

Bcl-2, which protects cells from non-Fas-mediated apoptosis,<sup>116,117</sup> is expressed in both T-lineage and B-lineage leukemias, but it is not yet known how this affects their ability to survive cytotoxic treatments. A related protein, Bax,<sup>118</sup> acts as an antagonist to Bcl-2 and may confer radiation sensitivity to cells.<sup>119</sup> In a recent CCG study, we found a marked variation in Bcl-2 expression by primary leukemic cells from 238 children with newly diagnosed ALL, including 52 patients with T-lineage ALL.<sup>120</sup> High-risk features, such as high WBC count, organomegaly, presence of MLL-AF4 or BCR-ABL fusion transcripts, or leukemic cell growth in SCID mice, were not associated with Bcl-2 expression in these patients. For patients with T-lineage ALL, high Bcl-2 expression was predictive of slow early response (ie, M3 day 14 marrow status). However, with limited follow-up and overall excellent outcome for patients, this correlation did not extend to EFS.

#### CLINICAL FEATURES AND TREATMENT OF T-LINEAGE ALL

T-lineage ALL is distinct from B-lineage ALL not only biologically, but also clinically. Although the basis for these differences is not well understood, clinical characteristics have been useful prognostic factors for guiding the use of experimental treatments. Below, we describe common presenting features, prognostic variables, and treatment outcome of patients with T-lineage ALL based on data accumulated over the last decade. We then focus on causes for treatment failure and discuss new strategies for improving outcome among subgroups of patients who remain at risk for relapse despite intensive therapy.

*Presenting features.* The relationship between T-lineage markers and unfavorable presenting characteristics was first noted by Borella, Sen, and others,<sup>121-124</sup> and numerous studies have now confirmed that compared to patients with B-lineage ALL, those with T-lineage ALL more frequently show the highest WBC range ( $\geq 50,000/\mu\text{L}$ ), are nonwhite, older, exhibit marked enlargement of the spleen, liver, and lymph nodes, and have a mediastinal mass.<sup>5,7,9,10,125</sup>

Modal chromosome number is often abnormal among patients with ALL, with hyperdiploidy ( $>50$  chromosomes) correlated with favorable outcome and pseudodiploidy associated with poor outcome.<sup>58,98,126-129</sup> The hyperdiploid karyotype is more often associated with pre-B or early pre-B immunophenotypes,<sup>129</sup> whereas the pseudodiploid karyotype is more often associated with the T-lineage immunophenotype.<sup>33</sup> Also, "near tetraploid" chromosome number ( $>65$ ) is more often associated with T-lineage ALL and poor outcome.<sup>130</sup> As described above, nonrandom translocations in T-lineage ALL preferentially occur in the TCR loci on chromosomes 7 and 14,<sup>33</sup> and those involving the TCR $\beta$  locus at 7q32-36 and the TCR $\alpha\delta$  14q11 collectively occur in approximately 20% of all T-lineage ALL cases.<sup>33</sup>

*Risk classification of T-lineage ALL.* In general, treatment protocols for childhood leukemias have relied on the known prognostic factors of age and WBC count, as well as organomegaly rather than immunophenotype for risk assessment. As a result, even though many patients with T-lineage ALL were

previously misclassified or not immunophenotyped, they were likely to receive treatment for high-risk ALL based on their other presenting features. In contemporary trials, various groups have used somewhat different criteria for classification, which has complicated comparisons of results between groups, but nevertheless has generally resulted in similar assignment of patients with T-lineage ALL to more intensive treatment protocols, such as Berlin-Frankfurt-Munster (BFM),<sup>131,132</sup> modified BFM,<sup>12</sup> and the New York (NY) regimen,<sup>13</sup> as well as those of the St Jude Children's Research Hospital<sup>133</sup> and Dana Farber Cancer Institute.<sup>11</sup>

From 1983 through 1993, children diagnosed with ALL who exhibited National Cancer Institute (NCI) standard risk features<sup>134</sup> were classified by the CCG as either low risk (ages 2 through 9 years and WBC <10,000/ $\mu$ L) or intermediate risk (ages 2 through 9 years and WBC <10,000 to 49,999/ $\mu$ L, or age 1 year and WBC <50,000/ $\mu$ L), whereas patients exhibiting NCI poor-risk characteristics were classified as follows: high risk, ages 1 through 9 years with WBC  $\geq$ 50,000/ $\mu$ L or age >10 years; infants, age <1 year; lymphomatous, patients with specific high-risk features, as described.<sup>135</sup> As shown in Table 1, patients with T-lineage ALL more frequently were assigned to the higher risk than to the lower risk protocols, which is consistent with their clinical features described above.

**Treatment outcome in T-lineage ALL.** As noted above, previous studies showed poorer outcomes for patients with T-lineage ALL compared with patients with B-lineage ALL. For example, in the BFM group, Henze et al<sup>136</sup> reported poor outcome for patients with T-lineage ALL who were treated on DAL (adapted from St Jude protocol VII), with 9-year probabilities of continuous complete remission (CCR) of 9%  $\pm$  9% and 41%  $\pm$  5%, for T-lineage and non-T-lineage, respectively. In contrast, patients treated on BFM achieved CCR of 52%  $\pm$  13% and 65%  $\pm$  5%, respectively, suggesting that BFM provided superior treatment for T-lineage ALL.

Investigators of the Pediatric Oncology Group<sup>7</sup> treated 53 patients with T-lineage ALL with a modified LSA<sub>2</sub>L<sub>2</sub> regimen that had been shown to be efficacious for treatment of T-cell non-Hodgkin's lymphoma.

Although complete remission was achieved for 88% of the patients, the projected overall 3-year EFS was only 40%

(SE = 8.3%). Moreover, for patients with WBC count <50,000, the projected 3-year EFS was 67%, whereas for patients with WBC count >50,000, 3-year EFS was only 19%. In a follow-up study, 253 children with T-lineage ALL treated by a modified LSA<sub>2</sub>L<sub>2</sub> regimen together with cranial radiation therapy and triple intrathecal therapy for presymptomatic treatment of central nervous system (CNS) disease achieved an overall 4-year EFS of 43% (SE = 4%).<sup>8</sup> Thus, although outcomes improved, the LSA<sub>2</sub>L<sub>2</sub> regimen remained ineffective for the majority of patients with T-lineage ALL. Similarly, in an analysis of data from St Jude studies X and XI, conducted from 1979 to 1983, 120 children with T-lineage ALL had a 5-year EFS of 46% (SE = 18%).<sup>137</sup> In a French trial, Garand et al<sup>10</sup> treated 88 pediatric patients with T-lineage ALL by protocols such as BFM or FRALLE,<sup>138</sup> and an EFS of approximately 58% was reported for a median follow-up of 30 months, suggesting that such therapy could improve outcome for these patients.

Although the studies described above generally found unfavorable outcomes for patients with T-lineage ALL, other recent studies have reported improved outcomes through the use of highly intensive treatment protocols. For example, using an intensive four-drug induction and multidrug continuation, including doxorubicin and prednisone together with prophylaxis for CNS disease and high-dose L-asparaginase, Clavell et al<sup>11</sup> reported improved outcome (4-year EFS of 71%) for high-risk patients, including those who had T-lineage ALL. More recently, in a study by Schorin et al<sup>14</sup> 20 patients with T-lineage ALL treated with multiagent chemotherapy together with cranial irradiation and intrathecal methotrexate for 2 years also had favorable outcomes (7-year EFS of 70%, SE = 10%).<sup>14</sup> The favorable outcome was attributed to the inclusion of L-asparaginase and doxorubicin in the treatment regimen.

Studies by the CCG also have shown improvements in EFS outcome for high-risk patients with ALL including those with the T-lineage immunophenotype. Steinherz et al<sup>13</sup> used an intensive multidrug chemotherapy (NY regimen) to treat 100 patients with characteristics previously correlated with a high risk for relapse. This patient population included 13 patients with T-lineage ALL (defined as E-rosette+). Four-year EFS for the entire cohort was 69% (SE = 5%), whereas 4-year EFS for patients with T-lineage ALL was 75%. Gaynon et al<sup>139</sup> used a modified BFM therapy involving four-drug induction and aggressive continuation therapy to treat high-risk children, including 60 who were E-rosette+. Overall 3-year EFS was 65% (SD = 3.5%); patients with WBC count >50,000 who were E-rosette+ had a 3-year EFS of 75% (SD = 6.9%), whereas those who were E-rosette- had an EFS of 51% (SD = 6.3%).

To investigate the outcome of patients with T-lineage ALL on these regimens more thoroughly, we recently analyzed data from the large cohort of patients enrolled on CCG studies conducted between 1983 and 1993.<sup>134</sup> Notably, we observed a significant improvement in outcome of patients with T-lineage ALL compared with those on earlier studies because of marked decreases in the incidences of induction failures, early bone marrow relapses, and CNS relapses when more aggressive therapy was given (Fig 1). The probability of 3-year survival for patients with T-lineage ALL increased from 56% in studies conducted between 1978 and 1983, to 65% in studies conducted

**Table 1. CCG and NCI Risk Group Classification of Children With B-Lineage and T-Lineage Acute Lymphoblastic Leukemia**

Risk Group	B-Lineage ALL N = 3,668		T-Lineage ALL N = 730	
	N	(%)*	N	(%)†
CCG-Low	705	(19.2)	58	(8.0)
CCG-Intermediate	1,575	(42.9)	71	(9.7)
CCG-High	1,059	(28.9)	169	(23.2)
CCG-Lymphomatous	216	(5.9)	425	(58.2)
CCG-Infant	113	(3.1)	7	(1.0)
Total	3,668	(100.0)	730	(100.0)
NCI-Standard	2,213	(60.3)	211	(28.9)
NCI-Poor	1,455	(39.7)	519	(71.1)
Total	3,668	(100.0)	730	(100.0)

\*Percentage of patients with B-lineage ALL classified into each risk group.

†Percentage of patients with T-lineage ALL classified into each risk group.




Fig 1. Improved EFS of patients with T-lineage ALL in the context of contemporary intensive chemotherapy programs. EFS for the entire cohort of patients with T-lineage and B-lineage ALL treated on the 1800 series and 100 series of CCG studies are shown. EFS values at designated points in follow-up are given in the text.

between 1983 and 1989, and to 78.8% in studies conducted between 1989 and 1993 (Table 2). Taken together, these various studies suggest that current risk for patients with T-lineage ALL treated by intensive therapeutic regimens is similar to that of patients with B-lineage ALL. Thus, a major improvement in treatment of T-lineage ALL has been achieved.

**Prognostic factors in T-lineage ALL.** A number of risk factors for T-lineage ALL were identified in the studies described above. For example, Dowell et al<sup>9</sup> and Shuster et al<sup>8</sup> reported that compared with patients with T-lineage ALL whose leukemic cells were CD10<sup>-</sup>, those whose cells were CD10<sup>+</sup> were more likely to achieve remission and have significantly improved EFS outcomes. In another study, Pui et al<sup>137</sup> reported that CD3 positivity in association with an abnormal karyotype was a significant adverse risk factor; 5-year EFS for patients with both of these characteristics was 35%. In contrast, Shuster et al<sup>8</sup> found no prognostic significance for CD3 expression; rather, the most important favorable prognostic factors for patients with low WBC count or high WBC count at diagnosis were CD5 positivity or expression of the THY antigen, respectively.

The findings that many patients with T-lineage ALL now can achieve a much improved outcome has motivated attempts to identify subgroups of patients within T-lineage ALL that may exhibit improved or reduced probabilities of survival. Two

previous CCG studies described above noted a favorable association between outcome and E-rosette (CD2) positivity among high-risk patients.<sup>13,139</sup> To determine comprehensively the clinical significance of CD2 expression in T-lineage ALL, we prospectively immunophenotyped leukemic cells from the large cohort of children enrolled on CCG studies between 1983 and 1993.<sup>140</sup> We noted a statistically significant correlation ( $P = .0006$ ) between the CD2 antigen expression frequency (ie, the average percentage of blasts that were positive for CD2) and EFS. Compared with patients with the highest CD2 expression level, patients with intermediate and low CD2 expression frequencies had relative hazard rates (RHR) of 1.27 and 2.01, indicating an increased risk of treatment failure. After 6 years of follow-up, the EFS estimates for the three CD2 expression groups (low expression frequency to high expression frequency) were 49.3%, 63.5%, and 72.2%, respectively. CD2 expression remained a significant predictor of EFS after adjustment for the effects of other covariates by multivariate regression. Expression of other antigens (CD3, CD5, CD10, or CD34) by leukemic cells was not correlated with EFS. Thus, the expression frequency of CD2 antigen is a powerful predictor of EFS that may be useful for risk classification or assignment to novel therapies aimed at improving patient outcome.

Maturation stage of the predominant leukemic clones also has been suggested as a means for subgrouping patients with T-lineage ALL. Crist et al<sup>4</sup> stratified 101 patients with T-lineage ALL into three maturation groups according to expression of T-lineage cell surface antigens, as follows: stage I, CD2<sup>+</sup>CD7<sup>+</sup>; stage II, CD2<sup>+</sup>CD7<sup>+</sup>CD1<sup>+</sup>CD4<sup>+</sup>CD8<sup>+</sup>; and stage III, CD2<sup>+</sup>CD7<sup>+</sup>CD1<sup>-</sup>(CD4<sup>+</sup> or CD8<sup>+</sup>)CD3<sup>+</sup>. Although the percentage of patients achieving remission following induction therapy was lower for patients with T-lineage ALL of the earliest maturation stage (79%, 100%, and 94% for stages I, II, and III, respectively), 4-year EFS was equally poor for all three groups (33%, 32%, and 38%, respectively).

Recently, we analyzed data from a large cohort of patients with T-lineage ALL treated on contemporary protocols of the CCG to further investigate the prognostic role of the apparent maturation stage of leukemic T-cell precursors.<sup>141</sup> Patients were immunophenotypically classified as follows: pro-thymocyte leukemia (pro-TL), CD7<sup>+</sup>CD2<sup>-</sup>CD5<sup>-</sup>; immature TL, CD7<sup>+</sup>(CD2<sup>+</sup> or CD5<sup>+</sup>)CD3<sup>-</sup>; and mature TL, CD7<sup>+</sup>CD2<sup>+</sup>CD5<sup>+</sup>CD3<sup>+</sup>. No group had a preponderance of favorable or unfavorable presenting characteristics. Four-year EFS was lower for patients with pro-TL (57.1%; SD = 8.4%) compared with patients with immature and mature TL (68.5%, SD = 3.5%; and 77.1%, SD = 4.0%; respectively) with an overall significance of  $P = .05$ . Highly significant differences were found for overall survival ( $P = .005$ ) as a result of the deaths of all patients with pro-TL who relapsed. Although CD2 also was a significant prognostic factor ( $P = .03$ ), RHRs of 2.11, 1.51, and 1.17 for patients with pro-TL, CD2<sup>-</sup> immature TL, and CD2<sup>+</sup> immature TL, respectively, suggested that the pro-TL maturation stage had added prognostic significance (Fig 2). Indeed, multivariate analysis indicated that the influence of ontogeny group was greater than that of CD2. Thus, leukemic cells of the pro-TL maturation stage identified a subgroup of patients with T-lineage ALL who have a significantly worse

Table 2. Outcome for Patients With T-Lineage ALL Treated During Three Consecutive CCG Treatment Eras

CCG Study Era	Years	Event-Free Survival (%)	
		3-Year	5-Year
CCG-160s	1978-1983	56.4	52.5
CCG-100s	1983-1989	65.8	61.0
CCG-1800s	1989-1993	78.2	75.2




Fig 2. EFS of patients with T-lineage ALL according to the apparent maturational stage of bone marrow leukemic blasts. EFS for (A) mature TL, (B) CD2<sup>+</sup> immature TL, (C) CD2<sup>-</sup> immature TL, and (D) pro-TL patients treated on the 1800 series of CCG protocols are shown. EFS values at designated points in follow-up are given in the text.

EFS outcome than patients whose leukemic cells correspond to a more mature stage of development.

The variant immunophenotype in which leukemic cells coexpress T-lineage- and myeloid-associated antigens represents a controversial prognostic factor. Although numerous investigators have reported that coexpression of myeloid antigens predicted an adverse risk for patients with T-lineage ALL,<sup>142,143</sup> others have found similar outcomes for myeloid antigen negative (My<sup>-</sup>) and myeloid antigen positive (My<sup>+</sup>) T-lineage ALL.<sup>144,145</sup> We recently evaluated the influence of myeloid antigen expression on treatment outcome in a large cohort of children with newly diagnosed ALL enrolled on risk-adjusted CCG studies.<sup>146</sup> Patients were classified as My<sup>-</sup> or My<sup>+</sup> T-lineage, according to expression of CD7, CD13, and CD33. Patients with My<sup>+</sup> T-lineage ALL were more likely than patients with My<sup>-</sup> T-lineage ALL to show favorable presenting features, but induction outcome and EFS outcome were similar for patients with My<sup>+</sup> and My<sup>-</sup> T-lineage ALL, with 4-year EFS of 72.7% (SD = 7.1%) and 70.1% (SD = 5.7%), respectively ( $P = .49$ ; Fig 3). These results show that regardless of treatment intensity, mixed myeloid-lymphoid phenotype was not an adverse prognostic factor for childhood T-lineage ALL.

#### IMPEDIMENTS TO EFFECTIVE TREATMENT

**Drug resistance.** Despite improvements in overall survival, relapse in the bone marrow, CNS, and other sites remains a significant problem for high-risk patients. Pieters et al<sup>147</sup> showed that patients with T-lineage ALL were particularly resistant to prednisone (PRED), daunorubicin, cytarabine, mafosfamide, and L-asparaginase, but wide ranges of resistance levels were observed within each immunophenotypic group.

For all patients, the probability of continuous complete remission decreased with increasing resistance to PRED. In a later study, these investigators reported that patients with T-lineage ALL were more resistant to a host of drugs including those mentioned above as well as teniposide, ifosfamide, vincristine, vindesine, and dexamethasone.<sup>148</sup> Lauer et al<sup>149</sup> found that a regimen of intensive rotating drug pairs was effective for prevention of drug resistance in high-risk patients with B-lineage, but not T-lineage ALL, again suggesting that immunophenotype plays a role in drug sensitivity. Others have attributed methotrexate (MTX) resistance in patients with T-lineage ALL to a decreased formation of MTX-polyglutamates, which is a determinant of toxicity.<sup>150,151</sup> Resistance to glucocorticoids is thought to be caused by low glucocorticoid receptor (GR) levels. However, the relationship between GR and outcome within the T-lineage immunophenotype is unclear. Qudus et al<sup>152</sup> reported that leukemic cell GR level did not predict outcome within the T-lineage group, whereas Costlow et al<sup>153</sup> reported that lower GR levels were correlated with unfavorable presenting features including T-lineage. Finally, although multidrug resistance is thought to be mediated by overexpression of P-glycoprotein, the product of the multidrug resistance gene MDR-1,<sup>154</sup> the specific significance of this phenomenon in T-lineage ALL has not been determined.

#### INNOVATIVE TREATMENT STRATEGIES FOR T-LINEAGE ALL

Current strategies for improving treatment of children with ALL have been aimed at maximizing efficacy of treatment according to risk. Reliable and accurate methods for predicting prognosis are required to achieve adequate treatment with the least intensive regimens. Identification of biological and clinical




Fig 3. Myeloid antigen expression in T-lineage ALL is not associated with poor EFS. EFS for My<sup>+</sup> TL and My<sup>-</sup> TL patients treated on the 1800 series of CCG protocols are shown. EFS values at designated points in follow-up are given in the text.

prognostic factors, as discussed above, has aided in stratifying patients according to risk. However, additional methods are required for identifying and more effectively treating subgroups of high-risk patients who are most likely to relapse despite intensive therapy.

Seventy-five percent of children with T-lineage ALL on CCG protocols fit within the NCI high-risk category based on presenting age and WBC count.<sup>134</sup> Patients with T-lineage ALL with standard risk represent less than 4% of patients with ALL and less than 6% of all standard-risk patients. Treatment of patients who have relapsed generally has consisted of intensive chemotherapy to achieve a second remission and subsequent use of either nonablative chemotherapy or ablative radiochemotherapy followed by bone marrow transplantation (BMT), and recurrence of leukemia is the major obstacle to the success of either approach. Intensification of cytotoxic therapy using conventional drugs will likely cause overlapping toxicities and may result in delays which may erode the intensity of therapy. Overall, the outcome for patients with relapsed T-lineage ALL is dismal because only a very small fraction can be saved with high-dose radiochemotherapy followed by BMT. Consequently, the development of new potent antileukemia drugs and the design of combinative treatment protocols using these new agents have emerged as exceptional focal points for research in modern therapy of relapsed T-lineage ALL.

*Immunotoxins and other targeted biotherapeutics.* Immunotoxins (MoAb-toxin conjugates) are a new class of immunopharmacologic agents that shows considerable promise for more effective treatment of T-lineage ALL. A vast number of MoAbs have been developed with the intent of specifically targeting cytotoxic agents to leukemia cells while limiting the deleterious effects on normal tissues. Immunoconjugates containing toxins such as pokeweed antiviral protein, ricin, Pseudomonas endotoxin, and diphtheria toxin directed against T-lineage-specific surface antigens have been developed for use as systemic therapy of T-lineage ALL.<sup>155-158</sup>

Murphy et al<sup>159</sup> as well as Kreitman et al<sup>160</sup> have pioneered the use of genetic engineering to redirect the lethal action of diphtheria toxin towards effective targeting of growth factor receptors on leukemic cells. In one example, researchers have developed a recombinant fusion toxin, DAB486IL-2, in which the native receptor binding domain of diphtheria toxin has been replaced with interleukin-2.<sup>161</sup>

*Deoxyguanosine analogs.* Another new and promising treatment program for T-lineage ALL is based on the potent antileukemia activity of deoxyguanosine analogs. The accumulation and the resulting toxicity of dGTP in T lymphocytes was first described in patients with a genetic deficiency for the enzyme purine nucleoside phosphorylase (PNP).<sup>162,163</sup> This observation led to the search for means by which cytotoxic levels of dGTP could be achieved in T-lineage leukemias. An analog of deoxyguanosine, Ara-G (9-β-D-arabinofuranosylguanine) accumulates in T cells and acts as a poor substrate for endogenous PNP, but is efficiently phosphorylated by deoxycytidine kinase<sup>164,165</sup>; in vitro studies have shown that Ara-G is selectively cytotoxic for T-cell lines and T-lineage leukemic cells.<sup>166,167</sup>

Recently, a water soluble pro-drug derivative of Ara-G, known as compound 506U/C-506 (2-amino-6-methoxypurine arabinoside), was developed for in vivo therapeutic applica-

tions.<sup>168</sup> Preliminary results of a Phase I trial of C-506 in adult T-cell malignancies suggested that daily infusion of C-506 could achieve and maintain cytotoxic levels of Ara-GTP.<sup>169</sup> These data indicate that C-506 warrants investigation as a new therapeutic drug for treatment of pediatric T-lineage ALL.

## CONCLUSIONS

The adverse risk previously associated with T-lineage ALL in children has progressively been surmounted by intensive chemotherapeutic regimens. Still, approximately 20% to 25% of children with T-lineage ALL continue to fail therapy. Further augmentation of the currently used intensive chemotherapeutic regimens may not be warranted because of the likelihood of significant adverse effects. Thus, the current challenge is to apply our expanding knowledge of biological regulation in leukemic cells to the development of novel biologic therapeutics, particularly those that specifically target leukemic cells. Such agents could theoretically be used either to trigger cell killing directly or to alter the leukemic cell's response to radiation or chemotherapeutics. Finally, the identification of prognostically distinct patient subgroups may lead to tailored and risk-adjusted therapies for children with T-lineage ALL. Use of these various strategies, singly and in combination, should allow further improvements in outcome for patients with ALL who remain at risk for treatment failure.

## REFERENCES

1. Poplack D: Acute lymphoblastic leukemia, in Pizzo P, Poplack D (eds): Principles and Practice of Pediatric Oncology (ed 2). Philadelphia, PA, Lippincott, 1993, p 431
2. Greaves MF: Differentiation-linked leukemogenesis in lymphocytes. *Science* 234:697, 1986
3. Smith LJ, Curtis JE, Messner HA, Senn JS, Furthmayr H, McCulloch EA: Lineage infidelity in acute leukemia. *Blood* 61:1138, 1983
4. Crist WM, Shuster JJ, Falletta J, Pullen DJ, Berard CW, Vietti TJ, Alvarado CS, Roper MA, Prasthofer E, Grossi CE: Clinical features and outcome in childhood T-cell leukemia-lymphoma according to stage of thymocyte differentiation: A Pediatric Oncology Group study. *Blood* 72:1891, 1988
5. Borowitz MJ, Dowell BL, Boyett JM, Pullen DJ, Crist WM, Qudus FM, Falletta JM, Metzgar RS: Clinicopathologic aspects of E rosette negative T cell acute lymphocytic leukemia: A Pediatric Oncology Group study. *J Clin Oncol* 4:170, 1986
6. Uckun F, Reaman G, Steinherz P, Arthur D, Sather H, Trigg M, Tubergen D, Gaynon P: Improved outcome for children with T-lineage acute lymphoblastic leukemia after contemporary chemotherapy: A children's cancer group study. *Leuk Lymphoma* 24:57, 1996
7. Pullen DJ, Sullivan MP, Falletta JM, Boyett JM, Humphrey GB, Starling KA, Land VJ, Dymont PG, Vats T, Duncan MH: Modified LSA2-L2 treatment in 53 children with E-rosette-positive T-cell leukemia: Results and prognostic factors (a Pediatric Oncology Group study). *Blood* 60:1159, 1982
8. Shuster JJ, Falletta JM, Pullen DJ, Crist WM, Humphrey GB, Dowell BL, Wharam MD, Borowitz M: Prognostic factors in childhood T-cell acute lymphoblastic leukemia: A Pediatric Oncology Group study. *Blood* 75:166, 1990
9. Dowell BL, Borowitz MJ, Boyett JM, Pullen DJ, Crist WM, Qudus FF, Russell EC, Falletta JM, Metzgar RS: Immunologic and clinicopathologic features of common acute lymphoblastic leukemia antigen-positive childhood T-cell leukemia. A Pediatric Oncology Group study. *Cancer* 59:2020, 1987

10. Garand R, Vannier JP, Bene MC, Faure G, Favre M, Bernard A: Comparison of outcome, clinical, laboratory, and immunological features in 164 children and adults with T-ALL. the Groupe D'Etude Immunologique Des Leucemies. *Leukemia* 4:739, 1990
11. Clavell LA, Gelber RD, Cohen HJ, Hitchcock Bryan S, Cassady JR, Tarbell NJ, Blattner SR, Tantravahi R, Leavitt P, Sallan SE: Four-agent induction and intensive asparaginase therapy for treatment of childhood acute lymphoblastic leukemia. *N Engl J Med* 315:657, 1986
12. Gaynon P, Steinherz P, Bleyer WA, Ablin A, Albo V, Finkelstein J, Grossman N, Littman P, Novak L, Pyesmany A, Reaman G, Sather H, Hammond D: Intensive therapy for children with acute lymphoblastic leukemia and unfavorable presenting features. *Lancet* 2:921, 1988
13. Steinherz PG, Gaynon P, Miller DR, Reaman G, Bleyer A, Finklestein J, Evans RG, Meyers P, Steinherz LJ, Sather H, Hammond D: Improved disease-free survival of children with acute lymphoblastic leukemia at high risk for early relapse with the New York regimen—A new intensive therapy protocol: A report from the Children's Cancer Study Group. *J Clin Oncol* 4:744, 1986
14. Schorin MA, Blattner S, Gelber RD, Tarbell NJ, Donnelly M, Dalton V, Cohen HJ, Sallan SE: Treatment of childhood acute lymphoblastic leukemia: Results of Dana-Farber Cancer Institute/Children's Hospital Acute Lymphoblastic Leukemia Consortium Protocol 85-01. *J Clin Oncol* 12:740, 1994
15. Blethen SL, Allen DB, Graves D, August G, Moshang T, Rosenfeld R: Safety of recombinant deoxyribonucleic acid-derived growth hormone: The National Cooperative Growth Study experience. *J Clin Endocrinol Metab* 81:1704, 1996
16. Lin YW, Kubota M, Wakazono Y, Hirota H, Okuda A, Bessho R, Usami I, Kataoka A, Yamanaka S, Akiyama Y, Furusho K: Normal mutation frequencies of somatic cells in patients receiving growth hormone therapy. *Mutat Res* 362:97, 1996
17. Rapaport R, Oberfield SE, Robison L, Salisbury S, David R, Rao J, Redmond GP: Relationship of growth hormone deficiency and leukemia. *J Pediatr* 126:759, 1995
18. Linet MS, Hatch EE, Kleinerman RA, Robison LL, Kaune WT, Friedman DR, Severson RK, Haines SM, Hartsock CT, Niwa S, Wacholder S, Tarone RE: Residential exposure to magnetic fields and acute lymphoblastic leukemia in children. *N Engl J Med* 337:1, 1997
19. Alexander FE: Space-time clustering of childhood acute lymphoblastic leukaemia: Indirect evidence for a transmissible agent. *Br J Cancer* 65:589, 1992
20. Alexander FE: Viruses, clusters and clustering of childhood leukemia: A new perspective? *Eur J Cancer* 29A:1424, 1993
21. Kinlen L: Evidence for an infective cause of childhood leukaemia: Comparison of a Scottish New Town with nuclear reprocessing sites in Britain. *Lancet* 2:1323, 1988
22. Petridou E, Revinthi K, Alexander FE, Haidas S, Kolioukas D, Kosmidis H, Piperopoulou F, Tzortzotou F, Trichopoulos D: Space-time clustering of childhood leukaemia in Greece: Evidence supporting a viral aetiology. *Br J Cancer* 73:1278, 1996
23. van Steensel Moll HA, Valkenburg HA, van Zanen GE: Childhood leukemia and parental occupation: A register-based case-control study. *Am J Epidemiol* 121:216, 1985
24. Wachsman W, Golde D, Chen I: HTLV and human leukemia: Perspectives. *Semin Hematol* 23:245, 1986
25. Williams D, Ragab A, McDougal J: HTLV-I antibodies in childhood leukemia. *JAMA* 253:2496, 1985
26. Lin KH, Su IJ, Chen RL, Lin DT, Tien HF, Chen BW, Lin KS: Peripheral T-cell lymphoma in childhood: A report of five cases in Taiwan. *Med Pediatr Oncol* 23:26, 1994
27. Toledano SR, Lange BJ: Ataxia-telangiectasia and acute lymphoblastic leukemia. *Cancer* 45:1675, 1980
28. Aurias A, Dutrillaux B, Buriot D, Lejeune J: High frequencies of inversions and translocations of chromosomes 7 and 14 in ataxia telangiectasia. *Mutat Res* 69:369, 1980
29. Heppell A, Butterworth SV, Hollis RJ, Kennaugh AA, Beatty DW, Taylor AM: Breakage of the T cell receptor alpha chain locus in non malignant clones from patients with ataxia telangiectasia. *Hum Genet* 79:360, 1988
30. Russo G, Isobe M, Pegoraro L, Finan J, Nowell PC, Croce CM: Molecular analysis of a t(7;14)(Q35;Q32) chromosome translocation in a T cell leukemia of a patient with ataxia telangiectasia. *Cell* 53:137, 1988
31. Hollis RJ, Kennaugh AA, Butterworth SV, Taylor AM: Growth of large chromosomally abnormal T cell clones in ataxiatelangiectasia patients is associated with translocation at 14q11: A model for other T cell neoplasia. *Hum Genet* 76:389, 1987
32. Champlin R, Gale RP: Acute lymphoblastic leukemia: Recent advances in biology and therapy [see comments]. *Blood* 73:2051, 1989
33. Raimondi SC, Behm FG, Roberson PK, Pui CH, Rivera GK, Murphy SB, Williams DL: Cytogenetics of childhood T-cell leukemia. *Blood* 72:1560, 1988
34. Kagan J, Finger LR, Letofsky J, Finan J, Nowell PC, Croce CM: Clustering of breakpoints on chromosome 10 in acute T-cell leukemias with the t(10;14) chromosome translocation. *Proc Natl Acad Sci USA* 86:4161, 1989
35. Zutter M, Hockett RD, Roberts CW, McGuire EA, Bloomstone J, Morton CC, Deaven LL, Crist WM, Carroll AJ, Korsmeyer SJ: The t(10;14)(q24;q11) of T-cell acute lymphoblastic leukemia juxtaposes the delta T-cell receptor with TCL3, a conserved and activated locus at 10q24. *Proc Natl Acad Sci USA* 87:3161, 1990
36. Begley CG, Aplan PD, Davey MP, Nakahara K, Tchorz K, Kurtzberg J, Hershfield MS, Haynes BF, Cohen DI, Waldmann TA, Kirsch IR: Chromosomal translocation in a human leukemic stem-cell line disrupts the T-cell antigen receptor delta-chain diversity region and results in a previously unreported fusion transcript. *Proc Natl Acad Sci USA* 86:2031, 1989
37. Boehm T, Baer R, Lavenir I, Forster A, Waters JJ, Nacheva E, Rabbitts TH: The mechanism of chromosomal translocation t(11;14) involving the T-cell receptor C delta locus on human chromosome 14q11 and a transcribed region of chromosome 11p15. *EMBO J* 7:385, 1988
38. Cheng JT, Yang CY, Hernandez J, Embrey J, Baer R: The chromosome translocation (11;14)(p13;q11) associated with T cell acute leukemia. Asymmetric diversification of the translocational junctions. *J Exp Med* 171:489, 1990
39. Garcia IS, Kaneko Y, Gonzalez Sarmiento R, Campbell K, White L, Boehm T, Rabbitts TH: A study of chromosome 11p13 translocations involving TCR beta and TCR delta in human T cell leukaemia. *Oncogene* 6:577, 1991
40. Carroll AJ, Crist WM, Link MP, Amylon MD, Pullen DJ, Ragab AH, Buchanan GR, Wimmer RS, Vietti TJ: The t(1;14)(p34;q11) is nonrandom and restricted to T-cell acute lymphoblastic leukemia: A Pediatric Oncology Group study. *Blood* 76:1220, 1990
41. Finger LR, Kagan J, Christopher G, Kurtzberg J, Hershfield MS, Nowell PC, Croce CM: Involvement of the TCL5 gene on human chromosome 1 in T-cell leukemia and melanoma. *Proc Natl Acad Sci USA* 86:5039, 1989
42. Chen Q, Yang CY, Tsan JT, Xia Y, Ragab AH, Peiper SC, Carroll A, Baer R: Coding sequences of the tal-1 gene are disrupted by chromosome translocation in human T cell leukemia. *J Exp Med* 172:1403, 1990
43. Begley CG, Aplan PD, Denning SM, Haynes BF, Waldmann TA, Kirsch IR: The gene SCL is expressed during early hematopoiesis and encodes a differentiation-related DNA-binding motif. *Proc Natl Acad Sci USA* 86:10128, 1989
44. Aplan PD, Lombardi DP, Ginsberg AM, Cossman J, Bertness



- VL, Kirsch IR: Disruption of the human SCL locus by "illegitimate" V-(D)-J recombinase activity. *Science* 250:1426, 1990
45. Aplan PD, Lombardi DP, Reaman GH, Sather HN, Hammond GD, Kirsch IR: Involvement of the putative hematopoietic transcription factor SCL in T-cell acute lymphoblastic leukemia. *Blood* 79:1327, 1992
  46. Bash RO, Crist WM, Shuster JJ, Link MP, Amylon M, Pullen J, Carroll AJ, Buchanan GR, Smith RG, Baer R: Clinical features and outcome of T-cell acute lymphoblastic leukemia in childhood with respect to alterations at the TAL1 locus: A Pediatric Oncology Group study. *Blood* 81:2110, 1993
  47. Porcher C, Swat W, Rockwell K, Fujiwara Y, Alt FW, Orkin SH: The T cell leukemia oncoprotein SCL/tal-1 is essential for development of all hematopoietic lineages. *Cell* 86:47, 1996
  48. Pulford K, Lecointe N, Leroy Viard K, Jones M, Mathieu Mahul D, Mason DY: Expression of TAL-1 proteins in human tissues. *Blood* 85:675, 1995
  49. Elwood NJ, Cook WD, Metcalf D, Begley CG: SCL, the gene implicated in human T-cell leukaemia, is oncogenic in a murine T-lymphocyte cell line. *Oncogene* 8:3093, 1993
  50. Dube ID, Raimondi SC, Pi D, Kalousek DK: A new translocation, t(10;14)(q24;q11), in T cell neoplasia. *Blood* 67:1181, 1986
  51. Kennedy MA, Gonzalez Sarmiento R, Kees UR, Lampert F, Dear N, Boehm T, Rabbitts TH: HOX11, a homeobox-containing T-cell oncogene on human chromosome 10q24. *Proc Natl Acad Sci USA* 88:8900, 1991
  52. Hatano M, Roberts CW, Minden M, Crist WM, Korsmeyer SJ: Deregulation of a homeobox gene, HOX11, by the t(10;14) in T cell leukemia. *Science* 253:79, 1991
  53. Dube ID, Kamel Reid S, Yuan CC, Lu M, Wu X, Corpus G, Raimondi SC, Crist WM, Carroll AJ, Minowada J, Baker JB: A novel human homeobox gene lies at the chromosome 10 breakpoint in lymphoid neoplasias with chromosomal translocation t(10;14). *Blood* 78:2996, 1991
  54. Dear TN, Sanchez Garcia I, Rabbitts TH: The HOX11 gene encodes a DNA-binding nuclear transcription factor belonging to a distinct family of homeobox genes. *Proc Natl Acad Sci USA* 90:4431, 1993
  55. Yamamoto H, Hatano M, Iitsuka Y, Mahyar NS, Yamamoto M, Tokuhisa T: Two forms of Hox11, a T cell leukemia oncogene, are expressed in fetal spleen but not in primary lymphocytes. *Mol Immunol* 32:1177, 1995
  56. Roberts CW, Shutter JR, Korsmeyer SJ: Hox11 controls the genesis of the spleen. *Nature* 368:747, 1994
  57. Salvati PD, Ranford PR, Ford J, Kees UR: HOX11 expression in pediatric acute lymphoblastic leukemia is associated with T-cell phenotype. *Oncogene* 11:1333, 1995
  58. Williams DL, Look AT, Melvin SL, Roberson PK, Dahl G, Flake T, Stass S: New chromosomal translocations correlate with specific immunophenotypes of childhood acute lymphoblastic leukemia. *Cell* 36:101, 1984
  59. Ribeiro RC, Raimondi SC, Behm FG, Cherrie J, Crist WM, Pui CH: Clinical and biologic features of childhood T-cell leukemia with the t(11;14). *Blood* 78:466, 1991
  60. Zalcborg IQ, Silva ML, Abdelhay E, Tabak DG, Ornellas MH, Simoes FV, Pucheri W, Ribeiro R, Seunaz HN: Translocation 11;14 in three children with acute lymphoblastic leukemia of T-cell origin. *Cancer Genet Cytogenet* 84:32, 1995
  61. McGuire EA, Hockett RD, Pollock KM, Bartholdi MF, O'Brien SJ, Korsmeyer SJ: The t(11;14)(p15;q11) in a T-cell acute lymphoblastic leukemia cell line activates multiple transcripts, including Ttg-1, a gene encoding a potential zinc finger protein. *Mol Cell Biol* 9:2124, 1989
  62. Royer Pokora B, Fleischer B, Ragg S, Loos U, Williams D: Molecular cloning of the translocation breakpoint in T-ALL 11;14 (p13;q11): Genomic map of TCR alpha and delta region on chromosome 14q11 and long-range map of region 11p13. *Hum Genet* 82:264, 1989
  63. Boehm T, Foroni L, Kaneko Y, Perutz MF, Rabbitts TH: The rhombotin family of cysteine-rich LIM-domain oncogenes: Distinct members are involved in T-cell translocations to human chromosomes 11p15 and 11p13. *Proc Natl Acad Sci USA* 88:4367, 1991
  64. Royer Pokora B, Loos U, Ludwig WD: TTG-2, a new gene encoding a cysteine-rich protein with the lim motif, is overexpressed in acute T-cell leukaemia with the t(11;14)(P13;Q11). *Oncogene* 6:1887, 1987
  65. Wilkinson DA, Neale GA, Mao S, Naeve CW, Goorha RM: Elf-2, a rhombotin-2 binding ets transcription factor: Discovery and potential role in T cell leukemia. *Leukemia* 11:86, 1997
  66. Raimondi SC, Pui CH, Behm FG, Williams DL: 7q32-q36 translocations in childhood T cell leukemia: Cytogenetic evidence for involvement of the T cell receptor beta-chain gene. *Blood* 69:131, 1987
  67. Cleary ML, Mellentin JD, Spies J, Smith SD: Chromosomal translocation involving the beta T cell receptor gene in acute leukemia. *J Exp Med* 167:682, 1988
  68. Mellentin JD, Smith SD, Cleary ML: lyl-1, a novel gene altered by chromosomal translocation in T cell leukemia, codes for a protein with a helix-loop-helix DNA binding motif. *Cell* 58:77, 1989
  69. Ellisen LW, Bird J, West DC, Soreng AL, Reynolds TC, Smith SD, Sklar J: TAN-1, the human homolog of the Drosophila notch gene, is broken by chromosomal translocations in T lymphoblastic neoplasms. *Cell* 66:649, 1991
  70. Burnett RC, David JC, Harden AM, Le Beau MM, Rowley JD, Diaz MO: The Lck gene is involved in the t(1;7)(P34;Q34) in the T-cell acute lymphoblastic leukemia derived cell line, Hsb-2. *Gene Chromosom Cancer* 3:461, 1991
  71. Tycko B, Smith SD, Sklar J: Chromosomal translocations joining lck and trcb loci in human T cell leukemia. *J Exp Med* 174:867, 1991
  72. Abraham KM, Levin SD, Marth JD, Forbush KA, Perlmutter RM: Thymic tumorigenesis induced by overexpression of P56lck. *Proc Natl Acad Sci USA* 88:3977, 1991
  73. Wildin RS, Garvin AM, Pawar S, Lewis DB, Abraham KM, Forbush KA, Ziegler SF, Allen JM, Perlmutter RM: Developmental regulation of lck gene expression in T lymphocytes. *J Exp Med* 173:383, 1991
  74. Finver SN, Nishikura K, Finger LR, Haluska FG, Finan J, Nowell PC, Croce CM: Sequence analysis of the MYC oncogene involved in the t(8;14)(q24;q11) chromosome translocation in a human leukemia T-cell line indicates that putative regulatory regions are not altered. *Proc Natl Acad Sci USA* 85:3052, 1988
  75. Erikson J, Williams DL, Finan J, Nowell PC, Croce CM: Locus of the alpha-chain of the T-cell receptor is split by chromosome translocation in T-cell leukemias. *Science* 229:784, 1985
  76. Finger LR, Huebner K, Cannizzaro LA, McLeod K, Nowell PC, Croce CM: Chromosomal translocation in T-cell leukemia line HUT 78 results in a MYC fusion transcript. *Proc Natl Acad Sci USA* 85:9158, 1988
  77. Elledge SJ: Cell cycle checkpoints: Preventing an identity crisis. *Science* 274:1664, 1996
  78. Boise LH, Thompson CB: Hierarchical control of lymphocyte survival. *Science* 274:67, 1996
  79. Stillman B: Cell cycle control of DNA replication. *Science* 274:1659, 1996
  80. Lowsky R, DeCoteau JF, Reitmair AH, Ichinohasama R, Dong WF, Xu Y, Mak TW, Kadin ME, Minden MD: Defects of the mismatch repair gene MSH2 are implicated in the development of murine and human lymphoblastic lymphomas and are associated with the aberrant expression of rhombotin-2 (Lmo-2) and Tal-1 (SCL). *Blood* 89:2276, 1997

81. Weinberg RA: Tumor Suppressor Genes. *Science* 254:1138, 1991
82. Hermanson M, Liu Y, Zabarovsky E, Grandt D, Gahrton G, Juliusson G, Rasool M, Wu X, Buys C, Yankovsky N, et al: Chromosome 13q14 deletions in lymphoid malignancies (meeting abstract). *Proc Am Assoc Cancer Res* 36:104, 1995, (abstr 3095)
83. Zhou M, Zaki SR, Ragab AH, Findley HW: Frequent alteration of Rb tumor-suppressor gene in childhood acute lymphoblastic leukemia (meeting abstract). *Proc Am Assoc Cancer Res* 34:104, 1993 (abstr 616)
84. Harper JW, Adami GR, Wei N, Keyomarsi K, Elledge SJ: The p21 Cdk-interacting protein Cip1 is a potent inhibitor of G1 cyclin-dependent kinases. *Cell* 75:805, 1993
85. Hannon GJ, Beach D: p15<sup>ink4b</sup> is a potential effector of Tgf-beta-induced cell cycle arrest [see comments]. *Nature* 371:257, 1994
86. Guan KL, Jenkins CW, Li Y, Nichols MA, Wu X, CL OK, Matera AG, Xiong Y: Growth suppression by p18, a p16<sup>INK4</sup>/MTS1- and p14<sup>INK4B</sup>/MTS2-related CDK6 inhibitor, correlates with wild-type pRb function. *Genes Dev* 8:2939, 1994
87. Chan FK, Zhang J, Cheng L, Shapiro DN, Winoto A: Identification of human and mouse p19, a novel CDK4 and CDK6 inhibitor with homology to p16<sup>ink4</sup>. *Mol Cell Biol* 15:2682, 1995
88. Matsuoka S, Edwards MC, Bai C, Parker S, Zhang P, Baldini A, Harper JW, Elledge SJ: p57<sup>kip2</sup>, a structurally distinct member of the p21<sup>cip1</sup> Cdk inhibitor family, is a candidate tumor suppressor gene. *Genes Dev* 9:650, 1995
89. Polyak K, Kato JY, Solomon MJ, Sherr CJ, Massague J, Roberts JM, Koff A: p27<sup>kip1</sup>, a cyclin-Cdk inhibitor, links transforming growth factor-beta and contact inhibition to cell cycle arrest. *Genes Dev* 8:9, 1994
90. Rasool O, Heyman M, Brandtner LB, Liu Y, Grandt D, Soderheall S, Einhorn S: p15<sup>ink4b</sup> and p16<sup>ink4</sup> gene inactivation in acute lymphocytic leukemia. *Blood* 85:3431, 1995
91. Ohnishi H, Kawamura M, Ida K, Sheng XM, Hanada R, Nobori T, Yamamori S, Hayashi Y: Homozygous deletions of p16/MTS1 gene are frequent but mutations are infrequent in childhood T-cell acute lymphoblastic leukemia. *Blood* 86:1269, 1995
92. Okuda T, Shurtleff SA, Valentine MB, Raimondi SC, Head DR, Behm F, Curcio Brint AM, Liu Q, Pui CH, Sherr CJ, Beach D, Look AT, Downing JR: Frequent deletion of p16<sup>INK4a</sup>/MTS1 and p15<sup>INK4b</sup>/MTS2 in pediatric acute lymphoblastic leukemia. *Blood* 85:2321, 1995
93. Guidal Giroux C, Gerard B, Cave H, Duval M, Rohrlrich P, Elion J, Vilmer E, Grandchamp B: Deletion mapping indicates that MTS1 is the target of frequent deletions at chromosome 9p21 in paediatric acute lymphoblastic leukaemias. *Br J Haematol* 92:410, 1996
94. Fizzotti M, Cimino G, Pisegna S, Alimena G, Quartarone C, Mandelli F, Pelicci PG, Lo Coco F: Detection of homozygous deletions of the cyclin-dependent kinase 4 inhibitor (p16) gene in acute lymphoblastic leukemia and association with adverse prognostic features. *Blood* 85:2685, 1995
95. Batova A, Diccianni MB, Yu JC, Nobori T, Link MP, Pullen J, Yu AL: Frequent and selective methylation of p15 and deletion of both p15 and p16 in T-cell acute lymphoblastic leukemia. *Cancer Res* 57:832, 1997
96. Chilcote RR, Brown E, Rowley JD: Lymphoblastic leukemia with lymphomatous features associated with abnormalities of the short arm of chromosome 9. *N Engl J Med* 313:286, 1985
97. Uckun FM, Gajl Peczkalska KJ, Provisor AJ, Heerema NA: Immunophenotype-karyotype associations in human acute lymphoblastic leukemia. *Blood* 73:271, 1989
98. Bloomfield CD, Secker Walker LM, Goldman AI, Van Den Berghe H, de la Chapelle A, Ruutu T, Alimena G, Garson OM, Golomb HM, Rowley JD, Kaneko Y, Whang-Peng J, Prigogina E, Philip P, Sandberg AA, Lawler SD, Mittleman F: Six-year follow-up of the clinical significance of karyotype in acute lymphoblastic leukemia. *Cancer Genet Cytogenet* 40:171, 1989
99. Sherr CJ: Cancer cell cycles. *Science* 274:1672, 1996
100. Harvey M, Vogel H, Morris D, Bradley A, Bernstein A, Donehower LA: A mutant p53 transgene accelerates tumour development in heterozygous but not nullizygous p53-deficient mice. *Nat Genet* 9:305, 1995
101. Yeargin J, Cheng J, Haas M: Role of the p53 tumor suppressor gene in the pathogenesis and in the suppression of acute lymphoblastic T-cell leukemia. *Leukemia* 6 Suppl 3:85s, 1992
102. Hsiao M, Low J, Dorn E, Ku D, Pattengale P, Yeargin J, Haas M: Gain-of-function mutations of the p53 gene induce lymphohematopoietic metastatic potential and tissue invasiveness. *Am J Pathol* 145:702, 1994
103. Savitsky K, Bar Shira A, Gilad S, Rotman G, Ziv Y, Vanagaite L, Tagle DA, Smith S, Uziel T, Sfez S, et al: A single ataxia telangiectasia gene with a product similar to PI-3 kinase [see comments]. *Science* 268:1749, 1995
104. Painter RB, Young BR: Radiosensitivity in ataxia-telangiectasia: A new explanation. *Proc Natl Acad Sci USA* 77:7315, 1980
105. Xu Y, Baltimore D: Dual roles of ATM in the cellular response to radiation and in cell growth control [see comments]. *Genes Dev* 10:2401, 1996
106. Xu Y, Ashley T, Brainerd EE, Bronson RT, Meyn MS, Baltimore D: Targeted disruption of ATM leads to growth retardation, chromosomal fragmentation during meiosis, immune defects, and thymic lymphoma [see comments]. *Genes Dev* 10:2411, 1996
107. Barlow C, Hirotsune S, Paylor R, Liyanage M, Eckhaus M, Collins F, Shiloh Y, Crawley JN, Ried T, Tagle D, Wynshaw Boris A: Atm-deficient mice: A paradigm of ataxia telangiectasia. *Cell* 86:159, 1996
108. Taylor A, Metalfe J, Thick J, Mak Y: Leukemia and lymphoma in ataxia telangiectasia. *Blood* 87:423, 1996
109. Spector B, Filipovich A, Perry G, Kersey K: Epidemiology of cancer in ataxia telangiectasia, in Bridges B, Harned D (eds): *A Cellular and Molecular Link Between Cancer, Neuropathology, and Immune Deficiency*. Chichester, UK, Wiley, 1982, p1
110. Collyn dHooghe M, Galieue Zouitina S, Szymiczek D, Lantoine D, Quief S, Loucheux Lefebvre MH, Kerckaert JP: Quantitative and qualitative variation of ETS-1 transcripts in hematologic malignancies. *Leukemia* 7:1777, 1993
111. Georgopoulos K, Moore DD, Derfler B: Ikaros, an early lymphoid-specific transcription factor and a putative mediator for T cell commitment. *Science* 258:808, 1992
112. Winandy S, Wu P, Georgopoulos K: A dominant mutation in the Ikaros gene leads to rapid development of leukemia and lymphoma. *Cell* 83:289, 1995
113. Debatin KM, Goldman CK, Waldmann TA, Krammer PH: APO-1-induced apoptosis of leukemia cells from patients with adult T-cell leukemia. *Blood* 81:2972, 1993
114. Debatin KM, Krammer PH: Resistance to APO-1 (CD95) induced apoptosis in T-ALL is determined by a BCL-2 independent anti-apoptotic program. *Leukemia* 9:815, 1995
115. Lucking Famira KM, Daniel PT, Moller P, Krammer PH, Debatin KM: APO-1 (CD95) mediated apoptosis in human T-ALL engrafted in SCID mice. *Leukemia* 8:1825, 1994
116. Cory S: Regulation of lymphocyte survival by the bcl-2 gene family. *Annu Rev Immunol* 13:513, 1995
117. Conroy LA, Alexander DR: The role of intracellular signalling pathways regulating thymocyte and leukemic T cell apoptosis. *Leukemia* 10:1422, 1996
118. Oltvai ZN, Milliman CL, Korsmeyer SJ: Bcl-2 heterodimerizes in vivo with a conserved homolog, Bax, that accelerates programmed cell death. *Cell* 74:609, 1993
119. Kitada S, Krajewski S, Miyashita T, Krajewska M, Reed JC: Gamma-radiation induces upregulation of Bax protein and apoptosis in radiosensitive cells in vivo. *Oncogene* 12:187, 1996

120. Uckun FM, Yang Z, Sather HN, Steinherz P, Nachman J, Bostrom B, Crotty L, Sarquis M, Ek O, Zren T, Tubergen D, Reaman G, Gaynon P: Cellular expression of anti-apoptotic BCL-2 oncoprotein in newly diagnosed childhood acute lymphoblastic leukemia. *Blood* 89:3769, 1997
121. Sen L, Borella L: Clinical importance of lymphoblasts with T markers in childhood acute leukemia. *N Engl J Med* 292:828, 1975
122. Borella L, Sen L: T- and B-lymphocytes and lymphoblasts in untreated acute lymphocytic leukemia. *Cancer* 34:646, 1974
123. Williams DL, Raimondi S, Rivera G, George S, Berard CW, Murphy SB: Presence of clonal chromosome abnormalities in virtually all cases of acute lymphoblastic leukemia [letter]. *N Engl J Med* 313:640, 1985
124. Pui CH, Crist WM, Look AT: Biology and clinical significance of cytogenetic abnormalities in childhood acute lymphoblastic leukemia. *Blood* 76:1449, 1990
125. Crist W, Boyett J, Pullen J, van Eys J, Vietti T: Clinical and biologic features predict poor prognosis in acute lymphoid leukemias in children and adolescents: A Pediatric Oncology Group review. *Med Pediatr Oncol* 14:135, 1986
126. Williams DL, Harber J, Murphy SB, Look AT, Kalwinsky DK, Rivera G, Melvin SL, Stass S, Dahl GV: Chromosomal translocations play a unique role in influencing prognosis in childhood acute lymphoblastic leukemia. *Blood* 68:205, 1986
127. Bloomfield CD, Goldman AI, Alimena G, Berger R, Borgstrom GH, Brandt L, Catovsky D, de la Chapelle A, Dewald GW, Garson OM, et al: Chromosomal abnormalities identify high-risk and low-risk patients with acute lymphoblastic leukemia. *Blood* 67:415, 1986
128. Look AT, Roberson PK, Williams DL, Rivera G, Bowman WP, Pui CH, Ochs J, Abromowitch M, Kalwinsky D, Dahl GV, et al: Prognostic importance of blast cell DNA content in childhood acute lymphoblastic leukemia. *Blood* 65:1079, 1985
129. Jackson JF, Boyett J, Pullen J, Brock B, Patterson R, Land V, Borowitz M, Head D, Crist W: Favorable prognosis associated with hyperdiploidy in children with acute lymphocytic leukemia correlates with extra chromosome 6: A Pediatric Oncology Group Study. *Cancer* 66:1183, 1990
130. Pui CH, Carroll AJ, Head D, Raimondi SC, Shuster JJ, Crist WM, Link MP, Borowitz MJ, Behm FG, Land VJ, et al: Near-triploid and near-tetraploid acute lymphoblastic leukemia of childhood. *Blood* 76:590, 1990
131. Zintl F, Plenert W, Malke H: Results of acute lymphoblastic leukemia therapy in childhood with a modified BFM protocol in a multicenter study in the German Democratic Republic. *Hamatol Bluttransfus* 30:471, 1987
132. Riehm H, Reiter A, Schrappe M, Berthold F, Dopfer R, Gerein V, Ludwig R, Ritter J, Stollmann B, Henze G: Corticosteroid-dependent reduction of leukocyte count in blood as a prognostic factor in acute lymphoblastic leukemia in childhood (therapy study ALL-BFM 83). *Klin Padiatr* 199:151, 1987
133. Dahl GV, Rivera GK, Look AT, Hustu HO, Kalwinsky DK, Abromowitch M, Mirro J, Ochs J, Murphy SB, Dodge RK, et al: Teniposide plus cytarabine improves outcome in childhood acute lymphoblastic leukemia presenting with a leukocyte count greater than or equal to  $100 \times 10^9/L$ . *J Clin Oncol* 5:1015, 1987
134. Smith M, Arthur D, Camitta B, Carroll AJ, Crist W, Gaynon P, Gelber R, Heerema N, Korn EL, Link M, Murphy S, Pui CH, Pullen J, Reamon G, Sallan SE, Sather H, Shuster J, Simon R, Trigg M, Tubergen D, Uckun F, Ungerleider R: Uniform approach to risk classification and treatment assignment for children with acute lymphoblastic leukemia [see comments]. *J Clin Oncol* 14:18, 1996
135. Steinherz PG, Siegel SE, Bleyer WA, Kersey J, Chard R, Jr., Coccia P, Leiken S, Lukens J, Neerhout R, Nesbit M, Miller DR, Reaman G, Sather H, Hammond D: Lymphomatous presentation of childhood acute lymphoblastic leukemia. *Cancer* 68:751, 1991
136. Henze G, Langermann HJ, Kaufmann U, Ludwig R, Schellong G, Stollmann B, Riehm H: Thymic involvement and initial white blood count in childhood acute lymphoblastic leukemia. *Am J Pediatr Hematol Oncol* 3:369, 1981
137. Pui CH, Behm FG, Singh B, Schell MJ, Williams DL, Rivera GK, Kalwinsky DK, Sandlund JT, Crist WM, Raimondi SC: Heterogeneity of presenting features and their relation to treatment outcome in 120 children with T-cell acute lymphoblastic leukemia. *Blood* 75:174, 1990
138. Schaison G, Leverger G, Bancillon A, Marty M, Olive D, Cornu G, Griselli C, Lemerle S, Harrousseau J, Bonnet M, Freycon F, Duffillot D, Demeocq F, Bateurs F, Lamagnere J, Taboureau O: Intermediate risk childhood acute lymphoblastic leukemias: Anascrine + cytosine arabinoside versus intermediate dose methotrexate for consolidation, and 6 mercaptopurine + methotrexate + vincristine versus monthly pulses for maintenance. *Hamatol Bluttransfus* 30:461, 1987
139. Gaynon PS, Bleyer WA, Steinherz PG, Finklestein JZ, Littman PS, Miller DR, Reaman GH, Sather HN, Hammond GD: Modified BFM therapy for children with previously untreated acute lymphoblastic leukemia and unfavorable prognostic features. Report of Children's Cancer Study Group Study CCG-193P. *Am J Pediatr Hematol Oncol* 10:42, 1988
140. Uckun F, Steinherz P, Sather H, Trigg M, Arthur D, Tubergen D, Gaynon P, Reaman G: CD2 antigen expression on leukemic cells as a predictor of event-free survival after chemotherapy for T-lineage acute lymphoblastic leukemia: A Children's Cancer Group Study. *Blood* 88:4288, 1996
141. Uckun FM, Gaynon P, Sensel M, Nachman J, Trigg M, Steinherz P, Bostrom B, Sather H, Reaman G: Clinical features and treatment outcome of childhood T-lineage acute lymphoblastic leukemia according to the apparent maturational stage of T-lineage leukemic blasts: A Children's Cancer Group Study. *J Clin Oncol* 15:2214, 1997
142. Wiersma SR, Ortega J, Sobel E, Weinberg KI: Clinical importance of myeloid-antigen expression in acute lymphoblastic leukemia of childhood [see comments]. *N Engl J Med* 324:800, 1991
143. Kurec AS, Belair P, Stefanu C, Barrett DM, Dubowy RL, Davey FR: Significance of aberrant immunophenotypes in childhood acute lymphoblastic leukemia. *Cancer* 67:3081, 1991
144. Pui CH, Behm FG, Singh B, Rivera GK, Schell MJ, Roberts WM, Crist WM, Mirro J Jr: Myeloid-associated antigen expression lacks prognostic value in childhood acute lymphoblastic leukemia treated with intensive multiagent chemotherapy. *Blood* 75:198, 1990
145. Bradstock KF, Kirk J, Grimsley PG, Kabral A, Hughes WG: Unusual immunophenotypes in acute leukaemias: Incidence and clinical correlations. *Br J Haematol* 72:512, 1989
146. Uckun FM, Sather HN, Gaynon P, Arthur D, Trigg M, Tubergen D, Nachman J, Steinherz P, Sensel M, Reaman G: Clinical features and treatment outcome of children with myeloid antigen positive acute lymphoblastic leukemia: A report from the Children's Cancer Group. *Blood* 90:28, 1997
147. Pieters R, Kaspers GJ, Klumper E, Veerman AJ: Clinical relevance of in vitro drug resistance testing in childhood acute lymphoblastic leukemia: The state of the art. *Med Pediatr Oncol* 22:299, 1994
148. Kaspers GJ, Pieters R, Van Zantwijk CH, Van Wering ER, Veerman AJ: Clinical and cell biological features related to cellular drug resistance of childhood acute lymphoblastic leukemia cells. *Leuk Lymphoma* 19:407, 1995
149. Lauer SJ, Camitta BM, Leventhal BG, Mahoney DH Jr, Shuster JJ, Adair S, Casper JT, Civin CI, Graham M, Kiefer G, Pullen J, Steuber CP, Kamen B: Intensive alternating drug pairs for treatment of high-risk childhood acute lymphoblastic leukemia. A Pediatric Oncology Group pilot study. *Cancer* 71:2854, 1993
150. Goker E, Lin JT, Trippett T, Elisseyeff Y, Tong WP, Niedzwiecki

D, Tan C, Steinherz P, Schweitzer BI, Bertino JR: Decreased polyglutamylation of methotrexate in acute lymphoblastic leukemia blasts in adults compared to children with this disease. *Leukemia* 7:1000, 1993

151. Barredo JC, Synold TW, Laver J, Relling MV, Pui CH, Priest DG, Evans WE: Differences in constitutive and post-methotrexate folypolyglutamate synthetase activity in B-lineage and T-lineage leukemia. *Blood* 84:564, 1994

152. Quddus FF, Leventhal BG, Boyett JM, Pullen DJ, Crist WM, Borowitz MJ: Glucocorticoid receptors in immunological subtypes of childhood acute lymphocytic leukemia cells: A Pediatric Oncology Group study. *Cancer Res* 45:6482, 1985

153. Costlow ME, Pui CH, Dahl GV: Glucocorticoid receptors in childhood acute lymphocytic leukemia. *Cancer Res* 42:4801, 1982

154. Gros P, Ben Neriah YB, Croop JM, Housman DE: Isolation and expression of a complementary DNA that confers multidrug resistance. *Nature* 323:728, 1986

155. Laurent G, Frankel AE, Hertler AA, Schlossman DM, Casellas P, Jansen FK: Treatment of leukemia patients with T101 ricin a chain immunotoxins. *Cancer Treat Res* 37:483, 1988

156. Kreitman RJ, Chaudhary VK, Waldmann TA, Hanchard B, Cranston B, FitzGerald DJ, Pastan I: Cytotoxic activities of recombinant immunotoxins composed of pseudomonas toxin or diphtheria toxin toward lymphocytes from patients with adult T-cell leukemia. *Leukemia* 7:553, 1993

157. Waurzyniak B, Schneider E, Yanishevski Y, Gunther R, Chelstrom LM, Wendorf H, Myers DE, Irvin JD, Messinger Y, Ek O, Seren T, Langlie M, Evans WE, Uckun FM: In vivo toxicity, pharmacokinetics, and antileukemic activity of TXU (anti-CD7)-pokeweed antiviral protein (PAP) immunotoxin. *Clin Cancer Res* 3:881, 1997

158. Uckun FM, Reaman GH: Immunotoxins for treatment of leukemia and lymphoma. *Leuk Lymphoma* 18:195, 1995

159. Murphy JR, Bishai W, Borowski M, Miyanohara A, Boyd J, Nagle S: Genetic construction, expression, and melanoma-selective cytotoxicity of a diphtheria toxin-related alpha-melanocyte-stimulating hormone fusion protein. *Proc Natl Acad Sci USA* 83:8258, 1986

160. Kreitman RJ, Chang CN, Hudson DV, Queen C, Bailon P,

Pastan I: Anti-Tac(Fab)-PE40, a recombinant double-chain immunotoxin which kills interleukin-2-receptor-bearing cells and induces complete remission in an in vivo tumor model. *Int J Cancer* 57:856, 1994

161. LeMaistre CF, Craig FE, Meneghetti C, McMullin B, Parker K, Reuben J, Boldt DH, Rosenblum M, Woodworth T: Phase I trial of a 90-minute infusion of the fusion toxin DAB486IL-2 in hematological cancers. *Cancer Res* 53:3930, 1993

162. Giblett ER: ADA and PNP deficiencies: How it all began. *Ann NY Acad Sci* 451:1, 1985

163. Kredich NM, Hershfield MS: Immunodeficiency diseases caused by adenosine deaminase deficiency and purine nucleoside phosphorylase deficiency, in Stanbury JB, Wyngaarden JB, Goldstein JL, Brown MS (eds): *The Metabolic Basis of Inherited Disease*. New York, NY, McGraw-Hill, 1983, p 1157

164. Ullman B, Martin DW Jr: Specific cytotoxicity of arabinosylguanine toward cultured T lymphoblasts. *J Clin Invest* 74:951, 1984

165. Verhoef V, Fridland A: Metabolic basis of arabinonucleoside selectivity for human leukemic T- and B-lymphoblasts. *Cancer Res* 45:3646, 1985

166. Hebert ME, Greenberg ML, Chaffee S, Gravatt L, Hershfield MS, Elion GB, Kurtzberg J: Pharmacologic purging of malignant T cells from human bone marrow using 9-beta-D-arabinofuranosylguanine. *Transplantation* 52:634, 1991

167. Gravatt LC, Chaffee S, Hebert ME, Halperin EC, Friedman HS, Kurtzberg J: Efficacy and toxicity of 9-beta-D-arabinofuranosylguanine (araG) as an agent to purge malignant T cells from murine bone marrow: Application to an in vivo T-leukemia model. *Leukemia* 7:1261, 1993

168. Lambe CU, Averett DR, Paff MT, Reardon JE, Wilson JG, Krenitsky TA: 2-Amino-6-methoxypurine arabinoside: An agent for T-cell malignancies. *Cancer Res* 55:3352, 1995

169. Plunkett W, Gandhi V, Nowak B, Du M, Rodriguez CO, Keating MJ: Pharmacokinetics of compound 506, a soluble prodrug of arabinosylguanine, in adult leukemias (meeting abstract). *Proc Am Assoc Cancer Res* 37:125, 1996