

REVIEW ARTICLE

DNA Methylation

By Rakesh Singal and Gordon D. Ginder

SINCE ITS FIRST recognition in 1948, the fifth base of human DNA, 5-methylcytosine (5-mC) has generated much interest and considerable controversy during attempts to understand its significance (for review, see Weissbach¹). DNA methylation in eukaryotes involves addition of a methyl group to the carbon 5 position of the cytosine ring (Fig 1). This reaction is catalyzed by DNA methyltransferase in the context of the sequence 5'-CG-3', which is also referred to as a CpG dinucleotide. It is the most common eukaryotic DNA modification and is one of the many epigenetic (alteration in gene expression without a change in nucleotide sequence) phenomena. Although extensive in plants and mammals, the absence of detectable DNA methylation in some eukaryotes such as *Drosophila*² and *Saccharomyces cerevisiae*³ has raised doubts about its significance in normal development and tissue-specific gene expression. However, recent studies showing abnormal development and embryonic lethality in transgenic mice expressing decreased but not completely absent DNA Methyltransferase (MTase) activity after DNA-MTase gene knockout⁴ lends support to a critical role for DNA methylation in developmental gene regulation. Others have proposed that the control of intragenomic parasites is the primary function of DNA methylation in mammalian cells.⁵

In this review, we will discuss the fundamental aspects of DNA methylation and its role in transcription repression, neoplasia, and transgene silencing and during development. In the interest of brevity, the role of DNA methylation in genomic imprinting and X chromosome inactivation are not discussed here, but have been reviewed recently elsewhere.⁶⁻⁸

DISTRIBUTION OF METHYLATED CYTOSINES AND CpG ISLANDS

Eukaryotic genomes are not methylated uniformly but contain methylated regions interspersed with unmethylated domains.⁹ During evolution, the dinucleotide CpG has been progressively eliminated from the genome of higher eukaryotes and is present at only 5% to 10% of its predicted frequency.¹⁰⁻¹² Cytosine methylation appears to have played a major role in this process, because most CpG sites lost represent the conversion through deamination of methylcytosines to thymines. Approximately 70% to 80% of the remaining CpG sites contain methylated cytosines in most vertebrates, including humans.^{10,12} These methylated regions are typical of the bulk chromatin that represents the late replicating DNA with its attendant histone composition and nucleosomal configuration and is relatively inaccessible to transcription factors.¹³ In contrast to the rest of the genome, smaller regions of DNA, called CpG islands, ranging from 0.5 to 5 kb and occurring on average every 100 kb, have distinctive properties. These regions are unmethylated, GC rich (60% to 70%), have a ratio of CpG to GpC of at least 0.6, and thus do not show any suppression of the frequency of the dinucleotide CpG.^{10,14} Chromatin containing CpG islands is generally heavily acetylated, lacks histone H1, and includes a nucleosome-free region.¹³ This so called open

chromatin configuration may allow, or be a consequence of, the interaction of transcription factors with gene promoters.¹⁴

The patterns of DNA methylation reflect two types of gene 5' regulatory regions in the genome. Approximately half of all genes in mouse and humans (ie, 40,000 to 50,000 genes) contain CpG islands.¹⁰ These are mainly housekeeping genes that have a broad tissue pattern of expression, but approximately 40% of genes with a tissue-restricted pattern of expression are also represented.¹⁴ Promoter region CpG islands are usually unmethylated in all normal tissues, regardless of the transcriptional activity of the gene. The main exceptions include nontranscribed genes on the inactive X-chromosome and imprinted autosomal genes where one of the parental alleles may be methylated.¹⁵ Tissue-specific genes without CpG islands are variably methylated, often in a tissue-specific pattern, and usually methylation is inversely correlated with the transcriptional status of the genes (for review, see Bird¹⁶ and Cedar¹⁷).

DNA METHYLTRANSFERASES

The enzymes that transfer methyl groups to the cytosine ring, cytosine 5-methyltransferases, or DNA methyltransferases (DNA-MTase) have been characterized in a number of eukaryotes.¹⁸ The target site for DNA-MTase in DNA is the dinucleotide palindrome CG (commonly referred to as CpG, with p denoting the phosphate group). The first eukaryotic DNA-MTase gene cloned from mouse almost a decade ago¹⁹ is now referred to as *Dnmt1*. This gene is highly conserved among eukaryotes. *Dnmt1* orthologs have been identified in various species, including humans (*DNMT1*).²⁰ Interestingly, *Dnmt1* orthologs have not been found in organisms lacking DNA methylation, such as *Saccharomyces cerevisiae*, *Caenorhabditis elegans*, and *Drosophila melanogaster*.

Mammalian *Dnmt1 methyltransferase* has a high affinity for hemimethylated substrates but is also capable of performing de novo methylation of unmethylated substrates in vitro. The de novo activity of mammalian *Dnmt1 methyltransferase* has been shown to be stimulated by aberrant DNA structures²¹ and 5-mC residues in single-stranded²² or double-stranded DNA substrates.²³ After it was found that the human and mouse *Dnmt1* genes contain additional 5' transcribed sequences with protein-

From the Department of Medicine, Feist-Weiller Cancer Center and Overton Brooks VA Medical Center, LSU Medical Center, Shreveport, LA; and the Departments of Internal Medicine and Human Genetics, Massey Cancer Center, Virginia Commonwealth University, Richmond, VA.

Submitted January 21, 1999; accepted March 26, 1999.

Supported by National Institutes of Health Grant No. DK29902 and by the Massey Cancer Center (G.D.G.) and the Feist-Weiller Cancer Center (R.S.).

Address reprint requests to Gordon D. Ginder, MD, Professor of Internal Medicine and Human Genetics, Director, Massey Cancer Center, Virginia Commonwealth University, 401 College St, PO Box 980037, Richmond, VA 23298-0037; e-mail: gginder@mcc1.mcc.vcu.edu.

© 1999 by The American Society of Hematology.

0006-4971/99/9312-0043\$3.00/0




Fig 1. Schematic representation of the biochemical pathways for cytosine methylation, demethylation, and mutagenesis of cytosine and 5-mC.

encoding potential,²⁴ it became apparent that previous biochemical and cell culture assays had relied on an incomplete *Dnmt1* cDNA sequence that codes for a truncated protein. Pradhan et al²⁵ demonstrated that full-length *Dnmt1* MTase synthesized in Baculovirus was equally active on unmethylated and hemimethylated substrates, in part by virtue of an inhibitory effect of the additional N-terminal amino acids on activity with hemimethylated DNA. These results are evidence that the *Dnmt1* MTase protein has inherent de novo methylating activity that may be altered by protein-protein interactions and enhanced by aberrant structures or 5-mC residues in the substrate DNA.

Evidence for an additional mammalian DNA-MTase capable of de novo methylation came from the experiments of Lei et al.²⁶ These investigators generated a null mutation of the *Dnmt1* gene through homologous recombination in mouse embryonic stem cells. *Dnmt1* null embryonic stem cells were viable and contained low but stable levels of methylcytosine and methyltransferase activity. Interestingly, integrated provirus DNA in Moloney murine leukemia virus (MoMuLV)-infected homozygous *Dnmt1* knockout ES cells exhibited de novo methylation to a similar extent as in wild-type cells. Although a second DNA-MTase has recently been identified in mouse (*Dnmt2*)²⁷ and in humans (*DNMT2*),²⁸ it is not yet clear if it is the long sought after de novo methyltransferase present in *Dnmt1* deficient ES cells.

TECHNIQUES TO STUDY DNA METHYLATION

Early techniques to study site-specific DNA methylation relied primarily on the inability of methylation-sensitive type II restriction enzymes to cleave sequences containing one or more methylated CpG sites, combined with Southern hybridization.²⁹⁻³¹ This method requires large amounts of high molecular weight DNA, detects methylation only if more than a few percent of alleles are methylated, and only provides information about those CpG sites found within the recognition sequence of

methylation-sensitive restriction enzymes. Singer-Sam et al³² improved the sensitivity of methylation detection by combining the use of methylation-sensitive restriction enzymes and polymerase chain reaction (PCR). After cleaving the DNA with methylation-sensitive restriction endonucleases, eg, *Hpa* II, PCR amplification with specific primers flanking the restriction site will only occur if DNA cleavage has been prevented by methylation. However, this method, like Southern-based approaches, can only monitor CpG methylation in methylation-sensitive restriction sites. Also, a false-positive result may be obtained due to incomplete restriction enzyme digestion of cellular DNA.

Genomic sequencing protocols that have previously been used to study DNA methylation use Maxam and Gilbert chemical cleavage reactions performed on genomic DNA³³ with linker-mediated PCR (LMPCR) to enhance the signal.³⁴ These methods are based upon the fact that 5-mC is not cleaved during the standard Maxam and Gilbert cytosine cleavage reaction.³⁵ Thus, 5-mC is identified in a sequencing gel by the lack of a band that corresponds to a cleavage product of a cytosine degradation reaction. This assay is technically demanding and is subject to both false-positive and false-negative results.

Frommer et al³⁶ introduced a procedure based on bisulfite-induced oxidative deamination of genomic DNA under conditions in which cytosine is converted to uracil and 5-mC remains unchanged. The target sequence is then amplified by PCR using strand-specific primers. Upon sequencing of the amplified DNA, all uracil and thymine residues become detectable as thymine and only 5-mC residues amplify as cytosines. This method, as shown in Fig 2, is presently the method of choice for the detailed analysis of 5-mC in any given genomic target sequence.

A number of rapid methods to detect 5-mC have been developed based on the bisulfite deamination reaction in combination with PCR amplification. These are suitable for






Fig 2. (A) Illustration of the steps involved in determining the methylation status of cytosines in a known DNA sequence by the bisulfite conversion method. (B) In vivo methylation of CpG dinucleotides of rho promoter in 5-day and adult chicken erythroid cells using bisulfite conversion technique. Arrows indicate methylated cytosines and are only seen in adult erythroid cells. Positions indicated are relative to the transcription start site. Cytosines that are not associated with CpG dinucleotides (sequence shown in [C]) have all been converted to thymidines in both 5-day and adult erythroid cells. (C) Graphic representation of the in vivo methylation of CpG dinucleotides of rho promoter in 5-day and adult chicken erythroid cells using bisulfite conversion technique. Arrows indicate methylated cytosines and are clearly seen in (A) (data not shown for CpG dinucleotide at position -15). Primers R and F indicate the sequence of rho globin promoter used for designating internal primers and for sequencing.

examining limited numbers of CpG dinucleotides that are either found within or immediately adjacent to the PCR primer sequences^{37,38} or within a restriction enzyme recognition sequence.³⁹

DNA METHYLATION AND TRANSCRIPTIONAL REPRESSION

A role for DNA methylation in the differential regulation of gene expression was hypothesized many years ago.^{40,41} The potential mechanism was suggested by a number of early observations in which site-specific cytosine methylation within or adjacent to genes was found to correlate with transcriptional repression.^{31,42-45} Subsequently, this inverse relationship between cytosine methylation and transcription has been observed in a large number of genes, although not universally.

Despite the long held view that DNA methylation might act as a negative regulator of transcription, the precise mechanism involved has remained elusive. Numerous reports have shown the ability of promoter DNA methylation to inhibit transcription of a wide variety of genes in in vitro transfection assays, and in some cases, such methylation corresponds to the inactive state



of the gene under study in vivo (reviewed in Bird¹⁶ and Razin and Cedar⁴⁶).

Three possible mechanisms have been proposed to account for transcriptional repression by DNA methylation, and each is shown in Fig 3. The first mechanism involves direct interference with the binding of specific transcription factors to their recognition sites in their respective promoters. Several transcription factors, including AP-2, c-Myc/Myn, the cyclic AMP-dependent activator CREB, E2F, and NF- κ B, recognize sequences that contain CpG residues, and binding to each has been shown to be inhibited by methylation. In contrast, other transcription factors (eg, Sp1 and CTF) are not sensitive to methylation of their binding sites,⁴⁷ and many factors have no CpG dinucleotide residues in their binding sites.

A second potential mechanism for methylation induced silencing is through the direct binding of specific transcriptional repressors to methylated DNA. Two such factors, MeCP-1 and MeCP-2 (methyl cytosine binding proteins 1 and 2), have been identified and shown to bind to methylated CpG residues in any sequence context. Although in vertebrates DNA methylation has been posited to inhibit transcription initiation, methylation

has also been shown to block transcription elongation in *Neurospora* through a mechanism that may be mediated through MeCP-1 and/or MeCP-2.⁴⁸

MeCP-1 binds to DNA containing multiple symmetrically methylated CpG sites, as opposed to hemimethylated CpGs, and manifests as a large complex on electrophoretic mobility shift assay.⁴⁹ Repression of transcription from densely methylated genes can be mediated by MeCP-1, and cells deficient in MeCP-1 show much reduced repression of methylated genes.⁵⁰ In a further study, it was demonstrated that sparse methylation could repress transfected genes completely, but the inhibition was fully overcome by the presence in *cis* of an SV40 enhancer. However, densely methylated genes could not be reactivated by the strong enhancer. It was proposed that sparsely methylated genes form an unstable complex with MeCP-1 that prevents transcription when the promoter is weak. This complex can be disrupted by a strong promoter, thereby allowing the methylated gene to be transcribed.⁵¹

We have recently shown that a complex with electrophoretic mobility similar to MeCP-1 forms efficiently with the methylated but not with unmethylated embryonic rho-globin gene promoter sequences, and the complex can be detected using nuclear extracts from the same primary avian erythroid cells in which methylation-mediated transcriptional inhibition was demonstrated.⁵² These results, in conjunction with the demonstration of a role for methylation in silencing rho-globin gene transcription in vivo in normal adult avian erythroid cells,^{53,54} suggest a role for MeCP-1 or a similar complex in developmental silencing of embryonic globin genes during normal erythropoiesis.

A component of the MeCP-1 complex, PCMI, has a methyl-CpG binding domain (MBD) and two cysteine-rich domains (CXXC) that are found in animal DNA methyltransferases and in the mammalian HRX proteins, MLL and ALL-1. Although the functional significance of the CXXC domain is not known, there is evidence that it is a part of a transcriptional repression domain.^{55,56} PCMI has been shown to repress transcription in vitro in a methylation-dependent manner.⁵⁷

MeCP-2 is more abundant than MeCP-1 in the cell and is able to bind to DNA containing a single methylated CpG pair.⁵⁸ MeCP-2, like DNA methyltransferase, is dispensable for the viability of embryonic stem cells, but is essential for embryonic development.⁵⁹ MeCP-2 has two domains: a methyl-CpG binding domain that is essential for chromosomal localization and a transcriptional repressor domain (TRD) that can inhibit transcription from a promoter at a distance, suggesting that MeCP-2 interacts with the transcriptional machinery or the initiation complex.⁶⁰ Recently, a region of MeCP-2 that localizes with the TRD was shown to associate with a corepressor complex containing the transcriptional repressor mSin3A and histone deacetylases. Transcriptional repression in vivo was relieved by the deacetylase inhibitor trichostatin A, suggesting that two global mechanisms of gene regulation, DNA methylation and histone deacetylation, can be linked by MeCP-2.^{61,62} However, in some instances DNA methylation has been shown to play a dominant role over histone deacetylation in transcriptional repression.^{53,54} It has been proposed that MeCP-2 might contribute to the assembly of a more stable repressive chromatin structure.⁶³ However, MeCP-2 can repress transcription of

naked DNA in a cell-free in vitro assay,⁶⁰ suggesting that chromatin formation is not necessary for its repressive action.

A third mechanism by which methylation may mediate transcriptional repression is by altering chromatin structure. Keshet et al⁶⁴ transfected mouse L cells with M13 plasmid constructs containing the human β -globin gene, as well as several other eukaryotic genes, after enzymatic methylation. Unmethylated DNA sequences, after integration and stable propagation in cell culture, were all detected in active chromatin, as measured by DNase I-sensitivity, in contrast to DNA sequences that were methylated in vitro before transfection that were contained in DNase I-resistant, transcriptionally inactive chromatin. Experiments using microinjection of certain methylated and nonmethylated gene templates into nuclei have shown that methylation inhibits transcription only after chromatin is assembled.⁶⁵ Even a strong transcriptional activator, GAL4-VP16, cannot counteract the effect of chromatin once it has assumed the inactive state induced by DNA methylation.⁶⁵ Therefore, in addition to stabilizing the inactive state, methylation also prevents activation by blocking the access of transcription factors.^{63,66} Whether this chromatin effect of methylation is mediated solely by MeCP-2-associated histone deacetylase remains to be determined.

One important issue regarding DNA methylation and transcriptional silencing has been whether methylation is a primary control mechanism or a secondary effect of gene activity. In the case of some genes with sparse CpG nucleotides, gain of methylation occurs after transcriptional silencing⁶⁷; in other cases, loss of methylation occurs after transcriptional activation.⁶⁸ At the same time, in other systems it has been shown that despite optimum nuclear conditions for transcription, including DNase I-sensitive chromatin, transcription can be tightly repressed by CpG methylation.^{52,53} It appears that methylation, particularly of CpG-rich genes, may serve as a locking off mechanism that may follow or precede other events that turn a gene off, but that once in place can prevent activation despite an optimum nuclear environment for transcription.

DNA DEMETHYLATION DURING DEVELOPMENT AND TISSUE-SPECIFIC DIFFERENTIATION

A critical aspect of the overall regulatory role of DNA methylation is the process of demethylation. Using methylation-sensitive restriction enzymes to monitor the general level of DNA methylation, it was shown that, during early development, a dramatic reduction in methylation levels occurs in the preimplantation embryo.⁶⁹ This is followed by a wave of de novo methylation involving most CpG residues but leaving the CpG islands unmethylated at the time of implantation.⁷⁰ After implantation, most of the genomic DNA is methylated, whereas tissue-specific genes undergo demethylation in their tissues of expression.⁴⁶ Thus, a means for both global and tissue- or gene-specific removal of methylcytosines from DNA must exist.

The mechanism underlying the process of demethylation has not been fully elucidated. In some cases, demethylation could be a passive process, ie, inhibition of methylation after DNA replication.⁴⁵ Despite the tight coordination of DNA methylation and replication that has been demonstrated in mammalian cells,⁷¹ the mechanism of methylation inhibition in the presence

of DNA replication remains to be defined. The existence of an active demethylation process not involving DNA replication in mammalian cells has been supported by a number of observations (reviewed in Szyf⁷²). In transient transfection assays using myoblast cells, a methylated α -actin gene was shown to have transcriptional activity similar to unmethylated template secondary to an active demethylation process.⁷³ Using transient transfection assays, Szyf et al⁷⁴ showed that an in vitro methylated SK-plasmid bearing no sequence homology to eukaryotic genes became fully demethylated between 1 and 2 days after transfection into the mouse embryonal carcinoma cell line P19. This demethylating activity was not sequence-selective and was independent of DNA replication.⁷⁴ During B-cell development, both the heavy and light chain Ig genes undergo demethylation.⁷⁵ Specific *cis*-acting elements, including both the intronic and 3' enhancers in the case of the κ light chain gene, have been shown to direct demethylation.^{75,76} Further analysis of the intronic enhancers suggested the involvement of the NF- κ B family of proteins as *trans*-acting factors in inducing demethylation. Both *cis*-acting elements and *trans*-acting factors, therefore, appear to direct the demethylation machinery to its target locus.

The biochemical mechanism underlying the process of demethylation remains unclear. However, extracts from chicken embryos have been demonstrated to have sequence nonspecific demethylating activity in vitro.⁷⁷ This enzymatic activity was limited to demethylation of hemimethylated DNA and involved removal of the methylated cytosine and subsequent repair of the resulting apyrimidinic acid residue.⁷⁸ In the case of rat myoblasts, demethylation activity is performed by an active component whose activity is to excise methylated nucleotides from the DNA template and replace them with an unmodified form.⁷⁹ Interestingly, in the presence of a myoblast extract, demethylation of the α -actin but not of the adenine phosphoribosyl transferase (APRT) CpG island fragment was observed. Although this demethylating activity was initially felt to be dependent on an RNA component, subsequent investigation of a more purified active fraction demonstrated complete resistance of the activity to RNase treatment.⁸⁰

It has long been supposed that direct removal of the methyl group by breaking the carbon-to-carbon bond of 5-mC would be so energetically unfavorable as to be an unlikely event. However, a gene encoding an enzyme capable of directly removing the methyl group from methylated CpG-containing DNA in cell free reactions as well as in transfected cells has been reported recently.⁸¹ If the corresponding endogenous gene product can be shown to have the same activity, this will provide a remarkable new avenue for studying the control of demethylation during normal differentiation and development as well as in oncogenesis.

It has been proposed that the core demethylase in cells is kept in an inactive form by interaction with a protein inhibitor. Demethylation could then come about by the removal or modification of the inhibitor. Such an inhibitor could either affect the entire genome resulting in a pattern similar to that seen in the early preimplantation embryo or be confined to the specific sequence influenced by local *cis*-acting elements and *trans*-acting factors, as in the case of the κ Ig gene example discussed above.⁸²

DNA METHYLATION AND CANCER

A role for DNA methylation in oncogenesis has been hypothesized for many years. Numerous studies have suggested aberrations in DNA methyltransferase activity in tumor cells.⁸³⁻⁸⁵ Transformed cells often have increased total DNA-MTase activity, widespread loss of methylation from normally methylated sites, and more regional areas of hypermethylated DNA.⁸⁶ The potential contribution of DNA methylation to oncogenesis appears to be mediated by one or more of the following mechanisms that are shown in Fig 4.

Signature C→T mutation in cancer cells. The high mutation rate of cytosine residues within the dinucleotide CpG, the target site of mammalian DNA-MTase, can be accounted for by an increased rate of cytosine to thymine transitions, which are, in turn, a consequence of hydrolytic deamination of 5-mC⁸⁷ (Fig 1). This mechanism of mutation was first recognized in prokaryotic systems.⁸⁸

Unmethylated cytosine can also undergo deamination to yield uracil, but the well-characterized Uracil-DNA glycosylase efficiently repairs G:U but not G:T mismatch.⁸⁹ However, DNA-MTase may block this repair, contributing to C→U→T transition. DNA-MTase may also mediate 5-mC→T transition at CpG dinucleotides under conditions that lead to increased DNA-MTase expression or decreased cellular S-adenosylmethionine levels.⁹⁰

A striking example of how this process may lead to oncogenesis is shown by the tumor-suppressor gene p53. Mutations in the p53 tumor-suppressor gene occur in more than 50% of human solid tumors.⁹¹ An estimated 24% of these mutations are C→T transitions at CpG dinucleotides, suggesting that DNA methylation may contribute to these mutations.⁹²

DNA hypomethylation in cancer. Decreased levels of overall genomic methylation are common findings in tumorigenesis.⁹³ This decrease in global methylation appears to begin early and before the development of frank tumor formation.^{94,95} Apart from the overall genomic hypomethylation, specific oncogenes have been observed to be hypomethylated in human tumors. A good inverse correlation between methylation and gene expression was observed in the antiapoptotic bcl-2 gene in B-cell chronic lymphocytic leukemia⁹⁶ and for the k-ras proto-oncogene in lung and colon carcinomas.⁹⁷

Hypermethylation of tumor-suppressor genes. In addition to point mutations or gene deletions, transcriptional repression by hypermethylation of promoter sequences suggests an alternative means for the inactivation of tumor-suppressor genes in cancer. This may result from the increased DNA-MTase levels that have been demonstrated in various cancers⁸³ or it could occur as a result of some other transient event that silences tumor-suppressor gene transcription. The retinoblastoma gene (Rb) was the first classic tumor-suppressor gene in which CpG island hypermethylation was detected. Three of 21 sporadic cases in one study⁹⁸ and 5 of 32 sporadic cases in another study⁹⁹ had hypermethylated CpG islands at the 5' end of the Rb gene. Subsequently, it was also shown that in vitro methylation of the promoter region of Rb directly blocked promoter activation.¹⁰⁰

Inactivation of the von Hippel-Lindau (VHL) gene by somatically acquired mutations in one copy of the VHL gene along with loss of the second allele has been implicated as the




Fig 3. Proposed mechanisms of transcriptional repression mediated by cytosine methylation.




Fig 4. Models for the different mechanisms through which cytosine methylation can promote oncogenesis.

initiating event in spontaneous cases of clear-cell renal carcinoma.¹⁰¹ Herman et al¹⁰² found hypermethylation of the VHL gene CpG island and concomitant lack of expression in 5 of 26 cases of the sporadic form of clear cell renal carcinoma that had no VHL intragenic mutations. Treatment of a clear cell renal carcinoma cell line with 5-deoxyazacytidine, a potent inhibitor of DNA-MTase, resulted in demethylation and expression of a previously silent VHL gene.

One of the most important cell cycle regulatory proteins is p16 (also known as MTS-1 for major tumor suppressor 1, INK4a for INhibitor of cyclin-dependent Kinase 4a, and CDKN2a for Cyclin-Dependent Kinase iNhibitor 2a). The major biochemical effect of p16 is to halt cell-cycle progression at the G1/S boundary, and the loss of p16 function may lead to cancer progression by allowing unregulated cellular proliferation.¹⁰³ A common genetic alteration in tumor cell lines and to a lesser extent in primary tumors is the loss of heterozygosity at chromosome 9p21, which contains both the related p16 and p15 genes. Among most solid tumors studied, a CpG island in the 5' region of p16 gene has been found to be frequently methylated, and treatment of cell lines carrying a hypermethylated p16 allele with 5-azacytidine results in transcriptional activation of the gene.^{104,105} The presence of either coding region deletions or promoter hypermethylation of p16 correlate inversely with the presence of Rb gene mutations in multiple tumor types (reviewed in Baylin et al⁸⁶). Interestingly, in some instances, p16 promoter hypermethylation may be the sole inactivating event for both alleles of the gene and may be the only lesion associated with loss of the cyclin D-Rb pathway.⁸⁶ In colon cancer, despite the lack of allelic loss of the 9p region, homozygous deletions of the p16 gene, or Rb gene mutations,¹⁰⁶ 30% to 40% demonstrate hypermethylated p16 alleles.¹⁰⁵ Hypermethylation-mediated inactivation of the p16 gene has been demonstrated in brain, breast, colon, head and neck, and non-small-cell lung cancer and in high grade non-Hodgkin's lymphoma (reviewed in Baylin et al⁸⁶).

P15 (INK4b) gene is an inhibitor of the cyclin-dependent kinases CDK4 and CDK6 and appears to play a role in transforming growth factor- β (TGF- β)-mediated growth inhibition responses.¹⁰⁷ Methylation-mediated inactivation of the p15 gene occurs predominantly in hematopoietic neoplasms such as acute myelogenous leukemia, acute lymphoblastic leukemia, and Burkitt's lymphoma.⁸⁶ Recently, hypermethylation of the p15 gene has been demonstrated in myelodysplastic syndromes¹⁰⁸ and hypermethylation of both p15 and p16 has been found in multiple myeloma.¹⁰⁹

Induction of chromosomal instability. Lengauer et al¹¹⁰ introduced exogenous CpG-rich sequences in the form of a retrovirally contained β -galactosidase gene into 10 colon cancer cell lines. Five of 10 cell lines failed to express the β -gal gene, and these lines were deficient in mismatch activity repair (MMR-), whereas other cell lines competent for mismatch repair (MMR+) expressed the gene. MMR- cell lines were found to be methylation proficient (MET+), and MMR+ cell lines were methylation deficient (MET-) based on Southern blot analysis and 5-azacytidine-induced reactivation of the β -gal gene. It was proposed that in mismatch repair proficient colon cells a methylation defect (MET-) directly facilitates the gain and loss of whole chromosomes, leading to the genomic

instability necessary for the development and progression of cancer. In contrast, MMR- cells have normal methylation proficiency and develop the required genomic instability by the alternative pathway of mismatch repair deficiency.¹¹¹ Consistent with a role for hypomethylation in chromosomal instability were the findings by Chen et al¹¹² in analyzing embryonic stem (ES) cells nullizygous for the DNA-MTase gene, Dnmt I. Gene deletions of selectable marker genes due to mitotic recombination or chromosomal loss were detected at a much higher frequency in the Dnmt 1-deficient ES cells compared with wild-type control cells.

Additional experimental support for the hypothesis linking DNA methylation to chromosomal integrity comes from earlier studies by Feinberg et al,¹¹³ who demonstrated an average of 8% and 10% reduction in genomic 5-mC content in colon adenomas and adenocarcinomas, respectively, with no significant difference between benign and malignant tumors. Interestingly, three patients with the highest 5-mC content in their normal colon appeared to have Lynch syndrome (HNPCC), which was subsequently shown to have the MMR- phenotype.¹¹³ In a recent study, the occurrence of genome-wide undermethylation, retroviral element amplification, and chromosome remodeling in an interspecific mammalian hybrid (*Macropus eugenii* \times *Wallabia bicolor*) was demonstrated. Atypically extended centromeres of *Macropus eugenii*-derived autosomes in the hybrid were composed primarily of unmethylated, amplified retroviral elements not detectable in either parental species, indicating that the failure of DNA methylation to occur and resultant mobile-element activity in the cell hybrids could facilitate rapid karyotypic evolution.¹¹⁴

ALTERED DNA-METHYLTRANSFERASE ACTIVITY IN CANCER

Several studies in the last few years have demonstrated an increase in DNA-MTase activity in neoplastic cells. Kautiainen and Jones⁸³ examined the levels of DNA methyltransferase in nuclei from 9 tumorigenic and 9 nontumorigenic cell lines. In all but 2 cases, the extractable methyltransferase activity was fourfold to 3,000-fold higher in tumorigenic than in nontumorigenic cells.⁸³ Increased DNA-MTase activity has been reported in colon cancers versus normal mucosa from the same patient.⁸⁴ Laird et al⁸⁵ bred normal mice heterozygous for deletion of the DNA-MTase gene (and having 50% of the DNA-MTase activity compared with wild-type) with mice having a mutant allele of the APC gene.⁸⁵ When combined with loss of the other APC allele, this mutation results in progressive formation of adenomas throughout the gastrointestinal tract of newborn animals. Intestinal adenomas were reduced by 50% in the offspring mice with both the APC and DNA-MTase gene mutations and were essentially eliminated when treatment with 5-deoxyazacytidine was combined with DNA-MTase allelic deletion.

The mechanism underlying the increased DNA-MTase activity in neoplastic cells has not yet been elucidated. MacLeod et al¹¹⁵ proposed that the activation of the *ras* signal transduction pathway causes increased activity by interacting with AP-1 sites in the presumed DNA-MTase promoter region in a murine adrenocortical tumor cell line. However, this apparent promoter region of the DNA-MTase gene was subsequently found to lie far downstream of the transcription start site.^{24,116} It has also

been demonstrated that overexpression of the *c-fos* gene results in cellular transformation through increased DNMTase activity.¹¹⁷ Further studies are required to understand the regulation of the DNA-MTase genes in normal and neoplastic cells.

The mechanisms responsible for maintaining normal and abnormal methylation patterns in normal and neoplastic cells remain unclear. A recent study demonstrated the binding of DNA-MTase to proliferating cell nuclear antigen (PCNA), an auxiliary factor for DNA replication and repair. This binding occurred in intact cells at foci of newly replicated DNA and did not alter DNA-MTase activity. A peptide derived from the cell cycle regulator p21(WAF1) was shown to disrupt the DNA-MTase-PCNA interaction, suggesting that p21(WAF1) may regulate methylation by blocking access of DNA-MTase to PCNA. The extent of expression of DNA-MTase and p21(WAF1) were found to be inversely related in both SV40-transformed and nontransformed cells.¹¹⁸ Based on these findings, it has been proposed that, in normal cells, the p21 protein negatively regulates DNA-MTase-PCNA interaction in early S phase and protects CpG islands from methylation, whereas diminished effects of p21 in late S phase result in the targeting of DNA-MTase to methylated DNA. In cancer cells, loss of p21 function allows DNA-MTase-PCNA interaction in early S phase, possibly facilitating aberrant increased methylation of CpG islands, whereas no change in decreased relative targeting of DNA-MTase to late S phase foci results in loss of normal methylation.¹⁵ This type of mechanism could explain the apparent paradox posed by the observation of both excessive and deficient DNA methylation during tumorigenesis.

CLINICAL AND THERAPEUTIC IMPLICATIONS OF DNA METHYLATION

Just as the vertebrate globin genes were among the first examples of an association between DNA methylation and transcriptional silencing,^{31,43,44,119} so too were they the first target for clinical intervention based on drugs that affect methylation.¹²⁰⁻¹²² Treatment with 5-azacytidine, an irreversible inhibitor of DNA-MTase, was shown to increase expression of the fetal γ -globin gene in nonhuman primates and subsequently in patients with β -thalassemia and sickle cell anemia. Because of its mutagenicity and the observation that other S-phase active cytotoxic agents that do not inhibit DNA methylation could induce similar increases in γ -globin gene expression,¹²³⁻¹²⁵ 5-azacytidine has not been widely used for this application. These experiences point to the limitations of attempting to alter gene expression through the use of global DNA methylation inhibitors that also possess other potent cellular effects and emphasize the need for a more complete understanding of the specificity of DNA methylation and demethylation control.

The recent advances in understanding of altered DNA methylation in cancer discussed above also have potential clinical implications. Because methylation of many involved genes may represent a process specific to neoplastic cells, this change may be a sensitive index of micrometastases.⁸⁶ It may also be of some prognostic value in certain situations. For instance, methylation of the *abl* promoter in chronic myeloid leukemia has been associated with disease of long-standing duration, most likely associated with a higher probability of imminent blast transformation.¹²⁶

The increased DNA-MTase activity seen in multiple cancers has prompted targeting of the inhibition of this enzyme as an anticancer strategy. The DNA-MTase inhibitors, 5-azacytidine and 5-azadeoxycytidine, have been used clinically for the treatment of patients with myelodysplastic syndromes or leukemia.^{127,128} Because, as noted, these agents have effects other than inducing demethylation, have many side effects, and need to be administered by continuous infusion, efforts to develop novel DNA-MTase inhibitors are clearly warranted. In one such approach, Ramchandani et al¹²⁹ demonstrated that intraperitoneal injection of DNA-MTase antisense oligonucleotides reduced the level of DNA-MTase and inhibited the growth of Y1 adrenocortical carcinoma in syngeneic LAF mice.

THE ROLE OF DNA METHYLATION IN EVOLUTION

Two hypotheses have been proposed for the evolutionary role of DNA methylation. The first hypothesis is derived from the perspective of gene numbers and biologic complexity.¹² The size of the genome in free-living organisms has increased from a few thousand genes in prokaryotes (eg, *Escherichia coli* with 4,000 genes) to 7,000 to 25,000 genes in nonvertebrate eukaryotes and 50,000 to 100,000 genes in vertebrates. Bird¹² has proposed that, if the number of tissue-specific genes is to increase during evolution, the efficiency of gene repression must be high. This repression, or transcriptional noise reduction, in eukaryotes has been attributed to two features (the nuclear envelope and histones) that are present in eukaryotes but not in prokaryotes. Vertebrates, in turn, have several fold more genes than the nonvertebrate eukaryotes, and one possible additional repression mechanism for vertebrates could be DNA methylation. A comparison of methylation patterns in invertebrates versus vertebrates shows some important differences. In the invertebrates, methylation of cytosine occurs at only a minor fraction of the CpG dinucleotides in the genome and in some cases, such as *Drosophila*, cannot be detected at all. It is likely that, in the vast majority of eukaryotes, DNA methylation functions as part of a system that silences potentially damaging DNA elements such as transposons, viral genomes, etc. However, vertebrates have the bulk of their DNA methylated, except for CpG islands. Bird¹² proposes that DNA methylation in vertebrates provides a novel layer of global repression, further reducing the transcriptional noise and thereby allowing vertebrates to accumulate and selectively use the extra genes that are crucial to their development.

The alternative theory proposed is that cytosine methylation in mammals is a nuclear host-defense system that evolved primarily to counter the threats posed by endogenous parasitic mobile genetic elements.⁵ Cytosine methylation inactivates the promoter of most viruses and transposons, including retroviruses and Alu elements, and such sequences are methylated in the DNA of differentiated cells. In fact, the large majority of 5-mCs in the genome lie within these elements. Demethylating drugs have been shown to activate transcription of endogenous transposons.¹³⁰ Also, cloned, unmethylated human L1 elements transpose at a higher rate in transfected human cells,¹³¹ and this transposition rate is far in excess of the rate of the identical but methylated endogenous elements. In an interspecific mammalian hybrid involving *Macropus eugenii* and *Wallabia bicolor*, the occurrence of genome-wide undermethylation, retroviral

element amplification, and chromosome remodeling has been demonstrated.¹¹⁴ Yoder et al⁵ also argue that there is little compelling evidence for a role for reversible promoter methylation in developmental gene control. However, our laboratory has shown that methylation does appear to play a role in specific developmental gene control.⁵²⁻⁵⁴

METHYLATION AND FOREIGN GENE SILENCING

In both cultured cells transfected with foreign DNA and transgenic organisms, the newly integrated foreign DNA frequently becomes de novo methylated.¹³² It has been proposed that de novo methylation constitutes a cellular defense mechanism to silence integrated foreign DNA or genes.¹³²

Evidence over the past several years suggests that DNA methylation is involved in the inactivation of virally introduced genes in vivo. The MoMuLV has been shown to be completely inactive in embryonic stem cell and embryonic carcinoma cell lines, and the inactivity is accompanied by de novo methylation of the proviral sequences.^{133,134} Orend et al¹³⁵ have shown that, upon integration, de novo methylation spreads from the center of the integrated collinear viral DNA. Methylation of specific sites in the adenoviral promoter results in promoter inactivation.¹³⁶ Herpes virus also undergoes de novo methylation in mammalian cells.¹³⁷ Epstein-Barr virus DNA has been found to be methylated in the normal lymphocytes of healthy volunteers.¹³⁸

In transduced murine hematopoietic cells, transcriptional inactivation of the proviral MoMuLV-LTR was shown to be associated with methylation of the proviral-LTR.¹³⁹ We have recently demonstrated that an in vitro-methylated retroviral LTR fragment containing 13 CpG dinucleotides was able to compete in binding assays for an MeCP complex that has a mobility similar to MeCP-1,¹⁴⁰ although the role of methylation in LTR silencing remains unresolved.

Methylation-mediated inactivation of foreign gene expression in specific cell types has important therapeutic and pharmacological implications in that inhibition of the methylation of therapeutically introduced genes might enhance gene therapy significantly by preventing transcriptional silencing.

CONCLUSION

DNA methylation clearly plays an important regulatory role in vertebrates, as evidenced by its vital role in embryonic development. Whether the primary evolutionary role of DNA methylation is through transcriptional silencing or as a host defense system against endogenous and exogenous parasitic sequence elements remains to be fully determined. However, there is substantial evidence that DNA methylation plays a critical role in silencing specific genes during development and cell differentiation. The intrinsic mutagenicity of 5-mC, activation of proto-oncogenes through hypomethylation, transcriptional inactivation of tumor-suppressor genes through hypermethylation, and defects in chromosomal segregation due to failure of de novo methylation may all contribute to neoplasia. Selective modulation of DNA methylation may therefore have important clinical implications for the prevention and treatment of cancer. To develop safe and effective strategies for therapeutic alteration of DNA methylation, the factors that regulate the specificity of both the methylation and demethylation processes

must be more fully understood. Likewise, understanding the factors involved in DNA methylation-induced gene silencing will facilitate attempts to selectively affect gene expression. Recent studies have linked two global mechanisms of gene regulation, DNA methylation, and histone deacetylation.^{61,62} Further investigations are necessary to understand the complex links between the methyltransferases, demethylases, methyl cytosine binding proteins, histone acetylation, and the transcriptional activity of genes.

ACKNOWLEDGMENT

The authors acknowledge the helpful assistance of Catharine W. Tucker in preparing their manuscript and thank Dr Steven Snyder for comments.

REFERENCES

1. Weissbach A: A chronicle of DNA methylation (1948-1975). *EXS* 64:1, 1993
2. Urieli-Shoval S, Gruenbaum Y, Sedat J, Razin A: The absence of detectable methylated bases in *Drosophila melanogaster* DNA. *FEBS Lett* 146:148, 1982
3. Proffitt JH, Davie JR, Swinton D, Hattman S: 5-Methylcytosine is not detectable in *Saccharomyces cerevisiae* DNA. *Mol Cell Biol* 4:985, 1984
4. Li E, Bestor TH, Jaenisch R: Targeted mutation of the DNA methyltransferase gene results in embryonic lethality. *Cell* 69:915, 1992
5. Yoder JA, Walsh CP, Bestor TH: Cytosine methylation and the ecology of intragenomic parasites. *Trends Genet* 13:335, 1997
6. Bartolomei MS, Tilghman SM: Genomic imprinting in mammals. *Annu Rev Genet* 31:493, 1997
7. Reik W, Walter J: Imprinting mechanisms in mammals. *Curr Opin Genet Dev* 8:154, 1998
8. Heard E, Clerc P, Avner P: X-chromosome inactivation in mammals. *Annu Rev Genet* 31:571, 1997
9. Bird AP: CpG-rich islands and the function of DNA methylation. *Nature* 321:209, 1986
10. Antequera F, Bird A: Number of CpG islands and genes in human and mouse. *Proc Natl Acad Sci USA* 90:11995, 1993
11. Antequera F, Bird A: CpG islands. *EXS* 64:169, 1993
12. Bird AP: Gene number, noise reduction and biological complexity. *Trends Genet* 11:94, 1995
13. Tazi J, Bird A: Alternative chromatin structure at CpG islands. *Cell* 60:909, 1990
14. Cross SH, Bird AP: CpG islands and genes. *Curr Opin Genet Dev* 5:309, 1995
15. Baylin SB: Tying it all together: Epigenetics, genetics, cell cycle, and cancer. *Science* 277:1948, 1997
16. Bird A: The essentials of DNA methylation. *Cell* 70:5, 1992
17. Cedar H: DNA methylation and gene activity. *Cell* 53:3, 1988
18. Bestor TH, Verdine GL: DNA methyltransferases. *Curr Opin Cell Biol* 6:380, 1994
19. Bestor T, Laudano A, Mattaliano R, Ingram V: Cloning and sequencing of a cDNA encoding DNA methyltransferase of mouse cells. The carboxyl-terminal domain of the mammalian enzymes is related to bacterial restriction methyltransferases. *J Mol Biol* 203:971, 1988
20. Yen RW, Vertino PM, Nelkin BD, Yu JJ, el-Deiry W, Kumaraswamy A, Lennon GG, Trask BJ, Celano P, Baylin SB: Isolation and characterization of the cDNA encoding human DNA methyltransferase. *Nucleic Acids Res* 20:2287, 1992
21. Laayoun A, Smith SS: Methylation of slipped duplexes, snapbacks and cruciforms by human DNA(cytosine-5)methyltransferase. *Nucleic Acids Res* 23:1584, 1995

22. Christman JK, Sheikhejad G, Marasco CJ, Sufrin JR: 5-Methyl-2'-deoxycytidine in single-stranded DNA can act in cis to signal de novo DNA methylation. *Proc the Natl Acad Sci USA* 92:7347, 1995
23. Tollefsbol TO, Hutchison CA 3rd: Control of methylation spreading in synthetic DNA sequences by the murine DNA methyltransferase. *J Mol Biol* 269:494, 1997
24. Yoder JA, Yen RWC, Vertino PM, Bestor TH, Baylin SB: New 5' regions of the murine and human genes for DNA (cytosine-5)-methyltransferase. *J Biol Chem* 271:31092, 1996
25. Pradhan S, Talbot D, Sha M, Benner J, Hornstra L, Li E, Jaenisch R, Roberts RJ: Baculovirus-mediated expression and characterization of the full-length murine DNA methyltransferase. *Nucleic Acids Res* 25:4666, 1997
26. Lei H, Oh SP, Okano M, Juttermann R, Goss KA, Jaenisch R, Li E: De novo DNA cytosine methyltransferase activities in mouse embryonic stem cells. *Development* 122:3195, 1996
27. Okano M, Xie S, Li E: Dnmt2 is not required for de novo and maintenance methylation of viral DNA in embryonic stem cells. *Nucleic Acids Res* 26:2536, 1998
28. Yoder JA, Bestor TH: A candidate mammalian DNA methyltransferase related to pmt1p of fission yeast. *Hum Mol Genet* 7:279, 1998
29. Bird AP, Southern EM: Use of restriction enzymes to study eukaryotic DNA methylation: I. The methylation pattern in ribosomal DNA from *Xenopus laevis*. *J Mol Biol* 118:27, 1978
30. Waalwijk C, Flavell RA: MspI, an isoschizomer of hpaII which cleaves both unmethylated and methylated hpaII sites. *Nucleic Acids Res* 5:3231, 1978
31. McGhee JD, Ginder GD: Specific DNA methylation sites in the vicinity of the chicken beta-globin genes. *Nature* 280:419, 1979
32. Singer-Sam J, Grant M, LeBon JM, Okuyama K, Chapman V, Monk M, Riggs AD: Use of a HpaII-polymerase chain reaction assay to study DNA methylation in the Pcg-1 CpG island of mouse embryos at the time of X-chromosome inactivation. *Mol Cell Biol* 10:4987, 1990
33. Church GM, Gilbert W: Genomic sequencing. *Proc Natl Acad Sci USA* 81:1991, 1984
34. Pfeifer GP, Steigerwald SD, Mueller PR, Wold B, Riggs AD: Genomic sequencing and methylation analysis by ligation mediated PCR. *Science* 246:810, 1989
35. Maxam AM, Gilbert W: Sequencing end-labeled DNA with base-specific chemical cleavages. *Methods Enzymol* 65:499, 1980
36. Frommer M, McDonald LE, Millar DS, Collis CM, Watt F, Grigg GW, Molloy PL, Paul CL: A genomic sequencing protocol that yields a positive display of 5-methylcytosine residues in individual DNA strands. *Proc Natl Acad Sci USA* 89:1827, 1992
37. Herman JG, Graff JR, Myohanen S, Nelkin BD, Baylin SB: Methylation-specific PCR: A novel PCR assay for methylation status of CpG islands. *Proc Natl Acad Sci USA* 93:9821, 1996
38. Gonzalgo ML, Jones PA: Rapid quantitation of methylation differences at specific sites using methylation-sensitive single nucleotide primer extension (Ms-SNuPE). *Nucleic Acids Res* 25:2529, 1997
39. Xiong Z, Laird PW: COBRA: A sensitive and quantitative DNA methylation assay. *Nucleic Acids Res* 25:2532, 1997
40. Riggs AD: X inactivation, differentiation, and DNA methylation. *Cytogenet Cell Genet* 14:9, 1975
41. Holliday R, Pugh JE: DNA modification mechanisms and gene activity during development. *Science* 187:226, 1975
42. Ginder GD, McGhee JD: DNA Methylation in the Chicken Adult β -Globin Gene: A Relationship With Gene Expression: Organization and Expression of Globin Genes. New York, NY, Liss, 1981, p 191
43. Shen CK, Maniatis T: Tissue-specific DNA methylation in a cluster of rabbit beta-like globin genes. *Proc Natl Acad Sci USA* 77:6634, 1980
44. van der Ploeg LH, Flavell RA: DNA methylation in the human gamma delta beta-globin locus in erythroid and nonerythroid tissues. *Cell* 19:947, 1980
45. Razin A, Riggs AD: DNA methylation and gene function. *Science* 210:604, 1980
46. Razin A, Cedar H: DNA methylation and gene expression. *Microbiol Rev* 55:451, 1991
47. Tate PH, Bird AP: Effects of DNA methylation on DNA-binding proteins and gene expression. *Curr Opin Genet Dev* 3:226, 1993
48. Rountree MR, Selker EU: DNA methylation inhibits elongation but not initiation of transcription in *Neurospora crassa*. *Genes Dev* 11:2383, 1997
49. Meehan RR, Lewis JD, McKay S, Kleiner EL, Bird AP: Identification of a mammalian protein that binds specifically to DNA containing methylated CpGs. *Cell* 58:499, 1989
50. Boyes J, Bird A: DNA methylation inhibits transcription indirectly via a methyl-CpG binding protein. *Cell* 64:1123, 1991
51. Boyes J, Bird A: Repression of genes by DNA methylation depends on CpG density and promoter strength: Evidence for involvement of a methyl-CpG binding protein. *EMBO J* 11:327, 1992
52. Singal R, Ferris R, Little JA, Wang SZ, Ginder GD: Methylation of the minimal promoter of an embryonic globin gene silences transcription in primary erythroid cells. *Proc Natl Acad Sci USA* 94:13724, 1997
53. Ginder GD, Whitters MJ, Pohlman JK: Activation of a chicken embryonic globin gene in adult erythroid cells by 5-azacytidine and sodium butyrate. *Proc Natl Acad Sci USA* 81:3954, 1984
54. Burns LJ, Glauber JG, Ginder GD: Butyrate induces selective transcriptional activation of a hypomethylated embryonic globin gene in adult erythroid cells. *Blood* 72:1536, 1988
55. Zeleznik-Le NJ, Harden AM, Rowley JD: 11q23 translocations split the "AT-hook" cruciform DNA-binding region and the transcriptional repression domain from the activation domain of the mixed-lineage leukemia (MLL) gene. *Proc Natl Acad Sci USA* 91:10610, 1994
56. Prasad R, Yano T, Sorio C, Nakamura T, Rallapalli R, Gu Y, Leshkowitz D, Croce CM, Canaani E: Domains with transcriptional regulatory activity within the ALL1 and AF4 proteins involved in acute leukemia. *Proc Natl Acad Sci USA* 92:12160, 1995
57. Cross SH, Meehan RR, Nan X, Bird A: A component of the transcriptional repressor MeCP1 shares a motif with DNA methyltransferase and HRX proteins. *Nat Genet* 16:256, 1997
58. Meehan R, Lewis J, Cross S, Nan X, Jeppesen P, Bird A: Transcriptional repression by methylation of CpG. *J Cell Sci Suppl* 16:9, 1992
59. Tate P, Skarnes W, Bird A: The methyl-CpG binding protein MeCP2 is essential for embryonic development in the mouse. *Nat Genet* 12:205, 1996
60. Nan X, Campoy FJ, Bird A: MeCP2 is a transcriptional repressor with abundant binding sites in genomic chromatin. *Cell* 88:471, 1997
61. Nan XS, Ng HH, Johnson CA, Laherty CD, Turner BM, Eisenman RN, Bird A: Transcriptional repression by the methyl-CpG-binding protein MeCP2 involves a histone deacetylase complex. *Nature* 393:386, 1998
62. Jones PL, Veenstra GJ, Wade PA, Vermaak D, Kass SU, Landsberger N, Strouboulis J, Wolffe AP: Methylated DNA and MeCP2 recruit histone deacetylase to repress transcription. *Nat Genet* 19:187, 1998
63. Kass SU, Pruss D, Wolffe AP: How does DNA methylation repress transcription? *Trends Genet* 13:444, 1997
64. Keshet I, Lieman-Hurwitz J, Cedar H: DNA methylation affects the formation of active chromatin. *Cell* 44:535, 1986
65. Kass SU, Landsberger N, Wolffe AP: DNA methylation directs a time-dependent repression of transcription initiation. *Curr Biol* 7:157, 1997
66. Siegfried Z, Cedar H: DNA methylation: A molecular lock. *Curr Biol* 7:R305, 1997

67. Enver T, Zhang JW, Papayannopoulou T, Stamatoyannopoulos G: DNA methylation: A secondary event in globin gene switching? *Genes Dev* 2:698, 1988
68. Sullivan CH, Norman JT, Borras T, Grainger RM: Developmental regulation of hypomethylation of delta-crystallin genes in chicken embryo lens cells. *Mol Cell Biol* 9:3132, 1989
69. Monk M, Boubelik M, Lehnert S: Temporal and regional changes in DNA methylation in the embryonic, extraembryonic and germ cell lineages during mouse embryo development. *Development* 99:371, 1987
70. Kafri T, Ariel M, Brandeis M, Shemer R, Urven L, McCarrey J, Cedar H, Razin A: Developmental pattern of gene-specific DNA methylation in the mouse embryo and germ line. *Genes Dev* 6:705, 1992
71. Araujo FD, Knox JD, Szyf M, Price GB, Zannis-Hadjopoulos M: Concurrent replication and methylation at mammalian origins of replication. *Mol Cell Biol* 18:3475, 1998
72. Szyf M: The DNA methylation machinery as a target for anticancer therapy. *Pharmacol Therapeutics* 70:1, 1996
73. Paroush Z, Keshet I, Yisraeli J, Cedar H: Dynamics of demethylation and activation of the alpha-actin gene in myoblasts. *Cell* 63:1229, 1990
74. Szyf M, Theberge J, Bozovic V: Ras induces a general DNA demethylation activity in mouse embryonal P19 cells. *J Biol Chem* 270:12690, 1995
75. Lichtenstein M, Keini G, Cedar H, Bergman Y: B cell-specific demethylation: A novel role for the intronic kappa chain enhancer sequence. *Cell* 76:913, 1994
76. Kirillov A, Kistler B, Mostoslavsky R, Cedar H, Wirth T, Bergman Y: A role for nuclear NF-kappaB in B-cell-specific demethylation of the Igkappa locus. *Nat Genet* 13:435, 1996
77. Jost JP: Nuclear extracts of chicken embryos promote an active demethylation of DNA by excision repair of 5-methyldeoxycytidine. *Proc Natl Acad Sci USA* 90:4684, 1993
78. Jost JP, Siegmund M, Sun L, Leung R: Mechanisms of DNA demethylation in chicken embryos. Purification and properties of a 5-methylcytosine-DNA glycosylase. *J Biol Chem* 270:9734, 1995
79. Weiss A, Keshet I, Razin A, Cedar H: DNA demethylation in vitro: involvement of RNA. *Cell* 86:709, 1996
80. Swisher JF, Rand E, Cedar H, Marie Pyle A: Analysis of putative RNase sensitivity and protease insensitivity of demethylation activity in extracts from rat myoblasts. *Nucleic Acids Res* 26:5573, 1998
81. Bhattacharya SK, Ramchandani S, Cervoni N, Szyf M: A mammalian protein with specific demethylase activity for mCpG DNA. *Nature* 397:579, 1999
82. Weiss A, Cedar H: The role of DNA demethylation during development. *Genes Cells* 2:481, 1997
83. Kautiainen TL, Jones PA: DNA methyltransferase levels in tumorigenic and nontumorigenic cells in culture. *J Biol Chem* 261:1594, 1986
84. Issa JP, Vertino PM, Wu J, Sazawal S, Celano P, Nelkin BD, Hamilton SR, Baylin SB: Increased cytosine DNA-methyltransferase activity during colon cancer progression. *J Natl Cancer Inst* 85:1235, 1993
85. Laird PW, Jackson-Grusby L, Fazeli A, Dickinson SL, Jung WE, Li E, Weinberg RA, Jaenisch R: Suppression of intestinal neoplasia by DNA hypomethylation. *Cell* 81:197, 1995
86. Baylin SB, Herman JG, Graff JR, Vertino PM, Issa JP: Alterations in DNA methylation: A fundamental aspect of neoplasia. *Adv Cancer Res* 72:141, 1998
87. Laird PW, Jaenisch R: The role of DNA methylation in cancer genetic and epigenetics. *Annu Rev Genet* 30:441, 1996
88. Coulondre C, Miller JH, Farabaugh PJ, Gilbert W: Molecular basis of base substitution hotspots in *Escherichia coli*. *Nature* 274:775, 1978
89. Gonzalgo ML, Liang G, Spruck CHR, Zingg JM, Rideout WMr, Jones PA: Identification and characterization of differentially methylated regions of genomic DNA by methylation-sensitive arbitrarily primed PCR. *Cancer Res* 57:594, 1997
90. Bender CM, Zingg JM, Jones PA: DNA methylation as a target for drug design. *Pharmaceutical Res* 15:175, 1998
91. Greenblatt MS, Bennett WP, Hollstein M, Harris CC: Mutations in the p53 tumor suppressor gene: Clues to cancer etiology and molecular pathogenesis. *Cancer Res* 54:4855, 1994
92. Magewu AN, Jones PA: Ubiquitous and tenacious methylation of the CpG site in codon 248 of the p53 gene may explain its frequent appearance as a mutational hot spot in human cancer. *Mol Cell Biol* 14:4225, 1994
93. Gama-Sosa MA, Slagel VA, Trewyn RW, Oxenhandler R, Kuo KC, Gehrke CW, Ehrlich M: The 5-methylcytosine content of DNA from human tumors. *Nucleic Acids Res* 11:6883, 1983
94. Christman JK, Sheikhnejad G, Dizik M, Abileah S, Wainfan E: Reversibility of changes in nucleic acid methylation and gene expression induced in rat liver by severe dietary methyl deficiency. *Carcinogenesis* 14:551, 1993
95. Pogribny IP, Miller BJ, James SJ: Alterations in hepatic p53 gene methylation patterns during tumor progression with folate/methyl deficiency in the rat. *Cancer Lett* 115:31, 1997
96. Hanada M, Delia D, Aiello A, Stadtmauer E, Reed JC: bcl-2 gene hypomethylation and high-level expression in B-cell chronic lymphocytic leukemia. *Blood* 82:1820, 1993
97. Feinberg AP, Vogelstein B: Hypomethylation distinguishes genes of some human cancers from their normal counterparts. *Nature* 301:89, 1983
98. Greger V, Debus N, Lohmann D, Hopping W, Passarge E, Horsthemke B: Frequency and parental origin of hypermethylated RB1 alleles in retinoblastoma. *Hum Genet* 94:491, 1994
99. Sakai T, Toguchida J, Ohtani N, Yandell DW, Rapaport JM, Dryja TP: Allele-specific hypermethylation of the retinoblastoma tumor-suppressor gene. *Am J Hum Genet* 48:880, 1991
100. Ohtani-Fujita N, Fujita T, Aoike A, Osifchin NE, Robbins PD, Sakai T: CpG methylation inactivates the promoter activity of the human retinoblastoma tumor-suppressor gene. *Oncogene* 8:1063, 1993
101. Gnarr JR, Tory K, Weng Y, Schmidt L, Wei MH, Li H, Latif F, Liu S, Chen F, Duh FM, Lobensky I, Duan DR, Florence C, Pozzatti R, Walther MM, Bander NH, Grossman HB, Brauch H, Pomer S, Brooks JD, Isaacs WB, Lerman MI, Zbar B, Linehan WM: Mutations of the VHL tumour suppressor gene in renal carcinoma. *Nat Genet* 7:85, 1994
102. Herman JG, Latif F, Weng Y, Lerman MI, Zbar B, Liu S, Samid D, Duan DS, Gnarr JR, Linehan WM, Baylin SB: Silencing of the VHL tumor-suppressor gene by DNA methylation in renal carcinoma. *Proc Natl Acad Sci USA* 91:9700, 1994
103. Liggett WH Jr, Sidransky D: Role of the p16 tumor suppressor gene in cancer. *J Clin Oncol* 16:1197, 1998
104. Merlo A, Herman JG, Mao L, Lee DJ, Gabrielson E, Burger PC, Baylin SB, Sidransky D: 5' CpG island methylation is associated with transcriptional silencing of the tumour suppressor p16/CDKN2/MTS1 in human cancers. *Nat Med* 1:686, 1995
105. Herman JG, Merlo A, Mao L, Lapidus RG, Issa JP, Davidson NE, Sidransky D, Baylin SB: Inactivation of the CDKN2/p16/MTS1 gene is frequently associated with aberrant DNA methylation in all common human cancers. *Cancer Res* 55:4525, 1995
106. Cairns P, Polascik TJ, Eby Y, Tokino K, Califano J, Merlo A, Mao L, Herath J, Jenkins R, Westra W, Rutter JL, Buckler A, Gabrielson E, Tockman M, Cho KR, Hedrick L, Bova GS, Isaacs W, Koch W, Schwab D, Sidransky D: Frequency of homozygous deletion at p16/CDKN2 in primary human tumours. *Nat Genet* 11:210, 1995
107. Hannon GJ, Beach D: p15INK4B is a potential effector of TGF-beta-induced cell cycle arrest. *Nature* 371:257, 1994
108. Uchida T, Kinoshita T, Nagai H, Nakahara Y, Saito H, Hotta T,

- Murate T: Hypermethylation of the p15INK4B gene in myelodysplastic syndromes. *Blood* 90:1403, 1997
109. Ng MH, Chung YF, Lo KW, Wickham NW, Lee JC, Huang DP: Frequent hypermethylation of p16 and p15 genes in multiple myeloma. *Blood* 89:2500, 1997
110. Lengauer C, Kinzler KW, Vogelstein B: DNA methylation and genetic instability in colorectal cancer cells. *Proc Natl Acad Sci USA* 94:2545, 1997
111. Jones PA, Gonzalgo ML: Altered DNA methylation and genome instability: A new pathway to cancer? *Proc Natl Acad Sci USA* 94:2103, 1997
112. Chen RZ, Pettersson U, Beard C, Jackson-Grusby L, Jaenisch R: DNA hypomethylation leads to elevated mutation rates. *Nature* 395:89, 1998
113. Feinberg AP, Gehrke CW, Kuo KC, Ehrlich M: Reduced genomic 5-methylcytosine content in human colonic neoplasia. *Cancer Res* 48:1159, 1988
114. Oneill RJW, Oneill MJ, Graves JAM: Undermethylation associated with retroelement activation and chromosome remodelling in an interspecific mammalian hybrid. *Nature* 393:68, 1998
115. MacLeod AR, Rouleau J, Szyf M: Regulation of DNA methylation by the Ras signaling pathway. *J Biol Chem* 270:11327, 1995
116. Tucker KL, Talbot D, Lee MA, Leonhardt H, Jaenisch R: Complementation of methylation deficiency in embryonic stem cells by a DNA methyltransferase minigene. *Proc Natl Acad Sci USA* 93:12920, 1996
117. Bakin AV, Curran T: Role of DNA 5-methylcytosine transferase in cell transformation by fos. *Science* 283:387, 1999
118. Chuang LS, Ian HI, Koh TW, Ng HH, Xu G, Li BF: Human DNA-(cytosine-5) methyltransferase-PCNA complex as a target for p21WAF1. *Science* 277:1996, 1997
119. Busslinger M, Hurst J, Flavell RA: DNA methylation and the regulation of globin gene expression. *Cell* 34:197, 1983
120. DeSimone J, Heller P, Hall L, Zwiers D: 5-Azacytidine stimulates fetal hemoglobin synthesis in anemic baboons. *Proc Natl Acad Sci USA* 79:4428, 1982
121. Ley TJ, DeSimone J, Anagnou NP, Keller GH, Humphries RK, Turner PH, Young NS, Keller P, Nienhuis AW: 5-azacytidine selectively increases gamma-globin synthesis in a patient with beta+ thalassemia. *N Engl J Med* 307:1469, 1982
122. Charache S, Dover G, Smith K, Talbot CC Jr, Moyer M, Boyer S: Treatment of sickle cell anemia with 5-azacytidine results in increased fetal hemoglobin production and is associated with nonrandom hypomethylation of DNA around the gamma-delta-beta-globin gene complex. *Proc Natl Acad Sci USA* 80:4842, 1983
123. Letvin NL, Linch DC, Beardsley GP, McIntyre KW, Nathan DG: Augmentation of fetal-hemoglobin production in anemic monkeys by hydroxyurea. *N Engl J Med* 310:869, 1984
124. Papayannopoulou T, Torrealba de Ron A, Veith R, Knitter G, Stamatoyannopoulos G: Arabinosylcytosine induces fetal hemoglobin in baboons by perturbing erythroid cell differentiation kinetics. *Science* 224:617, 1984
125. Platt OS, Orkin SH, Dover G, Beardsley GP, Miller B, Nathan DG: Hydroxyurea enhances fetal hemoglobin production in sickle cell anemia. *J Clin Invest* 74:652, 1984
126. Ben-Yehuda D, Krichevsky S, Rachmilewitz EA, Avraham A, Palumbo GA, Frassoni F, Sahar D, Rosenbaum H, Paltiel O, Zion M, Ben-Neriah Y: Molecular follow-up of disease progression and interferon therapy in chronic myelocytic leukemia. *Blood* 90:4918, 1997
127. Silverman LR, Holland JF, Weinberg RS, Alter BP, Davis RB, Ellison RR, Demakos EP, Cornell CJ Jr, Carey RW, Schiffer C: Effects of treatment with 5-azacytidine on the in vivo and in vitro hematopoiesis in patients with myelodysplastic syndromes. *Leukemia* 7:21, 1993 (suppl 1)
128. Willemze R, Archimbaud E, Muus P: Preliminary results with 5-aza-2'-deoxycytidine (DAC)-containing chemotherapy in patients with relapsed or refractory acute leukemia. The EORTC Leukemia Cooperative Group. *Leukemia* 7:49, 1993 (suppl 1)
129. Ramchandani S, MacLeod AR, Pinard M, von Hofe E, Szyf M: Inhibition of tumorigenesis by a cytosine-DNA, methyltransferase, antisense oligodeoxynucleotide. *Proc Natl Acad Sci USA* 94:684, 1997
130. Schmid CW: Alu: Structure, origin, evolution, significance and function of one-tenth of human DNA. *Prog Nucleic Acid Res Mol Biol* 53:283, 1996
131. Moran JV, Holmes SE, Naas TP, DeBerardinis RJ, Boeke JD, Kazanian HH Jr: High frequency retrotransposition in cultured mammalian cells. *Cell* 87:917, 1996
132. Doerfler W, Schubert R, Heller H, Kammer C, Hilger-Eversheim K, Knoblauch M, Remus R: Integration of foreign DNA and its consequences in mammalian systems. *Trends Biotechnol* 15:297, 1997
133. Harbers K, Schnieke A, Stuhlmann H, Jahner D, Jaenisch R: DNA methylation and gene expression: Endogenous retroviral genome becomes infectious after molecular cloning. *Proc Natl Acad Sci USA* 78:7609, 1981
134. Jahner D, Jaenisch R: Retrovirus-induced de novo methylation of flanking host sequences correlates with gene inactivity. *Nature* 315:594, 1985
135. Orend G, Kuhlmann I, Doerfler W: Spreading of DNA methylation across integrated foreign (adenovirus type 12) genomes in mammalian cells. *J Virol* 65:4301, 1991
136. Kruczek I, Doerfler W: Expression of the chloramphenicol acetyltransferase gene in mammalian cells under the control of adenovirus type 12 promoters: Effect of promoter methylation on gene expression. *Proc Natl Acad Sci USA* 80:7586, 1983
137. Barletta J, Greer SB: Methylation of HSV-1 DNA as a mechanism of viral inhibition: Studies of an analogue of methyldeoxycytidine: Trifluoromethyldeoxycytidine (F3mdCyd). *Antiviral Res* 18:1, 1992
138. Robertson KD, Ambinder RF: Methylation of the Epstein-Barr virus genome in normal lymphocytes. *Blood* 90:4480, 1997
139. Challita PM, Kohn DB: Lack of expression from a retroviral vector after transduction of murine hematopoietic stem cells is associated with methylation in vivo. *Proc Natl Acad Sci USA* 91:2567, 1994
140. Osborne C, Pasceri P, Singal R, Ginder GR: Amelioration of retroviral vector silencing in locus control region β -globin transgenic mice and trasduced F9 embryonic cells. *J Virol* (in press)