

# CXC chemokine receptor 3 expression on CD34<sup>+</sup> hematopoietic progenitors from human cord blood induced by granulocyte-macrophage colony-stimulating factor: chemotaxis and adhesion induced by its ligands, interferon $\gamma$ -inducible protein 10 and monokine induced by interferon $\gamma$

Tan Jinquan, Sha Quan, Henrik H. Jacobi, Chen Jing, Anders Millner, Bettina Jensen, Hans O. Madsen, Lars P. Ryder, Arne Svejgaard, Hans-Jørgen Malling, Per S. Skov, and Lars K. Poulsen

CXC chemokine receptor 3 (CXCR3), which is known to be expressed predominately on memory and activated T lymphocytes, is a receptor for both interferon  $\gamma$ (IFN- $\gamma$ )-inducible protein 10 ( $\gamma$ IP-10) and monokine induced by IFN- $\gamma$  (Mig). We report the novel finding that CXCR3 is also expressed on CD34<sup>+</sup> hematopoietic progenitors from human cord blood stimulated with granulocyte-macrophage colony-stimulating factor (GM-CSF) but not on freshly isolated CD34<sup>+</sup> progenitors. Freshly isolated CD34+ progenitors expressed low levels of CXCR3 messenger RNA, but this expression was highly up-regulated by GM-CSF, as indicated by a real-time quantitative reverse transcriptase-polymerase chain reaction technique. yIP-10 and Mig induced chemo-

## taxis of GM-CSF-stimulated CD34+ progenitors by means of CXCR3, since an anti-CXCR3 monoclonal antibody (mAb) was found to block yIP-10-induced and Mig-induced CD34<sup>+</sup> progenitor chemotaxis. These chemotactic attracted CD34<sup>+</sup> progenitors are colony-forming unitsgranulocyte-macrophage. yIP-10 and Mig also induced GM-CSF-stimulated CD34+ progenitor adhesion and aggregation by means of CXCR3, a finding confirmed by the observation that anti-CXCR3 mAb blocked these functions of $\gamma$ IP-10 and Mig but not of chemokine stromal cellderived factor 1a. vIP-10-induced and Mig-induced up-regulation of integrins (CD49a and CD49b) was found to play a crucial role in adhesion of GM-CSFstimulated CD34+ progenitors. Moreover,

yIP-10 and Mig stimulated CXCR3 redistribution and cellular polarization in GM-CSF-stimulated CD34<sup>+</sup> progenitors. These results indicate that CXCR3-yIP-10 and CXCR3-Mig receptor-ligand pairs, as well as the effects of GM-CSF on them, may be especially important in the cytokine/chemokine environment for the physiologic and pathophysiologic events of differentiation of CD34+ hematopoietic progenitors into lymphoid and myeloid stem cells, subsequently immune and inflammatory cells. These processes include transmigration, relocation, differentiation, and maturation of CD34+ hematopoietic progenitors. (Blood. 2000;96:1230-1238)

© 2000 by The American Society of Hematology

## Introduction

The trafficking of hematopoietic progenitor cells occurs during mobilization and homing, which play a key role in subsequent proliferation, differentiation, and maturation of these cells. However, little is known about the mechanisms and molecules that regulate the homing, retention, and migration of hematopoietic progenitor cells in hematopoietic organs. Analogous with the processes in mature leukocytes, these processes likely involve chemoattractant molecules and their receptors, which are known to regulate the trafficking of leukocytes under both physiologic and pathologic conditions. The CXC chemokine receptor (CXCR) 3, a G-protein-coupled 7-transmembrane receptor, is expressed at high levels on activated and memory T cells, B cells, natural killer cells, and plasmacytoid monocytes.<sup>1-3</sup> It has also been shown to bind interferon (IFN)  $\gamma$ -inducible protein 10 ( $\gamma$ IP-10) and monokine induced by IFN- $\gamma$  (Mig), with  $K_i$  values of 0.14 and 4.9 nmol/L, respectively.

Submitted December 30, 1999; accepted April 10, 2000.

yIP-10 and Mig are 2 members of the CXC chemokine superfamily whose expression is dramatically up-regulated by IFN-y. Both chemokines have been shown to be functional agonists of CXCR3.4 The CC chemokine 6Ckine and IFNinducible T-cell  $\alpha$  chemoattractant (I-TAC) were identified as ligands for CXCR3.5-7 yIP-10 and Mig induce rapid and transient adhesion of human interleukin (IL) 2-stimulated T lymphocytes to immobilized integrin ligands through their receptor CXCR3, which is selectively expressed on activated T cells.8 Immature T cells expressed only CXCR4, whereas most memory and activated T cells expressed CXCR3, and a small proportion expressed CC chemokine receptor (CCR) 3 and CCR5.9 CXCR3 was found to be expressed at high levels on T-helper (Th) 0 and Th1 lymphocytes and at low levels on Th2 lymphocytes.<sup>3</sup> In contrast, CCR3 and CCR4 were found on Th2 lymphocytes.<sup>9</sup> Circulating blood T cells, B cells, and natural killer cells also express CXCR3.3

**Reprints:** Tan Jinquan or Lars K. Poulsen, Laboratory of Medical Allergology, National University Hospital, 9 Blegdamsvej, DK-2200 Copenhagen Ø, Denmark; e-mail: tan@rh.dk or lkpallgy@inet.uni2.dk.

The publication costs of this article were defrayed in part by page charge payment. Therefore, and solely to indicate this fact, this article is hereby marked "advertisement" in accordance with 18 U.S.C. section 1734.

© 2000 by The American Society of Hematology

From the Laboratory of Medical Allergology, Allergy Unit, and the Laboratory for Tissue Typing, Department of Clinical Immunology, National University Hospital, Denmark; and the Department of Immunology, Anhui Medical University, People's Republic of China.

Supported by the Danish Allergy Research Center (T.J., S.Q., and A.M.), Directionen af Hovedstadens Sygehusfællensskab, Denmark (C.J.), Novo Nordisk A/S, Denmark (S.Q.), and the National Science Foundation of China (39870674).

Blood T cells expressing CXCR3 were mostly CD45RO<sup>+</sup> and generally expressed high levels of  $\beta$ 1 integrins. CXCR3 and CCR5 are markers for T cells associated with certain inflammatory reactions, particularly Th1-type reactions such as rheumatoid arthritis. CXCR3 and CCR5 appear to identify subsets of T cells in blood with a predilection for homing to these sites.<sup>3</sup> Interestingly, Mig was reported to promote tumor necrosis in vivo.<sup>10</sup> However, it is still unclear whether CXCR3 is expressed on CD34<sup>+</sup> hematopoietic progenitors and whether CXCR3 and its ligands ( $\gamma$ IP-10 and Mig) play any role in differentiation of CD34<sup>+</sup> hematopoietic progenitors.

### Materials and methods

#### Purification of CD34<sup>+</sup> hematopoietic progenitor cells

CD34<sup>+</sup> hematopoietic progenitor cells were purified as described elsewhere.<sup>11</sup> Briefly, umbilical cord blood samples were collected in accordance with institutional guidelines, and CD34<sup>+</sup> hematopoietic progenitors were isolated from mononuclear cells from cord blood. A positive-selection procedure using beads (Dynabeads M-450; Dynal A/S, Dynal, Norway) coated with anti-CD34 monoclonal antibody (mAb) was performed by following the manufacturer's instructions. The purity of the CD34<sup>+</sup> hematopoietic progenitors ranged from 90% to 99%, as determined by flow cytometry.

#### Flow cytometry

As previously described,<sup>12</sup> for detection of CXCR3, CD34<sup>+</sup> hematopoietic progenitors, either freshly isolated or stimulated with cytokines, were first incubated with a mouse antihuman CXCR3 mAb labeled with fluorescein isothiocyanate, conjugated (FITC) (49801.111; R&D Systems Europe Ltd, Abingdon, UK) or FITC-labeled CXCR4 mAb (12G5 [5 µg/mL]; R&D Systems Europe Ltd) as the positive control or matched isotype mouse IgG1 (5 µg/mL; Dako, Glostrup, Denmark) to which secondary FITC-labeled bovine antimouse mAb was consequently added. The incubation was done for 20 minutes in phosphate-buffered saline (PBS) containing 2% bovine serum albumin (BSA) and 0.1% sodium azide. The cells were then washed twice in staining buffer and incubated with a mouse antihuman CD34 class I phosphatidylethanolamine (PE)-labeled mAb (Immu409 [5 µg/mL]; Coulter-Immunoteck Co, Margency, France) in PBS containing 2% BSA and 0.1% sodium azide for 20 minutes and subsequently washed twice. For detection of integrins, CD34<sup>+</sup> hematopoietic progenitors stimulated with granulocytemacrophage colony-stimulating factor (GM-CSF) or with chemokine (for 8 hours) were incubated with a mouse antihuman antibody (5 µg/mL of either anti-CD29, 4B4; anti-CD49a, HP2B6; anti-CD49b, Gi9; anti-CD49c, C3VLA3; anti-CD49d, HP2/1; anti-CD49e, SAM1; or anti-CD49f, GoH3; Coulter-Immunoteck Co) for 20 minutes and then incubated with FITCconjugated bovine antimouse mAb (1:250 vol/vol, SeroTec Inc, Oxford, England) for 20 minutes. Subsequently, the cells were labeled with mouse antihuman CD34 class I PE-labeled mAb (Immu409) for 20 minutes as described above. All procedures were carried out at 4°C. The cells were then fixed with 1% paraformaldehyde. Analyses were performed with a flow cytometer (Coulter XL; Coulter Corporation, Miami, FL).

## Real-time quantitative reverse transcription–polymerase chain reaction (RT-PCR) assay

All real-time quantitative RT-PCR reactions were performed as described elsewhere.<sup>13-15</sup> Briefly, total RNA from CD34<sup>+</sup> hematopoietic progenitor cells ( $1 \times 10^6$ ; purity > 99%) was prepared by using a Quick Prep total RNA extraction kit (Pharmacia Biotech, Uppsala, Sweden), and any potential contaminating chromosomal DNA was digested with DNAase I according to the manufacturer's instructions. The RNA was reverse transcribed by using oligo(dT)<sub>12-18</sub> and Superscript II reverse transcriptase (Life Technologies, Grand Island, NY) according to the manufacturer's instructions. Reverse transcription was performed for 60 minutes at 37°C,

and any potential contaminating protein was denatured by incubation for 10 minutes at 95° C. The real-time quantitative PCR was performed in special optical tubes in a 96-well microtiter plate (Perkin Elmer Applied Biosystems, Foster City, CA) by using a sequence detector system (ABI Prism 7700; Perkin Elmer Applied Biosystems) according to the manufacturer's instructions. An SYBR green PCR core reagents kit (Perkin Elmer Applied Biosystems) was used to generate fluorescence during each PCR cycle by means of the 5' to 3' endonuclease activity of AmpliTaq Gold<sup>13</sup> to provide real-time quantitative PCR information. The CXCR3 genes were generated by connecting the following sequences of the specific primers (DNA Technology, Aarhus, Denmark): sense, 5'-GGAGCTGCTCAGAGTAAAT-CAC-3'; and antisense, 5'-GCACGAGTCACTCTCGTTTTC-3'.

All unknown complementary DNAs (cDNAs) were diluted to contain equal amounts of  $\beta$ -actin cDNA. The standards, "no template" controls, and unknown samples were added in a total volume of 50  $\mu$ L per reaction. PCR conditions were 2 minutes at 50°C, 10 minutes at 95°C, 40 cycles of 15 seconds at 95°C, and 60 seconds at 60°C for each amplification. Potential contaminants of the PCR product were digested by uracil-*N*-glycosylase (UNG), since deoxythymidine triphosphate is substituted with deoxyuridine triphosphate.<sup>14</sup> All PCR experiments were performed with a hot start. In the reaction system, UNG and AmpliTaq Gold (Perkin Elmer Applied Biosystems) were used according to the manufacturer's instructions.<sup>13,14</sup> To analyze data from the PCR products, 2 terms were used to express the results:  $\delta R_n$ , representing the normalized reporter signal minus the baseline signal established in the first few cycles of the PCR; and C<sub>T</sub> (threshold cycle), representing the PCR cycle in which an increase in reporter fluorescence signal above baseline levels could first be detected.

#### Northern blot analysis

As previously described,<sup>16</sup> total RNA from peripheral cells was prepared by using a Quick Prep total RNA extraction kit. Five micrograms of total RNA from each sample was electrophoresed under denaturing conditions, blotted on Nytran membranes (Schleicher & Schuell Inc, Keene, NH), and cross-linked by UV irradiation. cDNA probes were labeled with  $\alpha$ -phosphorus 32 (<sup>32</sup>P) deoxycytidine triphosphate. A CXCR3 cDNA probe was obtained by PCR amplification of the sequence shown above from total RNA of peripheral CD3<sup>+</sup> T lymphocytes from healthy adults. The membranes were hybridized overnight with 10<sup>6</sup> cpm/mL of <sup>32</sup>P-labeled probe and then washed with 0.2 × SSC (1 × SSC = 0.15 mol/L sodium chloride (NaCl) and 0.015 mol/L sodium citrate [pH 7.0]) and 0.1% sodium dodecyl sulfate before being autoradiographed.

#### In vitro chemotaxis assay

As described by Kim and Broxmeyer,17 chemotaxis and chemokinesis were assayed by a modification of the checkerboard assay. Fifty microliters of chemotaxis buffer (RPMI 1640, 0.5% BSA, and antibiotics) suspended with  $2 \times 10^5$  cells/mL was added to the upper well of the chamber, which was separated from the lower well by a polycarbonate, polyvinylpyrrolidonefree membrane (5 µm pore size) without a collagen coating (Nucleopore, Pleasanton, CA). Chemotaxis buffer was added to the lower chamber. Various amounts of chemoattractants were added to the chemotaxis buffer in the upper chamber, the lower chamber, or both, to form various chemoattractant concentration gradients to form various chemotactic gradients (positive gradient [0/+], negative gradient [+/0], and zero gradient [+/+ or 0/0]). All tests were performed in triplicate. Chambers were incubated at 37°C in 5% carbon dioxide (CO2) for 4 hours. Cells that migrated into the 3 lower chambers were collected and counted by using either a flow cytometer (Coulter XL) for 20 seconds at a high flow rate or a light microscope, by which identical results could be obtained. The cell migration was determined by calculating the percentage of input cells that migrated into the lower chamber.

#### Colony-forming cell assays

As described by Broxmeyer et al,<sup>18</sup> migrated or input cells were plated at concentrations of about 300 CD34<sup>+</sup> cells/mL in 35-mm plastic tissueculture dishes (Nunc, Roskilde, Denmark) containing 1 U/mL recombinant human erythropoietin (EPO; Sigma Chemical Co, St Louis, MO), 100 U/mL GM-CSF, and 100 U/mL IL-3, with and without various chemokines, in 1% methylcellulose culture medium containing 30% fetal-calf serum (FCS). After 7 to 10 days of incubation, burst-forming units–erythroid (BFU-E), colony-forming units–granulocyte-macrophage (CFU-GM), and mixed-cell units (colony-forming units–granulocyte-erythroid-macrophage-megakaryocyte [GEMM]), which respectively identify erythrocyte, granulocyte-macrophage, and multipotential progenitor cells, were scored in situ with an inverted microscope by using standard criteria for their identification.<sup>19</sup>

#### Adhesion assays

Adhesion assays were performed as described previously.12 Briefly, 96-well microtiter plates were coated with laminin (20 µg/mL; Sigma Chemical Co) in PBS for 1 hour at 37°C in a humidified atmosphere. The plates were washed with PBS and incubated with medium containing 0.2% BSA for 1 hour in 5% CO<sub>2</sub> to block nonspecific adhesion. Thereafter, single-cell suspensions were prepared in RPMI 1640 medium with 0.2% BSA ( $4 \times 10^5$ cells/mL), and yIP-10, Mig, or chemokine stromal cell-derived factor 1a (SDF-1a) was added (100 ng/mL). The cell suspension was added in triplicate to 96-well plates (100 µL per well) and incubated for 60 minutes at 37°C. To remove nonadherent cells, an 8-tipped manifold was used to aspirate all but about 50 µL of liquid from the wells by suspending the manifold at a uniform distance from the bottom of each well. The wells were then washed by carefully directing a stream of 0.2% BSA in PBS along the sides of the wells with the 8-tipped manifold after aspiration. Subsequently, the adherent cells were fixed with 1% formaldehyde and stained with 1% crystal violet. The crystal violet was then extracted by adding a 1:1 mixture of 0.1 mol/L sodium citrate (pH 4.2) and ethanol, and absorbency was read at 540 nm. Cells bound to collagen I (10 µg/mL) on separate wells were used to represent 100% attachment. Background cell adhesion to 2% BSA-coated wells was subtracted from all readings. For inhibition assays, cells were preincubated with different antibodies at 4°C for 30 minutes before the assay.

#### Aggregation assays

Aggregation assays were performed as described previously.<sup>12,20</sup> Briefly, the cells were added at a concentration of  $10^{6}$ /mL to culture plates with 24 wells. The chemokines were added to RPMI 1640 culture medium with 10% FCS. After 24 hours, the cells were observed, scored, and photographed by using a Leitz microphotography system. The following semiquantitative scoring method<sup>12,20</sup> was used: 0, no aggregation; 1+, less than 10% of cells aggregated; 2+, more than 50% of cells aggregated; 3+, up to 90% of cells small, loose clusters; and 4+, more than 90% of cells aggregated in large clusters.

# Polarization assay and immunofluorescence digital confocal microscopy

Immunofluorescence experiments were performed as described elsewhere.<sup>21</sup> Briefly, chemokines (100 ng/mL) were used in RPMI 1640 culture medium with 10% FCS in chemotaxis assays. The GM-CSF-stimulated CD34<sup>+</sup> hematopoietic progenitor cells were added at a concentration of 106/mL to a chemotactic chamber. After assay, the migrated GM-CSFstimulated CD34+ cells were collected and spun down on a slide, fixed with a mixture of methanol and acetone, and immersed in 1% BSA blocking buffer for 10 minutes to avoid nonspecific binding. Primary antibody, either anti-CXCR3 mAb (49801.111 [10 µg/mL]; R&D Systems Europe Ltd) or with isotype  $IgG_1$  (10  $\mu$ g/mL), was added, and after incubation overnight at 4°C, secondary FITC-labeled donkey antimouse antibody (1:250 vol/vol; Jackson ImmunoResearch Laboratories Inc, West Grove, PA) was added. Cells were observed with a fluorescence microscope (BX60; Olympus, Japan). Confocal microscopy analysis was performed with a confocal laser scanning system and an inverted microscope (LSMSIO; Zeiss, Germany). Images of serial cellular sections were acquired with a graphical user interface (Comos; Bio-Rad, Hercules, CA) as described previously.21

## Results

# Expression of CXCR3 on CD34 $^+$ progenitors is induced by GM-CSF

Flow cytometric analyses (Figure 1) showed only rare CXCR3<sup>+</sup> cell fractions in freshly isolated CD34<sup>+</sup> progenitors from cord blood (<3%; Figure 1B). After 36 hours of incubation with cytokine-free medium, there was no significant change in the CXCR3<sup>+</sup> cell fraction (data not shown). Interestingly, 10 ng/mL of GM-CSF significantly up-regulated the expression of CXCR3 on CD34<sup>+</sup> progenitors up to 97.6% (Figure 1C). Moreover, CXCR4 was constantly expressed in both freshly isolated CD34<sup>+</sup> progenitors (96.9%; Figure 1E) and GM-CSF–stimulated CD34<sup>+</sup> progenitors (98.2%; Figure 1F). Figures 1A and 1B show the results with the isotype controls for anti-CXCR3 and anti-CXCR4 antibodies, respectively.

In the kinetic study shown in Figure 2, a slightly increased expression (about 15.7%) of CXCR3 on GM-CSF-stimulated CD34<sup>+</sup> progenitors (Figure 2B) compared with freshly isolated cells (Figure 2A) was observed after 6 hours. The fractions of CXCR3<sup>+</sup> progenitors were 37.1%, 67.6%, 98.9%, and 99.4% after 16 hours (Figure 2C), 24 hours (Figure 2D), 36 hours (Figure 2E), and 48 hours (Figure 2F), respectively, of stimulation with GM-CSF. CXCR3 expression was not significantly changed in medium-cultured CD34<sup>+</sup> progenitors compared with freshly isolated cells at the different assessment times (data not shown).

#### CXCR3 messenger RNA (mRNA) expression in CD34<sup>+</sup> progenitors is up-regulated by GM-CSF

As shown in Figure 3, mRNA of CXCR3 was detected at low levels in freshly isolated CD34<sup>+</sup> hematopoietic progenitors from human umbilical blood. Compared with the amplification of standard DNA template ( $2.0 \times 10^4$  copies) with a housekeeping gene ( $\beta$ -actin), there were approximately  $6.3 \times 10^1$  copies for CXCR3 in the tested samples of freshly isolated CD34<sup>+</sup> progenitors. GM-CSF stimulation significantly up-regulated the expression of CXCR3 mRNA in CD34<sup>+</sup> progenitors. There were approximately  $4.3 \times 10^4$  copies for CXCR3 in the tested samples of



Figure 1. Double-color flow cytometric analysis of the distribution and modulation of CXC chemokine receptor (CXCR) 3 or CXCR4 on CD34<sup>+</sup> hematopoietic progenitors. The CD34<sup>+</sup> cells were either freshly isolated (A, B, D, and E) or stimulated with granulocyte-macrophage colony-stimulating factor (GM-CSF) for 36 hours (C and F). Panels A and D show results with isotype controls. The data are from a single experiment representative of 4 similar experiments performed.



Figure 2. Double-color flow cytometric analysis of the kinetics of CXCR3 expression on human cord blood CD34<sup>+</sup> hematopoietic progenitors. The cells were either freshly isolated (A) or stimulated with GM-CSF (10 ng/mL) for 6 hours (B), 16 hours (C), 24 hours (D), 36 hours (E), or 48 hours (F). The cells were then stained with fluorescein isothiocyanate, conjugated (FITC)–labeled anti-CXCR3 monoclonal antibody (mAb). The data are from a single experiment representative of 2 similar experiments performed.

GM-CSF-stimulated CD34<sup>+</sup> progenitors at 36 hours, whereas there were only about  $1.6 \times 10^2$  copies for CXCR3 in the tested samples of cultured nonstimulated CD34<sup>+</sup> progenitors at 36 hours (data not shown). A linear relation between C<sub>T</sub> and the log starting quantity of standard DNA template or target cDNA (CXCR3) was detected (data not shown). In all experiments, the correlation coefficient was approximately 0.94. CXCR3 mRNA expression in freshly isolated or GM-CSF-stimulated CD34<sup>+</sup> progenitors was confirmed by Northern blot assessment (Figure 3B). The upper panel in Figure 3B shows that freshly isolated CD34<sup>+</sup> progenitors expressed CXCR3 mRNA at low levels, whereas GM-CSFstimulated CD34<sup>+</sup> progenitors abundantly expressed CXCR3 mRNA. The lower panel shows that comparable total RNA amounts from different cells were added.



Figure 3. Assessment of messenger RNA (mRNA) expression in CD34<sup>+</sup> progenitors. (A) Plots of real-time detection and amplification of mRNA of CXCR3 in freshly isolated and GM-CSF-stimulated CD34<sup>+</sup> progenitors from human cord blood. Red plots represent the amplification of mRNA of CXCR3 in freshly isolated CD34<sup>+</sup> progenitors; blue plots represent the amplification of CXCR3 mRNA of GM-CSFstimulated CD34<sup>+</sup> progenitors; and black plots represent the amplification of standard DNA template ( $2.0 \times 10^4$  copies) with a housekeeping gene ( $\beta$ -actin). C<sub>T</sub>s were 23.5 for standard DNA template; 30.1 for mRNA of CXCR3 in freshly isolated CD34<sup>+</sup> progenitors; and 22.3 for mRNA of CXCR3 in GM-CSF-stimulated CD34<sup>+</sup> progenitors. The plots shown are representative of 2 similar experiments conducted. (B) CXCR3 mRNA Northern blot analysis of freshly isolated and GM-CSF-stimulated CD34<sup>+</sup> progenitors. Total RNA from different cells was isolated, electrophoresed, and blotted. The hybridization signals for CXCR3 mRNA from the cells are shown in the upper panel. Results with 28S recombinant RNAs (lower panel) confirmed that there were comparable amounts of loaded total RNA.

# $\gamma$ IP-10 and Mig induce GM-CSF–stimulated CD34+ progenitor chemotaxis by means of CXCR3

We examined the ability of  $\gamma$ IP-10 and Mig to induce chemotaxis in GM-CSF-stimulated CD34<sup>+</sup> progenitors and found that they induced significant chemotactic migration in the stimulated cells. As shown in Figure 4A (left),  $\gamma$ IP-10 and Mig did not induce chemotactic migration in freshly isolated CD34<sup>+</sup> progenitors. SDF-1 $\alpha$  was used as the positive control because it induces a strong



Figure 4. Assessments of migration of CD34<sup>+</sup> hematopoietic progenitors. (A) Migration of freshly isolated (left) and GM-CSF-stimulated (right) CD34+ hematopoietic progenitors toward interferon (IFN) γ-inducible protein 10 (γIP-10) (open bars), monokine induced by IFN-y (Mig) (black bars), and chemokine stromal cell-derived factor 1  $\alpha$  (SDF-1  $\alpha$ ) (gray bars). All results are expressed as the mean  $\pm$  SD percentage of input cells that migrated (n = 5) and are based on triplicate determinations of chemotaxis for each concentration of chemokines. An asterisk indicates a significant difference from the medium control (P < .03, except in one case, in which P < .04). (B) Checkerboard analysis for GM-CSF-stimulated CD34<sup>+</sup> hematopoietic progenitors toward vIP-10 (right) and Mig (right). Chemokines were applied in the upper chamber, the lower chamber, or both. The concentrations of chemokines applied in the lower chambers were 0, 1, 10, 100, and 1000 ng/mL (indicated as 1, 2, 3, 4, and 5, respectively). The data are from one representative experiment of 2 performed. (C) Specific chemotactic activity of vIP-10 (right), Mig (middle), and SDF-1 $\alpha$  (right) for subtypes of colony-forming, GM-CSF-stimulated CD34<sup>+</sup> hematopoietic progenitors from cord blood. CD34+ cells attracted to different chemokines at the indicated concentrations were assayed for colony-forming cells and the results expressed as the mean  $\pm$  SD percentage of input cells that migrated (n = 3). Open bars, black bars, and gray bars indicate burst-forming units-erythroid, colony-forming units-granulocyte-macrophage, and colony-forming units-granulocyte-erythroidmacrophage-megakaryocyte, respectively. An asterisk indicates a significant difference from the medium control (P < .04). (D) Flow cytometry analysis for CD34<sup>+</sup> cells before (left) and after (right) migration toward vIP-10. The cells were dual stained: percentages indicate positive or negative cells.

CD34<sup>+</sup> progenitor chemotactic migration. As shown in Figure 4A (right),  $\gamma$ IP-10 and Mig induced chemotactic migration in GM-CSF– stimulated CD34<sup>+</sup> progenitors, yielding typical bell-shaped, dosedependent chemotaxis response curves. The optimal chemotactic concentration of both  $\gamma$ IP-10 and Mig was 100 ng/mL (56.4% ± 8.5% and 51.7% ± 9.8% of input cells, respectively). SDF-1 $\alpha$ , the positive control, induced a similar GM-CSF–stimulated CD34<sup>+</sup> progenitor chemotactic migration (75.6% ± 9.6% of input cells). Spontaneous migration (medium control) of different CD34<sup>+</sup> progenitors was less then 5% (Figure 3A).  $\gamma$ IP-10 and Mig did not induce significant chemotactic migration of CD34<sup>+</sup> progenitor cells cultured in medium alone (data not shown).

To confirm that the observed GM-CSF-stimulated CD34<sup>+</sup> progenitor chemotaxis was indeed induced by  $\gamma$ IP-10 and Mig by means of CXCR3, we used anti-CXCR3 mAb to block the GM-CSF-stimulated CD34<sup>+</sup> progenitor chemotactic activity of  $\gamma$ IP-10 and Mig. The anti-CXCR3 mAb completely blocked the chemotaxis of GM-CSF-stimulated CD34<sup>+</sup> progenitors toward  $\gamma$ IP-10 and Mig (data not shown), whereas it had no effect on the chemotaxis of GM-CSF-stimulated CD34<sup>+</sup> progenitors toward SDF-1 $\alpha$  (data not shown). The isotype antibody had no blocking effect (data not shown).

To address the question of whether enhanced migration of GM-CSF–stimulated CD34<sup>+</sup> progenitors toward  $\gamma$ IP-10 and Mig is due to directional migration (chemotaxis), random migration (chemokinesis), or both,<sup>17</sup> we performed checkerboard chemotaxis assays in the presence of  $\gamma$ IP-10 or Mig in the lower chamber, the upper chamber, or both, in a combined manner. As the  $\gamma$ IP-10 or Mig concentration in the upper chamber increased, migration of GM-CSF–stimulated CD34<sup>+</sup> progenitors toward the lower chamber decreased (Figure 4B). When the chemotactic gradient became negative or zero, migration of GM-CSF–stimulated CD34<sup>+</sup> progenitors toward  $\gamma$ IP-10 (Figure 4B, left) or Mig (Figure 4B, right) to the lower chamber decreased to background levels, demonstrating that enhanced motility of the cells toward both chemokines was due to chemotaxis rather than chemokinesis.

To determine whether  $\gamma$ IP-10, Mig, and SDF-1 $\alpha$  had specificity to attract certain types of CD34<sup>+</sup> progenitors with respect to colony formation, we assayed the migrated CD34<sup>+</sup> progenitors in methylcellulose medium containing growth factors (GM-CSF, IL-3, EPO, and various chemokines). Interestingly, at a concentration of 100 ng/mL, yIP-10 (Figure 4C, left) and Mig (Figure 4C, middle) attracted mainly (and in significant numbers) CFU-GM in cord blood CD34<sup>+</sup> progenitors. The chemotactic activity for other cord blood progenitors, such as BFU-E and the more immature CFU-GEMM, was weak and not significant. However, the attractions of CFU-GM and BFU-E to SDF-1a (Figure 4C, right) were significant and equal, whereas the attraction of CFU-GEMM was weak and not significant. This observation can be explained by the conditions in our experiments.<sup>19</sup> We also examined the purity of GM-CSF-stimulated CD34<sup>+</sup> progenitors before and after migration toward yIP-10 (Figure 4D) and Mig (data not shown). The identical purity of GM-CSF-stimulated CD34<sup>+</sup> cells before (left) and after (right) migration excludes the possibility that contaminated T lymphocytes preferably migrated toward yIP-10.

# $\gamma \text{IP-10}$ and Mig induce GM-CSF–stimulated CD34+ progenitor adhesion by means of CXCR3

We examined the ability of  $\gamma$ IP-10 and Mig to induce GM-CSF– stimulated CD34<sup>+</sup> progenitor adhesion. Figure 5A shows that neither  $\gamma$ IP-10 nor Mig induced adhesion in freshly isolated CD34<sup>+</sup> progenitors (< 15% for both). SDF-1 $\alpha$  was used as the



Figure 5. Adhesion of CD34<sup>+</sup> hematopoietic progenitors from human cord blood induced by  $\gamma$ IP-10 (indicated by 1), Mig (2), or SDF-1 $\alpha$  (3). The CD34<sup>+</sup> progenitor cells were either freshly isolated (A), stimulated with GM-CSF (B), or stimulated with GM-CSF and incubated with anti-CXCR3 mAb (C). The cells were either freshly isolated from cord blood from a normal birth, preincubated with GM-CSF for 36 hours only, or preincubated with GM-CSF and then incubated with CXCR3 mAb (5  $\mu$ g/mL) for 60 minutes at room temperature before the adhesion assay. All chemokines were applied at a concentration of 100 ng/mL. The data represent mean values from at least 3 experiments performed. Results are expressed as the mean  $\pm$  SD percentage of adherent cells and are based on triplicate determinations of adhesion for each chemokine used.

positive control because it induces a strong CD34<sup>+</sup> progenitor adhesion (about 70%). Figure 5B shows that  $\gamma$ IP-10 and Mig induced a strong adhesion in GM-CSF–stimulated CD34<sup>+</sup> progenitors. About 68% and 75% of cells, respectively, had adhered 60 minutes after the addition of  $\gamma$ IP-10 or Mig for (100 ng/mL). SDF-1 $\alpha$ , the positive control, induced a similar GM-CSF– stimulated CD34<sup>+</sup> progenitor adhesion (65%).  $\gamma$ IP-10 and Mig did not induce significant adhesion of medium-cultured CD34<sup>+</sup> progenitor cells (data not shown).

To confirm that the observed GM-CSF–stimulated CD34<sup>+</sup> progenitor adhesion was induced by  $\gamma$ IP-10 and Mig by means of CXCR3, we used anti-CXCR3 mAb to block the GM-CSF–stimulated CD34<sup>+</sup> progenitor adhesive activity of  $\gamma$ IP-10 and Mig. The mAb completely blocked the adhesion of GM-CSF–stimulated CD34<sup>+</sup> progenitors induced by  $\gamma$ IP-10 and Mig (Figure 5C) (< 12%, respectively), whereas it had no effect on the adhesion of GM-CSF–stimulated CD34<sup>+</sup> progenitors induced by SDF-1 $\alpha$  (about 50%). The isotype antibody had no blocking effect (data not shown).

# $\gamma IP$ -10 and Mig induce GM-CSF–stimulated CD34 $^+$ progenitor aggregation by means of CXCR3

To examine whether  $\gamma$ IP-10 and Mig play an additional role in GM-CSF–stimulated CD34<sup>+</sup> progenitor adhesion, we performed aggregation tests of CD34<sup>+</sup> progenitors stimulated with  $\gamma$ IP-10, Mig, or SDF-1 $\alpha$ . The results shown in Figure 6 indicate that  $\gamma$ IP-10 and Mig do not induce aggregation in freshly isolated CD34<sup>+</sup> progenitors (Figure 6A and 6B, scored as 0). SDF-1 $\alpha$  was used as the positive control because it induces a strong CD34<sup>+</sup> progenitor aggregation (Figure 6C, scored as 4+).  $\gamma$ IP-10 and Mig induced a strong aggregation in GM-CSF–stimulated CD34<sup>+</sup> progenitors (Figure 6D, scored as 3+).  $\gamma$ IP-10 and Mig did not induce significant aggregation of medium-cultured CD34<sup>+</sup> progenitor cells (data not shown).

To confirm that the observed GM-CSF–stimulated CD34<sup>+</sup> progenitor aggregation was induced by  $\gamma$ IP-10 and Mig by means of CXCR3, we used anti-CXCR3 mAb to block the GM-CSF– stimulated CD34<sup>+</sup> progenitor aggregation activity of  $\gamma$ IP-10 and Mig. The mAb completely blocked the aggregation of GM-CSF– stimulated CD34<sup>+</sup> progenitors induced by  $\gamma$ IP-10 and Mig (Figure 6G and 6H, scored as 0), whereas it had no effect on the aggregation of GM-CSF–stimulated CD34<sup>+</sup> progenitors induced by SDF-1 $\alpha$  (Figure 6I, scored as 2+). The isotype antibody had no blocking effect (Figure 6J, scored as 2+; Figure 6K, scored as 4+; and Figure 6L, scored as 3+).

# Role of integrins in GM-CSF–stimulated CD34<sup>+</sup> progenitor adhesion induced by $\gamma$ IP-10 and Mig

We next examined the expression of some integrins on GM-CSF– stimulated CD34<sup>+</sup> progenitors induced by  $\gamma$ IP-10 or Mig. Because



Figure 6. Aggregation of CD34<sup>+</sup> hematopoietic progenitors. Aggregation induced by  $\gamma$ IP-10 (A, D, G, and J), Mig (B, E, H, and K), or SDF-1 $\alpha$  (C, F, I, and L) is shown. CD34<sup>+</sup> hematopoietic progenitors from human cord blood were either freshly isolated (A-C), stimulated with GM-CSF (D-F), incubated with anti-CXCR3 mAb and stimulated with GM-CSF (G-I), or incubated with isotype mAb and stimulated with GM-CSF (J-L). Arrows indicate the aggregation clusters. The cells were photographed at ×100 magnification. Aggregation in A, B, G, and H was estimated to be 0; that in D, E, F, I, J, and L to be 2+ to 3+; and that in C and K to be 4+. The photographs are representative of 3 similar experiments.



Figure 7. Double-color flow cytometric analysis of the distribution of different integrins on CD34<sup>+</sup> hematopoietic progenitors. The CD34<sup>+</sup> cells were either stimulated with GM-CSF for 36 hours (A-G) or stimulated with GM-CSF and then with  $\gamma$ IP-10 for 8 hours (H-N). The cells were isolated from cord blood from a normal birth and either preincubated with GM-CSF (10 ng/mL) for 36 hours only or preincubated with GM-CSF and then stimulated with  $\gamma$ IP-10 (100 ng/mL) for 8 hours. Subsequently, the cells were stained with the indicated anti-integrin mAbs. The data are from a single experiment representative of 4 similar experiments performed.

we had already demonstrated that  $\gamma$ IP-10 and Mig each produced a significant induction of GM-CSF-stimulated CD34<sup>+</sup> progenitor adhesion and aggregation, we conducted additional experiments to investigate whether yIP-10 or Mig would enhance the expression of integrins on GM-CSF-stimulated CD34<sup>+</sup> progenitors. The results of the flow cytometric analyses shown in Figure 7 indicate that yIP-10 can significantly increase the expression of certain integrins on GM-CSF-stimulated CD34<sup>+</sup> progenitors. On average, there were  $32.5\% \pm 5.1\%$  CD49a<sup>+</sup> cells on GM-CSF-stimulated CD34<sup>+</sup> progenitors (Figure 7B),  $31.5\% \pm 2.2\%$  CD49c<sup>+</sup> cells (Figure 7C), 36.2%  $\pm$  7.8% CD49d^+ cells (Figure 7D), 42%  $\pm$ 10.2% CD49e<sup>+</sup> cells (Figure 7E), 41.5% ± 12.5% CD49f<sup>+</sup> cells (Figure 7F), and 39%  $\pm$  7.7% CD49b<sup>+</sup> cells (Figure 7G). After stimulation with  $\gamma$ IP-10 (100 ng/mL) for 8 hours, the expression of CD49a and CD49b was selectively and substantially up-regulated. On average, there were 84%  $\pm$  2.8% CD49a<sup>+</sup> cells on  $\gamma$ IP-10– cultured GM-CSF-stimulated CD34<sup>+</sup> progenitors (Figure 7I) and 74.2%  $\pm$  4.7% CD49b<sup>+</sup> cells (Figure 7J). Statistical analysis for the 4 experiments showed a significant difference in CD49a and CD49b expression in GM-CSF-stimulated CD34<sup>+</sup> cells compared with GM-CSF-stimulated CD34<sup>+</sup> cells or  $\gamma$ IP-10-stimulated CD34<sup>+</sup> cells (both P < .003; n = 4). On average, there were 45.7%  $\pm$  4.5% CD49c^+ cells on  $\gamma IP\text{-}10\text{-}cultured,$  GM-CSFstimulated CD34<sup>+</sup> progenitors (Figure 7K), 45.5%  $\pm$  4.7% CD49d<sup>+</sup> cells (Figure 7L),  $45.2\% \pm 3.8\%$  CD49e<sup>+</sup> cells (Figure 7M), and  $46.2\% \pm 5.1\%$  CD49f<sup>+</sup> cells (Figure 7N). There was no significant difference in CD49c, CD49d, CD49e, or CD49f cell expression in GM-CSF-stimulated CD34+ cells compared with GM-CSFstimulated and  $\gamma$ IP-10-stimulated CD34<sup>+</sup> cells (all P > .05; n = 4).



Figure 8. Adhesion of CD34<sup>+</sup> hematopoietic progenitors from human cord blood. Adhesion induced by  $\gamma$ IP-10 (A), Mig (B), and SDF-1 $\alpha$  (C) is shown. The cells were either stimulated with GM-CSF alone (indicated by 1) or stimulated with GM-CSF and then incubated with anti-CD49a mAb (2), anti-CD49b mAb (3), anti-CD49a and arti-CD49b. The cells were preincubated with different mAbs (5  $\mu$ g/mL) for 60 minutes at room temperature before the adhesion assay. All chemokines were applied at a concentration of 100 ng/mL. The data represent mean values from at least 3 experiments performed. Results are expressed as the mean  $\pm$  SD percentage of adherent cells and are based on a triplicate determination of adhesion for each antibody used.

Figures 7A and 7H show the results with isotype controls for a-CD49a and a-CD49b.

We also observed that Mig (100 ng/mL) selectively and substantially up-regulated the expression of CD49a and CD49b on GM-CSF-stimulated CD34<sup>+</sup> progenitors (data not shown). In addition, we examined the expression of integrins on freshly isolated CD34<sup>+</sup> hematopoietic progenitors. There was no significant difference between freshly isolated and GM-CSF-stimulated CD34<sup>+</sup> hematopoietic progenitors in the expression of certain integrins (data not shown).

Because we found a selective and substantial up-regulation of expression of CD49a and CD49b on GM-CSF–stimulated CD34<sup>+</sup> progenitors induced by  $\gamma$ IP-10 or Mig, we explored the role of these integrins in the adhesion of GM-CSF–stimulated CD34<sup>+</sup> progenitors induced by IP-10 or Mig. As shown in Figure 8A,  $\gamma$ IP-10 induced significant adhesion in GM-CSF–stimulated CD34<sup>+</sup>

progenitors in the absence of anti-integrin antibodies (about 90%). Anti-CD49a or anti-CD49b mAb significantly inhibited the adhesion of GM-CSF-stimulated CD34<sup>+</sup> progenitors induced by yIP-10 (about 20% for each). However, the combination of anti-CD49a mAb and anti-CD49b mAb almost completely abolished the adhesion of GM-CSF-stimulated CD34<sup>+</sup> progenitors induced by  $\gamma$ IP-10 (< 10%), whereas isotype IgG1 had no inhibitory effect on adhesion (about 85%). A similar blocking effect on Mig-induced adhesion of GM-CSF-stimulated CD34+ progenitors was observed by using a combination of anti-CD49a mAb and anti-CD49b mAb (Figure 8B), whereas isotype IgG1 had no inhibitory effect on adhesion. In some experiments in which fibronectin replaced laminin as the substratum, similar blocking effects of anti-CD49a mAb and anti-CD49b mAb were observed (data not shown). Interestingly, neither anti-CD49a mAb, anti-CD49b mAb, nor isotype IgG1 significantly prohibited the adhesion of GM-CSFstimulated CD34<sup>+</sup> progenitors induced by SDF-1 $\alpha$  (Figure 8C). The results shown in Figure 8 indicate that  $\gamma$ IP-10 or Mig induces adhesion of GM-CSF-stimulated CD34+ progenitors by means of up-regulation of some adhesion molecules (certain integrins, CD49a and CD49b). CD49a and CD49b may be important ligands in the pathway of this reaction.

#### γIP-10 and Mig induce CXCR3 polarization on GM-CSF-stimulated CD34<sup>+</sup> progenitors

Because cell migration induced by chemoattractant gradients requires cell polarization and relative receptor redistribution, we assessed the cellular distribution of CXCR3 during stimulation of GM-CSF-stimulated CD34<sup>+</sup> progenitors with yIP-10, Mig, or SDF-1a. As shown in Figure 9, CXCR3 was evenly distributed throughout the GM-CSF-stimulated CD34<sup>+</sup> progenitors (Figure 9A). On stimulation of GM-CSF-stimulated CD34<sup>+</sup> progenitors with yIP-10 (Figure 9B) or Mig (Figure 9C), a rapid redistribution of CXCR3 took place that resulted in movement of the receptors to the leading edge of the cells. The insets in Figure 9 show the clusters of CXCR3 oriented in a certain direction. Figure 9D shows the CXCR3 evenly distributed throughout the GM-CSF-stimulated CD34<sup>+</sup> progenitors, although the cells migrated through the gradient of SDF-1a, indicating that the polarization of CXCR3 on the GM-CSF-stimulated CD34<sup>+</sup> progenitors is receptor-ligand ( $\gamma$ IP-10 and Mig) specific.



Figure 9. Assessment of cellular distribution of CXCR3 after stimulation. (A) CXCR3 distribution on GM-CSF–stimulated CD34<sup>+</sup> hematopoietic progenitors before the chemotaxis assay. After stimulation, the CXCR3 receptors redistributed to the leading edge of the polarized migrating GM-CSF–stimulated CD34<sup>+</sup> hematopoietic progenitors, toward  $\gamma$ IP-10 (B), Mig (C), or SDF-1 $\alpha$  (D). The CD34<sup>+</sup> cells were stimulated with GM-CSF for 36 hours and then subjected to the chemotaxis assay. The migrating cells were collected and stained and then photographed under epifluorescence conditions (original magnification ×1200; bar, 12 µm).

## Discussion

The mechanisms and specific molecules involved in the mobilization of hematopoietic progenitor cells from the hematopoietic organs into peripheral blood, in the homing of hematopoietic progenitor cells, and in their trafficking through the hematopoietic organs during maturation are unclear. Hematopoietic progenitor cells express CXCR422 and migrate toward a gradient of the chemotactic factor SDF-1a.<sup>23</sup> CXCR3 is not broadly expressed in cells of the immune system: 40% of resting T lymphocytes and low numbers of B cells and natural killer cells were found to stain positive for CXCR3.24 Interestingly, pretreatment with IL-2 resulted in cultures of fully responsive, CXCR3-positive T lymphocytes.24 CXCR3 has been identified as the receptor for yIP-10, Mig, secondary lymphoidtissue chemokine (SLC) (6Ckine and I-TAC.5,6,25 CXCR3 and CCR5 are preferentially expressed on human Th1 lymphocytes, whereas CCR4 and CCR3 are preferentially expressed on Th2 lymphocytes.26 Lymphocytic cells and cerebrospinal fluid T cells from patients with active multiple sclerosis are significantly enriched with cells expressing CXCR3 or CCR5.27 yIP-10 plays a role in the epidermotropism of cutaneous T-cell lymphoma.<sup>28</sup> yIP-10 and Mig appear to target stimulated lymphocytes specifically, but have no activity on either neutrophils or monocytes.<sup>29</sup> The gene for CXCR3 is on human chromosome Xq13, suggesting unique functions for this receptor and its ligands beyond their established role in T-cell-dependent immunity.24

The function of adhesion molecules on the surface of leukocytes is critically regulated by activating events triggered by chemoattractants binding to specific receptors on the leukocytes.<sup>30</sup> Hematopoietic progenitor cells home to the extravascular compartment of the bone marrow during transplantation. Multipotential and self-renewing hematopoietic cells migrate to the bone marrow from the fetal liver during fetal development. In the other direction, hematopoietic progenitor cells are mobilized from the bone marrow to the peripheral blood in response to injected cytokines such as GM-CSF, granulocyte colony-stimulating factor (G-CSF), and Steel factor (SLF).<sup>31</sup>

How do multiple chemoattractants cooperate in directing the migration of hematopoietic progenitor cells for homing and peripheral blood mobilization? SDF-1 and SLF were found to cooperate in attracting MO7e cells, cord blood cells, and bone marrow CD34<sup>+</sup> cells.<sup>32</sup> SDF-1 and SLF, along with other unidentified bone marrow chemoattractants, may be involved cooperatively in the migration of hematopoietic progenitor cells to the bone marrow and in preventing spontaneous mobilization of hematopoietic progenitor cells out of the bone marrow.<sup>32</sup> Chemokines play a direct role in mobilization of hematopoietic stem cells, although cytokines have other functions, such as proliferation, modulation of adhesion molecules, and alteration of the blood–bone marrow barrier. For example, G-CSF and GM-CSF appear to have no chemotactic or chemokinetic effects on hematopoietic stem cells, whereas SLF, IL-3, and IL-11 have been reported to chemoattract murine hematopoietic progenitor cells.<sup>33</sup>

In the current study, we demonstrated that  $\gamma$ IP-10 and Mig activate hematopoietic progenitors to chemotaxis and adhesion on induction of CXCR3 by GM-CSF. To our knowledge, this is the first report of CXCR3 expression on human hematopoietic progenitors induced by a growth factor–like cytokine, and it provides the first direct evidence of biologic activity of hematopoietic progenitors induced by CXC chemokines  $\gamma$ IP-10 and Mig. A rather complex picture is beginning to form regarding how hematopoietic progenitors selectively enter different hematopoietic organs, nest on sites, differentiate, and further transmigrate into functional destinations to cause physiologic and pathophysiologic events during continuous interaction with chemokines and cytokines. Our finding that  $\gamma$ IP-10 and Mig do not induce freshly isolated CD34<sup>+</sup> progenitor chemotaxis and adhesion are in agreement with the results of Kim and Broxmeyer,<sup>34</sup> who observed that the CXCR3 ligands  $\gamma$ IP-10 and Mig did not induce chemotaxis of freshly isolated CD34<sup>+</sup> hematopoietic progenitors but that another CXCR3 ligand, SLC, is a potential trafficking factor for hematopoietic progenitors. Interestingly, we found that GM-CSF is a strong inducer of expression of CXCR3 on progenitors at the mRNA and protein levels. Only GM-CSF–stimulated CD34<sup>+</sup> progenitors had chemotactic activity toward  $\gamma$ IP-10 and Mig. These results indicate that hematopoietic progenitors express different chemokine receptors at different stages of maturation and these receptors have different affinity to their ligands at the different stages. During the maturation of hematopoietic progenitors, GM-CSF plays a important role in the induction of expression of chemokine receptors and increases their affinity.

Transplanted human hematopoietic cells must retain specific adhesive capacity to interact with the vascular endothelium of the bone marrow.35 Murine hematopoietic progenitor cells interact in vivo with both P-selectin and E-selectin on vascular endothelium of bone marrow.33 Optimal recruitment of hematopoietic progenitor cells to the bone marrow requires the combined action of both selectins and vascular cell adhesion molecule 1.36,37 The rolling of hematopoietic progenitor cells on bone marrow endothelium may be accompanied by a coordinated sequence of adhesive and activation events leading to cell arrest, a critical step in the successful extravasation of blood-borne cells to extravascular sites.38,39 Leukocyte functionassociated molecule 1 (LFA-1) and very late antigen (VLA) 4 are constitutively expressed by cord blood CD34<sup>+</sup> hematopoietic progenitors in inactive forms.40 LFA-1 binding to endothelial intercellular adhesion molecule 1 is the principal adhesive interaction that mediates the firm arrest of leukocytes on vascular endothelium.40-42 Murine hematopoietic progenitors roll in vivo along bone marrow microvessels that display selectins and integrins.<sup>43</sup> SDF-1 is expressed at high levels on bone marrow endothelium and stimulates a firm integrin-dependent adhesion of circulating CD34+ progenitors to the bone marrow microvasculature.44

We demonstrated that up-regulation of  $\gamma$ IP-10–induced and Mig-induced integrins (CD49a and CD49b, also called VLA-1 and VLA-2) plays a crucial role in the adhesion of GM-CSF–stimulated CD34<sup>+</sup> progenitors. This finding suggests that  $\gamma$ IP-10, Mig, and their receptor, CXCR3, take part in the multistep mobilization of hematopoietic cells of migration to or from the bone marrow during their development, as well as mutual-direction mobilization between the bone marrow and the peripheral blood during immune reactions. Adhesion molecules, including integrins, are important elements in these processes.

In summary, we found that  $\gamma$ IP-10 and Mig, by means of CXCR3 induced by GM-CSF, activate CD34<sup>+</sup> progenitor cells to chemotaxis and adhesion. This finding provides a useful insight into novel mechanisms of action of  $\gamma$ IP-10 and Mig that may be especially important in the cytokine/chemokine environment for the physiologic and pathophysiologic events of mobilization and homing during proliferation, differentiation, and maturation of hematopoietic progenitor cells.

## Acknowledgments

We thank Gitte Pedersen, Anne Corfitz, Ulla Minuva, and Tina Mortensen for excellent technical assistance and Dr Fritz von Bülow, Institute of Anatomy, Panum Institute, University of Copenhagen, Denmark, for assistance in immunofluorescence digital confocal microscopy.

## References

- Trentin L, Agostini C, Facco M, et al. The chemokine receptor CXCR3 is expressed on malignant B cells and mediates chemotaxis. J Clin Invest. 1999;104:115.
- Cella M, Jarrossay D, Facchetti F, et al. Plasmacytoid monocytes migrate to inflamed lymph nodes and produce large amounts of type I interferon. Nat Med. 1999;5:919.
- Qin S, Rottman JB, Myers P, et al. The chemokine receptors CXCR3 and CCR5 mark subsets of T cells associated with certain inflammatory reactions. J Clin Invest. 1998;101:746.
- Weng Y, Siciliano SJ, Waldburger KE, et al. Binding and functional properties of recombinant and endogenous CXCR3 chemokine receptors. J Biol Chem. 1998;273:18288.
- Soto H, Wang W, Strieter RM, et al. The CC chemokine 6Ckine binds the CXC chemokine receptor CXCR3. Proc Natl Acad Sci U S A. 1998;95: 8205.
- Jenh CH, Cox MA, Kaminski H, et al. Species specificity of the CC chemokine 6Ckine signaling through the CXC chemokine receptor CXCR3: human 6Ckine is not a ligand for the human or mouse CXCR3 receptors. J Immunol. 1999;162: 3765.
- Piali L, Weber C, LaRosa G, et al. The chemokine receptor CXCR3 mediates rapid and shear-resistant adhesion-induction of effector T lymphocytes by the chemokines IP10 and Mig. Eur J Immunol. 1998;28:961.
- Sallusto F, Lenig D, Mackay CR, Lanzavecchia A. Flexible programs of chemokine receptor expression on human polarized T helper 1 and 2 lymphocytes. J Exp Med. 1998;187:875.
- Sgadari C, Farber JM, Angiolillo AL, et al. Mig, the monokine induced by interferon-gamma, promotes tumor necrosis in vivo. Blood. 1997;89: 2635.
- Caux C, Massacrier C, Vanbervliet B, et al. CD34<sup>+</sup> hematopoietic progenitors from human cord blood differentiate along two independent dendritic cell pathways in response to granulocyte-macrophage colony-stimulating factor plus tumor necrosis factor alpha. II. Functional analysis. Blood. 1997;90:1458.
- Jinquan T, Quan S, Feili G, Larsen CG, Thestrup-Pedersen K. Eotaxin activates T cells to chemotaxis and adhesion only if induced to express CCR3 by IL-2 together with IL-4. J Immunol. 1999;162:4285.
- Heid CA, Stevens J, Livak KJ, William PM. Real time quantitative PCR. Genome Res. 1996;6:986.
- Kruse N, Pette M, Toyka K, Rieckmann P. Quantification of cytokine mRNA expression by RT PCR in samples of previously frozen blood. J Immunol Methods. 1997;210:195.
- Jinquan T, Quan S, Jacobi HH, et al. Expression of the nuclear factors of activated T cells in eosinophils: regulation by IL-4 and IL-5. J Immunol. 1999;163:21.

- Sica A, Saccani A, Borsatti A, et al. Bacterial lipopolysaccharide rapidly inhibits expression of C-C chemokine receptors in human monocytes. J Exp Med. 1997;185:969.
- Kim CH, Broxmeyer HE. In vitro behavior of hematopoietic progenitor cells under the influence of chemoattractants: stromal cell-derived factor-1, Steel factor, and the bone marrow environment. Blood. 1998;91:100.
- Broxmeyer HE, Hangoc G, Cooper S, et al. Growth characteristics and expansion of human umbilical cord blood and estimation of its potential for transplantation in adults. Proc Natl Acad Sci U S A. 1992;89:4109.
- Kim CH, Pelus LM, White JR, Broxmeyer HE. Macrophage-inflammatory protein-3 beta/EBI1ligand chemokine/CK beta-11, a CC chemokine, is a chemoattractant with a specificity for macrophage progenitors among myeloid progenitor cells. J Immunol. 1998;161:2580.
- Bernard G, Zoccola D, Ticchioni M, Breittmayer JP, Aussel C, Bernard A. Engagement of the CD45 molecule induces homotypic adhesion of human thymocytes through a LFA-1/ICAM-3-dependent pathway. J Immunol. 1994;152:5161.
- Serrador JM, Alonso-Lebrero JL, del Pozo MA, et al. Moesin interacts with the cytoplasmic region of intercellular adhesion molecule-3 and is redistributed to the uropod of T lymphocytes during cell polarization. J Cell Biol. 1997;138:1409.
- Aiuti A, Turchetto L, Cota M, et al. Human CD34<sup>+</sup> cells express CXCR4 and its ligand stromal cellderived factor-1. Implications for infection by Tcell tropic human immunodeficiency virus. Blood. 1999;94:62.
- Aiuti A, Webb IJ, Bleul C, Springer T, Gutierrez-Ramos JC. The chemokine SDF-1 is a chemoattractant for human CD34<sup>+</sup> hematopoietic progenitor cells and provides a new mechanism to explain the mobilization of CD34<sup>+</sup> progenitors to peripheral blood. J Exp Med. 1997;185:111.
- Loetscher M, Loetscher P, Brass N, Meese E, Moser B. Lymphocyte-specific chemokine receptor CXCR3: regulation, chemokine binding and gene localization. Eur J Immunol. 1998;28:3696.
- Jenh CH, Cox MA, Kaminski H, et al. Species specificity of the CC chemokine 6Ckine signaling through the CXC chemokine receptor CXCR3: human 6Ckine is not a ligand for the human or mouse CXCR3 receptors. J Immunol. 1999;162: 3765.
- Bonecchi R, Bianchi G, Bordignon PP, et al. Differential expression of chemokine receptors and chemotactic responsiveness of type 1 T helper cells (Th1s) and Th2s. J Exp Med. 1998;187:129.
- Sorensen TL, Tani M, Jensen J, et al. Expression of specific chemokines and chemokine receptors in the central nervous system of multiple sclerosis patients. J Clin Invest. 1999;103:807.
- Sarris AH, Daliani D, Ulmer R, et al. Interferoninducible protein 10 as a possible factor in the pathogenesis of cutaneous T-cell lymphomas. Clin Cancer Res. 1997;3:169.
- 29. Farber JM. Mig and IP-10: CXC chemokines that target lymphocytes. J Leukoc Biol. 1997;61:246.
- 30. Dieu MC, Vanbervliet B, Vicari A, et al. Selective recruitment of immature and mature dendritic

cells by distinct chemokines expressed in different anatomic sites. J Exp Med. 1998;188:373.

- Donahue RE, Kirby MR, Metzger ME, et al. Peripheral blood CD34<sup>+</sup> cells differ from bone marrow CD34<sup>+</sup> cells in Thy-1 expression and cell cycle status in nonhuman primates mobilized or not mobilized with colony-stimulating factor and/or stem cell factor. Blood. 1996;87:1644.
- Kim CH, Qu CK, Hangoc G, et al. Abnormal chemokine-induced responses of immature and mature hematopoietic cells from motheaten mice implicate the protein tyrosine phosphatase SHP-1 in chemokine responses. J Exp Med. 1999;190: 681.
- Okumura N, Tsuji K, Ebihara Y, et al. Chemotactic and chemokinetic activities of stem cell factor on murine hematopoietic progenitor cells. Blood. 1996;87:4100.
- Kim CH, Broxmeyer HE. SLC/exodus2/6Ckine/ TCA4 induces chemotaxis of hematopoietic progenitor cells: differential activity of ligands of CCR7, CXCR3, or CXCR4 in chemotaxis vs. suppression of progenitor proliferation. J Leukoc Biol. 1999;66:455.
- Quesenberry PJ, Becker PS. Stem cell homing: rolling, crawling, and nesting. Proc Natl Acad Sci U S A. 1998;95:15155.
- Mazo IB, Gutierrez-Ramos J-C, Frenette PS, Hynes RO, Wagner DD, von Andrian UH. Hematopoietic progenitor cell rolling in bone marrow microvessels: parallel contributions by endothelial selectins and vascular cell adhesion molecule 1. J Exp Med. 1998;188:465.
- Frenette PS, Subbarao S, Mazo IB, von Andrian UH, Wagner DD. Endothelial selectins and vascular cell adhesion molecule-1 promote hematopoietic progenitor homing to bone marrow. Proc Natl Acad Sci U S A. 1998;95:14423.
- Springer TA. Traffic signals for lymphocyte recirculation and leukocyte emigration: the multistep paradigm. Cell. 1994;76:301.
- Butcher EC. Leukocyte-endothelial cell recognition: three (or more) steps to specificity and diversity. Cell. 1991;6:1033.
- Bargatze RF, Jutila MA, Butcher EC. Distinct roles of L-selectin and integrins alpha 4 beta 7 and LFA-1 in lymphocyte homing to Peyer's patch-HEV in situ: the multistep model confirmed and refined. Immunity. 1995;3:99.
- Bargatze RF, Butcher EC. Rapid G protein-regulated activation event involved in lymphocyte binding to high endothelial venules. J Exp Med. 1993;178:367.
- Schweitzer KM, Drager AM, van der Valk P, et al. Constitutive expression of E-selectin and vascular cell adhesion molecule-1 on endothelial cells of hematopoietic tissues. Am J Pathol. 1996;148: 165.
- Peled A, Petit I, Kollet O, et al. Dependence of human stem cell engraftment and repopulation of NOD/SCID mice on CXCR4. Science. 1999;283: 845.
- Peled A, Grabovsky V, Habler L, et al. The chemokine SDF-1 stimulates integrin-mediated arrest of CD34<sup>+</sup> cells on vascular endothelium under shear flow. J Clin Invest. 1999;104:1199.