

Expression of C/EBP β from the *C/ebp α* gene locus is sufficient for normal hematopoiesis in vivo

Letetia C. Jones, Meng-Liang Lin, Shih-Shun Chen, Utz Krug, Wolf-K. Hofmann, Stephen Lee, Ying-Hue Lee, and H. Phillip Koeffler

CCAAT/enhancer-binding proteins (C/EBPs) are critical transcriptional regulators of differentiation of hematopoietic cells. Previous studies have shown that targeted disruption of the *C/ebp α* gene results in a lack of granulocytic differentiation with an arrest at the stage of immature myeloblasts. By using a gene replacement strategy in which C/EBP β was expressed from the *C/ebp α* gene locus of C/EBP α -null mice, we have evaluated the ability of C/EBP β to func-

tion for C/EBP α in directing differentiation along the granulocytic pathway. We show that the morphology and the differential cell counts of the bone marrow and peripheral blood cells from C/EBP β knockin mice are indistinguishable from those of their wild-type littermates, indicating that hematopoiesis occurs normally in these animals. Additionally, we analyzed expression of 21 myeloid-specific genes, including markers for distinct stages of granulocytic differen-

tiation, and found no significant differences in their levels of expression in the bone marrow of C/EBP β knockin and wild-type mice. These results imply that C/EBP β can substitute for C/EBP α during hematopoiesis when expressed from the *C/ebp α* gene locus. (Blood. 2002;99:2032-2036)

© 2002 by The American Society of Hematology

Introduction

CCAAT/enhancer-binding proteins (C/EBPs) are a family of structurally related transcription factors made up of 6 members (C/EBP α , β , γ , δ , ϵ , and ζ).¹⁻⁷ All C/EBPs share conserved C-terminal regions that contain leucine-zipper dimerization motifs adjacent to basic DNA-binding domains.⁸ Their N-terminal regions are more diverse and contain transcriptional activation domains. Dimerization within the C/EBP family or with other transcription factors is a prerequisite for DNA binding and subsequent transactivation. With the exception of C/EBP ϵ , proteins in the C/EBP family are expressed in partially overlapping patterns in multiple tissues.⁹ However, targeted inactivation of C/EBP family genes in mice has demonstrated their individual contributions to cellular differentiation.

Knockout mice models have defined a critical role for C/EBPs in hematopoietic tissues.¹⁰⁻¹⁴ Two members, C/EBP α and C/EBP β , play key roles in determining the fate of differentiating hematopoietic cells. For example, C/EBP α is expressed in early myeloid cells,⁹ and its absence in C/EBP α ^{-/-} mice leads to a complete lack of granulocytic differentiation with an arrest at the stage of immature myeloblasts.¹⁰ Although mature neutrophils and eosinophils are absent in C/EBP α -null mice, other hematopoietic lineages are not affected. C/EBP β , however, appears to be a critical signaling molecule for more mature myeloid cells as well as for B lymphocytes because its expression is dramatically induced during macrophage differentiation^{9,15} and lymphopoiesis.¹² Targeted dele-

tion of C/EBP β in mice results in impaired macrophage function, lymphoproliferative disorders, and defective B lymphopoiesis. Differential expression of these 2 C/EBPs in hematopoietic tissues underscores their individual roles in the development of mature blood cells.

In addition to their roles in the hematopoietic system, C/EBP α and C/EBP β are important for normal development of liver^{16,17} and adipose tissue.^{4,18} Mice lacking C/EBP α die within 8 hours of birth because of a severe loss of liver function. However, a gene replacement approach in which C/EBP β is knocked into the *C/ebp α* gene locus of C/EBP α -null mice restores liver function and, consequently, their viability.¹⁹ These mutant mice, *C/ebp α ^{β/β}* , lack C/EBP α but have a concomitant gain of C/EBP β in tissues. In the current study, we evaluate the ability of C/EBP β functionally to replace C/EBP α in the hematopoietic system of *C/ebp α ^{β/β}* mice. We find that bone marrow and peripheral blood cells from C/EBP β knockin mice are indistinguishable from those of their wild-type littermates, indicating that hematopoiesis occurs normally in these animals. We confirm this finding on a molecular level by analyzing the expression of 21 myeloid-specific genes, including markers for distinct stages of granulocytic differentiation. Our results reveal no significant differences in the levels of expression of these genes in bone marrow of C/EBP β knockin and wild-type mice, thus implying that C/EBP β can substitute for C/EBP α during hematopoiesis when expressed from the *C/ebp α* gene locus.

From the Division of Hematology and Oncology, Department of Medicine, and the Department of Pathology, Cedars-Sinai Medical Center, University of California-Los Angeles School of Medicine; Laboratory of Molecular Pathology, Institute of Molecular Biology, Academia Sinica, Taipei, Taiwan; School of Medical Technology, China Medical College, Taichung, Taiwan.

Submitted August 2, 2001; accepted November 14, 2001.

Supported by grants from the National Institutes of Health (H.P.K.), the C. and H. Koeffler Fund, Parker Hughes Trust, Brian Harvey Fund, Frederick P. Begell Foundation, and the Joseph Troy Leukemia Foundation. L.C.J. is supported by grants from the American Cancer Society (PF-99-127-01-CNE) and the National Institutes of Health (T32 CA-75956). W.K.H. is a recipient of a

scholarship from the Deutsche Forschungsgemeinschaft (HO2207/1-1). H.P.K. is a member of the Jonsson Comprehensive Cancer Center and holds the endowed Mark Goodson Chair of Oncology Research at Cedars-Sinai Medical Center/UCLA School of Medicine.

Reprints: Letetia C. Jones, Division of Hematology and Oncology, Cedars-Sinai Medical Center, UCLA School of Medicine, 8700 Beverly Blvd, Suite BM-1, Rm 109, Los Angeles, CA 90048; e-mail: Letetia.Jones@cshs.org.

The publication costs of this article were defrayed in part by page charge payment. Therefore, and solely to indicate this fact, this article is hereby marked "advertisement" in accordance with 18 U.S.C. section 1734.

© 2002 by The American Society of Hematology

Materials and methods

Mice

C/ebp α ^{B β} mice were generated by using gene-targeting technology and the Cre/loxP DNA recombination system as reported previously.¹⁹ Mice were maintained in a specific pathogen-free animal facility at the Institute of Molecular Biology, Academia Sinica, Taipei. Five- to 6-week-old littermates from the heterozygote interbreedings were used in this study.

Bone marrow morphologic analysis

Bone marrow was collected from the femoral bone of wild-type and C/EBP β knockin mice. Smears were prepared and stained with Wright-Giemsa. Bone marrow smears were analyzed by light microscopy.

Differential cell counts

Differential white cell counts for peripheral blood and bone marrow were determined manually by 2 independent investigators. Percentages were calculated according to the cell morphology of a total of 600 cells per peripheral blood smear and 700 cells per bone marrow smear. Values are given as mean \pm SD from 3 independent animals.

RNA isolation and reverse transcription–polymerase chain reaction

Total RNA was isolated from mononuclear bone marrow cells of 4 wild-type and 4 C/EBP β knockin mice by using TRIzol (Life Technologies, Grand Islands, NY). Two micrograms of DNase I-treated RNA was reverse transcribed by using Moloney murine leukemia virus reverse transcriptase (Life Technologies), and 50 ng of the resulting complementary DNAs (cDNAs) was used as templates for polymerase chain reaction (PCR). Amplification was carried out by using HotStarTaq DNA polymerase (Qiagen, Valencia, CA) under the following conditions: an initial denaturation step at 95°C for 15 minutes followed by 35 cycles of 95°C for 30 seconds, 60°C for 30 seconds, and 72°C for 1 minute. The specificity of primer pairs used for amplification was confirmed by Southern blot by using internal oligonucleotides as probes. Reaction products were visualized on ethidium bromide-stained agarose gels, and images of C/EBP α , C/EBP β , and 18S DNAs were captured by using AlphaImager 2000 Gel Documentation software. Reverse transcriptase(RT)–PCR results were confirmed by varying input cDNA concentration and cycle number or by real-time PCR. For the latter, RT-PCR reactions were carried out by using HotStarTaq DNA polymerase (Qiagen), 50 ng cDNA for myeloid-specific genes (500–5 ng in serial dilutions for standard curves) or 1 pg for 18S (10–0.1 pg for standard curve), and SYBRGreenI nucleic acid gel staining solution in a 1:60 000 dilution. PCR conditions were as follows: a 95°C initial activation for 15 minutes followed by 45 cycles of 95°C for 15 seconds, 60°C for 15 seconds, and 72°C for 30 seconds, and fluorescence determination at the melting temperature of the product for 20 seconds on an ICycler detection system (BioRad, Hercules, CA).

Results

Construction of a C/EBP β knockin targeting vector and generation of homozygous *C/ebp α ^{B β}* mice were described previously.¹⁹ These mice carry a mutant *C/ebp α* allele in which the protein-coding region of *C/ebp α* was deleted and replaced with that of *C/ebp β* . *C/ebp α ^{B β}* mice are viable, fertile, and grossly normal and exhibit growth rates that are identical to their wild-type littermates. Furthermore, they show none of the liver abnormalities found in the C/EBP α -null mice, implying that C/EBP β can functionally replace C/EBP α in the liver when expressed from the *C/ebp α* gene locus.

The functional redundancy of C/EBP α and C/EBP β in liver raised the question of their redundancy in hematopoietic cells. Previous studies¹⁰ suggest that C/EBP α is indispensable for differentiation along the granulocytic pathway because C/EBP α -deficient mice lack mature neutrophils and eosinophils. Therefore, we evaluated the ability of C/EBP β to compensate for C/EBP α in the hematopoietic system of *C/ebp α ^{B β}* mice. Analysis of peripheral blood from *C/ebp α ^{B β}* mice and their wild-type littermates (Table 1) showed similar differential cell counts for all hematopoietic lineages, including myeloid elements from myeloblasts to mature neutrophils. Additionally, no differences in morphology were observed in Wright-Giemsa-stained bone marrow cells from C/EBP β knockin and wild-type mice (Figure 1). C/EBP β knockin and wild-type murine bone marrow also stained equally for the myeloid-specific, azurophilic protein myeloperoxidase and for Sudan black, a marker of myeloid progenitors (data not shown). The white blood cell counts of the C/EBP β knockin mice were in the normal range (mean, $4.4 \times 10^9/\mu\text{L}$), and for unclear reasons the wild-type mice had a slightly elevated white blood cell count (mean, $7.8 \times 10^9/\mu\text{L}$).²⁰ The morphology and number of neutrophils in the peripheral blood (Table 1 and data not shown) of the C/EBP β knockin animals were normal,²⁰ indicating that expression of C/EBP β from the *C/ebp α* gene locus overcomes the block in granulocytic differentiation observed in the C/EBP α -null mice. To rule out the possibility that this rescue is due to in vivo compensatory mechanisms other than the expression of C/EBP β , bone marrow cells from wild-type and C/EBP β knockin mice were used for in vitro colony assays in the presence of granulocyte colony-stimulating factor. The number of granulocyte colonies formed from bone marrow cells from knockin mice was not different from that seen when using bone marrow cells from wild-type animals (data not shown). The granulocyte colonies contained similar percentages of neutrophils in assays of bone marrow cells from knockin and wild-type mice (85.7 ± 3.1 and 82.3 ± 3.5 , respectively). These data imply that differentiation

Table 1. Analysis of peripheral blood and bone marrow from wild-type and C/EBP β knockin mice

	% \pm SD	
	<i>C/ebpα^{+/+}</i>	<i>C/ebp$\alphaB\beta$}</i>
Differential cell counts		
Peripheral blood		
Lymphocytes	77.7 \pm 3.1	81.7 \pm 6.5
Neutrophils	18.3 \pm 2.1	17.3 \pm 7.5
Monocytes	4.0 \pm 3.6	1.0 \pm 1.0
Bone marrow		
Myeloblasts & promyelocytes	4.3 \pm 0.6	4.7 \pm 2.1
Myelocytes & metamyelocytes	8.0 \pm 1.0	6.7 \pm 2.3
Neutrophils	47.0 \pm 3.5	51.0 \pm 2.6
Nucleated red blood cells	23.0 \pm 2.6	20.3 \pm 4.5
Lymphocytes	17.3 \pm 2.3	17.5 \pm 2.4
White blood cell count, $10^9/\mu\text{L}$ *	7.8	4.4
Red blood cell count, $10^{12}/\text{L}$	9.0	8.8
Platelet count, $10^9/\mu\text{L}$ *	771.3	691.0

For differential counts, percentages were based on cell morphology of a total of 600 cells per genotype in peripheral blood and 700 cells per genotype in bone marrow. Each value represents the mean \pm SD from 3 independent animals. Student *t* test showed a *P* value $>$.1 for all numbers. Total white cell, red cell, and platelet counts were determined from 2 independent *C/ebp α ^{B β}* and *C/ebp α ^{+/+}* mice.

*Normal reported counts²¹: white blood cells, 4.0 – $5.6 \times 10^9/\mu\text{L}$; red blood cells, 9.0 – $9.9 \times 10^{12}/\text{L}$; platelets, 800 – $1100 \times 10^9/\mu\text{L}$.




Figure 1. Wright-Giemsa–stained bone marrow cells from wild-type and *C/EBPα*^β knockin mice. Bone marrow smears from wild-type (A) and *C/ebpα*^{β/β} (B) mice show maturation of the myeloid lineage to mature granulocytes (indicated by G). Magnification × 100.

along the granulocytic pathway is due to functional replacement of *C/EBPα* with *C/EBPβ* and is not likely to be attributable to *C/EBPα*-independent pathways.

The presence of a mature granulocytic population in *C/ebpα*^{β/β} mice implies that genes necessary for differentiation are appropriately expressed. To verify this at the molecular level, we evaluated the expression of 21 myeloid-specific genes in bone marrow cells from *C/EBPβ* knockin mice and their wild-type littermates. We first confirmed the lack of *C/EBPα* expression in the bone marrow of *C/ebpα*^{β/β} mice by RT-PCR (Figure 2). The knockin mice showed increased expression of *C/EBPβ* compared with *C/ebpα*^{+/+} mice, representing transcripts from both the *C/ebpα* and *C/ebpβ* gene loci.

Other genes evaluated include (1) markers for different stages of maturation along the granulocytic pathway (primary and secondary granule proteins), (2) eosinophil-specific proteins, (3) colony-stimulating factor receptors and other cytokine-signaling proteins, and (4) proteins involved in phagocytosis (components of nicotinamide adenine dinucleotide phosphate oxidase and an antimicrobial protein). The individual genes analyzed are listed in Table 2. Our studies revealed no differences in the expression of these genes in *C/ebpα*^{β/β} and *C/ebpα*^{+/+} mice (Figure 3). This finding is consistent with the lack of morphologic differences between the 2 genotypes.




Figure 2. Expression of *C/EBPα* and *C/EBPβ* messenger RNAs in wild-type and *C/EBPβ* knockin mice. Gene expression was measured by RT-PCR using RNA from the bone marrow of wild-type and *C/EBPβ* knockin mice. After 35 (*C/EBPα* and *C/EBPβ*) and 15 (18S) cycles, amplification products were gel separated and stained with ethidium bromide.

For 7 of these genes, expression levels were quantified by using quantitative RT-PCR. RNA from the bone marrow of 4 wild-type and 4 *C/EBPβ* knockin mice were reverse transcribed, and the resulting cDNAs were analyzed by real-time PCR. The values for individual mice within each group were averaged, and expression values in *C/EBPβ* knockin mice are given in Table 3 relative to expression in wild-type animals (arbitrarily set at 1.0). Again, our results indicate that the levels of expression of these genes in *C/EBPβ* knockin mice do not significantly differ from those observed for the wild-type animals, suggesting that gene expression is normal in the knockin mice.

Table 2. Twenty-one myeloid-specific genes show no differences in expression in bone marrow cells of *C/ebpα*^{β/β} as compared with wild-type mice

Description	Gene
Myeloblastic & promyelocytic stages	Myeloperoxidase
	Neutrophil elastase
	B9
Myelocytic stage	Lactoferrin
	Neutrophil collagenase
	Neutrophil gelatinase
	CD11b (MAC-1)
Eosinophil-specific markers	Eosinophil peroxidase
	Major basic protein
Cytokines	MIP2
	TNF α
Cytokine receptors	G-CSF receptor
	GM-CSF receptor
	M-CSF receptor
	IL-6 receptor
	IL-8 receptor
Phagocytic proteins	p91phox
	p47phox
	MCLP
<i>C/EBP</i> family members	<i>C/EBPδ</i>
	<i>C/EBPε</i>

MIP2 indicates macrophage inflammatory protein-2; TNF α , tumor necrosis factor- α ; G-CSF, granulocyte colony-stimulating factor; GM-CSF, granulocyte-macrophage colony-stimulating factor; M-CSF, macrophage colony-stimulating factor; IL, interleukin; and MCLP, murine cathelinlike protein.




Figure 3. Expression of myeloid-specific genes in wild-type and C/EBP β knockin mice. Gene expression was measured by RT-PCR using RNA from the bone marrow of wild-type and C/EBP β knockin mice. After 15 (18S) and 35 (all other genes) cycles, amplification products were gel separated and stained with ethidium bromide. G-CSF R indicates granulocyte-colony stimulating factor receptor; MPO, myeloperoxidase.

Discussion

Mice with targeted deletion of C/EBP α die soon after birth from hypoglycemia, and analysis of their peripheral blood and bone marrow reveal hematopoietic abnormalities, including the absence of granulocytic differentiation.¹⁰ Thus, normal expression of C/EBP β does not compensate for the C/EBP α deficiency in the liver or hematopoietic tissue of C/EBP α -null mice. However, recent studies showed that expression of C/EBP β from the *C/ebp α* gene locus, in addition to its expression from the *C/ebp β* allele, restored liver function and overcomes the neonatal lethality of C/EBP α ^{-/-} mice.¹⁹ Similarly, in this report, we show that expression of C/EBP β from the *C/ebp α* locus restores normal hematopoiesis by overcoming the selective block in granulocytic differentiation observed in the C/EBP α -null mice.

The ability of C/EBP β to function for C/EBP α in hematopoietic cells of *C/ebp α ^{β/β}* mice is likely related to at least 2 aspects of expression from the *C/ebp α* gene locus: (1) the level of transcriptional activity and (2) the timing of expression. In hepatic tissues of *C/ebp α ^{β/β}* mice, C/EBP β messenger RNA expressed from the *C/ebp α* locus is significantly higher than expression of C/EBP β from its endogenous allele.¹⁹ Although our studies do not quantify the contribution of C/EBP β messenger RNA from each locus in hematopoietic cells, we do observe higher expression of C/EBP β in the knockin model, representing the combined expression from both loci. Radomska et al²¹ suggested that high levels of C/EBP α at

the stage of myeloid commitment is the molecular switch that directs myeloid precursors to the granulocytic pathway. Because myeloid progenitor cells express low levels of C/EBP β , the amount of C/EBP β in C/EBP α -null mice is possibly insufficient to transactivate genes whose expression is required for granulocytic differentiation.

Although it is low in early stages of myeloid differentiation, the level of C/EBP β expression increases dramatically at later stages of differentiation especially in maturing macrophages.⁹ C/EBP β ^{-/-} mice have defects in macrophage function and develop lymphoproliferative disorders as a result, but deletion of C/EBP β in mice does not adversely affect myeloid cell differentiation.¹³ Hence, its role in myelopoiesis is unclear. Expression of C/EBP β from the *C/ebp α* locus places the protein at the scene of hematopoietic differentiation earlier and probably at higher levels than when expressed from the *C/ebp β* gene locus. Therefore, the rescue of granulocytic differentiation in *C/ebp α ^{β/β}* mice likely reflects changes in the temporal expression of C/EBP β . Stages of differentiation in hematopoietic cells are driven not only by C/EBPs but also by other transcription factor groups including GATA-1^{22,23} and GATA-2,^{24,25} Myb,²⁶⁻²⁸ Ets,²⁹⁻³¹ and AML1.³²⁻³⁴ Perhaps high expression of C/EBP β early in myelopoiesis is sufficient to maintain the integrity of protein-protein interactions that direct myeloid progenitors toward mature granulocytes.

C/EBPs are highly homologous in their C-terminal dimerization and DNA-binding domains and are believed to bind the same recognition sites on DNA.³⁵ Therefore, C/EBP β conceivably can interact with dimerization partners of C/EBP α and bind C/EBP α -targeted promoters such as the receptors for granulocyte colony-stimulating factor³⁶ and interleukin 6,³⁷ myeloperoxidase,³⁸ and neutrophil elastase.³⁹ However, the N-terminal regions of C/EBPs are more diverse and mediate their transactivation functions. Given the complexity of transcriptional activation complexes, we are somewhat surprised that the transactivation domain of C/EBP β can recruit necessary cofactors to activate promoters normally directed by the transactivation domain of C/EBP α . Interestingly, C/EBP β does not substitute for C/EBP α in the expression of genes encoding adipocyte-specific factors adipin and leptin.¹⁹ Abnormalities in fat storage but normal liver development and hematopoiesis suggest that the redundancy of C/EBP α and C/EBP β may be tissue and gene specific. Previous studies implicate C/EBP α as a critical factor for granulocytic commitment of myeloid progenitor cells. Strain differences between mice may reconcile our findings with the strict requirement for C/EBP α in granulocytic differentiation reported previously. However, it is more likely that C/EBP α itself is less important than the timing and level of its expression.

Table 3. Real-time PCR analysis of gene expression in wild-type and C/EBP β knockin mice

Gene	WT \pm SD	KI \pm SD
G-CSF receptor	1.0 \pm 0.57	0.87 \pm 0.89
GM-CSF receptor	1.0 \pm 0.90	0.88 \pm 0.97
Myeloperoxidase	1.0 \pm 0.29	0.87 \pm 0.29
Neutrophil elastase	1.0 \pm 0.58	0.87 \pm 0.72
Neutrophil gelatinase	1.0 \pm 1.30	0.96 \pm 0.58
Eosinophil major basic protein	1.0 \pm 0.93	0.95 \pm 1.35
C/EBP ϵ	1.0 \pm 0.72	0.71 \pm 0.99

Expression values were calculated as previously described.²⁰ Mean \pm SD for 4 wild-type (WT) and 4 C/EBP β knockin (KI) mice are given. *P* values were $>$.1 in the Student *t* test for all genes tested. G-CSF indicates granulocyte colony-stimulating factor; GM-CSF, granulocyte-macrophage colony-stimulating factor.

References

- Landschulz WH, Johnson PF, Adashi EY, Graves BJ, McKnight SL. Isolation of a recombinant copy of the gene encoding C/EBP. *Genes Dev.* 1988;2:786-800.
- Akira S, Isshiki H, Sugita T, et al. A nuclear factor for IL-6 expression (NF-IL6) is a member of a C/EBP family. *EMBO J.* 1990;9:1897-1906.
- Roman C, Platero JS, Shuman JD, Calame K. Ig/EBP-1: a ubiquitously expressed immunoglobulin enhancer binding protein that is similar to C/EBP and heterodimerizes with C/EBP. *Genes Dev.* 1990;4:1401-1415.
- Cao Z, Umek RM, McKnight SL. Regulated expression of three C/EBP isoforms during adipose conversion of 3T3-L1 cells. *Genes Dev.* 1991;5:1538-1552.
- Antonson P, Stellan B, Yamanaka R, Xanthopoulos KG. A novel human CCAAT/enhancer binding protein gene, C/EBP ϵ , is expressed in cells of lymphoid and myeloid lineages and is located on chromosome 14q11.2 close to the T-cell receptor alpha/delta locus. *Genomics.* 1996;35:30-38.
- Chumakov AM, Grillier I, Chumakova E, Chih D, Slater J, Koeffler HP. Cloning of the novel human myeloid-cell-specific C/EBP ϵ transcription factor. *Mol Cell Biol.* 1997;17:1375-1386.
- Ron D, Habener JF. CHOP, a novel developmentally regulated nuclear protein that dimerizes with transcription factors C/EBP and LAP and functions as a dominant-negative inhibitor of gene transcription. *Genes Dev.* 1992;6:439-453.
- Landschulz WH, Johnson PF, McKnight SL. The DNA binding domain of the rat liver nuclear protein C/EBP is bipartite. *Science.* 1989;243:1681-1688.
- Scott LM, Civin CI, Rorth P, Friedman AD. A novel temporal expression pattern of three C/EBP family members in differentiating myelomonocytic cells. *Blood.* 1992;80:1725-1735.
- Zhang D, Zhang P, Wang N, Hetherington C, Darlington GJ, Tenen DG. Absence of granulocyte colony-stimulating factor signaling and neutrophil development in CCAAT enhancer binding protein α -deficient mice. *Proc Natl Acad Sci U S A.* 1997;94:569-574.
- Tanaka T, Akira S, Yoshida K, et al. Targeted disruption of the NF-IL6 gene discloses its essential role in bacteria killing and tumor cytotoxicity by macrophages. *Cell.* 1995;80:353-361.
- Chen X, Liu W, Ambrosino C, et al. Impaired generation of bone marrow B lymphocytes in mice deficient in C/EBP β . *Blood.* 1997;90:156-164.
- Scrapanti I, Romani L, Musiani P, et al. Lymphoproliferative disorder and imbalanced T-helper response in C/EBP beta-deficient mice. *EMBO J.* 1995;14:1932-1941.
- Yamanaka R, Barow C, Lekstrom-Himes J, et al. Impaired granulopoiesis, myelodysplasia and early lethality in C/EBP ϵ -deficient mice. *Proc Natl Acad Sci U S A.* 1997;94:13187-13192.
- Natsuka S, Akira S, Nishio Y, et al. Macrophage differentiation-specific expression of NF-IL6, a transcription factor for interleukin-6. *Blood.* 1992;79:460-466.
- Lee YH, Sauer B, Johnson PF, Gonzalez FJ. Disruption of the c/ebp α gene in adult mouse liver. *Mol Cell Biol.* 1997;17:6014-6022.
- Wang ND, Finegold MJ, Bradley A, et al. Impaired energy homeostasis in C/EBP α knockout mice. *Science.* 1995;269:1108-1112.
- Yeh WC, Cao Z, Classon M, McKnight SL. Cascade regulation of terminal adipocyte differentiation by three members of the C/EBP family of leucine zipper proteins. *Genes Dev.* 1995;9:168-181.
- Chen SS, Chen JF, Johnson PF, Muppala V, Lee YH. C/EBP β , when expressed from the C/ebp α gene locus, can functionally replace C/EBP α in liver but not in adipose tissue. *Mol Cell Biol.* 2000;20:7292-7299.
- Foster HL, Small JD, Fox JG. *The Mouse in Biomedical Research.* New York, NY: Academic Press; 1983:302-307.
- Radomska HS, Huettner CS, Zhang PU, Cheng T, Scadden DT, Tenen DG. CCAAT/enhancer binding protein α is a regulatory switch sufficient for induction of granulocytic development from bipotential myeloid progenitors. *Mol Cell Biol.* 1998;18:4301-4314.
- Pevny L, Simon M, Robertson E, et al. Erythroid differentiation in chimaeric mice blocked by a targeted mutation in the gene for transcription factor GATA-1. *Nature.* 1991;349:257-260.
- Simon M, Pevny L, Wiles M, Keller G, Costantini F, Orkin S. Rescue of erythroid development in gene targeted GATA-1 mouse embryonic stem cells. *Nat Genet.* 1992;1:92-98.
- Briegel K, Lim KC, Plank C, Beug H, Engel J, Zenke M. Ectopic expression of a conditional GATA-2/estrogen receptor chimera arrests erythroid differentiation in a hormone-dependent manner. *Genes Dev.* 1993;7:1097-1109.
- Tsai FY, Keller G, Kuo FC, et al. An early haematopoietic defect in mice lacking the transcription factor GATA-2. *Nature.* 1994;371:221-226.
- Clarke M, Kukowska-Latallo J, Westin E, Smith M, Prochownik E. Constitutive expression of a c-myc cDNA blocks Friend murine erythroleukemia cell differentiation. *Mol Cell Biol.* 1988;8:884-892.
- Gewirtz A, Calabretta B. A c-myc antisense oligodeoxynucleotide inhibits normal human hematopoiesis in vitro. *Science.* 1988;242:1303-1306.
- Graf T. Myb: a transcriptional activator linking proliferation and differentiation in hematopoietic cells. *Curr Opin Genet Dev.* 1992;2:249-255.
- Chen H, Zhang P, Voso M, et al. Neutrophils and monocytes express high levels of PU.1 (Spi-1) but not Spi-B. *Blood.* 1995;85:2918-2928.
- Klemsz M, McKercher S, Celada A, Beveren CV, Maki R. The macrophage and B cell-specific transcription factor PU.1 is related to the *ets* oncogene. *Cell.* 1990;61:113-124.
- Wotton D, Ghysdael J, Wang S, Speck N, Owen M. Cooperative binding of Ets-1 and core binding factor to DNA. *Mol Cell Biol.* 1994;14:840-850.
- Miyoshi H, Ohira M, Shimizu K, et al. Alternative splicing and genomic structure of the AML1 gene involved in acute myeloid leukemia. *Nucleic Acids Res.* 1995;23:2762-2769.
- Takahashi A, Satake M, Yamaguchi-Iwai Y, et al. Positive and negative regulation of granulocyte macrophage colony-stimulating factor promoter activity by AML1-related transcription factor, PEBP2. *Blood.* 1995;86:607-616.
- Tanaka T, Tanaka K, Ogawa S, et al. An acute myeloid leukemia gene, AML1, regulates hematopoietic myeloid cell differentiation and transcriptional activation antagonistically by two alternative spliced forms. *EMBO J.* 1995;14:341-350.
- Williams SC, Cantwell CA, Johnson PF. A family of C/EBP-related proteins capable of forming covalently linked leucine zipper dimers in vitro. *Genes Dev.* 1991;9:1553-1567.
- Smith LT, Hohaus S, Gonzalez DA, Dziennis SE, Tenen DG. PU.1 (Spi-1) and C/EBP α regulate the granulocyte colony-stimulating factor receptor promoter in myeloid cells. *Blood.* 1996;88:1234-1247.
- Zhang P, Iwama A, Datta MW, Darlington GJ, Link DC, Tenen DG. Upregulation of interleukin 6 and granulocyte colony-stimulating factor receptors by transcription factor CCAAT enhancer binding protein (C/EBP α) is critical for granulopoiesis. *J Exp Med.* 1998;6:1173-1184.
- Ford AM, Bennett CA, Healy LE, Towatari M, Greaves MF, Enver T. Regulation of the myeloperoxidase enhancer binding proteins PU.1, C/EBP α , β , and δ during granulocytic-lineage specification. *Proc Natl Acad Sci U S A.* 1996;93:10838-10843.
- Oelgeschlager M, Nuchprayoon I, Luscher B, Friedman AD. C/EBP, c-Myb, and PU.1 cooperate to regulate the neutrophil elastase promoter. *Mol Cell Biol.* 1996;16:4717-4725.