

Presence of a mobilizable intracellular pool of hepatocyte growth factor in human polymorphonuclear neutrophils

Alain Grenier, Sylvie Chollet-Martin, Bruno Crestani, Charlotte Delarche, Jamel El Benna, Anne Boutten, Valérie Andrieu, Geneviève Durand, Marie-Anne Gougerot-Pocidallo, Michel Aubier, and Monique Dehoux

Hepatocyte growth factor (HGF), a heparin-binding factor, is synthesized as a single-chain inactive precursor (pro-HGF), which is converted by proteolysis to an active heterodimer (mature HGF). HGF has pleiotropic activities and has been implicated in the regulation of mitogenesis, motogenesis, and morphogenesis of epithelial and endothelial cells. As polymorphonuclear neutrophils (PMNs) secrete numerous cytokines involved in the modulation of local inflammation, we investigated their ability to produce HGF. We found that HGF was stored in secretory vesicles and in gelatinase/specific

granules. This intracellular stock was rapidly mobilized by degranulation when neutrophils were stimulated with phorbol myristate acetate or N-formylmethionyl-leucyl-phenylalanine. Cycloheximide did not affect the release of HGF. Moreover, HGF messenger RNA and protein expression was found in bone marrow myeloid cells, suggesting that HGF synthesis likely occurs during PMN maturation. In mature circulating PMNs, intracellular HGF was in the pro-HGF form, whereas the HGF secreted by degranulation was the mature form. Furthermore, PMNs pretreated with diisopropyl fluorophosphate only re-

leased the pro-HGF form, suggesting that PMN-derived serine protease(s) are involved in the proteolytic process. We also obtained evidence that secreted mature HGF binds PMN-derived glycosaminoglycans (probably heparan sulfate). These findings suggest that PMNs infiltrating damaged tissues may modulate local wound healing and repair through the production of HGF, a major mediator of tissue regeneration. (Blood. 2002;99:2997-3004)

© 2002 by The American Society of Hematology

Introduction

Hepatocyte growth factor (HGF), a heparin-binding factor, was originally described as a potent mitogen for the growth of hepatocytes *in vitro* and was subsequently purified from rat platelets,¹ plasma of a patient with fulminant hepatic failure,² and normal human plasma.³ HGF has mitogenic, motogenic, and morphogenic effects on various epithelial and endothelial cell types.^{4,5} Many studies have demonstrated that embryogenesis, angiogenesis, hematopoiesis, as well as wound healing and organ regeneration, are critically controlled by HGF.⁵⁻⁷ HGF is secreted as a 92-kd single-chain pro-HGF that requires endoproteolytic processing. This processing is mediated by a serine protease. The HGF activator,⁸ blood coagulation factor XII,⁹ urokinase,¹⁰ and tissue-type plasminogen activator¹¹ are reported to activate HGF. Processing of HGF results in a bioactive form (mature HGF) consisting of a disulfide-linked 69-kd α chain and a 32-kd β chain.¹²

The biologic effects of HGF are mediated by the activation of its high-affinity binding site known as the tyrosine kinase receptor c-met.¹³ Besides c-met, HGF possesses lower-affinity/high-capacity binding sites corresponding to extracellular matrix molecules (glycosaminoglycans or collagen) or to cell surface-associated heparan sulfate^{14,15}; this correspondence creates a molecular reservoir of HGF on the cell surface, whereas HGF transfer to c-met initiates the cellular response.¹⁶ The broad range of HGF activities on many physiologic and pathologic processes is

reflected by c-met expression in a variety of organs and cell types. The finding that c-met receptor expression is induced by inflammatory cytokines¹⁷ suggests that the HGF/c-met system is involved in inflammatory responses to tissue injury.

Polymorphonuclear neutrophils (PMNs), the predominant tissue infiltrating cells during acute inflammatory process, participate in host defenses by generating reactive oxygen species and releasing proteolytic enzymes. Human PMNs can synthesize and secrete many proteins that influence the course of inflammatory responses and contribute to the modulation of local inflammation.^{18,19} After reports that human liver and bone marrow PMNs contain immunoreactive HGF,^{20,21} we investigated the regulation of HGF production by human blood PMNs, together with the subcellular localization and molecular form of HGF in these cells.

Materials and methods

Purification of blood PMNs and bone marrow myeloid cells

Human PMNs were purified as previously described.¹⁸ Briefly, blood PMNs from healthy volunteers were isolated in sterile conditions by sedimentation in medium containing 9% Dextran T-500 (Amersham Pharmacia Biotech, Uppsala, Sweden) and 38% Radioselectan (Schering, Lys-lez-Lannoy, France), and the leukocyte suspension was centrifuged on Ficoll-Paque medium (Amersham Pharmacia Biotech). The cell pellet was washed with phosphate-buffered saline (PBS), and erythrocytes were removed by

From the Laboratoire de Biochimie, Hôpital de Montfermeil, Montfermeil, France; and the Laboratoire de Biochimie, INSERM U 408, Laboratoire d'Hématologie et d'Immunologie et INSERM U 479, and Service de Pneumologie, Hôpital Bichat Assistance Publique-Hôpitaux de Paris, Paris, France.

Submitted January 10, 2001; accepted December 3, 2001.

Reprints: Monique Dehoux, Laboratoire de Biochimie and INSERM U408, Hôpital

Bichat, 46 rue Henri Huchard, 75877 Paris Cedex 18, France; e-mail: monique.dehoux@bch.ap-hop-paris.fr.

The publication costs of this article were defrayed in part by page charge payment. Therefore, and solely to indicate this fact, this article is hereby marked "advertisement" in accordance with 18 U.S.C. section 1734.

© 2002 by The American Society of Hematology

hypotonic lysis. PMNs were further purified by 20-minute incubation with pan antihuman HLA class II-coated magnetic beads (Dynal, Oslo, Norway) to deplete B lymphocytes, activated T lymphocytes, and monocytes. Nonspecific esterase staining always showed less than 0.5% of monocytes, and flow cytometry showed the absence of CD14⁺, CD3⁺, and CD19⁺ cells, confirming the recovery of highly purified PMNs. Two bone marrow samples were obtained from healthy allograft donors. Mononuclear cells had been removed by Ficoll density gradient centrifugation. These depleted samples were sedimented in medium containing 9% Dextran T-500 and 38% Radioselectan. The leukocyte-rich supernatant was centrifuged, and the cell pellet was washed with PBS. Erythrocytes were removed by hypotonic lysis. The distribution of neutrophil precursors was determined by differential cell counting of cytospin preparations from each donor sample, as follows: promyelocytes, 2% and 3%; myelocytes, 11% and 14%; metamyelocytes, 34% and 23%; band and segmented cells, 53% and 60%. In both preparations, myeloblasts, monocytes, and lymphocytes represented less than 1% of cells.

Immunocytochemical staining of HGF

Blood and bone marrow smears from healthy control subjects were air-dried for 24 hours and fixed in 4% formaldehyde in PBS for 20 minutes and permeabilized by incubation in PBS containing 0.1% Triton X-100 at room temperature for 30 minutes.²² Smears were first incubated with normal horse serum for 30 minutes to block nonspecific binding. The primary antibody, an antihuman-HGF monoclonal antibody (12.5 µg/mL) (MAB294, clone 24 612.111; R&D Systems, Abingdon, United Kingdom), was diluted in PBS containing 0.3% gelatin and added for 1 hour at room temperature. The slides were then incubated with a biotinylated antibody and then with alkaline phosphatase-labeled streptavidin according to the manufacturer's instructions (Vectastain ABC kit; Vector Laboratories, Burlingame, United Kingdom). Staining was completed by incubation with a substrate chromogen solution (Fast red substrate solution; DAKO, Carpinteria, CA). After washing, the slides were counterstained in Mayer hematoxylin, mounted, and examined with a light microscope. Positive staining developed as a red-colored reaction product. Smears incubated with a nonspecific mouse immunoglobulin of the same isotype (10 µg/mL; R&D Systems) served as negative controls.

Subcellular fractionation of blood PMNs

Specific and azurophilic granules were purified as previously described.²³ Briefly, purified neutrophils (100×10^6) in 5 mL ice-cold relaxation buffer (100 mM KCl, 3 mM NaCl, 1 mM Na₂ATP, 3.5 mM MgCl₂, 10 mM PIPES, pH 7.2) supplemented with EGTA and antiproteases were pressurized with N₂ for 20 minutes at 450 psi with constant stirring in a nitrogen bomb. The cavitate was then collected dropwise into EGTA, pH 7.4, sufficient for a final concentration of 1.25 mM. Nuclei and unbroken cells were pelleted by centrifugation of the cavitate at 400g for 15 minutes. The supernatant was decanted, loaded on top of a 2-layer Percoll gradient (1.05/1.12 g/mL), precooled to 4°C, and centrifuged at 4°C for 30 minutes at 40 000g. Three visible bands were identified: a lower band containing azurophilic granules, an intermediate band containing specific and gelatinase granules, and an upper band containing plasma membranes and secretory vesicles. The cytosol remained above the upper band. The different fractions were collected, and the purity of specific and azurophilic granule fractions was assessed by enzyme-linked immunosorbent assay (ELISA) measurement of their respective markers (lactoferrin and myeloperoxidase) in each fraction and in the total cavitate.

Degranulation experiments and purification of PMN-derived HGF

Pure PMNs (10⁷/mL) were resuspended in Hanks balanced salt solution (HBSS with Ca⁺⁺/Mg²⁺; Life Technologies, Cergy Pontoise, France). Part of the cells (unstimulated control PMNs) were kept for 20 minutes in medium alone, on ice, or at 37°C; the other part was preincubated at 37°C for 20 minutes with the following degranulating agents: lipopolysaccharide at 1 µg/mL (LPS from *Escherichia coli* 055:B5; Sigma, St Louis, MO),

10⁻⁶ M and 10⁻⁸ M N-formylmethionyl-leucyl-phenylalanine (fMLP; Sigma), or phorbol myristate acetate at 100 ng/mL (PMA; Sigma). In some experiments, to ensure total degranulation, PMNs were preincubated with 5 µg/mL cytochalasin B (Sigma) for 5 minutes, then with 10⁻⁶ M fMLP for 15 minutes at 37°C. In another set of experiments, to inhibit serine proteases, neutrophils were preincubated with 2 mM diisopropyl fluorophosphate (DFP; Sigma) for 15 minutes at 4°C and then treated as indicated above with cytochalasin B and fMLP. All tubes were then centrifuged, and the cell-free supernatants were stored at -80°C until HGF quantitation and Western blot analysis.

For HGF purification, the cell-free supernatant of PMNs stimulated with cytochalasin B + fMLP was applied to a Hi-Trap heparin affinity column (Amersham Pharmacia Biotech); the starting washing buffer was 0.02 M Tris, 0.2 M NaCl, pH 7.5, and the eluting buffer was 0.02 M Tris, 2 M NaCl, pH 7.5. Each harvested fraction (1 mL) was dialyzed against distilled water and assayed for HGF with an ELISA method.

Blood PMN culture

Pure PMNs (10⁷/mL) were cultured for 24 hours at 37°C with 5% CO₂ in 24-well tissue-culture plates (Costar, Cambridge, MA). The culture medium was RPMI 1640 (Sigma) supplemented with 2 mM glutamine and antibiotics. To avoid contaminating HGF and/or protease activities,²⁴ all culture media were free of fetal calf serum. Neutrophils were cultured with or without the following stimulating agents: LPS at 1 µg/mL, fMLP at 10⁻⁶ M, PMA at 100 ng/mL, and interleukin-1β (IL-1β) at 10 ng/mL (Immugenex, Los Angeles, CA). To evaluate de novo HGF synthesis, PMNs (10⁷/mL) were preincubated with or without 10 µg/mL cycloheximide (Sigma) for 30 minutes at 37°C and further incubated with the stimulating agents for 24 hours. Cell-free PMN culture supernatants were then collected and stored at -80°C until HGF assay.

HGF, lactoferrin, and myeloperoxidase assays

HGF was quantified by using a commercial ELISA kit (Quantikine; R&D Systems) with a detection limit of 40 pg/mL. As indicated by the manufacturer, both pro-HGF and mature HGF were recognized by this assay. Lactoferrin and myeloperoxidase were quantified by using ELISA kits (Oxis International, Portland, OR; detection limits 1 ng/mL), following the manufacturer's instructions.

Western blot analysis

Purified subcellular fractions of PMN granules, supernatants from cytochalasin B + fMLP-stimulated PMNs, or recombinant human HGF (rh-HGF, a mixture of pro-HGF and mature HGF according to the manufacturer; R&D Systems) were added to 2× Laemmli sample buffer, with or without 2-mercaptoethanol as reducing agent. Proteins were then resolved by sodium dodecyl sulfate-10% polyacrylamide gel electrophoresis (SDS-PAGE). After transfer to nitrocellulose membrane (Bio-Rad Laboratories, Hercules, CA), the proteins were immunoblotted. Nonspecific sites were blocked by overnight immersion in 10% nonfat dried milk, and the membranes were then probed with a mixture (1:1) of 2 monoclonal anti-HGF antibodies (0.5 µg/mL) both recognizing the α chain (MAB694 clone 24 516.1 and MAB294 clone 24 612.111; R&D Systems) or, in one set of experiments, with a monoclonal antiheparan sulfate antibody (5 µg/mL) (Seikagaku, Tokyo, Japan).²⁵ The immunoblots were developed by using an enhanced chemiluminescence method (Amersham Pharmacia Biotech) following the manufacturer's instructions. In selected experiments, before Western blot analysis, supernatants from cytochalasin B + fMLP-stimulated PMNs were incubated in 2 M NaCl, 0.02 M Tris, pH 7.5 for 30 minutes at 37°C to disrupt ionic interactions.²⁶ To check their specificity, the anti-HGF antibodies (5 µg) were preincubated with or without rh-HGF (1 µg) for 1 hour at room temperature before probing the membranes. The intensities of HGF bands (pro-HGF and α chain) decreased when the primary antibodies were preincubated with rh-HGF.

Treatment of pro-HGF from specific granules with serine proteases

Aliquots of purified specific granules (containing 1 ng pro-HGF as determined by Western blot analysis and ELISA) were incubated at 37°C for 10 and 60 minutes with either 0.1 to 100 IU urokinase-type plasminogen activator (uPA)/ng pro-HGF in 20 mM phosphate buffer pH 7.4,¹¹ or with 1 to 100 pmol elastase/ng pro-HGF in HBSS pH 7.5,²⁷ or with 1 to 100 pmol cathepsin G/ng pro-HGF in HBSS pH 7.5.²⁷ Elastase and cathepsin G were kindly provided by M. Chignard from Institut Pasteur-INSERM 285 (Paris, France); uPA was from Hoechst (Paris La Defense, France). After incubation, samples were immediately added to 2× Laemmli sample buffer with 2-mercaptoethanol as reducing agent and analyzed by Western blotting as detailed above.

Reverse transcriptase–polymerase chain reaction

Purified blood PMNs and bone marrow cells prepared as described above were harvested and analyzed by reverse transcriptase–polymerase chain reaction (RT-PCR) for HGF messenger RNA (mRNA) expression. In selected experiments, highly purified PMNs were incubated for 1 hour in 2 mL culture medium with or without 1 μg/mL LPS, 100 ng/mL PMA, 10⁻⁶M fMLP, or 10 ng/mL IL-1β as stimulating agents. Total cellular RNA was isolated with Trizol (Life Technologies) according to the manufacturer's instructions. Total RNA (4 μg) was reverse transcribed in reverse transcriptase reaction buffer (10 mM MgCl₂, 50 mM KCl, 50 mM Tris-HCl pH 8.3) with 25 μg/mL oligo-dT primer, 1 mM each deoxynucleotide triphosphate (Life Technologies), 1 U/μL Rnasin, and 0.5 U/μL AMV reverse transcriptase (Promega, Madison, WI). Samples were incubated for

Table 1. Distribution of hepatocyte growth factor among subcellular fractions from nitrogen-cavitated human neutrophils

	HGF	Lactoferrin	Myeloperoxidase
Cavitate	100	100	100
Plasma membrane	28	< 1	< 1
Cytosol	0	< 1	< 1
Specific granules	68	> 95	< 1
Azurophilic granules	4	< 1	> 95

Results are expressed as a percentage of the total cavitate content (arbitrarily 100%). Data are from a representative experiment of 3 independent experiments. HGF indicates hepatocyte growth factor.

5 minutes at 65°C and then at 42°C for 60 minutes. A specific pair of primers designed for HGF (Life Technologies) was then added to each complementary DNA reaction mix; the sense primer was 5'-TCCCCATCGC-CATCCCC-3' and the antisense primer was 5'-CACCATGGCCTCG-GCTGG-3'.²⁸ Amplification was performed in a solution containing 2 mM MgCl₂, 1× Taq polymerase buffer, 0.2 mM each deoxynucleotide triphosphate, 5% dimethyl sulfoxide (vol/vol), 25 U/mL Taq polymerase AmpliTaq Gold (Perkin Elmer, Foster City, CA), and 0.4 μmol/L specific primer, in an automated thermal cycler (Uno II; Biometra, Maidstone, United Kingdom) at 94°C for 30 seconds, 68°C for 30 seconds, and 72°C for 40 seconds. Amplification was stopped after 35 cycles. The MRC-5 human embryonic lung fibroblast line served as a positive control for HGF gene expression,²⁹ and water was used instead of RNA as a negative control of RT-PCR. The housekeeping gene porphobilinogen deaminase (PBGD) was amplified in each sample to confirm the integrity of RNA and the efficiency of complementary DNA synthesis. The expected PCR products of 749 (HGF) and 338 (PBGD) base pairs were detected by electrophoresis in 2% agarose gel containing ethidium bromide, along with molecular weight standards (Roche Diagnostics, Mannheim, Germany).

Results

PMNs contain an intracellular pool of pro-HGF

As shown in Figure 1, immunocytochemistry revealed the presence of intracellular HGF in unstimulated blood PMNs. To determine the precise localization of HGF, HGF, lactoferrin, and myeloperoxidase were determined in subcellular fractions. HGF was mainly located in the gelatinase/specific granules fraction and, to a lesser extent, the plasma membrane fraction (Table 1); the distribution of




Figure 1. Immunocytochemical staining of blood PMNs. (A) Negative control; no staining was observed with a control immunoglobulin. (B) Intracellular red staining was observed in PMNs with specific monoclonal anti-HGF antibodies; smears were examined by light microscopy at ×800; (Ly indicates lymphocytes; Mo, monocytes).




Figure 2. Western blot analysis of HGF in human neutrophil subcellular fractions. Human neutrophil subcellular fractions were prepared as described in "Materials and methods"; proteins from each fraction (azurophilic and specific granules (gr), membranes and cytosol) were separated by SDS-PAGE in reducing conditions and revealed with antihuman HGF monoclonal antibodies; rh-HGF (a mixture of pro-HGF and mature HGF) was loaded and blotted in parallel. The molecular masses of protein standards are indicated. These data are representative of one typical experiment of 5.

HGF (ELISA) matched that of lactoferrin but not that of myeloperoxidase. Western blot analysis confirmed the subcellular localization of HGF (Figure 2). We observed a distinct 92-kD band in the gelatinase/specific granules fraction and a weak band in the plasma membrane fraction, both migrating at the level of pro-HGF. HGF was not detected in azurophilic granules or in cytosol fractions. As Western blot analysis was conducted in reducing conditions, these results indicate that HGF was present in PMNs as the pro-HGF single-chain form.

PMNs release HGF by degranulation

As shown in Figure 3, PMN stimulated with 1 $\mu\text{g}/\text{mL}$ LPS, 10^{-8} and 10^{-6} M fMLP, the combination of cytochalasin B + 10^{-6} M fMLP, or 100 ng/mL PMA for 20 minutes at 37°C released a large amount of HGF (730 ± 120 , 741 ± 142 , $1\,234 \pm 165$, $2\,322 \pm 141$, and $2\,202 \pm 246$ pg/mL/ 10^7 PMN, respectively) in comparison with unstimulated control cells maintained at 37°C (261 ± 30 pg/mL/ 10^7 PMNs) or at 4°C (< 40 pg/mL).

To investigate the ability of PMNs to synthesize HGF de novo, purified blood PMNs were stimulated, in the presence or absence of cycloheximide, with LPS, PMA, fMLP, or IL-1, which induce cytokine synthesis in PMNs.³⁰ After 24 hours, unstimulated control PMNs cultured in medium alone showed spontaneous HGF release that was not affected by the presence of cycloheximide (Table 2). Each stimulating agent slightly increased HGF concentrations in PMN supernatants as compared with unstimulated PMNs. However, whatever the stimulus, cycloheximide treatment did not significantly modify the amounts of HGF detected after 24 hours of stimulation, suggesting that the HGF released in the culture supernatants was mainly derived from degranulation induced by LPS, PMA, fMLP, or IL-1. Similar results were obtained when PMNs were cultured with tumor necrosis factor α (100 U/mL) or granulocyte-macrophage colony-stimulating factor (100 U/mL) (data not shown).

In agreement with the absence of active HGF synthesis, a very low level of HGF mRNA was detected in unstimulated mature PMNs (Figure 4), and none of the stimulating agents (LPS, PMA, fMLP, or IL-1) modified the HGF mRNA level compared with unstimulated PMNs (densitometric analysis, in comparison with the housekeeping gene PBGD; data not shown).

Expression of HGF in human bone marrow myeloid cells

As most of the proteins stored in PMN granules are actively synthesized during granulocyte maturation, we investigated whether HGF protein and mRNA could be detected in bone marrow




Figure 3. Effect of degranulating agents on HGF secretion by human PMNs. Secreted HGF was measured in supernatants of PMNs ($10^7/\text{mL}$) incubated for 20 minutes at 37°C with 1 $\mu\text{g}/\text{mL}$ LPS, 100 ng/mL PMA, 10^{-8} M, and 10^{-6} M fMLP. When indicated, PMNs were preincubated with 5 $\mu\text{g}/\text{mL}$ cytochalasin B (CYTB). Unstimulated control cells (CTRL) were incubated at 4°C or 37°C in the medium alone. Results are expressed as the mean \pm SEM of 3 independent experiments. ND indicates not detected (< 40 pg/mL).

Table 2. Effects of cycloheximide on stimulus-induced hepatocyte growth factor secretion

	Without CHX pretreatment	With CHX pretreatment
Control PMNs	846 ± 134	897 ± 189
LPS (1 $\mu\text{g}/\text{mL}$)	$1\,480 \pm 239$	$1\,239 \pm 192$
PMA (100 ng/mL)	$1\,621 \pm 212$	$1\,408 \pm 285$
fMLP (10^{-6} M)	$1\,347 \pm 540$	$1\,524 \pm 746$
IL-1 (10 ng/mL)	$1\,068 \pm 272$	$1\,128 \pm 360$

Polymorphonuclear neutrophils (PMN; $10^7/\text{mL}$) were preincubated with or without cycloheximide (CHX) and were then stimulated for 24 hours at 37°C with various agents at the concentrations indicated; unstimulated control PMNs were incubated in medium alone. Hepatocyte growth factor (HGF) was assayed in the cell-free supernatants. Results are expressed as the mean \pm SEM of HGF levels (pg/mL) in 3 experiments. LPS indicates lipopolysaccharide; PMA, phorbol myristate acetate; fMLP, N-formylmethionyl-leucyl-phenylalanine; IL-1, interleukin 1.

myeloid cells. We first performed immunocytochemical analysis of normal human bone marrow smears ($n = 3$). We found strong positive staining in myelocytes, metamyelocytes, band, and segmented cells in all samples. Weak to moderate staining was found in promyelocytes, whereas lymphocytes, monocytes, and megakaryocytes were negative for HGF (Figure 5).

Mononuclear cell-depleted bone marrow samples and circulating purified PMNs were subjected to RT-PCR in the same run. HGF mRNA expression was found in bone marrow samples but was only weakly positive in mature circulating PMNs (Figure 4). By using densitometric analysis in comparison with the housekeeping gene PBGD, we found that the relative intensity was higher in bone marrow cells than in mature circulating PMNs (data not shown).

Taken together, these results show that HGF mRNA and protein are present in neutrophil precursors and suggest that HGF synthesis occurs during PMN maturation.

PMNs process pro-HGF to mature HGF in vitro

Western blot analysis in reducing conditions was used to characterize the HGF form released by PMNs into the extracellular medium. As shown in Figure 6, HGF released by cytochalasin B + fMLP-stimulated PMNs was mainly the mature form (presence of the 69-kD α chain). As the processing of pro-HGF to mature HGF involves serine proteases, the same degranulating experiment was conducted with an additional preincubation step of 15 minutes with 2 mM DFP, used to irreversibly inhibit neutrophil serine proteases.




Figure 4. RT-PCR analysis of HGF mRNA expression by bone marrow cells and mature PMNs. Total RNA was extracted from bone marrow (BM) cells (previously depleted from their mononuclear cells) and from mature PMNs (PMN). After reverse transcription, PCR was carried out with specific pairs of primers designed for HGF and PBGD. Unstimulated MRC-5 human embryonic lung fibroblasts served as positive controls (+). The figure indicates the size of amplification products relative to molecular weight standards run in parallel (M) and the negative control (-).




Figure 5. Immunocytochemical staining of bone marrow cells. (A) Negative control; no staining was observed with a control immunoglobulin. (B) Intracellular red staining was observed with specific monoclonal anti-HGF antibodies; (Pr indicates promyelocyte; My, myelocyte; Mt, metamyelocyte; B, band cell; S, segmented cell). Smears were examined by light microscopy at $\times 800$.

HGF secreted by DFP-pretreated PMNs was exclusively in the pro-HGF form, as shown by the sole detection of the 92-kd band by Western blotting in reducing conditions (Figure 6). These results indicated that the pro-HGF contained in PMNs was processed to




Figure 6. Western blot analysis of HGF released after PMN degranulation: effect of DFP. PMNs were incubated with 2 mM DFP (DFP^+) or without DFP (DFP^-) for 15 minutes at 4°C then stimulated with cytochalasin B and fMLP at 37°C ; the supernatant was then subjected to Western blot analysis in reducing conditions and revealed with antihuman HGF monoclonal antibodies. These data are representative of one experiment of 3.




Figure 7. Effects of serine proteases on pro-HGF from specific granules. Aliquots of purified specific granules were incubated at 37°C for 10 and 60 minutes with 10 pmol elastase/ng pro-HGF (E10, E60), 10 pmol cathepsin G/ng pro-HGF (C10, C60), or 10 IU uPA/ng pro-HGF (U10, U60); specific granules (gr) incubated in the medium alone for 60 minutes served as control. Proteins were then subjected to Western blot analysis in reducing conditions and revealed with antihuman HGF monoclonal antibodies. These data are representative of one experiment of 3.

mature HGF by neutrophil serine protease(s) during degranulation. We, thus, investigated the effects of elastase and cathepsin G, which are the main PMN serine proteases; uPA, which is present in specific granules³¹ and processes pro-HGF,¹¹ was also tested. As shown in Figure 7, none of these agents were able to catalyze the conversion of pro-HGF from specific granules, whatever the incubation period (from 10 to 60 minutes) or concentration (elastase and cathepsin G, from 1 to 100 pmol/ng pro-HGF; uPA, from 0.1 to 100 IU/ng pro-HGF).

Secreted HGF binds PMN-derived heparan sulfate

As HGF is a heparin-binding factor,³² we investigated whether HGF secreted by PMNs could link to glycosaminoglycans. First, in an attempt to purify PMN-derived HGF, the cell-free supernatant of cytochalasin B + fMLP-stimulated PMNs was submitted to heparin affinity chromatography. Contrary to control rh-HGF, HGF secreted by PMNs was not retained on the heparin column (ELISA; data not shown), suggesting that HGF binding sites for heparin were not available. Second, contrary to HGF purified from specific granules, Western blot analysis in nonreducing conditions of HGF secreted by PMNs failed to yield a specific HGF band (Figure 8). This latter result suggests that HGF epitopes were inaccessible to anti-HGF antibodies because of the presence of masking substances. As a high salt concentration is commonly used to release HGF from polyanionic substances such as glycosaminoglycans,²⁶ the supernatant of cytochalasin B + fMLP-stimulated PMNs was




Figure 8. Western blot analysis of PMN-derived HGF: effect of NaCl pretreatment. PMNs were stimulated with cytochalasin B and fMLP at 37°C ; the supernatant was then incubated for 30 minutes at 37°C with 2 M NaCl (NaCl^+) or without NaCl (NaCl^-) and then subjected to Western blot analysis in nonreducing conditions and revealed with antihuman HGF monoclonal antibodies; rh-HGF and HGF derived from purified specific granules (gr) were loaded and blotted in parallel. Note that both pro-HGF and mature HGF yielded a 92-kd band (arrow) in nonreducing conditions. These data are representative of one experiment of 3.




Figure 9. Western blot analysis of HGF released after PMN degranulation: colocalization with neutrophil-derived heparan sulfate. PMNs were stimulated with cytochalasin B and fMLP at 37°C; the supernatant was then subjected to Western blot analysis in reducing conditions and revealed with either antihuman HGF (anti-HGF) or antiheparan sulfate (anti-HS) monoclonal antibodies, as indicated; rh-HGF was loaded and blotted in parallel. These data are representative of one experiment of 3.

then incubated with 2 M NaCl. Western blot analysis revealed a 92-kd band in nonreducing conditions (Figure 8). This finding suggested that HGF secreted by PMNs binds glycosaminoglycans. As heparan sulfate is known to interact with HGF,³² we performed Western blot analysis of PMN-derived HGF with antiheparan sulfate monoclonal antibodies.²⁵ A specific 69-kd band migrating at the level of the HGF α chain was revealed, whereas no band was found with rh-HGF (Figure 9). Taken together, our results suggest that HGF secreted by PMNs binds a PMN-derived heparinlike substance (likely heparan sulfate).

Discussion

PMNs synthesize inflammatory cytokines and growth factors,³⁰ but there are few reports of preformed intracellular stocks of either cytokines¹⁸ or growth factors.³³ Our results demonstrate that both secretory vesicles and gelatinase/specific granules of human blood PMNs contain a mobilizable stock of pro-HGF that is proteolytically processed to mature HGF by neutrophil serine protease(s) during degranulation. Furthermore, this mature HGF bound neutrophil-derived polyanionic substances (likely heparan sulfate).

The bulk of HGF was found in the gelatinase/specific granules fraction and colocalized with the lactoferrin-containing fraction. The efficiency of fractionation was confirmed by measuring the reference markers of specific and azurophil granules (lactoferrin and myeloperoxidase, respectively) in each fraction. Interestingly, vascular endothelial growth factor, another heparin-binding factor, has also been found in specific granules.^{33,34} Intracellular HGF was also present in the plasma membrane and/or secretory vesicles, constituting a readily mobilizable store. This localization demonstrated by the results in Table 1 and in Figure 1 was confirmed by the pattern of HGF release obtained with 10^{-8} M fMLP (Figure 3), a concentration known to mobilize the exocytosis of secretory vesicles.^{35,36} Altogether, our results argue for a dual subcellular localization of HGF, in both secretory vesicles and gelatinase/specific granules.

In addition to releasing numerous preformed mediators into the extracellular medium, PMNs can synthesize cytokines de novo in certain conditions.³⁰ For instance, we have previously demon-

strated that PMNs can release oncostatin M through a 2-step mechanism involving the release of a preformed stock followed by de novo protein synthesis.¹⁸ In the present study we found no evidence of active HGF synthesis by human blood PMNs, as cycloheximide did not affect HGF levels after 24 hours of stimulation. Therefore, our results show that acute HGF release by mature circulating PMNs does not depend on ongoing or activated synthesis but rather on the release of HGF stored in PMNs. Most of the proteins stored in PMN granules are actively synthesized during granulocyte maturation; indeed, we found that HGF mRNA was expressed by bone marrow cells at a higher level than in circulating PMNs. The timing of granule formation is well documented; specific granules along with their protein contents are mainly formed at the myelocyte and metamyelocyte stages.^{37,38} As expected, our immunocytochemistry experiments showed that HGF⁺ cells are essentially myelocytes, metamyelocytes, and segmented cells. Weak to moderate staining was found in promyelocytes. This latter result could be explained by a continuum during precursor maturation.³⁸ Inaba et al³⁹ have shown that the human promyelocytic leukemia cell line HL-60 produces HGF and that dibutyl cAMP (a granulocyte inducer) stimulates HGF release. In addition, Majka et al⁴⁰ have shown that highly purified human CD34⁺ cells express HGF transcripts and produce HGF. Altogether, our results suggest that the bulk of HGF synthesis occurs during PMN maturation, along with most other granule constituents.

As shown by Western blot analysis in reducing conditions, HGF was present as the pro-HGF form in both gelatinase/specific granules and plasma membrane fractions; this mobilizable intracellular pool of HGF was released as the mature disulfide-linked heterodimer into the extracellular medium, as shown by the presence of the 69-kd α chain. These results suggest that HGF is processed to its mature form during PMN degranulation. Furthermore, proteolysis of pro-HGF to mature HGF was completely inhibited in DFP-treated PMNs, suggesting that PMN-derived serine protease(s) is/are involved in this process. Indeed, serine proteases have been shown to convert pro-HGF to mature HGF,^{8-11,41} and the active form of HGF has been found exclusively in injured tissues.⁴² In response to a trigger such as tissue injury, it is conceivable that locally activated PMNs release pro-HGF along with serine protease(s) and that neutrophils infiltrating damaged tissues participate in the local maturation of HGF required for its biologic activities.⁴³ Urokinase, a protease known to be present in PMN-specific granules³¹ and to process pro-HGF,¹⁰ failed to exhibit any converting activity in our experimental conditions. These results are in agreement with those of Mizuno et al⁴¹ who found no *in vitro* properties for plasminogen activators such as urokinase and tissue-type plasminogen activator and with those of Shimomura et al⁹ who reported that the converting activities of these agents were more than 1000 times lower than that of the HGF activator. We also investigated the effects of elastase and cathepsin G, the main serine proteases of PMNs, used over a range of relevant concentrations,²⁷ neither agent had any effect on the processing of pro-HGF. Therefore, the nature of the HGF-converting enzyme(s) in PMNs remains to be determined.

In addition to its specific signaling receptor c-met, HGF interacts with glycosaminoglycans, especially heparan sulfate.³² We infer that HGF secreted by PMNs into the extracellular medium binds PMN-derived heparan sulfate. Indeed, PMNs contain sulfated glycosaminoglycans (eg, heparan sulfate) in their azurophil granules,⁴⁴ which may bind to HGF during the degranulating process. Antiheparan sulfate antibodies revealed the existence of a 69-kd band comigrating with the HGF α chain. This finding is in

keeping with reports that HGF binds heparan sulfate by the N-terminal hairpin loop and the second kringle domain, both of which are located on the HGF α chain.²⁶ Unexpectedly, the apparent molecular mass of HGF bound to heparan sulfate was not modified. However, sulfated oligosaccharides as small as hexasaccharides are known to bind to HGF with high affinity.³⁴ Heparan sulfate potentiates HGF biologic activities; for instance, highly sulfated oligosaccharides increase HGF binding to the c-met receptor, as well as c-met dimerization and subsequent activation.³⁴

These results have functional implications for the role of PMNs in wound repair. First, the existence of a mobilizable stock of HGF would allow PMNs to respond rapidly to changes in the extracellular environment. PMNs represent the majority of infiltrating cells in injured tissues, at least in the early phases, and may, therefore, be an important source of HGF, a critical factor for tissue repair and regeneration. Indeed, HGF acts as a mitogen and morphogen for various epithelial cells and has been shown to play an essential part in liver⁴⁵ and lung⁷ regeneration after damage. HGF may exert its protective action by stimulating proliferation and angiogenesis⁴⁶ and by inhibiting apoptosis.⁴⁷ Various effects of HGF on peripheral blood cells have been described: HGF primes neutrophil oxidative

response,⁴⁸ triggers PMN transmigration,⁴⁹ and stimulates T-cell adhesion and migration,⁵⁰ as well as monocyte cytokine production (eg, IL-6) and monocyte motility and invasiveness.^{51,52} In monocytes, c-met expression is up-regulated in conditions mimicking inflammation (stimulation with IL-1 or LPS),^{52,53} making these cells accessible to the functional effects of HGF. Therefore, the autocrine/paracrine effects of HGF may play a role in the regulation of inflammatory process.

Finally, PMN-derived metalloproteases (eg, MMP-8 and MMP-9) can mobilize growth factors, including HGF, tethered to extracellular matrix proteoglycans and collagen,^{14,15} further supporting the role of PMNs in tissue repair.

In conclusion, our findings point to a new role of PMNs in tissue repair, through the release of HGF.

Acknowledgments

We thank A. Barnier, F. Hochedez, S. Laribe, V. Leçon-Malas, and J. Moreau for expert technical assistance and C. Poüs and J. B. Stern for helpful discussions.

References

- Nakamura T, Teramoto H, Ichihara A. Purification and characterization of a growth factor from rat platelets for mature parenchymal hepatocytes in primary cultures. *Proc Natl Acad Sci U S A*. 1986; 83:6489-6493.
- Gohda E, Tsubouchi H, Nakayama H, et al. Purification and partial characterization of hepatocyte growth factor from plasma of a patient with fulminant hepatic failure. *J Clin Invest*. 1988;81:414-419.
- Zarnegar R, Michalopoulos G. Purification and biological characterization of human hepatopoinetin A, a polypeptide growth factor for hepatocytes. *Cancer Res*. 1989;49:3314-3320.
- Bussolino F, Di Renzo MF, Ziche M, et al. Hepatocyte growth factor is a potent angiogenic factor which stimulates endothelial cell motility and growth. *J Cell Biol*. 1992;119:629-641.
- Zarnegar R, Michalopoulos GK. The many faces of hepatocyte growth factor: from hepatopoiesis to hematopoiesis. *J Cell Biol*. 1995;129:1177-1180.
- Nagaïke M, Hiraïo S, Tajima H, et al. Renotropic functions of hepatocyte growth factor in renal regeneration after unilateral nephrectomy. *J Biol Chem*. 1991;266:22781-22784.
- Yanagita K, Matsumoto K, Sekiguchi K, Ishibashi H, Niho Y, Nakamura T. Hepatocyte growth factor act as a pulmotrophic factor on lung regeneration after acute lung injury. *J Biol Chem*. 1993;268:21212-21217.
- Miyazawa K, Shimomura T, Kitamura A, Kondo J, Morimoto Y, Kitamura N. Molecular cloning and sequence analysis of the cDNA for a human serine protease responsible for activation of hepatocyte growth factor. *J Biol Chem*. 1993;268:10024-10028.
- Shimomura T, Miyazawa K, Komiyama Y, et al. Activation of hepatocyte growth factor by two homologous proteases, blood coagulation factor XIIa and hepatocyte growth factor activator. *Eur J Biochem*. 1995;229:257-261.
- Naldini L, Vigna E, Bardelli A, Follenzi A, Galimi F, Comoglio PM. Biological activation of pro-HGF (hepatocyte growth factor) by urokinase is controlled by a stoichiometric reaction. *J Biol Chem*. 1995;270:603-611.
- Mars WM, Zarnegar R, Michalopoulos GK. Activation of hepatocyte growth factor by the plasminogen activators uPA and tPA. *Am J Pathol*. 1993;143:949-958.
- Nakamura T, Nishizawa T, Hagiya M, et al. Molecular cloning and expression of human hepatocyte growth factor. *Nature*. 1989;342:440-443.
- Trusolino L, Pugliese L, Comoglio PM. Interactions between scatter factors and their receptors: hints for therapeutic applications. *FASEB J*. 1998; 12:1267-1280.
- Taipale J, Keski-Oja J. Growth factors in the extracellular matrix. *FASEB J*. 1997;11:51-59.
- Schuppan D, Schmid M, Somasundaram R, et al. Collagens in the liver extracellular matrix bind hepatocyte growth factor. *Gastroenterology*. 1998;114:139-152.
- Ruoslahti E, Yamaguchi Y. Proteoglycans as modulators of growth factor activities. *Cell*. 1991; 64:867-869.
- Moghul A, Lin A, Beedle A, et al. Modulation of c-met proto-oncogene (HGF receptor) mRNA abundance by cytokines and hormones: evidence for rapid decay of the 8 kb c-met transcript. *Oncogene*. 1994;9:2045-2052.
- Grenier A, Dehoux M, Boutten A, et al. Oncostatin M production and regulation by human polymorphonuclear neutrophils. *Blood*. 1999;93:1413-1421.
- Du Bois RM, Bernaudin JF, Paakko P, et al. Human neutrophils express the 1-antitrypsin gene and produce 1-antitrypsin. *Blood*. 1991;77:2724-2730.
- Sakaguchi H, Seki S, Tsubouchi H, Daikuhara Y, Niitani Y, Kobayashi K. Ultrastructural location of human hepatocyte growth factor in human liver. *Hepatology*. 1994;19:1157-1163.
- Wolf HK, Zarnegar R, Michalopoulos GK. Localization of hepatocyte growth factor in human and rat tissues: an immunohistochemical study. *Hepatology*. 1991;14:488-494.
- Sorensen O, Arnljots K, Cowland JB, Bainton DF, Borregaard N. The human antibacterial cathelicidin, hCAP-18, is synthesized in myelocytes and metamyelocytes and localized to specific granules. *Blood*. 1997;90:2796-2803.
- Borregaard N, Heiple JM, Simons ER, Clark RA. Subcellular localization of the b-cytochrome component of the human neutrophil microbicidal oxidase: translocation during activation. *J Cell Biol*. 1983;97:52-61.
- Shimomura T, Ochiai M, Kondo J, Morimoto Y. A novel protease obtained from FBS-containing culture supernatant, that processes single chain form hepatocyte growth factor to two chain form in serum-free culture. *Cytotechnology*. 1992;8: 219-229.
- David G, Mei Bai X, Van Der Schueren B, Cassiman JJ, Van Den Berghe H. Developmental changes in heparan sulfate expression: in situ detection with mAbs. *J Cell Biol*. 1992;119:961-975.
- Mizuno K, Inoue H, Hagiya M, et al. Hairpin loop and second kringle domain are essential sites for heparin binding and biological activity of hepatocyte growth factor. *J Biol Chem*. 1994;269:1131-1136.
- Renesto P, Chignard M. Enhancement of cathepsin G-induced platelet activation by leukocyte elastase: consequence of the neutrophil-mediated platelet activation. *Blood*. 1993;82:139-144.
- Borset M, Lien E, Espevik T, Helseth E, Waage A, Sundan A. Concomitant expression of hepatocyte growth factor/scatter factor and the receptor c-met in human myeloma cell lines. *J Biol Chem*. 1996;271:24655-24661.
- Tamura M, Arakaki N, Tsubouchi H, Takada H, Daikuhara Y. Enhancement of human hepatocyte growth factor production by interleukin-1 alpha and -1 beta and tumor necrosis factor-alpha by fibroblasts in culture. *J Biol Chem*. 1993;268: 8140-8145.
- Cassatella MA. The production of cytokines by polymorphonuclear neutrophils. *Immunol Today*. 1995;16:21-26.
- Pedersen TL, Plesner T, Horn T, Hoyer-Hansen G, Sorensen S, Hansen NE. Subcellular distribution of urokinase and urokinase receptor in human neutrophils determined by immunoelectron microscopy. *Ultrastruct Pathol*. 2000;24:175-182.
- Lyon M, Deakin JA, Mizuno K, Nakamura T, Gallagher JT. Interaction of hepatocyte growth factor with heparan sulfate. *J Biol Chem*. 1994;269: 11216-11223.
- Gaudry M, Bregerie O, Andrieu V, El Benna J, Pocard MA, Hakim J. Intracellular pool of vascular endothelial growth factor in human neutrophils. *Blood*. 1997;90:4153-4161.
- Zioncheck TF, Richardson L, Liu J, et al. Sulfated oligosaccharides promote hepatocyte growth factor association and govern its mitogenic activity. *J Biol Chem*. 1995;270:16871-16878.
- Sengelov H, Kjedsen L, Borregaard N. Control of exocytosis in early neutrophil activation. *J Immunol*. 1993;150:1535-1543.

36. Sengelov H, Kjeldsen L, Kroeze W, et al. Secretory vesicles are the intracellular reservoir of complement receptor 1 in human neutrophils. *J Immunol*. 1994;153:804-810.
37. Borregaard N, Cowland JB. Granules of the human neutrophilic polymorphonuclear leukocyte. *Blood*. 1997;89:3503-3521.
38. Cowland JB, Borregaard N. The individual regulation of granule protein mRNA levels during neutrophil maturation explains the heterogeneity of neutrophil granules. *J Leukoc Biol*. 1999;66:989-995.
39. Inaba M, Koyama H, Hino M, et al. Regulation of release of hepatocyte growth factor from human promyelocytic leukaemia cells, HL-60, by 1,25-dihydroxyvitamin D₃, 12-*o*-tetradecanoylphorbol 13-acetate, and dibutyryl cyclic adenosine monophosphate. *Blood*. 1993;82:53-59.
40. Majka M, Janowska-Wieczorek A, Ratajczak J, et al. Numerous growth factors, cytokines, and chemokines are secreted by human CD34⁺ cells, myeloblasts, erythroblasts, and megakaryoblasts and regulate normal hematopoiesis in an autocrine/paracrine manner. *Blood*. 2001;97:3075-3085.
41. Mizuno K, Takehara T, Nakamura T. Proteolytic activation of a single-chain precursor of hepatocyte growth factor by extracellular serine-protease. *Biochem Biophys Res Commun*. 1992;189:1631-1638.
42. Miyazawa K, Shimomura T, Naka D, Kitamura N. Proteolytic activation of hepatocyte growth factor in response to tissue injury. *J Biol Chem*. 1994;269:8966-8970.
43. Naldini L, Tamagnone L, Vigna E, et al. Extracellular proteolytic cleavage by urokinase is required for activation of hepatocyte growth factor/scatter factor. *EMBO J*. 1992;11:4825-4833.
44. Parmley RT, Hurst RE, Takagi M, Spicer SS, Austin RL. Glycosaminoglycans in human neutrophils and leukemic myeloblasts: ultrastructural, cytochemical, immunologic, and biochemical characterization. *Blood*. 1983;61:257-266.
45. Ueki T, Kaneda Y, Tsutsui H, et al. Hepatocyte growth factor gene therapy of liver cirrhosis in rats. *Nat Med*. 1999;5:226-230.
46. Aoki M, Morishita R, Taniyama Y, et al. Angiogenesis induced by hepatocyte growth factor in non infarcted myocardium and infarcted myocardium: up-regulation of essential transcription factor for angiogenesis, ets. *Gene Ther*. 2000;7:417-427.
47. Fan S, Ma YX, Wang JA, et al. The cytokine hepatocyte growth factor/scatter factor inhibits apoptosis and enhances DNA repair by a common mechanism involving signaling through phosphatidylinositol 3' kinase. *Oncogene*. 2000;119:2212-2223.
48. Jiang W, Puntis CA, Nakamura T, Hallett MB. Neutrophil priming by hepatocyte growth factor, a novel cytokine. *Immunology*. 1992;77:147-149.
49. Mine S, Tanaka Y, Suematu M, et al. Hepatocyte growth factor is a potent trigger of neutrophil adhesion through rapid activation of lymphocyte function-associated antigen-1. *Lab Invest*. 1998;78:1395-1404.
50. Adams DH, Harvath L, Bottaro DP, et al. Hepatocyte growth factor and macrophage inflammatory protein α : structurally distinct cytokines that induce rapid cytoskeletal changes and subset-preferential migration in T-cells. *Proc Natl Acad Sci U S A*. 1994;91:7144-7148.
51. Beilmann M, Vande Woude GF, Dienes HP, Schirmacher P. Hepatocyte growth factor-stimulated invasiveness of monocytes. *Blood*. 2000;95:3964-3969.
52. Galimi F, Cottone E, Vigna E, et al. Hepatocyte growth factor is a regulator of monocyte-macrophage function. *J Immunol*. 2001;166:1241-1247.
53. Beilmann M, Odenthal M, Jung W, Vande Woude GF, Dienes HP, Schirmacher P. Neoexpression of the c-met/hepatocyte growth factor-scatter factor receptor gene in activated monocytes. *Blood*. 1997;90:4450-4458.