Combination of the low anticoagulant heparin CX-01 with chemotherapy for the treatment of acute myeloid leukemia

Tibor J. Kovacsovics,¹ Alice Mims,² Mohamed E. Salama,^{1,3} Jeremy Pantin,⁴ Narayanam Rao,¹ Ken M. Kosak,¹ Peter Ahorukomeye,¹ Martha J. Glenn,¹ Michael W. N. Deininger,¹ Kenneth M. Boucher,⁵ Linda M. Bavisotto,⁶ Gerardo Gutierrez-Sanchez,⁷ Thomas P. Kennedy,⁸ Stephen G. Marcus,⁹ and Paul J. Shami¹

¹Division of Hematology and Hematologic Malignancies, Huntsman Cancer Institute, University of Utah, Salt Lake City, UT; ²Hollings Cancer Center, Medical University of South Carolina, Charleston, SC; ³Department of Pathology and Associated Regional and University Pathologists Laboratories, University of Utah, Salt Lake City, UT; ⁴Division of Hematology/Oncology, Augusta University, Augusta, GA; ⁵Department of Internal Medicine and Huntsman Cancer Institute, University of Utah, Salt Lake City, UT; ⁶Porta Clinica PLLC, Seattle, WA; ⁷Complex Carbohydrate Research Center, University of Georgia, Athens, GA; ⁸Division of Pulmonary Medicine, Wake Forest University, Winston-Salem, NC; and ⁹Cantex Pharmaceuticals, Weston, FL

Key Points

- In a pilot study, the nonanticoagulant heparin derivative CX-01 was well tolerated when combined with chemotherapy for the treatment of AML.
- Preliminary results show encouraging complete remission rates and rapid platelet recovery.

Relapses in acute myelogenous leukemia (AML) are a result of quiescent leukemic stem cells (LSCs) in marrow stromal niches, where they resist chemotherapy. LSCs employ CXCL12/CXCR4 to home toward protective marrow niches. Heparin disrupts CXCL12-mediated sequestration of cells in the marrow. CX-01 is a low-anticoagulant heparin derivative. In this pilot study, we combined CX-01 with chemotherapy for the treatment of AML. Induction consisted of cytarabine and idarubicin (7 + 3) with CX-01. Twelve patients were enrolled (median age, 56 years; 3 women). Three, 5, and 4 patients had good-, intermediate-, and poor-risk disease, respectively. Day 14 bone marrows were available on 11 patients and were aplastic in all without detectable leukemia. Eleven patients (92%) had morphologic complete remission after 1 induction (CR1). Eight patients were alive at a median follow-up of 24 months (4 patients in CR1). Three patients received an allogeneic stem cell transplant in CR1. Median disease-free survival was 14.8 months. Median overall survival was not attained at the maximum follow-up time of 29.4 months. No CX-01-associated serious adverse events occurred. Median day to an untransfused platelet count of at least 20×10^9 /L was 21. CX-01 is well tolerated when combined with intensive therapy for AML and appears associated with enhanced count recovery and treatment efficacy.

Introduction

Acute myelogenous leukemia (AML) afflicted 21 380 patients in the United States in 2017, with 10 590 deaths and a 5-year survival of 26.9%.¹ The mainstay of therapy is variations on the combination of cytarabine and an anthracycline ("7 + 3").² Relapses are common because of quiescent leukemic stem cells (LSCs) in marrow stromal niches, where they are resistant to chemotherapy.³

Hematopoietic stem cells (HSCs) migrate to and are anchored by chemotactic signals produced in bone marrow stroma.^{4,5} Foremost among these signals is the CXC chemokine CXCL12 (stem cell-derived factor-1, or SDF-1), which ligates chemokine receptor 4 (CXCR4) on the HSC surface and activates migration toward, and attachment in, the marrow.^{6,7} CXCL12/CXCR4 also activates phosphatidylinositol-3-kinase and protein kinase B, enhancing survival.⁸ In the marrow, HSCs shuttle between an endosteal osteoblastic niche inducing quiescence and a sinusoidal endothelial vascular

Submitted 16 October 2017; accepted 21 January 2018. DOI 10.1182/ bloodadvances.2017013391.

^{© 2018} by The American Society of Hematology

niche promoting proliferation and differentiation.⁴ Quiescence in the endosteal niche is maintained by close association with megakaryocytes.^{9,10} Megakaryocytes negatively regulate HSC proliferation through secretion of platelet factor-4 (PF4), a cationic chemokine that also enhances integrin-mediated endothelial adhesion during marrow homing.¹¹

LSCs employ the same CXCL12-, CXCR4-dependent mechanisms as normal HSCs to home toward a marrow niche.^{3,4} CXCR4 is an unfavorable AML prognostic factor.¹² Disrupting CXCR4 inhibits transmigration, survival, and chemotherapy resistance in vitro and in vivo.^{13,14} The CXCR4 inhibitor plerixafor has been tested in AML.¹⁵ Plerixafor administered for up to 7 days for stem cell mobilization is well tolerated,¹⁶ but carries the possibility of increased secondary myeloid malignancies.¹⁷

Heparin avidly binds to a cluster of cationic amino acids on the CXCL12 surface that mediates attachment of the chemokine to heparan sulfate on cell surfaces.^{18,19} Heparin can disrupt CXCL12mediated sequestration of cells in marrow by competing with marrow heparan sulfate.²⁰ Heparin's competitive inhibition of CXCL12 binding to heparan sulfate is not diminished by removal of 2-O sulfates,²⁰ but is apparent only at concentrations exceeding therapeutic plasma levels for anticoagulation.²⁰⁻²² An alternative to disrupt the CXCL12/CXCR4 axis is use of a heparin analog with reduced anticoagulation. Low anticoagulant 2-O, 3-O desulfated heparin (ODSH, CX-01) is a porcine intestinal heparin derivative that retains many heparin anti-inflammatory properties.²³ CX-01 binds PF4 with affinity similar to heparin, but the CX-01/PF4 complex does not bind antibodies that mediate heparin-induced thrombocytopenia.^{24,25} Megakaryopoiesis is negatively regulated by PF4, and anti-PF4 blocking antibodies diminish chemotherapyinduced thrombocytopenia in mice.²⁶ By reversing the biologic activities of PF4, CX-01 is able to mitigate chemotherapy-induced thrombocytopenia in PF4-overexpressing transgenic mice.²⁷

We conducted a pilot study combining CX-01 with chemotherapy for AML treatment. Results show no added toxicity from CX-01. Complete remission (CR) rates and count recovery were encouraging. Exploratory in vitro studies suggest CX-01 may interfere with the CXCL12/CXCR4 axis.

Methods

Clinical trial

This was an open-label pilot study conducted at 3 academic centers. The protocol was approved by the University of Utah, Augusta University, and the Medical University of South Carolina respective Institutional Review Boards (ClinicalTrials.gov identifier: NCT02056782). The study was conducted in accordance with the Declaration of Helsinki. Adult patients with previously untreated AML based on World Health Organization criteria²⁸ with an Eastern Cooperative Oncology Group performance status of 0 to 2 and adequate organ function were enrolled. Patients with acute promyelocytic leukemia and acute megakaryoblastic leukemia were excluded. Patients needing anticoagulation were excluded.

Treatment protocol. Induction cycles consisted of cytarabine at 100 mg/m² as a continuous 24-hour infusion on days 1 through 7, and idarubicin at 12 mg/m² as a slow intravenous push on days 1 through 3. CX-01 was administered as a bolus of 4 mg/kg over the course of 30 minutes immediately after the first dose of idarubicin, followed by a continuous infusion at 0.25 mg/kg per hour for 24 hours on days 1 through 7. Patients who were aged 60 years or older received further therapy off study. Patients who were younger than 60 years remained on study and received up to 4 consolidation cycles consisting of cytarabine at a dose of 3 g/m² intravenously over the course of 3 hours every 12 hours on days 1, 3, and 5. For consolidation cycles, CX-01 was given as a bolus of 4 mg/kg over the course of 30 minutes after completion of the first dose of cytarabine, followed by a continuous infusion at 0.25 mg/kg per hour for 24 hours on days 1 through 5. Eligible patients underwent allogeneic stem cell transplant whenever ready after completion of induction therapy.

Supportive care. Red blood cell transfusions, antibiotics, and myeloid growth factors were given per institutional guidelines. Prophylactic platelet transfusions were given to maintain a platelet count of at least $10 \times 10^9/L$ for afebrile patients and of at least $20 \times 10^9/L$ for febrile patients. Otherwise, platelets were given as clinically indicated.

End points. Primary endpoints included safety and tolerability of CX-01 when combined with chemotherapy and the effect of CX-01 on platelet transfusion independence, defined as the first of 5 consecutive days with an untransfused platelet count of at least 20×10^9 /L. Secondary endpoints included rate of CR, platelet nadir, number of platelet transfusions, and toxicity using Common Terminology Criteria for Adverse Events version 4.0.²⁹ Responses were determined using International Working Group Criteria.³⁰

Statistical analysis. Platelet recovery and toxicity endpoints were analyzed using descriptive statistics with reporting of minimum, maximum, mean, median, and 95% confidence interval of the mean. CR rates were reported as a fraction of enrolled subjects together with an exact 95% binomial confidence interval constructed using the method of Clopper and Pearson.³¹ Planned enrollment was 10 evaluable patients. The target rate of unacceptable CX-01-related toxicity was 20% with the following decision rule: If 3 or more subjects experience unacceptable CX-01-related toxicity, an acceptable dose of CX-01 has been exceeded. If 0-2 patients experience unacceptable CX-01-related toxicity, an acceptable dose of CX-01 has not been exceeded. With this rule, there is a 7% chance of making an incorrect determination that an acceptable dose of CX-01 has been exceeded if the true rate of unacceptable CX-01-related toxicity is 10%. There is a 6% chance of making an incorrect determination that an acceptable dose of CX-01 has not been exceeded if the true rate of unacceptable CX-01-related toxicity is 50%.

Pharmacokinetic studies. Single-dose and steady-state pharmacokinetic sampling at 5 points was obtained during induction cycles for 7 patients. Duplicate samples for each point were analyzed by BioCascade Incorporated (Arlington, WI), using the Sta-Chrom anti-Xa assay,²³ calibrated to express CX-01 concentration in micrograms per milliliter.

In vitro studies

Surface plasmon resonance. Surface plasmon resonance analyses on immobilized CX-01 and heparin sodium USP. One hundred micrograms per milliliter NeutrAvidin (Thermo Scientific) in 10 mM sodium acetate at pH 4.5 was immobilized on 2 flow cells of a Series S CM5 (GE Healthcare) sensor chip, using the amine coupling method.^{32,33} The remaining activated

groups were blocked by ethanoline injection. For direct binding studies, biotinylated CX-01 and heparin sodium were then injected at a concentration of 50 μ g/mL for 2 min at 5 μ L/min, resulting in approximately 174 to 243 response units CX-01 or heparin sodium immobilized. Control flow cells were used for background subtraction, having biotin captured over the immobilized neutravidin. For direct binding analysis, CXCL12 α was diluted in HBS-EP (0.01 M HEPES pH 7.4, 0.15 M NaCl, 3 mM EDTA, 0.05% v/v Surfactant P20; GE Healthcare). Different dilutions of the protein were injected at a flow rate of 30 μ L/min for 180 seconds. A dissociation time of 300 seconds was set up, followed by regeneration with 3 M NaCl in HBS-EP buffer.

Competition for CXCL12 binding between immobilized heparin sodium and CX-01. For the competition experiment, CXCL12 α (final concentration, 100 nM) was mixed with various concentrations of CX-01 (0, 1, 5, 10, 50, 100, 1000, and 2000 nM) and then injected. The effect of CX-01 on CXCL12 α binding to immobilized heparin was monitored as relative response units, used to calculate 50% inhibitory concentration values.

Surface plasmon resonance data evaluation. Surface plasmon resonance (SPR) data were analyzed by global fitting either to a 1:1 Langmuir binding model or to a 2-state reaction binding model, using the Biacore T100 BIAevaluation software 2.0.4 (GE Health-care). For competition studies, data were plotted and the amount of CX-01 required to inhibit 50% of the CXCL12 binding to heparin was determined.

Cell migration studies. To determine the effect of CX-01 on the migration of AML cells toward CXCL12, we conducted in vitro migration assays using a cell migration kit from Cell Biolabs (San Diego, CA) with the U937 monocytic leukemia cell line (ATCC, Manassas, VA), after verification of CXCR4 expression by flow cytometry. U937 cells (0.2×10^6 cells/0.1 mL) were loaded in inserts with 5-µm pore membranes. Inserts were transferred to wells containing 0.5 mL OPTI-MEM media with CXCL12 (100 ng/mL), CX-01 (200 µg/mL), or both. To investigate the direct effect of CX-01 on leukemia cells in 1 variable, CXCL12 only was added to the wells, and U937 cells (2×10^6 /mL) were pretreated for 30 minutes with 500 µg/mL CX-01. Cells were then washed before loading into inserts. Plates were incubated overnight at 37°C in a 5% CO₂ atmosphere. Cell migration was quantitated with the CyQuant GR dye stain according to the manufacturer's protocol.

Results

Clinical trial

Patient characteristics. Enrollment was planned for 10 patients; 12 were enrolled (median age, 54 years; range, 22-74 years; 3 women). Two patients (patients 2001 and 1007) were taken off study because of adverse events unrelated to CX-01. They were replaced for the purposes of time to count recovery analysis and were not included in that analysis. They were included in the response and toxicity analysis. Patient 2001 was taken off study after 2 days of induction because of acute recurring angioedema-like allergic reactions that began before treatment and may have been from angiotensin-converting enzyme inhibitor therapy. Patient 1007 developed acute renal failure secondary to tumor lysis and vancomycin nephrotoxicity, and was taken off treatment after completing 5 of the 7 days of induction. All patients were evaluable for safety. Ten completed the induction cycle and were evaluable for

efficacy. Table 1 summarizes patient characteristics. Three, 5, and 4 patients had good-, intermediate-, and poor-risk disease, respectively.

Response to induction. Eleven (92%) of 12 patients obtained a morphologic CR after 1 induction cycle (CR1), including the 2 patients who did not complete induction. All patients with de novo AML (11/11) obtained a CR. Two patients in morphologic CR had minimal residual disease by molecular testing. Patient 1007 received 5 days of induction and had detectable inv(16) by polymerase chain reaction. Patient 1006 had a FLT-3/ITD mutation and an inv(3) that were detectable at the end of induction. The patient who did not obtain a CR (1003) had presented with extensive lymphadenopathy with a bone marrow showing chronic myelomonocytic leukemia-2, but was included in the study on the basis of evidence of granulocytic sarcoma on lymph node biopsy. It is worth noting that day 14 bone marrows were available on 11 patients and were aplastic in all without detectable leukemia.

Count recovery after induction. We analyzed count recovery in the 10 patients who completed a full induction cycle. They had a median time to recovery of an untransfused platelet count of at least 20×10^9 /L and of at least 50×10^9 /L of 21 (mean, 21.3 ± 1.7 ; range, 18-22) and 23.5 (mean, 23.1 ± 2.0 ; range, 19-25) days, respectively. Induction cycles required a median of 6.0 platelet transfusions (mean, 6.5 ± 3.9 ; range, 0-10 transfusions). Patient 1010 had a platelet count nadir of 11×10^9 /L and required no platelet transfusions. Patient 1002 had an induction course complicated by a rectal abscess and hematuria and received 10 platelet transfusions. The median platelet nadir count for induction cycles was 8×10^9 /L (mean, 8.4×10^9 /L $\pm 2.6 \times 10^9$ /L; range, 0.5-14 $\times 10^9$ /L).

Patient 1010 was the only patient who received filgrastim support at the discretion of the treating physician and not because of a delay in count recovery. As we aimed to obtain preliminary results on the effect of CX-01 on bone marrow recovery, patient 1010 was not evaluated for total white blood cell (WBC) or neutrophil recovery. For patients who did not receive filgrastim support, we analyzed time to recovery to a total WBC of at least 1×10^{9} /L and a neutrophil count of at least 0.5×10^9 /L. Eight patients were evaluable for total WBC recovery and had a median time to a WBC count of at least 1×10^{9} /L of 21 days (mean, 23 ± 4.8; range, 20-35 days). Patient 1009 was not included in the analysis for a total WBC recovery to a count of at least 1×10^9 /L because his WBC never went below 1×10^9 /L during the whole induction cycle. Five patients were evaluable for neutrophil recovery and had a median time to a neutrophil count of at least 0.5×10^9 /L of 22 days (mean, 22.6 \pm 2.4; range, 21-27 days). The other patients were discharged from the hospital with a rising absolute neutrophil count (ANC) before achieving the target neutrophil count, and were therefore not included in the analysis for the time to ANC recovery.

Frequency of platelet transfusions after consolidation. Four patients received 1 or more consolidation cycles on study, with median frequencies of platelet transfusions as follows (n = evaluable patients): for the first consolidation cycle (n = 4), 2 (mean, 2.0 \pm 0.7; range, 1-3) transfusions; for the second consolidation cycle (n = 3), 3 (mean, 2.3 \pm 0.9; range, 1-3) transfusions; for the third consolidation cycle, (n = 2), 2 (mean, 2.0 \pm 0.0; range, 2- 2) transfusions. Patient 1005 received a fourth consolidation cycle on study and had a prolonged hospitalization

able 1.	Patient	Char	acteristics and Outcomes				
Patient	Age	Sex	Karyotype and molecular findings	% Marrow blasts at diagnosis	Day 14 bone marrow	Induction outcome	Postinduction therapy and outcome
1002	63	Σ	+8, t(9:11), FLT3 TKD	03	<5% cellularity; NED	g	Three cycles of IDAC consolidation off study followed by allogeneic stem cell transplant in CR1.
1003	46	Σ	t(12,13)	6.4	<5% cellularity; NED	Failure	Allogeneic transplant after salvage therapy.
1004	74	Σ	add(8) t(8,19) FLT3 TKD	64	Suboptimal; NED	ర	Two cycles of IDAC consolidation and relapse 6 mo after diagnosis. Postrelapse therapy on an experimental protocol and died of cerebral hemorrhage.
1005	46	Σ	XY (SNP array)	26	Suboptimal; serous atrophy; NED	£	Four cycles of HIDAC + CX-01 consolidation on study. Relapse 16 m after diagnosis. Underwent salwage therapy and allogeneic transplant. Had extramedullary relapse posttransplant and underwent salvage therapy and donor lymphocyte infusion.
1006	5	ш	inv(3) FLT3 ITD	0 LG	<5% celularity; serous atrophy; NED	CR MRD	Two cycles of IDAC consolidation off study followed by allogeneic stem cell transplant in CR1. Relapse 9 mo after diagnosis, received further therapy on an experimental protocol and died of infectious complications.
1007	58	Σ	inv(16) +8, +22, +21	51	<5% cellularity serous atrophy; NED	CR MRD	Four cycles of HiDAC consolidation off study because of incomplete induction cycle.
1009	54	Σ	XY - NPM1 +	37	10% cellularity, NED	ଞ	Three cycles of consolidation with HiDAC + CX:01 on study One cycle of consolidation with HiDAC alone after withdrawing from study.
1010	52	Σ	XY - FLT3 ITD	30	<5% cellularity serous atrophy; NED	స	Developed line-associated deep vein thrombosis and went off study after induction. – Two cycles of HiDAC consolidation off study followed by allogenetic stem cell transplant in CR1.
2001	5	Σ	t(6,9) - FLT3 ITD	20	Not done	с	Received 2 d of induction and relapsed 7 wk after diagnosis. Underwent salvage therapy and allogeneic stem cell transplant. Subsequently died of transplant-related complications.
3001	22	ш	+ NPM1 +	21	<10% cellularity; NED	CR	One cycle of HiDAC + CX-01 consolidation on study and was lost to follow-up after withdrawing study. Relapsed 13.5 mo after diagnosis, received salvage therapy and underwent allogeneic stem cell transplant in second remission.
3002	67	Σ	dei(20) (q11.2) - FLT3 ITD (tested at relapse)	20	Hypocellular; NED	ర	Three cycles of consolidation therapy off study. Relapsed 11.5 mo after diagnosis. Underwent sakage therapy with transient response and relapse followed by allogeneic stem cell transplant.
3003	25	ш	XX	6 <u>6</u>	<20% cellularity; NED	S	Two cycles of HiDAC + CX-01 consolidation on study. Relapsed 6 mo after diagnosis and underwent salvage therapy and allogeneic stem cell transplant. Subsequently relapsed again and underwent salvage and second allogeneic stem cell transplant and died of transplant-related complications.
HiDAC,	high-dose) cytarab	aine; IDAC, intermediate-dose cytarabine; MRD, n	minimal residual disease; NED, no eviden	ice of disease.		



Figure 1. Kaplan Meier curves of overall survival (OS) and disease-free survival (DFS).

because of septic shock with multiorgan failure in the setting of neutropenia. He was therefore not evaluable for the number of platelet transfusions associated with that cycle. The starting platelet counts at the beginning of the first consolidation cycle were typically higher than for other cycles because most patients experienced some degree of rebound thrombocytosis on full hematologic recovery from induction. Platelet and WBC nadirs for consolidation cycles were not evaluated because subjects had blood count checks only 3 times per week on an outpatient basis on completion of chemotherapy.

Postinduction therapy and long-term follow-up. Of the 12 patients enrolled, 7 were not eligible to receive postinduction treatment on study because of age 60 years or older, induction failure, or incomplete induction cycles (Table 1). Of the remaining patients, 1 developed a line-associated deep venous thrombosis requiring anticoagulation and was taken off study before consolidation. The remaining 4 patients each received 1 or more cycles of consolidation on study as follows: patient 1005 completed all 4 cycles; patient 1009 received 3 cycles of consolidation and withdrew from study; patient 3001 completed 1 consolidation cycle, withdrew from study, and was lost to follow-up; patient 3003 received 2 consolidation cycles before relapsing. Three patients who completed induction received an allogeneic stem cell transplant in CR1 (patients 1002, 1006, 1010). Two of those patients were alive without evidence of disease at the time of final data analysis. Six of 8 patients who obtained a CR with induction therapy and did not receive a transplant in CR1 relapsed at a median time of 8 months. Among those were patient 2001, who had not completed induction and relapsed 7 weeks after diagnosis, and patient 3001, who received only 1 cycle of consolidation and relapsed 13.5 months after diagnosis. With a median follow-up of 24 months, 8 patients were alive. Median disease-free survival was 14.8 months. Median overall survival was not attained at the maximum follow-up time of 29.4 months (Figure 1).

Safety and toxicity. CX-01 was well tolerated and did not increase the risk of bleeding. Coagulation parameters in induction and consolidation were not significantly affected by CX-01. Anti-factor Xa



Figure 2. Peak and steady-state CX-01 plasma concentrations vs time curves in 7 patients with AML during induction. Shown are averages and standard errors of the mean for each time point.

activity before and during administration of CX-01 was not significantly changed from baseline and remained below published therapeutic anticoagulant reference ranges. Two patients experienced grade 3 to 4 and 3 patients experienced grade 1 to 2 bleeding episodes, all deemed to be unrelated to CX-01.

A total of 5 serious adverse events occurred in 5 patients. Grade 4 sepsis occurred in 2 patients during the first and fourth consolidation. These events were deemed to be possibly related and not related to CX-01, respectively. Another patient experienced febrile neutropenia not related to CX-01 during induction. Patient 1002 had a perirectal abscess during induction requiring surgery with platelet support, an event that was unrelated to CX-01. Patients 1007 and 2001 did not complete induction because of serious adverse events that were unrelated to CX-01 (see "Patient characteristics").

The most frequent nonserious adverse events were hematologic, followed by infectious and organ toxicity complications resulting from treatment and/or the underlying leukemia. Grade 3 and 4 hematologic toxicities were consistent with intensive AML chemotherapy. Transient asymptomatic low-grade elevations of liver transaminases were observed during chemotherapy administration in induction cycles. These were deemed by investigators to be possibly related to CX-01, irrespective of other potential causes of liver function abnormalities. Transient asymptomatic grade 3 to 4 liver transaminase elevations were observed during consolidation cycles in 2 patients and were deemed possibly related to CX-01. Serum bilirubin remained normal for all patients, except for an isolated grade 4 hyperbilirubinemia as a consequence of hepatic failure complicating septic shock in patient 1005 during the fourth consolidation.

Pharmacokinetics. Single-dose and steady-state pharmacokinetic plasma sampling was obtained during induction at 5 points for 7 patients (Figure 2). Peak levels of $38.5 \pm 7.5 \,\mu$ g/mL (average \pm SEM) were reached 15 minutes after CX-01 bolus injection. For unknown reasons, patients 3001 and 3002 had substantially lower peak levels of CX-01 than the remaining patients



Figure 3. Surface plasmon resonance assay of the binding of CX-01 to CXCL12 α . (A) BlAcore sensorgrams for the binding of 100 nM CXCL12 α after preincubation with different concentrations of CX-01 (0, 1, 5, 10, 50, 100, 1000, and 2000 nM). (B) Competition assay shows the CX-01 concentration (50% inhibitory concentration = 4.7 nM or 0.055 μ g/mL) that inhibits 50% of the maximum CXCL12 α attachment response to immobilized heparin.

(13.3 \pm 1.85 vs 48.6 \pm 5.25 $\mu g/mL).$ Steady state levels at day 6 were 6.5 \pm 1.5 $\mu g/mL.$

In vitro studies

To explore possible mechanisms of action of CX-01, we conducted in vitro studies to evaluate the effect of CX-01 on the CXCR4/ CXCL12 axis.

SPR. The interaction properties of CX-01 or heparin sodium with CXCL12 α were compared. CXCL12 α bound to immobilized CX-01 or heparin sodium with comparable association and dissociation constants, yielding apparent K_D values from 26 to 40 nM and 5 to 30 nM, respectively. To determine whether CX-01 inhibits binding of CXCL12 α to immobilized heparin, a constant amount of CXCL12 α was incubated with varying concentrations of CX-01 and injected over the immobilized heparin. The amount of residual binding was determined at each concentration of CX-01, and the corresponding amount of free CXCL12 α was determined. Competition experiments demonstrated that binding of 100 nM CXCL12 α to immobilized heparin was inhibited by progressively higher concentrations of CX-01 (Figure 3A), with 5 nM CX-01 inhibiting 46.8% of the interaction between CXCL12 α and heparin, for a 50% inhibitory concentration of 4.7 nM (0.055 µg/mL) (Figure 3B).

Cell migration studies. To test the effects of CX-01 on chemotaxis, we performed in vitro migration studies using U937 AML cells. CXCL12 at a concentration of 100 ng/mL induced chemotaxis of U937 AML cells. Addition of CX-01 (200 μ g/mL) to CXCL12 in the wells did not inhibit migration. However, pretreating U937 cells with CX-01 (500 μ g/mL) nullified the chemotactic effect of CXCL12, suggesting a direct effect of CX-01 on leukemic cells (Figure 4).

Discussion

In this pilot exploratory study, CX-01 in combination with standard intensive AML therapy was safe and well tolerated, with no unexpected serious adverse events, suggesting that the approach is feasible. The combination was associated with an encouraging CR

rate of 92%. Hematologic recovery was also encouraging, with a median of 22 days to reach an ANC larger than $0.5\times10^9/L$, and 21 days to reach a platelet count higher than $20\times10^9/L$ in induction cycles.

From a toxicity point of view, we observed transient asymptomatic low-grade elevations in liver transaminases during induction cycles.



Figure 4. Effect of CX-01 on U937 cell migration toward CXCL12. The assay was conducted using a cell migration assay kit from Cell Biolabs (San Diego, CA), as detailed in the Methods section. CXCL12 was added at a concentration of 100 ng/mL in the wells. CX-01 was added at a concentration of 200 μ g/mL in the wells. To investigate the direct effect of CX-01 on leukemia cells in 1 variable, CXCL12 only was added to the wells and U937 cells were pretreated for 30 minutes with 500 μ g/mL CX-01 before loading them into inserts. The graph shows averages and standard errors of the mean of 2 to 3 replicate wells in each group (the media blank variable had 1 well). The legend indicates CXCL12 and/or CX-01 added to the wells or pretreatment of U937 cells with CX-01. The graph is representative of 2 separate experiments. RFU, relative fluorescence unit.

Elevation of liver transaminases is a class effect for heparins.³⁴ It has also been described for idarubicin and cytarabine induction regimens in the absence of heparin.³⁵ Transient asymptomatic grade 3 to 4 liver transaminase elevations were also noted during consolidation cycles in 2 patients. Simultaneous coadministration of CX-01 with cytarabine in this study may confound attribution of causality for this effect, as hepatotoxicity has also been reported for cytarabine.³⁶ Although the CX-01 dose and schedule were identical for induction and consolidation cycles (except for the duration of the CX-01 continuous infusion), consolidation cycles employed a much higher dose of cytarabine than induction cycles, raising the possibility of multifactorial causation of elevated liver transaminase levels.

Pharmacokinetic studies showed a similar pattern of CX-01 clearance as in normal volunteers.²³ It should be noted that the doses we used were lower than doses use in the latter study.²³ These results suggest that CX-01 can be administered with cytarabine and idarubicin without significant drug-drug interaction. Hematologic recovery was rapid compared with published data, with a median of 22 days to reach an ANC higher than 0.5×10^9 /L, and 21 days to reach a platelet count higher than 20×10^9 /L in induction cycles. A randomized comparison will be required to confirm these findings.

The combination was associated with an encouraging CR rate of 92%, higher than the 71% CR rate reported by Vogler et al using a similar regimen of cytarabine and idarubicin.³⁷ None of our patients needed a second induction cycle to achieve CR, compared with 23% in the Vogler study.³⁷ Although our study is small, it is conceivable that the improvement over the expected response is the result of disrupting the CXCL12/CXCR4mediated homing of LSCs to the protective marrow niche.3,4 Intravenous boluses of sulfated polysaccharides mobilize normal stem cells from the niche into the circulation, suggesting that sulfated polysaccharides also mobilize LSCs.³⁸ Promoting cell division and permissive migration closer to the vascular compartment may expose leukemic cells to chemotherapy in a vulnerable state. Our hypothesis is supported by the similar binding affinity of CX-01 and unfractionated heparin for CXCL12 and the ability of CX-01 to inhibit binding of CXCL12 to immobilized heparin at concentrations below those achieved in the circulation of treated patients. This suggests similar competition in vivo of CX-01 with marrow glycosaminoglycans. Interestingly, when CX-01 was added to CXCL12 in U937 cell migration studies, it did not inhibit migration of the leukemic cells toward CXCL12. The discrepancy between the SPR data and the cell migration data may be a result of differences between a cell-free assay (SPR) and a cell-based assay (migration). However, pretreatment of U937 cells with CX-01 abrogated migration toward CXCL12, suggesting a direct effect on the leukemic cells to block the CXCL12/CXCR4 axis. As the CX-01 concentration in these experiments was higher than the measured drug concentrations in patients, further work will be required to determine whether that observation can be reproduced at lower CX-01 concentrations.

The mean days of recovery to a platelet count above $50 \times 10^9/L$ and to a WBC count above $1 \times 10^9/L$ were 23.1 and 23.0 days, respectively; a more rapid return of platelets (mean of 35.1 days for a platelet count higher than $50 \times 10^9/L$) and WBC (mean of 31.2 days for WBC count higher than $1 \times 10^9/L$) than previously reported

with this regimen.³⁷ Furthermore, the median time to recovery of neutrophils for induction cycles in CX-01-treated patients is essentially identical (22 days) to that of patients supported with myeloid growth factors during induction.³⁹ Although the number of patients in our study is small, and while acknowledging the inherent limitations of such comparisons, our results are encouraging with respect to WBC and neutrophil recovery.

PF4 levels inversely correlate with recovery of platelet counts after chemotherapy in children with acute lymphoblastic leukemia,⁴⁰ and PF4 suppresses the proliferation of normal HSCs.^{9,10} Because CX-01 inhibits the biologic action of PF4, our observations suggest that CX-01 might be abrogating the suppressive effect of PF4 on marrow recovery. Furthermore, the effects on platelet and neutrophil counts in our study are compatible with the recently appreciated role of megakaryocytes and PF4 in promoting quiescence of normal HSCs.^{9,10} Thus, PF4 might conceivably play a role in maintaining LSC quiescence, similar to normal HSCs. In these conditions, CX-01 (a polyanion that charge-inhibits critical cationic structural sequences of PF4) could remove any PF4 constraint on LSC activation, while simultaneously promoting megakaryocytic and myeloid recovery through neutralization of the inhibitory effect of PF4 on HSCs.

Results of this pilot study are preliminary and exploratory in nature. They require confirmation in large studies, as well as more extensive evaluation of our mechanistic hypotheses. However, if confirmed, CX-01 might offer a valuable adjunctive therapy for one of hematology's most challenging malignancies.

Acknowledgment

This study was supported with funding from Cantex Pharmaceuticals.

Authorship

Contribution: T.J.K. participated in study design, patient care, data and analysis; A.M. participated in study design and patient care; M.E.S. participated in pathologic review of bone marrows; J.P. participated in study design and patient care; N.R. participated in laboratory studies; K.M.K. participated in laboratory studies; P.A. participated in laboratory studies; M.J.G. participated in patient care; M.W.N.D. participated in patient care; K.M.B. participated in statistical design and analysis; L.M.B. participated in data analysis; G.G.-S. participated in laboratory studies; T.P.K. participated in data analysis; S.G.M. participated in study design and data analysis; P.J.S. participated in study design, patient care, and data analysis; and all authors participated in manuscript preparation.

Conflict-of-interest disclosure: T.P.K. is a stockholder in Cantex Pharmaceuticals. S.G.M. is employed by and is a stock holder in Cantex Pharmaceuticals. P.J.S. received research support from Cantex Pharmaceuticals. The remaining authors declare no competing financial interests.

The current affiliation for A.M. is Division of Hematology, Ohio State University, Columbus, OH.

Correspondence: Paul J. Shami, Huntsman Cancer Institute, Suite 2100, 2000 Circle of Hope, Salt Lake City, UT 84112; e-mail: paul.shami@utah.edu.

References

- 1. Siegel R, Ma J, Zou Z, Jemal A. Cancer statistics, 2014. CA Cancer J Clin. 2014;64(1):9-29.
- 2. Estey EH. Acute myeloid leukemia: 2014 update on risk-stratification and management. Am J Hematol. 2014;89(11):1063-1081.
- 3. Konopleva MY, Jordan CT. Leukemia stem cells and microenvironment: biology and therapeutic targeting. J Clin Oncol. 2011;29(5):591-599.
- 4. Blau O. Bone marrow stromal cells in the pathogenesis of acute myeloid leukemia. Front Biosci. 2014;19(1):171-180.
- Lapidot T, Kollet O. The essential roles of the chemokine SDF-1 and its receptor CXCR4 in human stem cell homing and repopulation of transplanted immune-deficient NOD/SCID and NOD/SCID/B2m(null) mice. *Leukemia*. 2002;16(10):1992-2003.
- 6. Peled A, Kollet O, Ponomaryov T, et al. The chemokine SDF-1 activates the integrins LFA-1, VLA-4, and VLA-5 on immature human CD34(+) cells: role in transendothelial/stromal migration and engraftment of NOD/SCID mice. *Blood*. 2000;95(11):3289-3296.
- Avigdor A, Goichberg P, Shivtiel S, et al. CD44 and hyaluronic acid cooperate with SDF-1 in the trafficking of human CD34⁺ stem/progenitor cells to bone marrow. *Blood.* 2004;103(8):2981-2989.
- 8. Sahin AO, Buitenhuis M. Molecular mechanisms underlying adhesion and migration of hematopoietic stem cells. Cell Adhes Migr. 2012;6(1):39-48.
- 9. Bruns I, Lucas D, Pinho S, et al. Megakaryocytes regulate hematopoietic stem cell quiescence through CXCL4 secretion. *Nat Med.* 2014;20(11): 1315-1320.
- Zhao M, Perry JM, Marshall H, et al. Megakaryocytes maintain homeostatic quiescence and promote post-injury regeneration of hematopoietic stem cells. Nat Med. 2014;20(11):1321-1326.
- 11. Zhang J, Lu SH, Liu YJ, Feng Y, Han ZC. Platelet factor 4 enhances the adhesion of normal and leukemic hematopoietic stem/progenitor cells to endothelial cells. *Leuk Res.* 2004;28(6):631-638.
- 12. Spoo AC, Lübbert M, Wierda WG, Burger JA. CXCR4 is a prognostic marker in acute myelogenous leukemia. Blood. 2007;109(2):786-791.
- Chen Y, Jacamo R, Konopleva M, Garzon R, Croce C, Andreeff M. CXCR4 downregulation of let-7a drives chemoresistance in acute myeloid leukemia. J Clin Invest. 2013;123(6):2395-2407.
- 14. Zeng Z, Shi YX, Samudio IJ, et al. Targeting the leukemia microenvironment by CXCR4 inhibition overcomes resistance to kinase inhibitors and chemotherapy in AML. *Blood.* 2009;113(24):6215-6224.
- Uy GL, Rettig MP, Motabi IH, et al. A phase 1/2 study of chemosensitization with the CXCR4 antagonist plerixafor in relapsed or refractory acute myeloid leukemia. Blood. 2012;119(17):3917-3924.
- 16. De Clercq E. The bicyclam AMD3100 story. Nat Rev Drug Discov. 2003;2(7):581-587.
- 17. Deol A, Abrams J, Masood A, et al. Long-term follow up of patients proceeding to transplant using plerixafor mobilized stem cells and incidence of secondary myelodysplastic syndrome/AML. Bone Marrow Transplant. 2013;48(8):1112-1116.
- Sadir R, Baleux F, Grosdidier A, Imberty A, Lortat-Jacob H. Characterization of the stromal cell-derived factor-1α-heparin complex. J Biol Chem. 2001; 276(11):8288-8296.
- Ziarek JJ, Veldkamp CT, Zhang F, et al. Heparin oligosaccharides inhibit chemokine (CXC motif) ligand 12 (CXCL12) cardioprotection by binding orthogonal to the dimerization interface, promoting oligomerization, and competing with the chemokine (CXC motif) receptor 4 (CXCR4) N terminus. *J Biol Chem.* 2013;288(1):737-746.
- Zhang S, Condac E, Qiu H, et al. Heparin-induced leukocytosis requires 6-O-sulfation and is caused by blockade of selectin- and CXCL12 proteinmediated leukocyte trafficking in mice. J Biol Chem. 2012;287(8):5542-5553.
- 21. Seeger FH, Rasper T, Fischer A, et al. Heparin disrupts the CXCR4/SDF-1 axis and impairs the functional capacity of bone marrow-derived mononuclear cells used for cardiovascular repair. Circ Res. 2012;111(7):854-862.
- 22. Hirsh J, Anand SS, Halperin JL, Fuster V; American Heart Association. Guide to anticoagulant therapy: Heparin : a statement for healthcare professionals from the American Heart Association. *Circulation*. 2001;103(24):2994-3018.
- 23. Rao NV, Argyle B, Xu X, et al. Low anticoagulant heparin targets multiple sites of inflammation, suppresses heparin-induced thrombocytopenia, and inhibits interaction of RAGE with its ligands. *Am J Physiol Cell Physiol*. 2010;299(1):C97-C110.
- Joglekar MV, Quintana Diez PM, Marcus S, et al. Disruption of PF4/H multimolecular complex formation with a minimally anticoagulant heparin (ODSH). Thromb Haemost. 2012;107(4):717-725.
- 25. Krauel K, Hackbarth C, Fürll B, Greinacher A. Heparin-induced thrombocytopenia: in vitro studies on the interaction of dabigatran, rivaroxaban, and lowsulfated heparin, with platelet factor 4 and anti-PF4/heparin antibodies. *Blood.* 2012;119(5):1248-1255.
- 26. Lambert MP, Rauova L, Bailey M, Sola-Visner MC, Kowalska MA, Poncz M. Platelet factor 4 is a negative autocrine in vivo regulator of megakaryopoiesis: clinical and therapeutic implications. *Blood.* 2007;110(4):1153-1160.
- Lambert MP, Sharma SS, Xiao L, et al. 2-O, 3-O-Desulfated Heparin (ODSH) mitigates chemotherapy-induced thrombocytopenia (CIT) by blocking the negative paracrine effect of platelet factor 4 (PF4) on megakaryopoiesis [abstract]. Blood. 2012;120(21). Abstract 386.
- 28. Vardiman JW, Harris NL, Brunning RD. The World Health Organization (WHO) classification of the myeloid neoplasms. Blood. 2002;100(7):2292-2302.
- US Department of Health and Human Services. Common Terminology Criteria for Adverse Events (CTCAE). Version 4.0. https://evs.nci.nih.gov/ftp1/ CTCAE/CTCAE_4.03_2010-06-14_QuickReference_5x7.pdf. Accessed 13 February 2013.

- 30. Cheson BD, Bennett JM, Kopecky KJ, et al; International Working Group for Diagnosis, Standardization of Response Criteria, Treatment Outcomes, and Reporting Standards for Therapeutic Trials in Acute Myeloid Leukemia. Revised recommendations of the International Working Group for diagnosis, standardization of response criteria, treatment outcomes and reporting standards for therapeutic trials in acute myeloid leukemia. J Clin Oncol. 2003; 21(24):4642-4649.
- 31. Clopper CJ, Pearson ES. The use of confidence or fiducial limits illustrated in the case of the binomial. Biometrika. 1934;26(4):404-413.
- 32. Osmond RI, Kett WC, Skett SE, Coombe DR. Protein-heparin interactions measured by BIAcore 2000 are affected by the method of heparin immobilization. *Anal Biochem.* 2002;310(2):199-207.
- Amara A, Lorthioir O, Valenzuela A, et al. Stromal cell-derived factor-1alpha associates with heparan sulfates through the first beta-strand of the chemokine. J Biol Chem. 1999;274(34):23916-23925.
- 34. Harrill AH, Roach J, Fier I, et al. The effects of heparins on the liver: application of mechanistic serum biomarkers in a randomized study in healthy volunteers. *Clin Pharmacol Ther.* 2012;92(2):214-220.
- 35. Wiernik PH, Banks PLC, Case DC Jr, et al. Cytarabine plus idarubicin or daunorubicin as induction and consolidation therapy for previously untreated adult patients with acute myeloid leukemia. *Blood.* 1992;79(2):313-319.
- 36. US National Library of Medicine. Drug record: cytarabine. https://livertox.nlm.nih.gov/Cytarabine.htm. Accessed 30 May 2016.
- 37. Vogler WR, Velez-Garcia E, Weiner RS, et al. A phase III trial comparing idarubicin and daunorubicin in combination with cytarabine in acute myelogenous leukemia: a Southeastern Cancer Study Group Study. J Clin Oncol. 1992;10(7):1103-1111.
- Sweeney EA, Lortat-Jacob H, Priestley GV, Nakamoto B, Papayannopoulou T. Sulfated polysaccharides increase plasma levels of SDF-1 in monkeys and mice: involvement in mobilization of stem/progenitor cells. *Blood.* 2002;99(1):44-51.
- Sierra J, Szer J, Kassis J, et al. A single dose of pegfilgrastim compared with daily filgrastim for supporting neutrophil recovery in patients treated for lowto-intermediate risk acute myeloid leukemia: results from a randomized, double-blind, phase 2 trial. BMC Cancer. 2008;8(1):195.
- 40. Lambert MP, Reznikov A, Grubbs A, et al. Platelet factor 4 platelet levels are inversely correlated with steady-state platelet counts and with platelet transfusion needs in pediatric leukemia patients. J Thromb Haemost. 2012;10(7):1442-1446.